

with free OH groups in the α and γ positions of the side chain.

LITERATURE CITED

1. Z. K. Sailov, L. S. Smirnova, and Kh. A. Abduazimov, Khim. Drevesiny, Vol. 2, 40 (1977).
2. N. A. Veksler, L. S. Smirnova, and Kh. A. Abduazimov, Khim. Priir. Soedin., 100 (1977).

COMPONENTS OF *Gomphocarpus fruticosus*

L. G. Kretsu

UDC 547.918

The preparation "Gomfotin," which is close in its pharmacological properties to strophanthin, is obtained from the epigeal mass of *Gomphocarpus fruticosus* (L) R. Br. The main active agents of this preparation are the cardiac glycosides that have been obtained previously from the seeds [1] and leaves [2] of *G. fruticosus*. One of the glycosides has been synthesized independently [3]. The other components of the plants have been studied.

We have investigated the leaves, stems, roots, and follicles of *G. fruticosus* grown on an experimental plot of the botanical garden of the Academy of Sciences of the Moldavian SSR, the material for analysis being taken in the fruit-bearing phase. The amounts of alkaloids in the various organs of the plant were determined by Van'kovskii's method [4]. It was established that the roots contained about 0.5%, the valves of the seeds 0.5%, and the leaves 0.15% of alkaloids.

Preliminary analysis with the aid of spot reactions, and also by paper and thin-layer chromatography on silica gel in various solvent systems showed that the epigeal part contained flavonoids (Shinoda's test) and coumarins (visualization on paper or thin-layer chromatograms in UV light before and after spraying with a 5% solution of alkali), while the roots contained tanning substances (test with 1% ferric chloride solution) and flavonoids. Cardiac glycosides were detected only in the epigeal organs of the plant (Legal and Baljet tests). Quantitative analysis of the combined glycosides calculated as Gomfotin performed by the method of Tsarenko and Shraiber [5] showed that the largest amount of them was present in the leaves and follicles (about 1%) while the stems contained 0.06%, i.e., the epigeal mass of the plant is of interest for obtaining the combined glycosides.

The sum of the flavonoids was obtained by the following scheme: the comminuted air-dry material (epigeal part and roots separately) was extracted with chloroform and then with hot methanol. The methanolic extract was extracted to small volume, diluted with water, and treated with ethyl acetate. The organic layer was diluted with three volumes of chloroform. The yellow amorphous precipitate that deposited was separated off; it consisted of a mixture of flavonoid compounds — 9% in the epigeal part and 4% in the roots. On the basis of Bryant's test [6], aglycones and glycosides were detected in the mixture. Crystals with mp 189–190°C deposited from ethanol. Hydrolysis gave an aglycone with mp 302–305°C, identical according to PC with quercetin. Glucose and rhamnose were detected in the mother solution of the hydrolysate by paper chromatography.

Thus, the crystals with mp 189–190°C were rutin, as was confirmed by a direct comparison of its chromatographic mobility and UV spectrum with those of an authentic sample [7].

On studying the chloroform extract of the epigeal parts of the plant, in addition to carotenoids and chlorophyll, a substance was detected which was colored pink by the Sanníe reagent. Column chromatography on silica gel in the hexane-ether (7:3) system yielded a compound with mp 135–136°C (ethanol), coinciding in chromatographic mobility with

Botanical Garden, Academy of Sciences of the Moldavian SSR, Kishinev. Translated from Khimiya Prirodnykh Soedinenii, No. 6, pp. 795–796, November–December, 1982. Original article submitted May 24, 1982.

β -sitosterol [8]. The acetate of the compound, after recrystallization, had mp 119°C. These substances were identified by direct comparison with authentic samples.

This is the first time that β -sitosterol, rutin, alkaloids, coumarins, and tanning substances have been detected in *Gomphocarpus fruticosus*.

LITERATURE CITED

1. V. T. Chernobai, Med. Promst. SSSR, No. 1, 36 (1957).
2. V. T. Chernobai and D. G. Kolesnikov, Med. Promst. SSSR, No. 3, 29 (1957).
3. V. T. Chernobai and N. F. Komissarenko, Khim. Prir. Soedin., 445 (1971).
4. A. I. Ermakov, Methods for the Biochemical Investigation of Plants [in Russian], Leningrad (1972), p. 340.
5. N. Ya. Tsarenko and M. S. Shraiber, Farmatsiya, No. 2, 39 (1970).
6. E. Bryant, J. Am. Pharm. Assoc., 39, 481 (1960).
7. M. A. Azimov and Z. N. Nazirov, Khim. Prir. Soedin., 271 (1970).
8. C. Bodea, Tratat de biochimie vegetală [Treatise on Plant Biochemistry; in Romanian], Bucharest, Part I (1964), p. 368.