

## DUALITY IN COHERENT CONFIGURATIONS

A. NEUMAIER

*Received July 1, 1986*

*Revised July 12, 1987*

By introducing a duality operator in coherent algebras (i.e. adjacency algebras of coherent configurations) we give a new interpretation to Delsarte's duality theory for association schemes. In particular we show that nonnegative matrices and positive semidefinite matrices,  $(0, 1)$ -matrices and distance matrices, regular graphs and spherical 2-designs, distance regular graphs and Delsarte matrices are pairs of dual objects. Several "almost dual" properties which are not yet fully understood are also reported.

In his thesis, Delsarte [8] observed a formal duality in the theory of association schemes which becomes an actual duality in the case that an abelian group is present. In this note we shall develop certain aspects of this duality in the slightly more general context of coherent algebras. In particular, it turns out that graphs and distance matrices, introduced in Neumaier [12], are dual objects.

A *coherent algebra* is an algebra  $V$  of square complex  $v \times v$ -matrices (indexed by a set  $X$  with  $v$  elements) closed under componentwise (Schur) multiplication  $\circ$  and under conjugate transposition  $*$  such that the identity matrix  $I$  and the all-one matrix  $J$  belong to  $V$ . It is not difficult to show that the partition of  $X \times X$  defined by the equivalence relation

$$(x, y) \equiv (x', y') \text{ iff } A_{xy} = A_{x'y'} \text{ for all } A \in V$$

turns  $X$  into a coherent configuration in the sense of Higman [10]; conversely, the adjacency algebra of a coherent configuration is a coherent algebra ([10], (2.8)). In particular, if  $V$  is commutative,  $X$  is an association scheme in the sense of Delsarte [8], and  $V$  is its Bose—Mesner algebra (cf. [5], where only the symmetric case is treated). See also Higman [21].

Many properties of coherent algebras and related configurations are formally dual and in some cases, this is due to the existence of a duality operator.

We call a coherent algebra  $V$  *selfdual* if there is a semi-linear duality operator  $\tau$  on  $V$  such that

$$A^\tau = vA \quad \text{for all } A \in V.$$

$$(\alpha A)^\tau = \bar{\alpha} A^\tau \quad \text{for all } \alpha \in \mathbb{C} \text{ and } A \in V.$$

$$(A^*)^\tau = (A^\tau)^* \quad \text{for all } A \in V.$$

$$(AB)^\tau = A^\tau \circ B^\tau \quad \text{for all } A, B \in V.$$

Note that then  $A^* = (AI)^* = A^* \circ I^*$ , so

$$I^* = J, \quad J^* = I^{**} = vI,$$

$$(A \circ B)^* = \frac{1}{v^2} (A^{**} \circ B^{**})^* = \frac{1}{v^2} (A^* B^*)^{**} = \frac{1}{v} A^* B^*;$$

in particular  $V$  is commutative.

**Example 1.** Let  $G$  be a finite abelian group of order  $v$ . A decomposition of  $G$  into a direct product  $G = C_1 \times \dots \times C_t$  of cyclic groups  $C_i$  of order  $e_i$  with generators  $g_i$  determines a multiplicative inner product

$$\langle x, y \rangle = \prod \zeta_i^{x_i y_i}, \quad \text{if } x = \prod g_i^{x_i}, \quad y = \prod g_i^{y_i},$$

(where  $\zeta_i$  is a primitive  $e_i$ -th root of unity) with the following properties:

$$\langle x, y \rangle = \langle x, y \rangle, \quad \langle x, y \rangle^{-1} = \langle x^{-1}, y \rangle,$$

$$\langle x, yz \rangle = \langle x, y \rangle \langle x, z \rangle,$$

$$\langle x, y \rangle = 1 \quad \text{for all } y \in G \Leftrightarrow x = 1.$$

$$\sum_{y \in G} \langle x, y \rangle = \begin{cases} v & \text{if } x = 1, \\ 0 & \text{otherwise,} \end{cases}$$

since

$$\sum_{y_1=0}^{e_1} \dots \sum_{y_t=0}^{e_t} \zeta_i^{x_i y_i} = 0.$$

In particular, the map  $x \mapsto \langle x, \cdot \rangle$  is an isomorphism between  $G$  and its character group  $\chi(G)$ .

The semi-linear map  $\tau$  obtained by extending the definition

$$x^* = \sum_{y \in G} \langle x, y \rangle y \quad \text{with} \quad \left( \sum_{x \in G} a_x x \right)^* = \sum_{x \in G} \bar{a}_x x^*$$

to the group algebra  $CG$  satisfies

$$x^{**} = \left[ \sum_y \langle x, y \rangle y \right]^* = \sum_y \overline{\langle x, y \rangle} y^* = \sum_y \langle x, y \rangle^{-1} y^* =$$

$$= \sum_y \sum_z \langle x^{-1}, y \rangle \langle y, z \rangle z = \sum_z \sum_y \langle x^{-1} z, y \rangle z = vx,$$

$$(xy)^* = \sum_z \langle xy, z \rangle z = \sum_z \langle x, z \rangle \langle y, z \rangle z = \sum_z \langle x, z \rangle z \circ \sum_z \langle y, z \rangle z = x^* \circ y^*,$$

$$1^* = \sum_y \langle 1, y \rangle y = \sum_y y.$$

Let  $V$  be the centralizer algebra of  $G$  (cf. Wielandt [18]) in its regular action (=the set of all matrices that are invariant under  $G$ ),  $V = \{A = (a_{xy})_{x,y \in G} | a_{xz}, yz = a_{x,y} \text{ for all } x, y, z \in G\}$  is isomorphic to the group algebra of  $G$  via the isomorphism  $a = \sum_x a_x x \mapsto \Pi(a) = (a_{xy^{-1}})_{x,y \in G}$ .  $V$  is a coherent algebra, and the map  $\tau$  defined by  $\Pi(a)^* = \Pi(a^*)$  is a duality operator on  $V$ .

**Example 2.** An association scheme is of Latin square type (negative Latin square type) if every  $(0, 1)$ -matrix with zero diagonal contained in the Bose—Mesner algebra  $V$  is the adjacency matrix of a pseudo Latin square graph (negative Latin square graph). Here a  $\frac{\text{pseudo}}{\text{negative}}$  Latin square graph is a strongly regular graph with

$\frac{\mu=s(s+1)}{\mu=r(r+1)}$  (cf. Cameron et al. [7]); in particular, imprimitive strongly regular graphs (i.e. disjoint unions of cliques and multipartite graphs) are degenerate pseudo Latin square graphs. E.g. the points of an affine plane (or more generally a net (Bruck [6])), together with a partition  $(\Sigma_1, \dots, \Sigma_s)$  of the line directions (points at infinity) defines an association scheme of Latin square type by calling  $x, y$   $i$ -th associates if  $x$  and  $y$  are on a line with direction  $\in \Sigma_i$ . See also [19, 20].

Define for  $A \in V$  the numbers  $k_A, d_A$  by

$$AJ = k_A J, \quad A \circ I = d_A I.$$

Then for any association scheme of Latin square type, the map  $\tau$  with

$$\bar{A}^* = \pm nA + (n \mp 1)^{-1}(k_A \mp nd_A)(nI \mp J), \quad n = \sqrt{v}$$

(upper sign for Latin square type, lower sign for negative Latin square type) is a duality operator.

**Proposition 1.** A commutative coherent algebra  $V$  has a unique basis  $D_0, \dots, D_s$  of  $(0, 1)$ -matrices such that  $D_0 = I, D_i^* \in \{D_0, \dots, D_s\}$  for all  $i, D_0 + D_1 + \dots + D_s = J$ .

**Proof.** Every finite dimensional commutative algebra has a basis of idempotents. Let  $D_0, \dots, D_s$  be such a basis with respect to the  $\circ$ -multiplication. Then each  $D_i$  is a  $(0, 1)$ -matrix, and since  $D_i \circ D_j = 0$  for  $i \neq j$ , the linear combination of the  $D_i$  representing  $J$  must be  $J = D_0 + \dots + D_s$ . Similarly,  $D_i^*$  and  $I$  are represented by a sum of  $D_j$  ( $j=0, \dots, s$ ) which is seen to contain only one term, and by a suitable permutation we may take  $D_0 = I$ . ■

From now on, let  $V$  be selfdual (and hence commutative).

**Proposition 2.** Let  $T = (\tau_{ij})_{i,j}$  be the matrix defined by

$$D_i^* = \sum_{j=0}^s \tau_{ij} D_j.$$

Then  $D_i$  has row sums  $k_i = \tau_{i0}$ , and we have the relations

$$T\bar{T} = vI,$$

$$\tau_{ji}k_i = \bar{\tau}_{ij}k_j^*.$$

**Proof.**  $(D_i^*)^* = \sum_{j=0}^s \bar{\tau}_{ij} D_j^* = \sum_{j=0}^s \sum_{l=0}^s \bar{\tau}_{ij} \tau_{jl} D_l$  but also  $D_i^{**} = vD_i$ , so  $\bar{T}T = vI$ . Next we have

$$D_i^* \circ I = \sum_{j=0}^s \tau_{ij} D_j \circ I = \tau_{i0} D_0 = \tau_{i0} I,$$

hence  $D_i J = \frac{1}{v} (D_i^\tau \circ J^\tau)^\tau = (D_i^\tau \circ I)^\tau = (\tau_{i0} I)^\tau = \tau_{i0} J$ , giving  $k_i = \tau_{i0}$ . Finally

$$D_i^\tau D_j \circ I = \sum_{l=0}^s \tau_{il} (D_l D_j \circ I) = \sum_{l=0}^s \tau_{il} (\delta_{lj} k_j I) = \tau_{ij} k_j I$$

since  $D_i D_j \circ I = \delta_{ij} k_j I$ , hence

$$\begin{aligned} \bar{\tau}_{ij} k_j J &= (\tau_{ij} k_j I)^\tau = (D_i^\tau D_j \circ I)^\tau = \frac{1}{v} (D_i^\tau D_j)^\tau I^\tau = \\ &= \frac{1}{v} (D_i^{\tau\tau} \circ D_j^\tau) J = (D_i \circ D_j) J = \tau_{ji} D_i J = \tau_{ij} k_i J, \end{aligned}$$

since  $D_i \circ D_j^\tau = D_i \circ \sum_l \tau_{jl} D_l = \sum_l \tau_{jl} D_i \circ D_l = \tau_{ji} D_i$ ; and we get  $\bar{\tau}_{ij} k_j = \tau_{ji} k_i$ . ■

**Proposition 3.**

- (i)  $A$  has constant row sum  $k \Leftrightarrow A^\tau$  has constant diagonal entries  $k$ .
- (ii)  $A$  is a  $(0, 1)$ -matrix of valency  $k \Leftrightarrow \frac{1}{v} A^\tau$  is idempotent of rank  $k$ .
- (iii)  $A$  is nonnegative  $\Leftrightarrow A^\tau$  is positive semidefinite.
- (iv) The  $\frac{1}{v} D_i^\tau$  are the irreducible idempotents of  $V$ .
- (v)  $D_i$  has the eigenvalues  $\bar{\tau}_{ij}$  with multiplicity  $k_j$  ( $j=0, \dots, s$ ).

**Proof.**

- (i)  $AJ = kJ \Leftrightarrow A^\tau \circ I = kI$ .
- (ii)  $A \circ A = A \Leftrightarrow \frac{1}{v} A^\tau \cdot \frac{1}{v} A^\tau = \frac{1}{v} A^\tau$ , trace  $\frac{1}{v} A^\tau = k$ , so rank  $\frac{1}{v} A^\tau = k$  (since the only eigenvalues are 0 and 1).
- (iii)  $A$  is nonnegative  $\Leftrightarrow A = \sum_{i=0}^s a_i D_i$ , and all  $a_i \geq 0 \Leftrightarrow A^\tau = \sum_{i=0}^s a_i D_i^\tau$ , and all  $a_i \geq 0 \Leftrightarrow A^\tau$  is positive semidefinite.
- (iv) We have  $\frac{1}{v} D_i^\tau \cdot \frac{1}{v} D_j^\tau = \frac{1}{v} (D_i \circ D_j)^\tau = \frac{1}{v} \delta_{ij} D_i^\tau$ . If  $\frac{1}{v} D_i^\tau = \frac{1}{v} E^\tau + \frac{1}{v} F^\tau$  is a decomposition of  $\frac{1}{v} D_i^\tau$  into idempotents then  $D_i = E + F$  is a decomposition into  $(0, 1)$ -matrices. But  $D_i$  is a basis element so this is impossible.
- (v)  $D_i^\tau = \sum_j \tau_{ij} D_j$  implies  $D_i = \sum_j \bar{\tau}_{ij} \frac{1}{v} D_j^\tau$ , and since  $\frac{1}{v} D_j^\tau$  is idempotent,  $D$  has eigenvalues  $\bar{\tau}_{ij}$  with multiplicity rank  $\left(\frac{1}{v} D_j^\tau\right) = \text{trace} \left(\frac{1}{v} D_j\right) = k_j$ . ■

The *eigenmatrices* of a commutative coherent algebra (equivalently, of a symmetric association scheme) are the matrices  $P = (P_i(j))_{i,j}$  and  $Q = (Q_i(j))_{i,j}$  relating the basis of  $(0, 1)$ -matrices  $D_i$  and the basis of idempotents  $E_i$  as follows

$$D_i = \sum_{j=0}^s P_i(j) E_j; \quad v E_i = \sum_{j=0}^s Q_i(j) D_j.$$

Of course  $PQ = vI$ , and by the above results, we have  $\bar{P} = Q = T$  for a self-dual algebra. Conversely, a commutative coherent configuration with  $\bar{P} = Q$  is self-dual, since the semi-linear map  $\tau$  with  $D_i^\tau = vE_i$  is a duality operator.

If  $V_0$  is a coherent subalgebra of a self-dual coherent algebra, then so is its dual  $V_0^\tau$ ; the valencies  $k_i'$ , multiplicities  $f_i'$ , and eigenmatrices  $P', Q'$  of  $V_0^\tau$  are related to the valencies  $k_i$ , multiplicities  $f_i$ , and eigenmatrices  $P, Q$  of  $V_0$  by

$$k_i' = f_i, \quad f_i' = k_i, \quad P' = \bar{Q}, \quad Q' = \bar{P}.$$

In particular, we have

**Theorem 1.** *If a self-dual coherent algebra contains the adjacency matrix of a strongly regular graph  $\Gamma$  with eigenvalues  $k, r, s$  with multiplicities  $1, f, v-1-f$ , then it also contains the adjacency matrix of a strongly regular graph  $\Gamma$  with the dual parameters:*

$$(*) \quad v' = v, \quad k' = f, \quad r' = \frac{v-k+s}{r-s}, \quad s' = \frac{s-k}{r-s}, \quad f' = k.$$

*In particular, the hypothesis can only be satisfied if either  $\Gamma$  is a conference graph, or  $r-s$  is an integer dividing  $v$  (then the known necessary conditions already imply that  $(r-s)|(s-k)$ ).*

**Proof.** The eigenmatrices of  $\Gamma$  are (cf. Lemma 2.2 of Cameron et al. [7])

$$P = \begin{pmatrix} 1 & 1 & 1 \\ k & r & s \\ l & -r-1 & -s-1 \end{pmatrix}, \quad Q = \begin{pmatrix} 1 & 1 & 1 \\ f & \frac{fr}{k} & -\frac{f(r+1)}{l} \\ g & \frac{gs}{k} & -\frac{g(s+1)}{l} \end{pmatrix},$$

where  $l = v-1-k$ ,  $g = v-1-f$ , and a similar formula holds for the dual. Now it is clear that  $v' = v$  and the relations  $P' = Q$ ,  $Q' = P$  give  $k' = f, f' = k$ , and  $r' = \frac{fr}{k}$ ,  $s' = \frac{-f(r+1)}{l}$ . Now, using e.g. the formulas in Lemma 2.1 of Cameron et al. [7], we find

$$s' = -\frac{f(r+1)}{l} = -\frac{(k(s+1)+ls)(r+1)}{(r-s)l} = -\frac{l(k+rs)+ls(r+1)}{(r-s)l} = \frac{s-k}{r-s}$$

and

$$r' - s' = \frac{f}{kl} (rl + (r+1)k) = \frac{fg}{kl} (r-s) = \frac{v}{r-s}. \quad \blacksquare$$

**Corollary 1** (Delsarte [8]) *If a strongly regular graph  $\Gamma$  has a regular abelian automorphism group then there is a dual strongly regular graph  $\Gamma'$  such that their parameters are related by A.*  $\blacksquare$

**Corollary 2** (Delsarte [8]). *The Bose-Mesner algebra  $V$  of a strongly regular graph  $\Gamma$  is self-dual iff  $v = (r-s)^2$  iff  $\Gamma$  is a conference graph, a pseudo Latin square graph, or a negative Latin square graph.*  $\blacksquare$

Next we relate the adjacency matrices of graphs in a self-dual coherent algebra with their dual, which are distance matrices (see Neumaier [12], [13], [14]). We shall also establish a relation with distance regular graphs (see Biggs [4]).

A *distance matrix* (Neumaier [12]) is a nonzero real symmetric  $v \times v$ -matrix  $C = (c_{xy})$  with nonnegative entries, indexed by a set  $X$  with  $v$  elements, such that the distance function  $d(x, y) := c_{xy}^{1/2}$  satisfies  $d(x, x) = 0$  and  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$ . If  $C$  has no repeated rows then this turns  $X$  into a metric space, and conversely, for every metric  $d$  on a finite set  $X$ , the matrix  $C := (d(x, y)^2)$  is a distance matrix. A distance matrix  $C$  has *degree*  $s$  if  $C$  has precisely  $s$  distinct nonzero entries, and *strength* (at least)  $t$  if for all nonnegative integers  $i, j$  with  $i + j \leq t$ , the  $(x, y)$ -entry of the product of  $C^{(i)} = (c_{xy}^i)$  and  $C^{(j)} = (c_{xy}^j)$  can be written as a polynomial  $f_{ij}$  of degree  $\leq \min(i, j)$  in  $c_{xy}$ ; we write this as  $C^{(i)}C^{(j)} = f_{ij} \circ C$ . A distance matrix  $C$  is *spherical* if  $G = \gamma J - C$  is positive semidefinite for  $\gamma > 0$ . Distance matrices are closely related to *spherical  $t$ -designs* defined by Delsarte et al. [9].

**Proposition 4.** *Let  $C$  be a distance matrix with  $v$  rows.*

- (i)  *$C$  has strength 1 iff*

$$(1) \quad CJ = kJ \text{ for some } k > 0.$$
- (ii)  *$C$  has strength 2 iff (1) holds and  $G := \frac{k}{v}J - C$  satisfies*

$$(2) \quad G^2 = nG \text{ for some } n \geq 0;$$
*in this case  $C$  is spherical.*
- (iii)  *$C$  has strength 3 iff (1), (2) hold, and*

$$(3) \quad (G \circ G)G = mG \text{ for some } m \geq 0.$$
- (iv)  *$C$  is the distance matrix of a spherical 1-design iff  $C$  is spherical and (1) holds, of a spherical 2-design iff (1) and (2) hold, and of a spherical 3-design iff (1), (2), and (3) hold with  $m = 0$ .*

**Proof.** See Neumaier [12]. ■

A distance matrix which has strength  $t$  for all  $t$  is called a *Delsarte matrix* ([12]).

A *distance regular graph* is a graph  $\Gamma$  such that the distance relation turns  $\Gamma$  into an association scheme. It is well-known (Biggs [4]) that a graph  $\Gamma$  of diameter  $\delta$  is distance regular iff for all  $i \leq \delta$  the distance- $i$ -adjacency matrix  $A_i$  is a polynomial of degree  $i$  in the (distance-1)-adjacency matrix  $A$  of  $\Gamma$ .

For symmetric matrices  $A \in V$  with constant row sums  $k = k_A$ , we define  $C_A := k_A J - A^*$ . If  $A$  is a symmetric  $(0, 1)$ -matrix, denote by  $\Gamma(A)$  the graph whose adjacency matrix is  $A$ .

**Theorem 2.** *Let  $A \in V$  be a symmetric matrix with nonnegative diagonal and constant row sums  $k$ . Then*

- (i)  *$C_A$  is a spherical distance matrix of strength 1 iff  $A$  is nonnegative.*
- (ii)  *$C_A$  has strength 2 iff  $A$  is a  $(0, 1)$ -matrix, i.e. the adjacency matrix of a regular graph of valency  $k$ .*
- (iii)  *$C_A$  has strength 3 iff  $\Gamma(A)$  is an edge-regular graph, i.e. every edge of  $\Gamma(A)$  is in a constant number of triangles.*

- (iv)  $C_A$  is the distance matrix of a spherical 3-design iff  $\Gamma(A)$  is a triangle-free graph.
- (v)  $C_A$  is a Delsarte matrix iff  $\Gamma(A)$  is distance regular.
- (vi)  $C_A$  has repeated rows iff  $\Gamma(A)$  is disconnected.

**Proof.**

- (i)  $A$  is nonnegative  $\Rightarrow A^r$  positive semidefinite  $\Rightarrow k_A J - C_A$  is positive semidefinite  $\Rightarrow C_A$  is a spherical 1-design. Conversely,  $C_A$  is a spherical 1-design  $\Rightarrow \gamma J - C_A$  is positive semidefinite for some  $r > 0 \Rightarrow (\gamma - k_A)J + A^r$  is positive semidefinite  $\Rightarrow (\gamma - k_A)I + A$  is nonnegative  $\Rightarrow A$  is nonnegative (since  $A$  has nonnegative diagonal).
- (ii)  $C_A$  has strength 2  $\Leftrightarrow A^r A^r = v A^r \Leftrightarrow A \circ A = A \Leftrightarrow A$  is a  $(0, 1)$ -matrix.
- (iii)  $C_A$  has strength 3  $\Leftrightarrow (A^r \circ A^r) A^r = \text{const.}$   $A_r \Leftrightarrow A^2 \circ A = \text{const.}$   $A \Leftrightarrow \Gamma(A)$  is edge regular.
- (iv) Same as (iii) with constant = 0.
- (v)  $C_A$  is a Delsarte matrix  
 $\Leftrightarrow E_i = Q_i \circ A$  (where  $Q_i$  is a polynomial of degree  $i$ )  
 $\Leftrightarrow D_i = Q_i(v k_A I - v A) = \varphi_i(A)$  for some polynomial  $\varphi_i$  of degree  $i \Leftrightarrow \Gamma(A)$  is distance regular (for the last equivalence see Biggs [4]).

$$(vi) \quad C_A = \sum_i a_i D_i = k_A J - A^r \Leftrightarrow A = k_A I - \sum_i \frac{a_i}{v} D_i \Leftrightarrow$$

$$\Leftrightarrow A \text{ has eigenvalues } k_A - \frac{a_i}{v},$$

so  $C_A$  has repeated rows iff  $C_A$  has some off diagonal elements zero iff some  $a_k \neq 0$  (for  $k \neq 0$ ) iff  $A$  has multiple eigenvalue  $k_A$  iff  $\Gamma(A)$  is disconnected. ■

**Remarks. 1.** Similarly (cf. [15]),  $C_A$  has strength 4 iff in  $\Gamma(A)$  two points at distance zero, one, or two have the same number  $k$ ,  $\lambda$ , or  $\mu$  of common neighbours, respectively.

**2.** The dual of a spherical distance matrix is a *Fiedler matrix*, i.e. a symmetric matrix with zero row sums and off-diagonal entries  $\leq 0$ . It is possible to dualize the concept of strength to arbitrary Fiedler matrixes (not only to those inside a self-dual coherent algebra). Then the Fiedler matrixes of unbounded strength are essentially the matrices  $kI - A$  where  $A$  is the adjacency matrix of a distance regular graph of valency  $k$ . See Neumaier [15].

**3.** In general, the actual duality in case of matrices inside a self-dual coherent algebra is replaced by a formal duality between ordinary matrix product and element-wise product, between ranks and valencies, between Gram matrices and nonnegative matrices, etc. There are more such properties which look like duals, but the analogies are not yet understood. We outline the basic analogies.

For the following facts see e.g. Bannai and Ito [3]. Let  $\Gamma$  be a graph of diameter  $s^* = \max_{x,y} d(x, y)$ , and girth  $t^* + 1 = \text{minimal size of a circuit}$ . Then  $t^* \leq 2s^*$ , and equality holds iff  $\Gamma$  is a Moore graph. (This can be taken as the definition of a Moore graph; cf. [11]). Every Moore graph is distance regular, and the Moore graphs of valency 2 are the polygons; there are only a few Moore graphs of valency  $> 2$ ,

and in particular, there is none for  $s^* \geq 3$ . A graph  $\Gamma$  of diameter  $s^*$  and valency  $k$  contains  $v \leq v_{s^*}(k) = \frac{k(k-1)^{s^*}-2}{k-2}$  points, with equality iff  $\Gamma$  is a Moore graph;  $v_{s^*}(k)$  is a polynomial of degree  $s^*$  in  $k$ . Finally, if the valencies of a primitive association scheme are ordered such that  $k_1 \leq k_2 \leq \dots \leq k_n$  then  $k_{i+1} \leq k_i(k_1-1)$ , and equality implies that the graph with valency  $k_1$  corresponding to the first relation has girth  $t^*+1$  with  $t^* \geq 4$ .

Compare this with properties of a spherical set  $X$  of points, discussed in Delsarte, Goethals and Seidel [9]. Suppose that  $X$  has  $s$  distances and is a spherical  $t$ -design. Then  $t \leq 2s$ , and equality holds iff  $X$  is a tight design. Every tight design is a Delsarte space; and the tight designs with  $f=2$  are the polygons; there are only a few tight designs with  $f > 2$ , and in particular, there is none for  $s \geq 12$ . (See Bannai and Damerell [1], [2]). A set  $X$  with  $s$  distances in the unit sphere of  $\mathbb{R}^f$  contains

$$w \leq w_s(f) = \binom{f+s-1}{s} + \binom{f+s-2}{s-1}$$

points, with equality iff  $\Gamma$  is a tight design;  $w_s(f)$  is a polynomial of degree  $s$  in  $f$ . Finally, if the multiplicities of a primitive association scheme are ordered such that  $f_1 \leq f_2 \leq \dots \leq f_n$  then it can be shown that  $f_2 \leq \frac{1}{v}(f_1+2)(f_1-1)$ , and equality implies that the set of points corresponding to the idempotent of rank  $f_1$  is spherical 4-design.

Some immediate questions arise: How to explain the analogies? Why are the formulae slightly different in the two cases? Is there a spherical analogue of  $k_{i+1} \leq k_i(k_1-1)$  for  $i \geq 2$ ? Is there a Delsarte matrix analogue of the inequalities  $k_{i+1}k_{i-1} \leq k_i^2$  obtained by Taylor and Levingston [17] for distance regular graphs?

I want to close with some remarks on Delsarte's setting for duality. The basic observation is that if  $G$  is a  $v \times v$ -matrix such that  $GG^* = G^*G = vI$ , and  $G = (G_0 | G_1 | \dots | G_s)$  is a partition of the columns then the matrices  $E_i = \frac{1}{v} G_i G_i^*$  are mutually orthogonal idempotents, and hence define an algebra. Hence we might call two algebras of  $v \times v$ -matrices *dual* if there is a matrix  $G$  with  $GG^* = G^*G = vI$  such that a suitable partition of the columns gives the first algebra and a suitable partition of the rows gives the second algebra. Finally, we call two association schemes *dual* (in Delsarte's sense) if their Bose—Mesner algebras are dual with respect to a matrix  $G$ , and if, in addition, the column partition is given by the associates of a suitable point of the first scheme. Then it can be shown that the parameters of a dual pair of association schemes again satisfies the relations  $k'_i = f_i$ ,  $f'_i = k_i$ ,  $P' = Q$  and  $Q' = P$ .

In the case of a regular abelian group, Delsarte's duality is equivalent with our duality, since we may take  $G = (\langle x, y \rangle)_{x, y \in X}$  and the spherical point defining the partition can be taken to be zero.

**Acknowledgements.** This paper is a polished account of one of a series of lectures I gave in 1980 at the Technische Hogeschool Eindhoven. I recall with pleasure the time I spent there, supported by a grant No. B—61—99 of the Netherlands Organization for the Advancement of Pure Research (Z.W.O.), the many discussions with



Prof. Seidel, Aart Blokhuis, and the audience of my lectures who encouraged me to organize my ideas on the subject to a coherent whole. I also want to thank René Klerx for providing a set of lecture notes [16], whose last chapter formed the basis for the present paper.

### References

- [1] E. BANNAI and R. M. DAMERELL, Tight spherical designs, *I. J. Math. Soc. Japan*, **31** (1979), 199—207.
- [2] E. BANNAI and R. M. DAMERELL, Tight spherical designs, *II. J. London Math. Soc. (2)*, **21** (1980), 13—30.
- [3] E. BANNAI and T. ITO, *Algebraic Combinatorics I: Association schemes*. Benjamin/Cummings, Menlo Park, Calif. 1984.
- [4] N. L. BIGGS, *Algebraic Graph Theory*. Cambridge University Press, Cambridge 1974.
- [5] R. C. BOSE and D. M. MESNER, On linear associative algebras corresponding to association schemes of partially balanced designs, *Ann. Math. Statist.* **30** (1959), 21—38.
- [6] R. H. BRUCK, Finite nets I: Numerical invariants, *Canad. J. Math.* **3** (1951), 94—107.
- [7] P. J. CAMERON, J. M. GOETHALS and J. J. SEIDEL, Strongly regular graphs having strongly regular subconstituents, *J. Algebra* **55** (1978), 257—280.
- [8] PH. DELSARTE, An algebraic approach to the association schemes of coding theory, *Philips Res. Rep. Suppl.* **10** (1973).
- [9] PH. DELSARTE, J. M. GOETHALS and J. J. SEIDEL, Spherical codes and designs, *Geom. Dedicata* **6** (1977), 363—388.
- [10] D. G. HIGMAN, Coherent configurations, part I: Ordinary representation theory, *Geom. Dedicata* **4** (1975), 1—32.
- [11] A. J. HOFFMAN and R. R. SINGLETON, On Moore graphs with diameters 2 and 3, *IBM J. Res. Develop.* **4** (1960), 497—504.
- [12] A. NEUMAIER, Distances, graphs, and designs, *Europ. J. Combinatorics* **1** (1980), 163—174.
- [13] A. NEUMAIER, Distance matrices and  $n$ -dimensional designs, *Europ. J. Combinatorics* **2** (1981), 165—172.
- [14] A. NEUMAIER, Distance matrices, dimension, and conference graphs, *Indagationes Math.* **43** (1981), 385—391.
- [15] A. NEUMAIER, Classification of graphs by regularity, *J. Combin. Theory (Ser. B)* **30** (1981), 318—331.
- [16] A. NEUMAIER, Combinatorial configurations in terms of distances, *Memorandum 81—09 (Wiskunde)*, TH Eindhoven 1981.
- [17] D. E. TAYLOR and R. LEVINGSTON, Distance-regular graphs, in: *Combinatorial Mathematics* (D. A. Holton and J. Seberry, eds.), *Springer Lecture Notes in Mathematics* **686** (1978), 313—323.
- [18] H. WIELANDT, *Finite permutation groups*, Academic Press, New York, 1964.
- [19] JA. JU. GOLDFAND and M. H. KLIN, Amorphic cellular rings I, *Investigations in algebraic theory of combinatorial objects*, Moscow, Inst. for Systems Studies, 1985, pp. 32—39.
- [20] J. L. HAYDEN, Association algebras for finite projective planes, *J. Combin. Theory (Ser. A)* **34** (1983), 360—374.
- [21] D. G. HIGMAN, Coherent algebras, *Linear Algebra Appl.* **93** (1987), 209—240.

A. Neumaier

Institut für Angewandte Mathematik  
Universität Freiburg i. Br.  
West-Germany