

## EXISTENCE OF POLYHEDRAL EMBEDDINGS OF GRAPHS

BOJAN MOHAR\*

Received July 6, 1998

It is proved that the decision problem about the existence of an embedding of face-width 3 of a given graph is **NP**-complete. A similar result is proved for some related decision problems. This solves a problem raised by Neil Robertson.

### 1. Introduction

Let  $C$  and  $C'$  be cycles in a graph  $G$ . We say that  $C$  and  $C'$  *meet properly* if the intersection of  $C$  and  $C'$  is either empty, a single vertex or an edge.

Let  $G$  be a 3-connected graph. A 2-cell embedding of  $G$  in some surface is *polyhedral* if every facial walk is a cycle and any two facial cycles meet properly. Equivalently, we require that the graph is 3-connected and that the embedding has face-width at least three [8] (cf. also [1, 5, 6]). Let us recall that the *face-width* (also called the *representativity*) of a (2-cell) embedded graph  $G$  is the minimum integer  $r$  such that  $G$  has  $r$  facial walks whose union contains a cycle which is noncontractible on the surface. (In the case when there are no noncontractible cycles, we let the face-width be  $\infty$ .)

At the Seventh Vermont Summer Workshop on Combinatorics and Graph Theory in 1995, Neil Robertson asked how difficult it is to see whether a given 3-connected graph admits a polyhedral embedding. In this note we answer his question by proving that the decision problem about the

---

*Mathematics Subject Classification (2000):* 05C10, 68Q17

\* Supported in part by the Ministry of Science and Technology of Slovenia, Research Project J1-0502-0101-98.

existence of polyhedral embeddings is **NP**-complete. The problem remains **NP**-complete even if we ask about polyhedral embeddings in orientable surfaces and require that the given graph is 6-connected. A similar problem where we ask about embeddings of face-width exactly 3 is also **NP**-complete. However, it is not known if existence of embeddings of face-width 4 or more is still **NP**-complete.

An indication that there is some nontriviality in polyhedral embeddings is that for a complete graph  $K_n$  ( $n \geq 5$ ), a polyhedral embedding is necessarily a triangulation, and a significant part of Ringel and Youngs' Map Color Theorem [7] was to determine which complete graphs have such embeddings. Our result is not that much of interest from the computational complexity point of view. Its main message is that any theory on polyhedral embeddings is rich and interesting.

It is worth mentioning that a similar problem concerning embeddings of face-width at least two may be polynomially solvable. This problem is easily reduced to 2-connected graphs (cf. [6, 8]), and there are two long standing conjectures which are closely related to the Cycle Double Cover Conjecture (cf. [4, 10]), and whose affirmative solution would give a trivial answer about existence of embeddings of face-width at least 2.

**Conjecture 1.1** (Haggard [3]). Every 2-connected graph has an embedding of face-width 2 or more.

**Conjecture 1.2** (Jaeger [4]). Every 2-connected graph has an orientable embedding of face-width 2 or more.

## 2. Embeddings and compatible cycles

Our treatment of graph embeddings follows essentially [6]. All graphs are simple, so there are no loops or multiple edges. We only consider 2-cell embeddings into closed surfaces which can be defined combinatorially as follows. An *embedding* of a connected graph  $G$  is a pair  $\Pi = (\pi, \lambda)$  where  $\pi = \{\pi_v \mid v \in V(G)\}$  is a collection of *local clockwise rotations*, i.e.,  $\pi_v$  is a cyclic permutation of the edges incident with  $v$  ( $v \in V(G)$ ), and  $\lambda: E(G) \rightarrow \{+1, -1\}$  is a *signature*. The local rotation  $\pi_v$  describes the cyclic clockwise order of edges incident with  $v$  on the surface, and the signature  $\lambda(uv)$  of the edge  $uv$  is positive if and only if the local rotations  $\pi_u$  and  $\pi_v$  both correspond to the clockwise (or both to anticlockwise) rotations when traversing the edge  $uv$  on the surface. An embedding of a graph  $G$  is *nonorientable* if  $G$  contains a cycle whose number of edges with negative signature is odd.

The embedding  $\Pi$  determines a set of  $\Pi$ -facial walks. If a  $\Pi$ -facial walk is a cycle, it is also called a  $\Pi$ -facial cycle. The underlying surface of the embedding  $\Pi$  is obtained by pasting discs along the  $\Pi$ -facial walks in  $G$ .

Let  $G$  be a graph. Two subgraphs  $H_1, H_2$  of  $G$  are said to be *compatible* if  $E(H_1) \cap E(H_2)$  is a matching in  $G$ . Equivalently, no two edges of  $H_1$  incident with the same vertex are both contained in  $H_2$ .

Let  $G$  be a  $\Pi$ -embedded graph and let  $v \in V(G)$ . Let  $H$  be the subgraph of  $G$  consisting of all neighbors of  $v$  and all edges  $uw$  such that  $vuw$  is a  $\Pi$ -facial cycle. Then  $H$  is called the *link* of  $v$ , and is denoted by  $\text{link}(v, G, \Pi)$ .

**Lemma 2.3.** *Let  $G$  be a  $\Pi$ -embedded graph and let  $u, v$  be distinct vertices of  $G$ . If no vertex adjacent to  $v$  is of degree 4 in  $G$ , then  $\text{link}(u, G, \Pi)$  and  $\text{link}(v, G, \Pi)$  are compatible subgraphs of  $G$  whose maximum degree is at most 2.*

**Proof.** Each edge  $vw$  is contained in at most two facial triangles. Therefore, the maximum degree in the link of  $v$  does not exceed 2. If  $\text{link}(u, G, \Pi)$  and  $\text{link}(v, G, \Pi)$  share two edges  $aw, bw$  incident with the same vertex  $w$ , then the link of  $w$  is the cycle  $avbu$  and  $w$  is of degree 4. This completes the proof. ■

Thomassen [9] proved:

**Theorem 2.4 (Thomassen [9]).** *The decision problem whether a given cubic bipartite graph contains two compatible Hamilton cycles is NP-complete.*

The cubic bipartite graphs  $G$  in Thomassen's proof of Theorem 2.4 in [9] are 2-connected and contain many edges which are contained in any Hamilton cycle of  $G$ . Such edges are easily discovered in  $G$ . This shows that the same problem is NP-complete also when the input graph is 2-connected and has three prescribed edges which are contained in every Hamilton cycle of  $G$ .

Let  $T$  be a tree of maximum degree  $d$ , and suppose that  $G_0$  is a graph and  $e_1, \dots, e_d$  are edges of  $G_0$ . Take a distinct copy  $G_t$  of  $G_0$  for each vertex  $t \in V(T)$ . Label each oriented edge  $tt'$  of  $T$  by a number in  $\{1, \dots, d\}$  so that the edges emanating from the same vertex receive distinct labels. Now, for each edge  $tt' \in E(T)$ , repeat the following operation. Let  $a$  and  $b$  be the labels of  $tt'$  and  $t't$ , respectively. Remove the edge  $e_a = xy$  from  $G_t$ , remove  $e_b = x'y'$  from  $G_{t'}$ , and add the edges  $xx'$  and  $yy'$ . Let  $G$  be any graph resulting from this operation.

**Lemma 2.5.** *Let  $G_0$  be a graph and let  $T$  be a tree of maximum degree  $d$ . Let  $G$  be a graph constructed as described above, and let  $e_1, \dots, e_d$  be the edges of  $G_0$  used in the construction. Then  $G$  contains two compatible Hamilton cycles if and only if  $G_0$  contains two compatible Hamilton cycles each of which contains all edges  $e_1, \dots, e_d$ .*

**Proof.** Suppose first that  $G$  contains two compatible Hamilton cycles  $H_1, H_2$ . We shall use the notation introduced in the definition of  $G$ . Suppose that  $t$  is a vertex of degree  $d$  in  $T$ . The removal of the edges  $xx'$  and  $yy'$  disconnects the graph  $G$ . Let  $G'$  be the component of  $G - xx' - yy'$  which contains  $V(G_t)$ . Clearly,  $xx'$  and  $yy'$  are both contained in  $H_1$  and in  $H_2$ . Therefore,  $H'_1 = (H_1 \cap G') + xy$  and  $H'_2 = (H_2 \cap G') + xy$  are compatible Hamilton cycles of  $G' + xy$ . By repeating such a reduction for all edges incident with  $t$  in  $T$ , we obtain two compatible Hamilton cycles of  $G_t$  (and hence of  $G_0$ ) which contain all edges  $e_1, \dots, e_d$ .

Suppose now that  $G_0$  contains two compatible Hamilton cycles  $H_1^\circ$  and  $H_2^\circ$  each of which contains all edges  $e_1, \dots, e_d$ . We shall prove by induction on  $|V(T)|$  that  $G$  admits two compatible Hamilton cycles  $H_1, H_2$  such that all edges  $e_1, \dots, e_d$  in each copy  $G_t$  ( $t \in V(T)$ ) which remain in  $G$  are contained in  $H_1$  and in  $H_2$ . (Here we allow that  $d$  is larger than the maximum degree in  $T$ .) This is clear if  $|V(T)| = 1$ . Otherwise, let  $t$  be a leaf of  $T$ , and let  $t'$  be the neighbor of  $t$  in  $T$ . Let  $G' = (G - V(G_t)) + x'y'$ . Then  $G'$  is obtained from  $G_0$  and  $T - t$  in the same way as described before the lemma. By the induction hypothesis,  $G'$  has two compatible Hamilton cycles  $H'_1, H'_2$  which contain  $e_b = x'y'$  (and all other edges  $e_1, \dots, e_d$  in each copy  $G_s$ ,  $s \in V(T - t)$ , which remain in  $G'$ ). Let  $H_j = (H'_j - x'y') \cup (H_j^\circ - xy) + xx' + yy'$ ,  $j = 1, 2$ . Then  $H_1$  and  $H_2$  are compatible Hamilton cycles in  $G$  with the desired property. ■

**Lemma 2.6.** *Let  $G_0$ ,  $T$ , and  $G$  be as in Lemma 2.5. Suppose that  $T$  has more than  $4k$  leaves where  $k$  is a positive integer. If  $G$  contains two compatible spanning subgraphs  $H_1, H_2$  such that for  $i = 1, 2$ , the maximum degree in  $H_i$  is at most two and such that the number of connected components of  $H_i$  is  $\leq k$ , then  $G_0$  contains two compatible Hamilton cycles.*

**Proof.** Note that both  $H_1$  and  $H_2$  are disjoint unions of isolated vertices, paths, and cycles. Let  $U$  be the vertex set of  $G$  containing all vertices of degree less than 2 in  $H_1$  or in  $H_2$ , and containing one vertex of each cycle in  $H_1$  or in  $H_2$ . Then  $|U| \leq 4k$ , and hence there is a leaf  $t$  of  $T$  such that  $U \cap V(G_t) = \emptyset$ . Then  $H_1 \cap G_t$  and  $H_2 \cap G_t$  give rise to two compatible Hamilton cycles in  $G_0$ . ■

### 3. Reduction

**Theorem 3.7.** *The decision problem “Does a given graph  $G$  have a polyhedral embedding” is **NP**-complete. The problem remains **NP**-complete also if we ask about polyhedral embeddings in orientable surfaces and require that  $G$  is 6-connected.*

**Proof.** Let  $G_0$  be an arbitrary 2-connected cubic bipartite graph, and let  $e_1, e_2, e_3 \in E(G_0)$  be distinct edges of  $G_0$  such that every Hamilton cycle of  $G_0$  contains each of them. By Theorem 2.4 (and the remark following it), it is **NP**-complete to decide if  $G_0$  has two compatible Hamilton cycles. Thus, Theorem 3.7 will follow if we prove that one can construct in polynomial time a 6-connected graph  $G_1$  which has a polyhedral embedding if and only if  $G_0$  has two compatible Hamilton cycles and, moreover, if  $G_1$  has a polyhedral embedding, then it also has an orientable polyhedral embedding.

Let  $T$  be a cubic tree (i.e., each vertex of  $T$  is of degree 3 or 1) of order 104, so that  $T$  has 53 leaves. Construct the graph  $G$  as described before Lemma 2.5. Clearly,  $G$  is cubic, bipartite and 2-connected. Let  $V(G) = V_1 \cup V_2$  be the bipartition of  $G$ . Now, define the graph  $G_1$  which is obtained from  $G$  as follows. First, replace each vertex  $v \in V_2$  by two mutually adjacent vertices  $v', v''$  which are both adjacent to the same three vertices in  $V_1$  as  $v$ . Let  $V' = \{v' \mid v \in V_2\}$  and  $V'' = \{v'' \mid v \in V_2\}$ . Finally, add four new vertices  $a', b', a'', b''$  where  $a'$  and  $b'$  are adjacent to all vertices in  $V_1 \cup V'$ , and  $a'', b''$  are adjacent to all vertices in  $V_1 \cup V''$ .

The resulting graph  $G_1$  is 6-connected. To see this, one considers each pair  $x, y$  of vertices and shows that there are 6 internally disjoint paths joining  $x$  and  $y$ . The details are rather straightforward and are left to the reader.

We claim that  $G_0$  contains two compatible Hamilton cycles if and only if  $G_1$  has a polyhedral embedding. First, assume that  $G_0$  admits two compatible Hamilton cycles. Since every Hamilton cycle of  $G_0$  contains  $e_1, e_2$ , and  $e_3$ , Lemma 2.5 shows that  $G$  has two compatible Hamilton cycles, say  $H_1$  and  $H_2$ . For  $i = 1, 2$ , let  $H'_i$  (resp.  $H''_i$ ) be the cycle in  $G_1$  obtained from  $H_i$  by replacing each vertex  $v \in V_2$  by the vertex  $v' \in V'$  (resp.  $v'' \in V''$ ). It is easy to see that  $G_1$  has (a unique) embedding in which all facial cycles are triangles such that the link of  $a'$  (resp.  $b', a'', b''$ ) is  $H'_1$  (resp.  $H'_2, H''_1, H''_2$ ). This embedding is clearly polyhedral. It is also orientable. To see this, orient the facial triangles as follows:  $a'v'v_1$  (if  $v'v_1 \in E(H'_1) \setminus E(H'_2)$ ),  $a'v_1v'$  (if  $v'v_1 \in E(H'_1) \cap E(H'_2)$ ), similarly around  $b'$ ,  $v_1v'v''$  (if  $v'v_1 \in E(H'_1)$ ),  $v_1v''v'$  (if  $v'v_1 \in E(H'_2)$ ), where  $v_1 \in V_1$ ,  $v' \in V'$ , and  $v'' \in V''$ . Similarly we orient

the triangles containing the edges of  $H_1''$  and  $H_2''$ . The details are left to the reader.

Conversely, let  $\Pi$  be a polyhedral embedding of  $G_1$ . Let us consider the  $\Pi$ -facial cycles containing  $a'$ . Each such facial cycle  $C = a'v_1v_2 \dots v_k$  is an induced cycle in  $G_1$ . (If  $C$  had a chord  $e$ , then a facial cycle containing  $e$  would meet improperly with  $C$ .) We say that  $C$  is *exceptional* if  $k > 2$ . It is *strongly exceptional* if  $V(C)$  contains at least one of the vertices  $b', a'', b''$ , and *weakly exceptional* otherwise. There are at most three strongly exceptional faces containing  $a'$  since no two strongly exceptional faces contain the same pair of (nonconsecutive) vertices  $\{a', x\}$ ,  $x \in \{b', a'', b''\}$ .

An exceptional face  $C = a'v_1 \dots v_k$  is induced. Therefore,  $v_1, v_k \in V_1 \cup V'$  and  $v_2, \dots, v_{k-1} \notin V_1 \cup V'$ . Similar conclusions hold for the exceptional faces at the vertices  $b', a''$ , and  $b''$ . This implies that at most 10 vertices of  $V''$  belong to strongly exceptional faces at the vertices  $a', b', a'', b''$ .

Suppose now that  $C$  is weakly exceptional. Then  $k = 3$  since  $v_2 \in V''$ , and hence  $v_3$  is a neighbor of  $a'$ . Let  $v \in V_2$  be the vertex such that  $v_2 = v''$ . If  $v_1 \in V'$ , then  $v_1 = v'$  and thus  $v_1v_3 \in E(G_1)$ , a contradiction. Hence  $v_1, v_3 \in V_1$ . As mentioned above, there are at most 10 vertices in  $V''$  contained in a strongly exceptional face. Therefore, there are at most 10 weakly exceptional facial cycles containing  $a'$  such that  $v_2$  is contained in some strongly exceptional facial cycle.

Suppose now that  $C$  is not such a face. Consider the  $\Pi$ -clockwise ordering around  $v''$ . If the edges  $v''a''$  and  $v''v'$  are consecutive in that ordering, the facial cycle containing these two edges is not induced (as we just proved above when considering the possibility that  $v_1 \in V'$ ). Similarly,  $v''b''$  and  $v''v'$  are not consecutive around  $v''$ . In particular,  $v_2 = v''$  belongs to a strongly exceptional face containing  $a''$  and  $b''$ , a contradiction. This implies that there are at most 10 weakly exceptional and at most 3 strongly exceptional faces containing  $a'$ . This shows that  $\text{link}(a', G_1, \Pi)$  is a subgraph of  $G_1$  of maximum degree at most 2 and with at most 13 connected components. The same holds for  $\text{link}(b', G_1, \Pi)$ . By [Lemma 2.3](#), these links are compatible subgraphs of  $G_1$ . Clearly, they give rise to compatible subgraphs in  $G$ . Since  $T$  has 53 leaves, [Lemma 2.6](#) implies that  $G_0$  (and hence also  $G$  by [Lemma 2.5](#)) contains two compatible Hamilton cycles. Additionally, as the previous paragraph shows,  $G_1$  admits an orientable polyhedral embedding determined by two compatible Hamilton cycles of  $G$ .

We reduced, in polynomial time, the **NP**-complete problem of [Theorem 2.4](#) to the existence of polyhedral embeddings of 6-connected graphs. Since the embedding of  $G_1$  obtained from two compatible Hamilton cycles in  $G_0$  (and in  $G$ ) is orientable, this completes the proof. ■

The proof of [Theorem 3.7](#) shows that in every embedding  $H$  of  $G_1$  of face-width at least 3, the link of  $a'$  determines a Hamilton path in one of the subgraphs  $G_t$  of  $G$  where  $t$  is some leaf of  $T$ . This can be used to show that  $G_1$  has no embeddings of face-width 4 or more. Hence, the same proof also shows:

**Corollary 3.8.** *The decision problem “Does a given 6-connected graph  $G$  have an (orientable) embedding of face-width exactly 3” is NP-complete.*

## References

- [1] D. BARNETTE: Polyhedral maps on 2-manifolds, in: *Convexity and Related Combinatorial Geometry* (Norman, Okla., 1980), Dekker, New York, 1982, 7–19.
- [2] J. A. BONDY, U. S. R. MURTY: *Graph Theory with Applications*, North-Holland, New York, 1981.
- [3] G. HAGGARD: Edmonds’ characterization of disc embeddings, *Congr. Numer.*, **19** (1977), 291–302.
- [4] F. JAEGER: A survey of the cycle double cover conjecture, in: *Cycles in Graphs* (B. Alspach and C. Godsil, Eds.), *Ann. Discrete Math.*, **27** (1985), 1–12.
- [5] B. MOHAR: Face-width of embedded graphs, *Math. Slovaca*, **47** (1997), 35–63.
- [6] B. MOHAR, C. THOMASSEN: *Graphs on Surfaces*, Johns Hopkins University Press, to appear.
- [7] G. RINGEL: *Map Color Theorem*, Springer-Verlag, Berlin, 1974.
- [8] N. ROBERTSON, R. P. VITRAY: Representativity of surface embeddings, in: *Paths, Flows, and VLSI-Layout* (B. Korte, L. Lovász, H. J. Prömel, and A. Schrijver, eds.), Springer-Verlag, Berlin, 1990, 293–328.
- [9] C. THOMASSEN: Triangulating a surface with a prescribed graph, *J. Combin. Theory Ser. B*, **57** (1993), 196–206.
- [10] C.-Q. ZHANG: *Integer Flows and Cycle Covers of Graphs*, Marcel Dekker, New York, 1996.

Bojan Mohar

*Department of Mathematics,*

*University of Ljubljana,*

*1111 Ljubljana, Slovenia*

[bojan.mohar@uni-lj.si](mailto:bojan.mohar@uni-lj.si)