

THE PARITY PROBLEM OF POLYMATROIDS WITHOUT
DOUBLE CIRCUITS*

MÁRTON MAKAI, JÁCINT SZABÓ

Received February 7, 2007

According to the present state of the theory of the matroid parity problem, the existence of a good characterization to the size of a maximum matching depends on the behavior of certain substructures, called double circuits. In this paper we prove that if a polymatroid has no double circuits then a partition type min-max formula characterizes the size of a maximum matching. Applications to parity constrained orientations and to a rigidity problem are given.

1. Introduction

Polymatroid parity (a.k.a. matroid parity or matroid matching) is an involved notion of combinatorial optimization concerning parity. Early known special cases of it are the matching problem of graphs and the matroid intersection problem, which in fact motivated Lawler to introduce this notion. Jensen and Korte [6], and Lovász [11] have shown that in general the matroid parity problem is of exponential complexity under the independence oracle framework. However, Lovász gave a good characterization to the size of a maximum matching and also a polynomial algorithm for linearly represented matroids [10, 11]. Lovász [9], and Dress and Lovász [1] manifest that the solubility of the linear case is due to the fact that these matroids can be embedded into a matroid sharing the so-called *double circuit property*, or *DCP* for short. It was also shown that full linear, full algebraic,

Mathematics Subject Classification (2000): 05B35

* Research is supported by OTKA grants K60802, TS049788 and by European MCRTN Adonet, Contract Grant No. 504438.

full graphic, and full transversal matroids are DCP matroids [1]. The subsequent results in the literature are so as to guarantee the DCP in particular matroid classes. The disadvantage of this approach is that, due to the embedding into a bigger matroid, in many cases the obtained min-max formula cannot be translated into a combinatorial form. However, the diversity and the importance of solvable special cases of the matroid parity problem force to search for techniques which yield combinatorial characterizations.

In this work we investigate the class of those polymatroids which have no non-trivial double circuits. We prove that in these polymatroids a partition type combinatorial formula characterizes the size of a maximum matching. Thereafter, two applications are presented. First we show that the parity constrained orientation problem of Király and Szabó [7] can be formulated as a polymatroid parity problem in such a way that the polymatroid in question has no non-trivial double circuits, yielding the partition type formula of [7]. Second, we deduce a result of Fekete [2] about adding a clique of minimum size to a graph resulting in a generic rigid graph in the plane.

To formulate our main result, some definitions are in order. We denote by \mathbb{R}_+ and \mathbb{Z}_+ the set of non-negative reals and non-negative integers resp. Let S be a finite ground set. A set-function $f: 2^S \rightarrow \mathbb{Z}$ is said to be *submodular* if

$$(1) \quad f(X) + f(Y) \geq f(X \cap Y) + f(X \cup Y)$$

holds whenever $X, Y \subseteq S$. The set-function f is said to be *non-decreasing* if $f(X) \leq f(Y)$ for every $\emptyset \neq X \subseteq Y \subseteq S$, and we say that f is *non-increasing* if $-f$ is non-decreasing. A non-decreasing submodular set-function $f: 2^S \rightarrow \mathbb{Z}_+$ with $f(\emptyset) = 0$ is called a *polymatroid function*. A vector $x \in \mathbb{Z}_+^S$ is said to be *even* if x_i is even for every $i \in S$. The even vectors $m \in \mathbb{Z}_+^S$ with $m(U) \leq f(U)$ for every $U \subseteq S$ are said to be the *matchings* of f . A matching m with maximum $m(S)$ is said to be a *maximum matching* of f , and we denote

$$\nu(f) = \max\{m(S)/2 : m \text{ is a matching of } f\}.$$

The *polymatroid parity* problem is to determine the value of $\nu(f)$ for a polymatroid function f . Our main result is as follows. The notion of a non-trivial compatible double circuit will be defined in [Section 2](#).

Theorem 1.1. *Let $f: 2^S \rightarrow \mathbb{Z}_+$ be a polymatroid function having no non-trivial compatible double circuits. Then*

$$\nu(f) = \min \sum_{j=1}^t \left\lfloor \frac{f(U_j)}{2} \right\rfloor,$$

where the minimum is taken over all partitions U_1, U_2, \dots, U_t of S .

Recently, Pap [13] developed a deep combinatorial algorithm for finding a maximum matching in a polymatroid without non-trivial compatible double circuits. His result serves as a completion of Theorem 1.1, as we do not concern algorithmic aspects in this paper.

2. Preliminaries and proof of the main theorem

We recollect some important notions and results from the theory of matroids, polymatroids and matroid parity. We omit the proofs of some well-known facts, as the details can be found e.g. in [14]. If $x \in \mathbb{R}^S$ and $U \subseteq S$ then we use the notations $x(U) = \sum_{u \in U} x(u)$ and

$$(x|_U)(i) = \begin{cases} x_i & \text{if } i \in U, \\ 0 & \text{if } i \notin U. \end{cases}$$

If $f: 2^S \rightarrow \mathbb{Z}_+$ is a function then we define the polyhedra

$$P(f) = \{x \in \mathbb{R}_+^S : x(U) \leq f(U) \text{ for every } U \subseteq S\}, \quad \text{and} \\ B(f) = \{x \in \mathbb{R}_+^S : x(U) \leq f(U) \text{ for every } U \subseteq S, \ x(S) = f(S)\}.$$

In the theory of submodular functions it is proved that if f is a polymatroid function then $P(f)$ and $B(f)$ are nonempty integer polyhedra. Recall that an even vector $m \in \mathbb{Z}_+^S$ is a matching of f if and only if $m \in P(f)$.

There is a close relation between polymatroid functions and matroids. First, if $M = (T, r)$ is a matroid and $\varphi: T \rightarrow S$ is a function then $f: 2^S \rightarrow \mathbb{Z}_+$, $X \mapsto r(\varphi^{-1}(X))$ is a polymatroid function, the *homomorphic image of M under φ* . Second, for any polymatroid function f it is possible to define a matroid M , the homomorphic image of which is f , in such a way that M is “most independent” in some sense. The ground set T of M is the disjoint union of sets T_i for $i \in S$ of size $|T_i| \geq f(\{i\})$. If $X \subseteq T$ then we define the vector $\chi^X \in \mathbb{Z}_+^S$ with $\chi_i^X = |X \cap T_i|$ for $i \in S$. With this notation, a set $X \subseteq T$ is defined to be independent in M if $\chi^X \in P(f)$. It is routine to prove that M is indeed a matroid with rank function

$$r(X) = \min_{Y \subseteq X} (|Y| + f(\varphi(X - Y))),$$

where $\varphi: T \rightarrow S$ maps t to i if $t \in T_i$. This M is called a *prematroid* of f . Note that a prematroid M is uniquely determined by f and by the sizes $|T_i|$, $i \in S$. If M is a matroid with rank function r then the prematroids of r are the parallel extensions of M . If we consider a prematroid M then we tacitly assume that $M = (T, r)$ and that the function $\varphi: T \rightarrow S$ is given with $t \mapsto i$ if $t \in T_i$.

If f is a polymatroid function and $x \in \mathbb{Z}_+^S$ then we define the *rank* of x as

$$(2) \quad r_f(x) = \min_{U \subseteq S} (x(U) + f(S - U)).$$

If $x \in \mathbb{Z}_+^S$ then $r_f(x) = x(S)$ if and only if $x \in P(f)$. Besides, if $M = (T, r)$ is a prematroid of f and $X \subseteq T$ then

$$(3) \quad r_f(\chi^X) = r(X).$$

The *span* of $x \in \mathbb{Z}_+^S$ is defined by $\text{sp}_f(x) = \{i \in S : r_f(x + \chi_i) = r_f(x)\}$. If M is a prematroid of f and $X \subseteq T$ then $\text{sp}_f(\chi^X) = \{i \in S : T_i \subseteq \text{sp}_M(X)\}$.

Let $M = (T, r)$ be a matroid. A set $C \subseteq T$ is said to be a *circuit* if $r(C) = |C| - 1$ and $r(C - x) = |C| - 1$ for every $x \in C$. A set $D \subseteq T$ is a *double circuit* if $r(D) = |D| - 2$ and $r(D - x) = |D| - 2$ for every $x \in D$. If D is a double circuit then the dual of $M|D$ is a matroid of rank 2 without loops, that is a line, showing that there exists a partition called *principal partition* $D = D_1 \dot{\cup} D_2 \dot{\cup} \dots \dot{\cup} D_d$, $d \geq 2$, such that the circuits of D are exactly the sets of the form $D - D_i$, $1 \leq i \leq d$. We say that D is *non-trivial* if $d \geq 3$, and *trivial* otherwise. A trivial double circuit is simply the direct sum of two circuits.

Analogously, we define circuits and double circuits of the polymatroid function $f : 2^S \rightarrow \mathbb{Z}_+$. For a vector $x \in \mathbb{R}_+^S$ let $\text{supp}(x) = \{i \in S : x_i > 0\}$. A vector $c \in \mathbb{Z}_+^S$ is a *circuit* of f if $r_f(c) = c(S) - 1$ and $r_f(c - \chi_i) = c(S) - 1$ for every $i \in \text{supp}(c)$. A vector $w \in \mathbb{Z}_+^S$ is a *double circuit* of f if $r_f(w) = w(S) - 2$ and $r_f(w - \chi_i) = w(S) - 2$ for every $i \in \text{supp}(w)$.

The exact relation between matroidal and polymatroidal double circuits is as follows, which can be seen easily from (3).

Lemma 2.1. *Let M be a prematroid of f , $D \subseteq T$ and $\chi^D = w$. Then D is a double circuit of M if and only if w is a double circuit of f .*

Let M be a prematroid of f and w be a double circuit of f such that there is a set $D \subseteq T$ with $\chi^D = w$. By Lemma 2.1, D is a double circuit of M , thus it has a principal partition $D = D_1 \dot{\cup} D_2 \dot{\cup} \dots \dot{\cup} D_d$. Due to the structure of prematroids it is easy to check that $\text{supp}(w)$ has a partition $W_0 \dot{\cup} W_1 \dot{\cup} \dots \dot{\cup} W_d$ (called the *principal partition* of w) with the property that each set D_j is either a singleton belonging to some T_i with $w_i \geq 2$ and $i \in W_0$, or is equal to $D \cap \bigcup_{i \in W_h} T_i$ for some $1 \leq h \leq d$. Note that a partition $W_0 \dot{\cup} W_1 \dot{\cup} \dots \dot{\cup} W_d$ of $\text{supp}(w)$ is the principal partition of w if and only if $w - \chi_i$ is a circuit of f and $w_i \geq 2$ whenever $i \in W_0$, moreover, $w|_{W - W_i}$ is a circuit of f for each $1 \leq i \leq d$. A double circuit w is said to be *compatible* if $W_0 = \emptyset$, and it is *non-trivial* if D is non-trivial.

We shortly mention what is the double circuit property (DCP). If $M = (T, r)$ is a prematroid of the polymatroid function f and $Z \subseteq T$ then $\varphi(M/Z)$ is called a *contraction* of f . A polymatroid function f is said to have the DCP if whenever w is a non-trivial compatible double circuit in a contraction f' of f with principal partition $W_1 \dot{\cup} \dots \dot{\cup} W_d$ then $f'(\bigcap_{1 \leq i \leq d} \text{sp}_{f'}(w|_{W-W_i})) > 0$, (Dress, Lovász [1]). A polymatroid function without non-trivial compatible double circuits has not necessarily the DCP, as its contractions may have many non-trivial compatible double circuits.

Note that every polymatroid function has double circuits, say $(f(\{i\}) + 2)\chi_i$ for some $i \in S$. However, these are not compatible, because $W_0 = \{i\}$.

Lemma 2.2. *If $w \in \mathbb{Z}_+^S$ is a double circuit of the polymatroid function $f : 2^S \rightarrow \mathbb{Z}_+$ with principal partition $W = W_0 \dot{\cup} W_1 \dot{\cup} \dots \dot{\cup} W_d$ then $f(W) = w(W) - 2$ and $f(W - W_i) = w(W - W_i) - 1$ for $1 \leq i \leq d$.*

Proof. We prove that if $x \in \mathbb{Z}_+^S$ is a vector with the property that $r_f(x) = r_f(x - \chi_i)$ for all $i \in \text{supp}(x)$ then $f(\text{supp}(x)) = r_f(x)$. By definition, $r_f(x) = x(U) + f(S - U)$ for some $U \subseteq S$. Note that $r_f(x - \chi_i) \leq (x - \chi_i)(U) + f(S - U) = r_f(x) - 1$ for all $i \in \text{supp}(x) \cap U$. Thus $\text{supp}(x) \cap U = \emptyset$. Concluding, $f(S - U) = r_f(x) \leq f(\text{supp}(x)) \leq f(S - U)$, because f is non-decreasing. If x is a circuit or a double circuit then $r_f(x) = r_f(x - \chi_i)$ for all $i \in \text{supp}(x)$, so we are done. ■

Next we explore how two polymatroid operations alter double circuits.

Translation

If $f : 2^S \rightarrow \mathbb{Z}$ is a function and $n \in \mathbb{Z}^S$ then define $f + n : 2^S \rightarrow \mathbb{Z}$ by $X \mapsto f(X) + n(X)$. If f is a polymatroid function and $n \in \mathbb{Z}_+^S$ then $f + n$ is clearly a polymatroid function, too.

Claim 2.3. *Let $n \in \mathbb{Z}^S$, and let f and $f + n$ be polymatroid functions. Then a vector $w \in \mathbb{Z}_+^S$ is a double circuit of f with $W = \text{supp}(w)$ if and only if $w + n|_W$ is a double circuit of $f + n$. In this case their principal partition coincide.*

Proof. Let $y \in \mathbb{Z}_+^S$ and assume that $y' := y + n|_{\text{supp}(y)} \in \mathbb{Z}_+^S$. Note that in (2) we can always choose U with $U \cup \text{supp}(y) = S$, thus $r_{f+n}(y') - y'(S) = r_f(y) - y(S)$. Hence by symmetry, it is enough to prove that if w is a double circuit of f with support W then $\text{supp}(w + n|_W) = W$. Otherwise $w_i + n_i \leq 0$ for some $i \in W$, thus by Lemma 2.2 we would have

$$w(W - i) \geq w(W) + n_i = f(W) + n_i + 2 \geq f(W - i) + 2,$$

which is impossible. ■

Deletion or upper bound

Let $u \in \mathbb{Z}_+^S$ be a bound vector and define $f \setminus u = \varphi(r_{M|Z})$ where M is a prematroid of f and $Z \subseteq T$ with $\chi^Z = u$. The matroid union theorem asserts that $(f \setminus u)(X) = \min_{Y \subseteq X} (u(Y) + f(X - Y))$ for $X \subseteq S$. If M is a matroid with rank function r then $r \setminus u$ is the rank function of $M|_{\text{supp}(u)}$.

Claim 2.4. *Let $u \in \mathbb{Z}_+^S$. If $w \in \mathbb{Z}_+^S$ is a double circuit of $f' := f \setminus u$ then w is either a double circuit of f with the same principal partition, or trivial, or non-compatible.*

Proof. Let $M = (T, r)$ be a prematroid of f and let $Z \subseteq T$ with $\chi^Z = u$. If $w \leq \chi^Z$ then w is a double circuit of f with the same principal partition by Lemma 2.1. Observe that $w_i \leq f'(\{i\}) + 2$ and $f'(\{i\}) \leq u_i$ for every $i \in S$. Thus if $w \not\leq \chi^Z$ then there exists an $i \in S$ such that $w_i - f'(\{i\}) \in \{1, 2\}$. If $w_i = f'(\{i\}) + 2$ then $r_{f'}(w_i \chi_i) = w_i - 2$, thus $W_0 = \text{supp}(w) = \{i\}$, implying that w is non-compatible. If $w_i = f'(\{i\}) + 1$ then $w_i \chi_i$ is a circuit of f' thus if $W_0 \neq \emptyset$ then w is non-compatible, and if $W_0 = \emptyset$ then w is trivial. ■

We are ready to prove Theorem 1.1.

Proof of Theorem 1.1. It is easy to see that $\nu(f) \leq \sum_{j=1}^t \lfloor \frac{f(U_j)}{2} \rfloor$ holds for every partition U_1, U_2, \dots, U_t of S . For the other direction we use a deep and important result of Lovász on 2-polymatroids, which can be translated to polymatroids as follows.

Theorem 2.5 (Lovász [9]). *If $f: 2^S \rightarrow \mathbb{Z}_+$ is a polymatroid function then at least one of the following cases holds.*

1. $f(S) = 2\nu(f) + 1$.
2. $S = S_1 \dot{\cup} S_2$ where $S_i \neq \emptyset$ and $\nu(f) = \nu(f|_{2^{S_1}}) + \nu(f|_{2^{S_2}})$.
3. There exists an $i \in S$ such that for all maximum matchings m we have $i \in \text{sp}_f(m)$.
4. There exists a certain substructure, called ν -double flower in f , which we do not define here, but which always contains a non-trivial compatible double circuit.

We argue by induction on the pair $(|S|, |K(f)|)$, where

$$K(f) = \{s \in S : s \in \text{sp}_f(m) \text{ for each maximum matching } m \text{ of } f\}.$$

If $S = \emptyset$ then the statement is trivial. If $K(f) = \emptyset$ then either 1. or 2. holds in Theorem 2.5. If 1. holds then the trivial partition will do, while if 2. holds then we can use our induction hypothesis applied to $f|_{2^{S_1}}$ and $f|_{2^{S_2}}$.

Next, let $K(f) \neq \emptyset$. We prove that if m is a maximum matching of $f + 2\chi_s$ then $m(s) \geq 2$. Indeed, assume that $m(s) = 0$. As m is a maximum matching, there exists a set $s \in U \subseteq S$ with $m(U) \geq (f + 2\chi_s)(U) - 1$. Thus $m(U - s) = m(U) \geq (f + 2\chi_s)(U) - 1 \geq f(U - s) + 1$, which is a contradiction. It is also clear that $m + 2\chi_s$ is a matching of $f + 2\chi_s$ for each matching m of f . Therefore, m is a maximum matching of f if and only if $m + 2\chi_s$ is a maximum matching of $f + 2\chi_s$.

Let $s \in K(f)$. Clearly, $\nu(f) \leq \nu(f + \chi_s) \leq \nu(f + 2\chi_s) = \nu(f) + 1$ and we claim that in fact, $\nu(f + \chi_s) = \nu(f)$ holds. Indeed, if $\nu(f + \chi_s) = \nu(f) + 1$ and m is a maximum matching of $f + \chi_s$ then m is also a maximum matching of $f + 2\chi_s$, thus $m(s) \geq 2$. Then $m - 2\chi_s$ is a maximum matching of f and, as $s \in \text{sp}_f(m - 2\chi_s)$, there exists a set $s \in U \subseteq S$ with $(m - 2\chi_s)(U) = f(U)$. This implies $m(U) = f(U) + 2$, contradicting to that m is a matching of $f + \chi_s$.

So if m is a maximum matching of f then m is a maximum matching of $f + \chi_s$, too, and clearly, $\text{sp}_f(m) \subseteq \text{sp}_{f+\chi_s}(m) - \{s\}$. Thus we have $K(f + \chi_s) \subseteq K(f) - s$. By [Claim 2.3](#), $f + \chi_s$ has no non-trivial compatible double circuits, so we can apply induction to $f + \chi_s$. This gives a partition U_1, U_2, \dots, U_t of S such that

$$\nu(f + \chi_s) = \sum_{j=1}^t \left\lfloor \frac{(f + \chi_s)(U_j)}{2} \right\rfloor.$$

But then,

$$\nu(f) = \nu(f + \chi_s) = \sum_{j=1}^t \left\lfloor \frac{(f + \chi_s)(U_j)}{2} \right\rfloor \geq \sum_{j=1}^t \left\lfloor \frac{f(U_j)}{2} \right\rfloor. \quad \blacksquare$$

3. A parity constrained orientation theorem

Frank, Jordán and Szigeti [3] proved that the existence of a k -rooted-connected orientation with prescribed parity of in-degrees can be characterized by a partition type condition. Recently, Király and Szabó [7] proved that the connectivity requirement in this parity constrained orientation problem can be given by a more general non-negative intersecting supermodular function. It is well-known that all these problems can be formalized as polymatroid parity problems. In this section we show that it is possible to formalize the problem of Király and Szabó in such a way that the arising polymatroid function has no non-trivial compatible double circuits. So [Theorem 1.1](#) can be applied to yield the result in [7].

$H = (V, \mathcal{E})$ is called a *hypergraph* if V is a finite set and $\emptyset \notin \mathcal{E}$ is a collection of multisets of V , the set of *hyperedges* of H . A *directed hyperedge* is obtained

from $h \in \mathcal{E}$ by *orienting* it, more precisely by designating a vertex $\text{head}(h) \in h$ as the *head vertex* of h . The other vertices of h are called *tail vertices*, which have therefore multiplicity function $h - \chi_{\text{head}(h)}$. By orienting each hyperedge of a hypergraph $H = (V, \mathcal{E})$ we get the *directed hypergraph* $D = (V, \mathcal{A})$. For a set $X \subseteq V$, let $\delta_D(X)$ denote the number of directed hyperedges *entering* X , that is the set of hyperedges with head in X and at least one vertex in $V - X$.

Let $p: 2^V \rightarrow \mathbb{Z}_+$ be a function with $p(\emptyset) = p(V) = 0$. An orientation D of a hypergraph $H = (V, \mathcal{E})$ *covers* p if $\delta_D(X) \geq p(X)$ for every $X \subseteq V$. In a *connectivity orientation problem* the question is the existence of an orientation covering p . When we are talking about *parity constrained orientations*, we are looking for connectivity orientations such that the out-degree at each vertex is of prescribed parity. For clarity, the in-degree of a vertex v is $|\{h \in \mathcal{E} : \text{head}(h) = v\}|$, while the out-degree of v is $(\sum_{h \in \mathcal{E}} \text{tail}(h))(v)$. Now define $b: 2^V \rightarrow \mathbb{Z}$ by

$$(4) \quad b(X) = \sum_{h \in \mathcal{E}} h(X) - |\mathcal{E}[X]| - p(X) \quad \text{for } X \subseteq V,$$

where $\mathcal{E}[X]$ denotes the set of hyperedges $h \in \mathcal{E}$ with $h \cap (V - X) = \emptyset$, and h equivalently stands for the hyperedge and its multiplicity function. It is clear that if $x: V \rightarrow \mathbb{Z}_+$ is the out-degree vector of an orientation covering p then $x \in B(b)$. The contrary is also easy to prove (see e.g. [4]).

Lemma 3.1. *Let $H = (V, \mathcal{E})$ be a hypergraph, $p: 2^V \rightarrow \mathbb{Z}_+$ be a function with $p(\emptyset) = p(V) = 0$, and $x: V \rightarrow \mathbb{Z}_+$. Then H has an orientation covering p such that the out-degree of each vertex $v \in V$ is $x(v)$ if and only if $x \in B(b)$.*

The function $b: 2^V \rightarrow \mathbb{Z}$ is said to be *intersecting submodular* if (1) holds whenever $X \cap Y \neq \emptyset$. Similarly, $p: 2^V \rightarrow \mathbb{Z}$ is *intersecting supermodular* if $-p$ is intersecting submodular. If $b: 2^V \rightarrow \mathbb{Z}_+$ is a non-negative, non-decreasing intersecting submodular function then we can define a polymatroid function $\widehat{b}: 2^V \rightarrow \mathbb{Z}_+$ by

$$\widehat{b}(X) = \min \left\{ \sum_{i=1}^t b(X_i) : X_1 \dot{\cup} X_2 \dot{\cup} \cdots \dot{\cup} X_t = X \right\} \quad \text{for } X \subseteq V,$$

which is called the *Dilworth truncation* of b .

Lemma 3.2. *If $p: 2^V \rightarrow \mathbb{Z}_+$ is intersecting supermodular with $p(V) = 0$ then p is non-increasing.*

Proof. If $\emptyset \neq X \subseteq Y \subseteq V$ then $p(Y) \leq p(Y) + p((V - Y) \cup X) \leq p(X) + p(V) = p(X)$. ■

The following theorem can be proved using basic properties of polymatroids (Frank [5]).

Theorem 3.3. *Let $H = (V, \mathcal{E})$ be a hypergraph and $p : 2^V \rightarrow \mathbb{Z}_+$ be an intersecting supermodular function with $p(\emptyset) = p(V) = 0$. Define b as in (4). Then H has an orientation covering p if and only if*

$$(5) \quad b(V) \leq \sum_{j=1}^t b(U_j)$$

holds for every partition U_1, U_2, \dots, U_t of V .

Proof. If there exists an orientation, then its out-degree vector satisfies

$$b(V) = x(V) = \sum_{j=1}^t x(U_j) \leq \sum_{j=1}^t b(U_j)$$

for every partition U_1, U_2, \dots, U_t of V . For the other direction, first, b is non-decreasing by Lemma 3.2. Second, if $\emptyset \subseteq U \subseteq V$ then $b(U) + b(V - U) \geq b(V)$ by (5), implying that $b(U) \geq b(V) - b(V - U) \geq 0$. Hence b is also non-negative, and we can define the Dilworth-truncation \hat{b} . Hence, (5) implies $\hat{b}(V) = b(V)$ and $B(\hat{b}) = B(b)$. As $B(\hat{b})$ is a non-empty integer polyhedron, taking any integer vector $x \in B(\hat{b})$, Lemma 3.1 gives the orientation. ■

Let $H = (V, \mathcal{E})$ be a hypergraph and $T \subseteq V$. Our goal is to find an orientation of H covering p , where the set of odd out-degree vertices is as close as possible to T .

Theorem 3.4 (Király and Szabó [7]). *Let $H = (V, \mathcal{E})$ be a hypergraph, $T \subseteq V$, $p : 2^V \rightarrow \mathbb{Z}_+$ be an intersecting supermodular function with $p(\emptyset) = p(V) = 0$, and assume that H has an orientation covering p . Define b as in (4). For an orientation D of H let $Y_D \subseteq V$ denote the set of odd out-degree vertices in D . Then*

$$(6) \quad \min \{|T \Delta Y_D| : D \text{ is an orientation of } H \text{ covering } p\} \\ = \max \left\{ b(V) - \sum_{j=1}^t b(U_j) + |\{j : b(U_j) \not\equiv |T \cap U_j| \pmod{2}\}| \right\},$$

where the maximum is taken over partitions U_1, U_2, \dots, U_t of V .

Our proof of Theorem 3.4 is based on the fact that the polymatroid of the out-degree vectors of orientations covering p have no non-trivial compatible double circuits:

Lemma 3.5. *Let $H = (V, \mathcal{E})$ be a hypergraph, $\alpha: V \rightarrow \mathbb{Z}_+$ and $p: 2^V \rightarrow \mathbb{Z}_+$ an intersecting supermodular function with $p(\emptyset) = 0$. Suppose moreover that $b: 2^V \rightarrow \mathbb{Z}$, defined by $b(X) = \alpha(X) - |\mathcal{E}[X]| - p(X)$ ($X \subseteq V$) is non-negative and non-decreasing. Then the polymatroid function $f := \widehat{b}$ has no non-trivial compatible double circuits.*

Proof. Assume that $w: V \rightarrow \mathbb{Z}_+$ is a non-trivial compatible double circuit of f with principal partition $W = W_1 \dot{\cup} W_2 \dot{\cup} \dots \dot{\cup} W_d$. Clearly,

$$b(W) \geq w(W) - 2.$$

Let $1 \leq i < j \leq d$ and $Z = W - W_i$. As $w|_Z$ is a circuit of f , Lemma 2.2 yields that $w(Z) - 1 = f(Z) = \min \sum \{b(X_i) : X_1, \dots, X_k \text{ partitions } Z\}$. However, if a non-trivial partition with $k \geq 2$ gave equality here, then we would have $f(Z) = \sum b(X_i) \geq \sum f(X_i) \geq \sum w(X_i) = w(Z) = f(Z) + 1$, using that $w|_{X_i} \in P(f)$. Thus

$$w(W - W_i) - 1 = b(W - W_i) \quad \text{and similarly,} \quad w(W - W_j) - 1 = b(W - W_j).$$

By applying intersecting submodularity to $W - W_i$ and $W - W_j$, and using that $w|_{W - W_i - W_j} \in P(f)$, we get

$$\begin{aligned} 0 &\geq b(W) - b(W - W_i) - b(W - W_j) + b(W - W_i - W_j) \\ &\geq (w(W) - 2) - (w(W - W_i) - 1) - (w(W - W_j) - 1) + w(W - W_i - W_j) = 0, \end{aligned}$$

so equality holds throughout. As a corollary, each hyperedge $e \in \mathcal{E}[W]$ is spanned by one of the W_i 's, and

$$\begin{aligned} (7) \quad \binom{d-1}{2} (b(W) + 2) &= \binom{d-1}{2} w(W) = \sum_{1 \leq i < j \leq d} w(W - W_i - W_j) \\ &= \sum_{1 \leq i < j \leq d} b(W - W_i - W_j). \end{aligned}$$

On the other hand,

$$\binom{d-1}{2} \alpha(W) = \sum_{1 \leq i < j \leq d} \alpha(W - W_i - W_j),$$

as α is modular, and

$$\binom{d-1}{2} p(W) \leq \sum_{1 \leq i < j \leq d} p(W - W_i - W_j),$$

as p is non-negative and non-increasing. Finally,

$$\binom{d-1}{2} |\mathcal{E}[W]| = \binom{d-1}{2} \sum_{i=1}^d |\mathcal{E}[W_i]| = \sum_{1 \leq i < j \leq d} |\mathcal{E}[W - W_i - W_j]|.$$

By the definition of b , the last 3 equalities together contradict (7). \blacksquare

Proof of Theorem 3.4. For every $v \in T$ add a loop $2\chi_v$ to \mathcal{E} , resulting in the hypergraph $H' = (V, \mathcal{E}')$. Define b' as in (4), w.r.t. H' . As there is a straightforward bijection between the orientations of H and H' , we have

$$\begin{aligned} \min\{|T \triangle Y_D| : D \text{ is an ori. of } H \text{ covering } p\} \\ = \min\{|Y_{D'}| : D' \text{ is an ori. of } H' \text{ covering } p\}. \end{aligned}$$

Furthermore,

$$\begin{aligned} b(V) - \sum_{j=1}^t b(U_j) + |\{j : b(U_j) \not\equiv |T \cap U_j| \pmod{2}\}| \\ = b'(V) - \sum_{j=1}^t b'(U_j) + |\{j : b'(U_j) \text{ is odd}\}|. \end{aligned}$$

Thus we can assume that $T = \emptyset$, and let us use again the original notations.

By Lemma 3.1, the integer vectors of $B(b)$ are exactly the out-degree vectors of the orientations of H covering p . Thus the \geq direction is easy to check. Now we prove the other direction. By Theorem 3.3, H has an orientation covering p , and we can define the polymatroid function $f = \widehat{b}$ as in the proof of Theorem 3.3. Now, we can use Lemma 3.5 with the choice $\alpha(v) = \sum_{h \in \mathcal{E}} h(v)$ for $v \in V$, stating that f has no non-trivial compatible double circuits. Then, by Theorem 1.1 we have

$$(8) \quad \nu(f) = \min \sum_{i=1}^s \left\lfloor \frac{f(V_i)}{2} \right\rfloor,$$

where the minimum is taken over all partitions V_1, V_2, \dots, V_s of V . Indeed, using the definition of the Dilworth-truncation and that $b(V) = f(V)$ by Theorem 3.3, we get

$$\begin{aligned} \min\{|Y_D| : D \text{ is an ori. of } H \text{ covering } p\} &= f(V) - 2\nu(f) \\ &= b(V) - \min \left\{ \sum_{i=1}^s f(V_i) - |\{i : f(V_i) \text{ is odd}\}| : V_1, \dots, V_s \text{ partitions } V \right\} \\ &\leq b(V) - \min \left\{ \sum_{j=1}^t b(U_j) - |\{j : b(U_j) \text{ is odd}\}| : U_1, \dots, U_t \text{ partitions } V \right\}. \blacksquare \end{aligned}$$

The next corollary of [Theorem 3.4](#) is formally very similar to the characterization in [Theorem 3.3](#).

Theorem 3.6. *Let $H = (V, \mathcal{E})$ be a hypergraph and $p: 2^V \rightarrow \mathbb{Z}_+$ an intersecting supermodular function with $p(\emptyset) = p(V) = 0$. Then a set $T \subseteq V$ arises as the set of odd out-degree vertices in an orientation of H covering p if and only if*

$$(9) \quad b(V) \leq \sum_{j=1}^t b(U_j) - |\{j : b(U_j) \not\equiv |T \cap U_j| \pmod{2}\}|$$

holds for every partition U_1, U_2, \dots, U_t of V .

Let us give an example showing that polymatroids without non-trivial compatible double circuits are not closed under taking contractions. Let $V = \{v_1, v_2, v_3, v_4\}$, $\mathcal{E} = \{v_1 v_i, v_i v_i : i \in \{2, 3, 4\}\}$, $p(\{v_1\}) = 1$ and $p(U) = 0$ for all other sets $U \subseteq V$. Then b is non-decreasing and non-negative, thus by [Lemma 3.5](#), \hat{b} has no non-trivial compatible double circuits. However, the polymatroid obtained from \hat{b} by contracting an element from the preimage of v_1 in the prematroid of \hat{b} has the non-trivial compatible double circuit $(1, 2, 2, 2)$, with the trivial principal partition.

4. A planar rigidity problem

If $G = (V, E)$ is a graph and $p: V \rightarrow \mathbb{R}^2$ is an embedding into the Euclidean plane then (G, p) is said to be a *framework*. We think of the edges of G as rigid bars with flexible joints at the vertices. An *infinitesimal motion* means an assignment of a velocity $x(v) \in \mathbb{R}^2$ to each vertex $v \in V$ such that the bar lengths are preserved, that is $(p(u) - p(v)) \perp (x(u) - x(v))$. The framework (G, p) is called *rigid* if all infinitesimal motions of (G, p) correspond to isometries of \mathbb{R}^2 . The question of finding a vertex set $Z \subseteq V$ of minimum size such that $(G + K_Z, p)$ is rigid was solved by Lovász in his seminal paper [9] about matroid parity. We say that $G = (V, E)$ is *generic rigid* if all frameworks (G, p) with *algebraically independent* coordinates p are rigid. The problem of finding a vertex set $Z \subseteq V$ of minimum size such that $G + K_Z$ is generic rigid is left open by [9], and it was solved recently by Fekete [2]. For more on the 2-dimensional rigidity see Laman [8] and Lovász and Yemini [12].

The setup of [2] puts the problem into a bit more general setting. Let $G = (V, E)$ be a graph, and for $l \in \{2, 3\}$ let $M_{2,l}$ be the matroid on ground set E such that $F \subseteq E$ is independent in $M_{2,l}$ if and only if $|F[X]| \leq 2|X| - l$ for all $X \subseteq V$, $|X| \geq 2$. It can be proved that M is really a matroid. For clarity, $M_{2,2}$ is two times the cycle matroid of G , and so G has two edge-

disjoint spanning trees if and only if $r_{2,2}(E) = 2|V| - 2$. As $M_{2,3}$ is the rigidity matroid of G , the graph G is generic rigid if and only if $r_{2,3}(E) = 2|V| - 3$. For $Z \subseteq V$ let $K_Z = (Z, E_Z)$ be the graph with vertex set Z having $4 - l$ parallel edges between any two vertices of Z . Our goal is to find a set $Z \subseteq V$ of minimum size such that $E + E_Z$ has rank $2|V| - l$ in $M_{2,l}$.

We assume that E is independent in $M_{2,l}$, since if E is replaced by one of its bases then the solution set does not change. Fekete [2] proved the following lemma. For $X \subseteq V$ let $e(X)$ denote the number of edges having at least one end vertex in X .

Lemma 4.1 ([2]). *Let $l \in \{2, 3\}$. Assume that E is independent in $M_{2,l}$ and that $r_{2,l}(E) < 2|V| - l$. Let $Z \subseteq V$. Then $r(E + E_Z) = 2|V| - l$ if and only if $e(Y) \geq 2|Y|$ for every $Y \subseteq V - Z$.*

Therefore, the goal is to find a set $Z \subseteq V$ of minimum size such that $e(Y) \geq 2|Y|$ for every $Y \subseteq V - Z$. Let $f : 2^V \rightarrow \mathbb{Z}_+$ be the polymatroid function with $f(X) = \min_{Y \subseteq X} 2|Y| + e(X - Y)$ for $X \subseteq V$. That is, f is obtained from the polymatroid function $X \mapsto e(X)$ by putting upper bound 2 on the elements (which is called above as deletion operation). Hence for $l = 2$ the value $|V| - \nu(f)$ is just the minimum size of a set Z whose contraction results in a graph with two edge-disjoint spanning trees, and for $l = 3$ it is the minimum size of a set Z such that $G + K_Z$ is generic rigid. In [9] the computation of $\nu(f)$ is reduced to the matching problem of graphs, yielding a partition type characterization. This characterization follows from the previous results of this paper, too. First, by Lemma 3.5 with the choice $p = 0$, the polymatroid function $X \mapsto e(X)$ has no non-trivial compatible double circuits. As f is obtained from $X \mapsto e(X)$ by deletion, Claim 2.4 yields that nor f has. Thus,

$$\nu(f) = \min \sum_{j=1}^t \left\lfloor \frac{f(U_j)}{2} \right\rfloor,$$

where the minimum is taken over all partitions U_1, U_2, \dots, U_t of V . By the definition of f , we get the following.

Theorem 4.2 (Fekete [2]). *Let $l \in \{2, 3\}$. Assume that E is independent in $M_{2,l}$ and that $r_{2,l}(E) < 2|V| - l$. Then the minimum size of a set $Z \subseteq V$ such that $r(E + E_Z) = 2|V| - l$ is $|V| - \nu(f)$, where*

$$\nu(f) = \min \left| V - \bigcup_{j=1}^t U_j \right| + \sum_{j=1}^t \left\lfloor \frac{e(U_j)}{2} \right\rfloor,$$

where the minimum is taken over all subpartitions U_1, U_2, \dots, U_t of V .

References

- [1] A. DRESS and L. LOVÁSZ: On some combinatorial properties of algebraic matroids, *Combinatorica* **7**(1) (1987), 39–48.
- [2] ZS. FEKETE: Source location with rigidity and tree packing requirements, Technical Report TR-2005-04, Egerváry Research Group (EGRES), Budapest (2005).
<http://www.cs.elte.hu/egres>
- [3] A. FRANK, T. JORDÁN and Z. SZIGETI: An orientation theorem with parity conditions, *Discrete Appl. Math.* **115**(1–3) (2001), 37–47. 1st Japanese–Hungarian Symposium for Discrete Mathematics and its Applications (Kyoto, 1999).
- [4] A. FRANK, T. KIRÁLY and Z. KIRÁLY: On the orientation of graphs and hypergraphs, *Discrete Appl. Math.* **131**(2) (Submodularity) (2003), 385–400.
- [5] ANDRÁS FRANK: On the orientation of graphs, *J. Combin. Theory Ser. B* **28**(3) (1980), 251–261.
- [6] P. M. JENSEN and B. KORTE: Complexity of matroid property algorithms, *SIAM J. Comput.* **11**(1) (1982), 184–190.
- [7] T. KIRÁLY and J. SZABÓ: A note on parity constrained orientations, Technical Report TR-2003-11, Egerváry Research Group, Budapest, (2003).
<http://www.cs.elte.hu/egres>
- [8] G. LAMAN: On graphs and rigidity of plane skeletal structures, *J. Engrg. Math.* **4** (1970), 331–340.
- [9] L. LOVÁSZ: Matroid matching and some applications, *J. Combin. Theory Ser. B* **28**(2) (1980), 208–236.
- [10] L. LOVÁSZ: Selecting independent lines from a family of lines in a space, *Acta Sci. Math. (Szeged)* **42**(1–2) (1980), 121–131.
- [11] L. LOVÁSZ: The matroid matching problem, in *Algebraic methods in graph theory, Vol. I, II (Szeged, 1978)*, volume **25** of *Colloq. Math. Soc. János Bolyai*, pages 495–517, North-Holland, Amsterdam, 1981.
- [12] L. LOVÁSZ and Y. YEMINI: On generic rigidity in the plane, *SIAM J. Algebraic Discrete Methods* **3**(1) (1982), 91–98.
- [13] GYULA PAP: *A constructive approach to matching and its generalizations*, PhD thesis, Eötvös University, Budapest, 2006.
- [14] ALEXANDER SCHRIJVER: *Combinatorial optimization. Polyhedra and efficiency*, volume **24** of *Algorithms and Combinatorics*. Springer-Verlag, Berlin, 2003.

Márton Makai, Jácint Szabó

MTA-ELTE Egerváry Research Group (EGRES)

Institute of Mathematics

Eötvös University

H-1117 Budapest, Pázmány P. s. 1/C

Hungary

{marci,jacint}@cs.elte.hu