

## WATER-EXCHANGE KINETICS IN LABILE AQUO AND SUBSTITUTED AQUO TRANSITION METAL IONS BY MEANS OF $^{17}\text{O}$ NMR STUDIES

J.P. HUNT

*Chemistry Department, Washington State University, Pullman, Washington 99163 (U.S.A.)*

(Received November 2nd, 1970)

### CONTENTS

A.	Introduction	1
B.	Consideration of experimental measurements, theory and interpretation	2
(i)	Theory of the NMR method for water-exchange kinetics	2
(ii)	Experimental aspects of the NMR measurements	3
(iii)	Treatment of data	3
(iv)	Chemical aspects of the measurements	3
C.	Results	4
(i)	General comments	4
(ii)	More specific consideration of the cobalt and nickel data	6
(iii)	Some special observations	8
D.	Conclusion	9
	References	10

### ABBREVIATIONS

Tiron	4,5 dihydroxybenzene-1,3-disulfonate	Terpy	2,2',2''-terpyridyl
IDA	iminodiacetate	bipy	2,2'-bipyridyl
SS	5-sulfosalicylate	dien	diethylenetriamine
phen	<i>ortho</i> -phenanthroline	tren	triethylenetriamine
mal	malonate	tren	$\beta, \beta', \beta''$ -triethylenetriaminoamine
en	ethylenediamine	EDTA or Y	ethylenediaminetetraacetate

### A INTRODUCTION

Several reviews have appeared which discuss the uses of NMR methods to study water exchange kinetics in aqueous metal ion solutions<sup>1</sup> and the importance of these results in attempting to understand net substitution processes in labile systems<sup>2</sup>. A moderate amount of NMR data, largely from our laboratories, is now available concerning the kinetics of water exchange in partially substituted aquo ions. It is the purpose of this review

to bring together the current results, consider correlating them, and to mention some related matters suggested by the NMR work. Inferences about water-exchange kinetics have, of course, been drawn from the large amount of work on net substitution reactions<sup>1-3</sup>, but these are not reconsidered in detail in this discussion.

## B. CONSIDERATION OF EXPERIMENTAL MEASUREMENTS, THEORY, AND INTERPRETATION

### (i) Theory of the NMR method for water-exchange kinetics

We are concerned here with relating measurements of line broadenings and shifts for  $^{17}\text{O}$  in solvent or bulk water, produced by paramagnetic (usually) water-containing metal ion complexes, to the kinetics of water exchange between solvent and bound water molecules in the first coordination sphere of a metal ion. Discussions of the general subject are numerous<sup>1</sup> and only a brief account will be given here. The procedures and equations used for the studies are based on the work of Swift and Connick<sup>4</sup>. Some modifications we have used are given by Dodgen et al.<sup>5</sup>. The basic equations for two sites, as we use them, are as follows where  $T_{2p'}$  is a molar line-broadening function and  $Q$  is the molar shift function

$$T_{2p'} \equiv \frac{2[M]}{\gamma\Delta'} = \frac{[M] \tau_{\text{H}_2\text{O}} [(1/T_{2M} + 1/\tau_M)^2 + \Delta\omega_M^2]}{[1/T_{2M}^2 + 1/T_{2M}\tau_M + \Delta\omega_M^2]}$$

$$Q \equiv \frac{TS[\text{H}_2\text{O}]}{[M]} = \frac{T[\text{H}_2\text{O}] \Delta\omega_M}{\omega_0 [M] \tau_{\text{H}_2\text{O}} \tau_M [(1/\tau_M + 1/T_{2M})^2 + \Delta\omega_M^2]}$$

The symbols used have the following meanings.  $[M]$  is the concentration of the aquo metal ion complex,  $[\text{H}_2\text{O}]$  is the concentration of solvent water,  $\tau_{\text{H}_2\text{O}}$  is the mean lifetime for water exchange in bulk water ( $\tau_{\text{H}_2\text{O}} = [\text{H}_2\text{O}]/\text{rate of exchange}$ ),  $T_{2M}$  is the spin-spin relaxation time for  $^{17}\text{O}$  bound to the metal ion,  $\tau_M$  is the mean lifetime for exchange of a bound water molecule ( $n[M]/\text{rate of exchange}$ , where  $n$  is the number of water molecules bound/metal ion),  $\Delta\omega_M = (\omega_M - \omega_{\text{obs}}) \simeq (\omega_M - \omega_0)$  for dilute solutions, where  $\omega_M$  is the NMR absorption frequency (rad/sec) for a bound water,  $\omega_{\text{obs}}$  is the frequency observed for bulk water in the given solution and  $\omega_0$  is the frequency observed for bulk water in the absence of the paramagnetic metal ion,  $\gamma$  is the magnetogyric ratio for  $^{17}\text{O}$  ( $3628 \text{ gauss}^{-1} \cdot \text{sec}^{-1}$ ),  $\Delta'$  is the observed line broadening ( $W_{\text{obs}} - W_0$ ) where  $W_{\text{obs}}$  is the full line width at half-maximum in the NMR absorption curve for the paramagnetic solution and  $W_0$  the same quantity for the reference or blank,  $T$  is the absolute temperature and occurs because  $\Delta\omega$  is assumed to follow Curie's Law and  $S = \Delta\omega_{\text{obs}}/\omega_0$  ( $\Delta\omega_{\text{obs}} = \omega_{\text{obs}} - \omega_0$ ). Thus,  $S$  and  $\Delta'$  can be thought of as the experimental observables in general. In some cases,  $T_{2M}$  and  $\Delta\omega_M$  may be directly measured and also used to help extract  $\tau_M$  or  $\tau_{\text{H}_2\text{O}}$  which contain the kinetic information.

It should be noted that the treatment based on Swift and Connick's equations is valid only for two-site or parallel two-site situations. Further, only solutions dilute in the paramagnetic ion ( $< 0.5 M$  perhaps) are precisely treated. The equations for the dilute solution

three-site situation have been presented<sup>6</sup>. In principle, the kinetic information can be obtained either from line broadening or shift measurements or a combination of both, which latter approach seems preferable. Limiting forms of the equations are useful, but it is desirable to fit the data finally using the complete equations

*(ii) Experimental aspects of the NMR measurements*

Various kinds of equipment and techniques of measurement are used. The use of derivative curves or side-band techniques is common with commercial equipment. Our experiments have been mainly done using a single-coil method<sup>5,7</sup>. In all cases, care is required to ensure the proper use of the spectrometer. Problems of field and frequency stability, phasing, modulation parameters, Rf power level (avoidance of saturation), good temperature control of sample etc., and others must be considered. Signal averaging can help to increase signal-noise ratios.

A reference solution is needed in making broadening and shift measurements at each temperature. The nature of this reference is less critical if one can work with relatively large line broadenings and shifts. If this is not possible, the reference solution should be chosen to resemble the working solution as closely as possible.

Possible outer-sphere line broadening and line shifts should be looked for and taken into account if found. These seem to be quite small in general when protons are not being studied.

*(iii) Treatment of data*

The parameters of kinetic interest usually have to be extracted from a rather complex function. Although the usual non-linear least squares computer treatment have been used, they are not ideally suited to the problem at hand. Variations in the individual parameters over a range of values should be tried to get a proper assessment of the influence of each on the results. The relative importance and precision of measurements of each parameter depends on the temperature. To date, we have preferred to use the computer to calculate curves which can be compared visually with the experimental curves, adjusting parameters systematically to give the best fit to the more precise data. Because of the inherent difficulties in the measurements, rate constants will rarely be reliable to better than  $\pm 5\%$  and  $\Delta H^\ddagger$  values to  $\pm 0.5$  kcal/mole even in favorable cases.

*(iv) Chemical aspects of the measurements*

In systems where several species exist in equilibrium, a knowledge of the formation constants and enthalpy of formation values is needed under prevailing conditions. Uncertainties in these quantities are reflected in the absolute errors present in the end results reported. Species present may change over the range of temperatures used and if so can cause misleading interpretations. Often the isomeric structures present are not known with any certainty.

## C. RESULTS

## (1) General comments

The available data are collected in Table 1. The rate constant  $k_1$  is defined by the relation  $R = nk_1[M]$ , where  $R$  is the observed rate of water exchange,  $n$  is the number of kinetically equivalent, exchanging water molecules and  $M$  refers to the particular metal complex species involved. The quantity  $A$ , from the electron-nuclear spin interaction term  $AI \cdot S$ , is reported as  $A/h$  ( $h$  = Planck's constant) and is calculated from

$$A/h = T(\Delta\omega_M/\omega_0)(3k/2\pi)(\gamma_N/S(S+1)g_{\text{eff}}\beta)$$

Here  $T$  is the absolute temperature,  $k$  the Boltzmann constant,  $\gamma_N$  the nuclear magnetogyric ratio,  $S$  the spin quantum number and  $g_{\text{eff}}\beta$  is consistent with  $\mu_{\text{eff}}$  for the metal complex with  $\mu_{\text{eff}} = g_{\text{eff}}[S(S+1)]^{1/2}$ . Here  $\mu_{\text{eff}}$  is the magnetic moment in Bohr magnetons,  $\beta$  is the Bohr magneton in cgs units and  $g_{\text{eff}}$  is the observed "g-factor". The other symbols have been defined previously. Errors indicated are estimates based on the precision of measurement and uncertainties in quantities used in the data treatment. They are thought to be conservatively assigned.

TABLE 1

Water exchange data

System <sup>a</sup>	$k_1$ (25°C) (sec <sup>-1</sup> )	$\Delta H^\ddagger$ (kcal/mol)	$A/h$ (Hz)	Ref
VO <sup>2+</sup> pH ca 1, $\mu < 0.3$ , ClO <sub>4</sub> <sup>-</sup> anion	Equatorial water $5 \times 10^2$	13.7	$3.8 \times 10^6$	8
VO (IDA), pH ca. 4, low $\mu$	$1.2 \times 10^5$	11.7	$2.8 \times 10^6$	8
VO (SS) <sup>-</sup> pH ca. 4, low $\mu$	$1.5 \times 10^5$	10.8	$4.5 \times 10^6$	8
VO (Tiron) <sup>2-</sup> , pH ca. 4, low $\mu$	$5.3 \times 10^5$	11.8	$4.9 \times 10^6$	8
Mn (H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> pH ca. 4, low $\mu$	$2.3 (\pm 0.2) \times 10^7$	$9 \pm 1$	$7 (\pm 0.1) \times 10^6$	21
Mn (H <sub>2</sub> O) <sub>4</sub> (phen) <sup>2+</sup> pH ca. 4, low $\mu$	$5.3 (\pm 0.7) \times 10^7$	$9 \pm 2$	$1 (\pm 0.5) \times 10^7$	21
Mn (H <sub>2</sub> O) <sub>2</sub> (phen) <sub>2</sub> <sup>2+</sup>	$1.2 (\pm 0.2) \times 10^8$	$9 \pm 2$	$2 (\pm 0.6) \times 10^7$	21
Co (H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> , 0.1 M Co (ClO <sub>4</sub> ) <sub>2</sub>	$2.6 (\pm 0.2) \times 10^6$	$11.9 \pm 0.7$	$1.35 \times 10^7$	22
Co (H <sub>2</sub> O) <sub>5</sub> Cl <sup>+</sup> dil. HCl	$1.7 (\pm 0.3) \times 10^7$	$13.8 \pm 0.7$	$1.35 \times 10^7$	22
Co (H <sub>2</sub> O) <sub>6</sub> <sup>2+</sup> , pH ca. 4, 2 M NH <sub>4</sub> NO <sub>3</sub>	$2.24 (\pm 0.05) \times 10^6$	$10.3 \pm 0.2$	$1.20 (\pm 0.02) \times 10^7$	14
Co (H <sub>2</sub> O) <sub>5</sub> NH <sub>3</sub> <sup>2+</sup> 2 M NH <sub>4</sub> NO <sub>3</sub>	$1.6 (\pm 0.2) \times 10^7$	$12.6 \pm 0.6$	$1.2 (\pm 0.1) \times 10^7$	14

TABLE 1 (continued)

$\text{Co}(\text{H}_2\text{O})_4(\text{NH}_3)_2^{2+}$ $2\text{M NH}_4\text{NO}_3$	$6.5 (\pm 1) \times 10^7$	$9.4 \pm 1.5$	$1.2 (\pm 0.1) \times 10^7$	14
$\text{Co}(\text{H}_2\text{O})_4 \text{ mal}^\circ$ , $\mu$ ca. 0.3	$2.2 (\pm 0.4) \times 10^7$	$12.9 \pm 0.8$	$1.2 (\pm 0.1) \times 10^7$	14
$\text{Co}(\text{H}_2\text{O})_2(\text{mal})_2^{2-}$ $\mu \approx 0.3$	$> 10^8$			14
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , 0.1 M $\text{HClO}_4$ , $\mu$ ca. 0.4	$2.7 \times 10^4$	11.6		23
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , 0.1 M $\text{HClO}_4$ $\mu$ ca. 0.2 - 0.7	$3.0 (\pm 0.3) \times 10^4$	$10.8 \pm 0.5$	$1.9 \times 10^7$	24
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , 6.5 M $\text{LiClO}_4$ pH ca. 7	$4.1 (\pm 0.5) \times 10^4$	$9.6 \pm 1$	$2.8 (\pm 0.5) \times 10^7$	19
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , $\mu = 0.5$ pH ca. 6	$4.4 (\pm 0.2) \times 10^4$	$10.3 \pm 0.5$	$2.2 (\pm 0.4) \times 10^7$	25
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , $\mu = 0.4 - 1.0$ pH ca. 3	$3.4 (\pm 0.1) \times 10^4$	$12.1 \pm 0.3$		26
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , 2 M $\text{NH}_4\text{NO}_3$ pH ca. 5	$3.6 (\pm 0.2) \times 10^4$	$12.3 \pm 0.5$	$2.3 (\pm 0.2) \times 10^7$	11
$\text{Ni}(\text{H}_2\text{O})_6^{2+}$ , $10^{-3}$ M $\text{HCl}$ $\mu = 0.3 \rightarrow 1.4$	$3.2 (\pm 0.2) \times 10^4$	$12.1 \pm 0.5$	$2.1 (\pm 0.1) \times 10^7$	27
$\text{Ni}(\text{H}_2\text{O})_4(\text{NH}_3)_2^{2+}$ 2 M $\text{NH}_4\text{NO}_3$	$2.5 (\pm 0.2) \times 10^5$	$10.6 \pm 0.5$	$2.5 (\pm 0.2) \times 10^7$	11
$\text{Ni}(\text{H}_2\text{O})_4(\text{NH}_3)_2^{2+}$ 2 M $\text{NH}_4\text{NO}_3$	$6.1 (\pm 0.6) \times 10^5$	$7.8 \pm 0.5$	$1.9 (\pm 0.2) \times 10^7$	11
$\text{Ni}(\text{H}_2\text{O})_3(\text{NH}_3)_3^{2+}$ 2 M $\text{NH}_4\text{NO}_3$	$2.5 (\pm 0.3) \times 10^6$	$10.2 \pm 0.7$	$2.6 (\pm 0.2) \times 10^7$	11
$\text{Ni}(\text{H}_2\text{O})_4\text{en}^{2+}$ , $\mu = 0.5$	$4.4 (\pm 0.2) \times 10^5$	$10.0 \pm 0.5$	$2.2 (\pm 0.2) \times 10^7$	25
$\text{Ni}(\text{H}_2\text{O})_2\text{en}_2^{2+}$ , $\mu = 0.5$	$5.4 (\pm 0.3) \times 10^6$	$9.1 \pm 0.5$	$2.3 (\pm 0.2) \times 10^7$	25
$\text{Ni}(\text{H}_2\text{O})_5\text{Cl}^+$ , 7 M $\text{LiCl}$	$1.4 (\pm 0.3) \times 10^5$	$8 \pm 1$	$3.5 (\pm 1) \times 10^7$	19
$\text{Ni}(\text{H}_2\text{O})_5(\text{NCS})_4^{1+}$ , 8 M $\text{KNCS}$	$1.1 (\pm 0.1) \times 10^6$	$6 \pm 0.6$	$2.0 (\pm 0.2) \times 10^7$	20
$\text{Ni}(\text{H}_2\text{O})_3(\text{terpy})^{2+}$ pH ca. 3, low $\mu$	$5.2 (\pm 0.4) \times 10^4$	$10.7 \pm 0.4$		26
$\text{Ni}(\text{H}_2\text{O})_4(\text{bipy})^{2+}$ $10^{-3}$ M $\text{HCl}$ , low $\mu$	$4.9 (\pm 0.5) \times 10^4$	$12.6 \pm 0.5$	$1.6 (\pm 0.2) \times 10^7$	27
$\text{Ni}(\text{H}_2\text{O})_2(\text{pipy})_2^{2+}$ $10^{-3}$ M $\text{HCl}$ , low $\mu$	$6.6 (\pm 0.8) \times 10^4$	$13.7 \pm 1$	$3.8 (\pm 0.5) \times 10^7$	27
$\text{Ni}(\text{H}_2\text{O})_3(\text{dien})^{2+}$ pH ca. 8, low $\mu$	$1.2 (\pm 0.1) \times 10^6$	$5.5 \pm 0.5$	$2.3 \times 10^7$	28
$\text{Ni}(\text{H}_2\text{O})_2(\text{tren})^{2+}$ pH ca. 8, low $\mu$	$2.9 (\pm 0.3) \times 10^6$	$7 \pm 0.5$	$2.1 \times 10^7$	28
$\text{Ni}(\text{H}_2\text{O})_2(\text{tren})^{2+}$ pH ca. 7, low $\mu$	$6.0 (\pm 0.6) \times 10^5$ $1.0 (\pm 0.1) \times 10^7$	$8 \pm 1$		28
$\text{Ni}(\text{H}_2\text{O})(\text{EDTA})^{2-}$	$7.0 (\pm 0.5) \times 10^{-2}$	$8.0 \pm 0.5$	$2.6 (\pm 0.2) \times 10^7$	29
$\text{Ni}(\text{H}_2\text{O})(\text{HEDTA})^-$	$2 (\pm 0.1) \times 10^5$	$9.8 \pm 0.3$	$2.3 (\pm 0.1) \times 10^7$	

The VO (IV) aquo species is unique in having one relatively non-labile oxygen, one rapidly exchanging water and four (equatorial) moderately labile water molecules. Labilizing effects of the ligands used are quite large. These are discussed in detail by Wuthrich and Connick<sup>8</sup> in terms of structures and electron donor effects and their discussion will not be repeated here. Similar arguments will be used for the Ni and Co systems discussed later in the review.

Very little data are available on the Mn(II) complexes. Exchange rates are near the upper limit of present day  $^{17}\text{O}$  NMR techniques and labilizing effects will be hard to measure. The phen ligand produces only a small effect as will be discussed later. It might be noted that the enthalpies of activation are fairly large though generally smaller than for Co and Ni. The aquo ion data are very much like those for  $\text{Mn}(\text{NH}_3)_6^{2+}$  in liquid ammonia<sup>9</sup>, an observation which is even more striking for Co and Ni examples<sup>5b</sup>.

Most of the available results refer to Ni(II) and Co(II) species. These will be discussed in the next section in a variety of ways. Comparison of these metals is of considerable theoretical interest. No detailed and systematic studies on effects of ionic strength ( $\mu$ ), added salts, solvent mixtures, and pH have been made. The data listed do suggest that at least small effects can be expected upon varying  $\mu$  and pH. Probably the Ni(II) systems warrant some detailed physico-chemical studies.

In general the bound water molecules behave as if they are kinetically equivalent with the nickel-tren and nitrilo triacetate<sup>10</sup> (NTA) systems being exceptions. Detection conditions for non-equivalence will vary considerably depending on the detailed NMR behavior found as well as on the equilibrium properties relating species and the kinetic parameters for the particular water molecules. Some of the results reported may be averages of similar values. One might generally expect to see clearly differences in  $k_1$  values of a factor of 3 and in  $\Delta H^\ddagger$  of 2–3 kcal/mole and smaller ones in favorable circumstances. In many cases no clear-cut evidence exists as to the isomer(s) present.

#### *(ii) More specific consideration of the cobalt and nickel data*

Considering the  $k_1$  values at 25°C, a simple correlation was found<sup>11</sup> when  $\log k_1$  (25°C) was plotted vs.  $\text{H}_2\text{O bound}/\text{M}^{2+}$ . In general,  $\log k_1$  (25°C) increases roughly linearly as  $\text{H}_2\text{O}/\text{M}^{2+}$  decreases for Ni(II) and Co(II) examples with the ligands  $\text{Cl}^-$ ,  $\text{NH}_3$ , malonate, en, dien, and tren as shown in Fig. 1. Some extrapolations in temperature are made and because of the variability in  $\Delta H^\ddagger$  the correlation typically works best at one temperature only. The ligands  $\text{NCS}^-$ , bipy, terpy, *o*-phen, and EDTA characteristically (considering the few examples) do not produce as large an increase in  $k_1$  (25°C) as the above correlation predicts (terpy, for example, is about 1/100 as effective as three  $\text{NH}_3$  groups). The small increases in  $\log k_1$  (25°C), however, are still roughly linear in  $\text{H}_2\text{O}/\text{Ni}$ . The above observations can be considered in conventional terms by regarding the complexes to be largely ionic in nature with electron-donating or withdrawing effects of the ligands altering the effective metal ion charge. These effects would be expected to be roughly additive. Good electron donors should enhance the rate of water exchange. Funahashi and Tanaka<sup>12</sup> have correlated some rates of substitution with the  $E_n$  (electron donating) parameter of Edwards<sup>13</sup>. In general,  $E_n$  values are not available for all the




Fig. 1. Semilogarithmic plot of  $\log k_1$  (25°C) vs.  $\text{H}_2\text{O}/\text{M}$  for cobalt and nickel systems. Right hand scale and  $\bullet$  refer to cobalt data while left hand scale and  $\circ$  refer to nickel. Y is EDTA

ligands of interest. If one assumes that  $\text{NH}_3$  and the other amines have the same  $E_n$  value, the correlation is only approximate. The more complex amines deviate by producing smaller effects, in general, than  $\text{NH}_3$ . Chloride is not consistent between cobalt and nickel, it should be about two-thirds as effective as  $\text{NH}_3$ . Using twice the  $E_n$  value of acetate for malonate predicts about the observed results.

The ligands bipy, terpy, and phen (in the  $\text{Mn(II)}$  case) produce little effect. These observations can be rationalized by postulating significant back  $\pi$ -bonding to the ligand, compensating for sigma donation. Thiocyanate may then be an intermediate case with somewhat less back-bonding.

The EDTA systems are interesting and will be described in a bit more detail under (iii). Based on the malonate result for cobalt, one might expect more rate enhancement than is observed. Protonation of the free acetate arm considerably reduces the enhancement. Comparison of these highly chelated systems with the simpler ones may not, of course, be particularly valid.

One might expect some spectral correlations if electron donating – withdrawing effects are involved. The spectra, however, are complex and it is not easy to obtain proper parameters. Simple correlations with  $10Dq$ , in general, do not work. Hoggard<sup>14</sup> has suggested the use of Jørgensen's "optical electronegativities". Although there are

many difficulties in application of these ideas, some success seems to be possible and perhaps warrants further investigation.

That single temperature data can be misleading is a well-known but often largely ignored fact. While the  $\Delta H^\ddagger$  values are not as precise as may be desired, they are not grossly in error and qualitative trends can be discussed. In the  $\text{VO}^{2+}$  system the ligand labilization of water is accompanied by some reduction in  $\Delta H^\ddagger$  although the greatest labilizing effect at 25°C is for the system with the least reduction in  $\Delta H^\ddagger$ . In the  $\text{Mn(II)}$  examples the  $\Delta H^\ddagger$  values are not precise but are similar as are the rates. In the cobalt(II) systems the  $\text{Cl}^-$ ,  $\text{NH}_3$ , and malonate ligands (mono-complexes) *increase* or do not change  $\Delta H^\ddagger$ , while bound water is being labilized. For the diammine  $\text{Co(II)}$  a significant reduction in  $\Delta H^\ddagger$  occurs. For  $\text{Ni(II)}$  the  $\Delta H^\ddagger$  values to some extent follow the trend in  $k_1$  though not in any regular way. Perhaps the main value of these parameters at this time is to emphasize how little is really understood in detail about the kinetic properties of similar species and the importance of entropy and solvation effects.

A comparison of the cobalt and nickel  $\Delta H^\ddagger$  values is of special interest. For the hexaquo ions cobalt has a lower  $\Delta H^\ddagger$  by ca. two kcal/mole (in 2M  $\text{NH}_4\text{NO}_3$ ). In the case of the hexammine species in liquid ammonia<sup>5b</sup> both have  $\Delta H^\ddagger = 11 \pm 0.5$  kcal/mole. In the other systems, cobalt has the *higher* value. Various theoretical considerations have suggested that the faster  $\text{Co(II)}$  rates (about 100X at 25°C) are due to a *lower*  $\Delta H^\ddagger$  for a common mechanism. Solvation effects are usually assumed to be identical for cobalt and nickel. If these ideas are correct (and they may not be), one may wish to consider that the mechanisms are indeed not the same for these metals, as was suggested earlier<sup>5b</sup>. The high  $\Delta H^\ddagger$  and positive  $\Delta S^\ddagger$  for cobalt may reflect greater dissociative character (four-coordinate cobalt?) for cobalt.

### (iii) Some special observations

The implications of ligand effects on solvent exchange kinetics for substitution processes have been recently discussed by Hewkin and Prince<sup>2</sup>. Although most of the data cited here were not available to them their general conclusions are not particularly altered. The observation of more than one water-exchange rate in the  $\text{Ni(II)-tren}$  and  $\text{NTA}$  systems does, of course, require some new thought concerning substitution in the relevant complexes.

Margerum et al.<sup>3</sup> have revised the substitution data using ammonia to replace bound water in nickel. Treating the observed substitution rate constant as a product of an outer-sphere constant and the rate constant for water exchange they estimate " $k_1$ " values which can be compared with the  $^{17}\text{O}$  NMR results. These are as follows ( $^{17}\text{O}$  NMR result given first): for  $\text{Ni(H}_2\text{O)}_4\text{en}^{2+}$   $4.4 \times 10^5$ ,  $1.8 \times 10^5$ ; for  $\text{Ni(H}_2\text{O)}_2\text{en}_2^{2+}$   $5.4 \times 10^6$ ,  $2.1 \times 10^8$ ; for  $\text{Ni(H}_2\text{O)}_3\text{dien}^{2+}$   $1.2 \times 10^6$ ,  $8.6 \times 10^5$ ; for  $\text{Ni(H}_2\text{O)}_2\text{tren}^{2+}$   $3.8 \times 10^6$ ,  $3.6 \times 10^6$ ; for  $\text{Ni(H}_2\text{O)}_3\text{tren}^{2+}$   $7 \times 10^5$  and  $6 \times 10^4$ ,  $7.8 \times 10^6$ . The agreement is about as good as usually found except for the  $\text{en}_2$  case. A similar result is found for terpyridine substituting on  $\text{Ni(H}_2\text{O)}_3\text{terpy}^{2+}$  where the usual estimate of  $k_1$  is 100–200 times too large<sup>15</sup>.

The shift data for the  $^{17}\text{O}$  resonance in bulk water for  $\text{Ni(II)}$  and  $\text{Co(II)}$ , but not  $\text{VO}^{3+}$ , solutions imply fairly constant  $A/h$  values (for a constant  $\mu_{\text{eff}}$ ) for metal–oxygen coupling regardless of substitution in octahedral species. The main exceptions appear to be




$\text{Ni}(\text{H}_2\text{O})_5\text{Cl}^+$  and  $\text{Ni}(\text{H}_2\text{O})_2(\text{bipy})_2^{2+}$ ; however, these values may not be well-determined. An important implication is that at least an approximate idea of the number of bound water molecules in a species can be obtained from such shift measurements. Earlier Morgan et al.<sup>16</sup> had made use of the assumption of constant  $A/h$  to obtain equilibrium data. The  $\text{Ni}(\text{H}_2\text{O})\text{EDTA}^{2-}$  species has received, in this way, strong support in our high pH studies. A shift in the  $^{17}\text{OH}_2$  resonance at high pH was observed and using Higginson and Samuels<sup>17</sup> results for the abundance of the species a "normal"  $A/h$  value is obtained suggesting an octahedral complex with one unprotonated acetate arm replaced by a water molecule.

In the nickel-tren system, the existence of two forms has been reported<sup>18</sup>. A minor species at 25°C suggested to be planar is postulated in addition to the normal octahedral one. In our work,  $^{17}\text{O}$  shift measurements at high temperature do show an abnormal decrease consistent with the endothermic formation of a slow-exchanging or diamagnetic species. Again, shift measurements have promise of revealing more detail on the nature of species present in solution.

The ligand tren appears to bind in four coordination positions to nickel(II). It is easily seen from models that the two remaining water molecules are not structurally equivalent. Discovery of two water-exchange rates is perhaps not surprising in view of the earlier observations<sup>10</sup> on nitrilotriacetate but is still a rarity. The substitution of  $\text{NH}_3$ <sup>3</sup> appears to be for the more rapidly exchanging water as would be expected.

A subject of some importance which has not received much experimental attention concerns equilibria such as



By use of NMR methods, exchange kinetics for both free  $\text{H}_2\text{O}$  and  $\text{Cl}^-$  can be determined<sup>19</sup>. If *only* the above process occurs, water and chloride exchange must follow the same kinetics in detail. In this particular case, and the similar  $\text{Ni}(\text{H}_2\text{O})_2(\text{NCS})_4^{2-}$  system<sup>20</sup> the situation is more complicated than a single replacement equilibrium permits. The question of direct ligand replacement can be studied if suitable systems can be found. Preferably, only a single metal ion species should be present.

#### D. CONCLUSION

A significant amount of water-exchange data now exists and it can be understood approximately in conventional terms. The mechanisms for exchange are not proven and the order with respect to solvent remains unknown. It clearly is desirable to obtain more precise activation parameters for exchange as well as for net substitution processes. Comparisons between different metals give promise of revealing some unexpected results. In addition to kinetic information, the  $^{17}\text{O}$  NMR methods can be very helpful in understanding the nature of partially substituted aquo ions and their solutions.

#### ACKNOWLEDGEMENTS

The studies made at Washington State University were made possible by generous sup-

port from the U.S. Atomic Energy Commission, the Washington State University Research Committee, and the National Science Foundation through an Institutional Grant.

## REFERENCES

- 1 T R. Stengle and C H Langford, *Coord. Chem. Rev.* 2 (1967) 349, E.F. Caldin, *Fast Reactions in Solution*, Blackwell, Oxford, 1964.
- 2 D J. Hewkin and R H Prince, *Coord. Chem. Rev.* 5 (1970) 45, A. McAuley and J. Hill, *Quart. Rev.*, 23 (1969) 18
- 3 J.P. Jones, E J Billo and D W Margerum, *J Amer. Chem. Soc.*, 92 (1970) 1875 and references therein.
- 4 T.J. Swift and R E Connick, *J Chem. Phys.*, 37 (1962) 307, 41 (1964) 2553
- 5 (a) R. Murray, H.W. Dodgen and J P Hunt, *Inorg. Chem.*, 3 (1964) 1576;  
(b) H H. Glaeser, H.W. Dodgen and J P Hunt, *Inorg. Chem.*, 4 (1965) 1061.
- 6 N.S. Angerman and R B. Jordan, *Inorg. Chem.*, 8 (1969) 1824
- 7 H.H. Glaeser, G A. Lo, H.W. Dodgen and J P. Hunt, *Inorg. Chem.*, 4 (1965) 206.
- 8 K. Wuthrich and R E Connick, *Inorg. Chem.*, 6 (1967) 583, 7 (1968) 1377.
- 9 M. Grant, H W. Dodgen and J P. Hunt, *J Amer. Chem. Soc.*, 91 (1969) 6318
- 10 R.E. Connick, private communication
- 11 A. G. Desai, H.W. Dodgen and J.P. Hunt, *J Amer. Chem. Soc.*, 92 (1970) 798
- 12 S. Funahashi and M. Tanaka, *Inorg. Chem.*, 8 (1969) 2159.
- 13 J.O. Edwards, *J. Amer. Chem. Soc.*, 78 (1956) 1819.
- 14 P. Hoggard, *Ph. D. Thesis*, Washington State University, 1970
- 15 R.H. Holyer, C D Hubbard, S F A. Kettle and R.G. Wilkins, *Inorg. Chem.*, 5 (1966) 622.
- 16 A H. Zeltman, N A Matwiyoff, and L O Morgan, *J. Phys. Chem.*, 72 (1968) 121.
- 17 W.C.E. Higginson and B. Samuel, *J. Chem. Soc.*, (1970) 1579
- 18 R.G. Wilkins, R. Yelin, D W. Margerum and D C Weatherburn, *J. Amer. Chem. Soc.*, 91 (1969) 4326
- 19 S.F. Lincoln, F. Aprile, H W. Dodgen and J P. Hunt, *Inorg. Chem.*, 7 (1968) 929.
- 20 R B Jordan, H W. Dodgen, and J P. Hunt, *Inorg. Chem.*, 5 (1966) 1906.
- 21 M. Grant, H.W. Dodgen and J.P. Hunt, *Inorg. Chem.*, 10 (1971) 71.
- 22 A.H. Zeltman, N A. Matwiyoff and L.O. Morgan, *J. Phys. Chem.*, 73 (1969) 2689.
- 23 T.J. Swift and R.E. Connick, *J. Chem. Phys.*, 37 (1962) 307.
- 24 R.E. Connick and D. Fiat, *J. Chem. Phys.*, 44 (1966) 4103.
- 25 A.G. Desai, H.W. Dodgen, and J.P. Hunt, *J. Amer. Chem. Soc.*, 91 (1969) 5001.
- 26 D. Rablen and G. Gordon, *Inorg. Chem.*, 8 (1969) 395.
- 27 M. Grant, H.W. Dodgen and J.P. Hunt, *J. Amer. Chem. Soc.*, 92 (1970) 2321.
- 28 D.P. Rablen, unpublished work in these laboratories.
- 29 M. Grant, H.W. Dodgen and J.P. Hunt, *Chem. Commun.*, (1970) 1446.