

CHROMATOGRAPHY OF COORDINATION COMPLEXES

LEONARD F. DRUDING*

Rutgers, The State University, Newark, New Jersey 07102 (U.S.A.)

GEORGE B. KAUFFMAN**

California State College at Fresno, Fresno, California 93726 (U.S.A.)

(Received March 25th, 1968)

CONTENTS

- A. Introduction
- B. Separation and identification of different elements by formation of coordination complexes
- C. Preparation and purification of coordination complexes
- D. Separation of different coordination complexes of the same element
 - (i) Complexes with different charges
 - (ii) Complexes with the same charge
- E. Separation and identification of geometric isomers
- F. Resolution of optically active coordination complexes
- G. Summary

A. INTRODUCTION

Although chromatography in its many forms has been applied mostly to the separation of organic and biological materials, recent years have seen a growing number of applications of the various chromatographic techniques to inorganic separations. Not only has chromatography been used for the qualitative and quantitative analysis of inorganic substances, but it has also been employed to prepare pure inorganic compounds, to follow inorganic reaction kinetics, and to determine such physical parameters as the configuration and charge of inorganic species. It

* This author gratefully acknowledges the financial support of the Rutgers University Research Council and the U.S. Educational Foundation in the Republic of China.

** This author gratefully acknowledges the financial support of the donors of the Petroleum Research Fund administered by the American Chemical Society (Grant 1152-B), and the California State College at Fresno Research Committee.

is in these latter areas of investigation that chromatography has been particularly useful to the coordination chemist. For the separation of ionic species, ion exchange chromatography has of course been the preferred technique. Yet the literature contains numerous examples of the separation of inorganic complexes by column, paper, thin-layer, and gas chromatography as well as by ion electrophoresis.

Most of the chromatographic separations of complexes have involved hexacoordinate octahedral species, but a few studies of tetracoordinate square planar complexes have been made^{39,69,71}. The following generalizations should be valid for all coordination numbers and configurations.

The type of separation to be effected largely determines the choice of technique. For example, in cases in which the preparation or isolation of a chemical species is the goal, a column, with its greater capacity, is generally preferred. However, for the qualitative identification of components in a mixture or for the study of properties such as charge or geometry, paper, gas, or thin-layer chromatography, techniques which are rapid and adaptable to micro amounts of sample, are often more attractive.

The type of species to be separated also influences the choice of chromatographic method, adsorbent, and solvent system. Most frequently, the separation involves ions of different charge density so that use of ion exchange resins with an aqueous solvent system is most effective. Differences in retention by the resin can then be related to the charge density or charge distribution of the ions or to the ease of complex formation with the eluent*.

* We must note here that in addition to the conventional ion exchange resins, certain other adsorbents such as silica gel and alumina should probably be classified as ion exchange types of solid substrates, at least as far as inorganic separations are concerned. A number of studies have been made of the adsorption of complex ions on silica gel¹²³, and considerable evidence has accumulated that silica gel can behave as a weak cation exchange resin. The gel surface consists of silanol (Si-O-H) sites whose hydrogen atoms may be exchanged for cations in a reasonably predictable manner. Dugger *et al.*⁴¹ have tabulated enthalpy, entropy, and free energy values for the exchange reaction between metal ions and the hydrogen atom of the silanol group of silica gel. They found that the bond energy was roughly proportional to the charge density of the unhydrated ion. This finding suggests that in the chromatography of cations, those of higher charge density should be retained more strongly and so have lower R_F values. The results of a thin-layer chromatographic investigation of some inorganic ions by Druding³⁸ do not entirely agree with this prediction, but they do suggest that retention is at least proportional to the charge density of the hydrated ion. In the separation of alkali metal ions and alkaline earth metal ions on silica gel in an acidic solvent, the order of R_F values was $\text{Li}^+ > \text{Na}^+ > \text{Rb}^+ > \text{Cs}^+$ and $\text{Be}^{2+} > \text{Mg}^{2+} > \text{Ca}^{2+} > \text{Sr}^{2+} > \text{Ba}^{2+}$.

An alternative explanation for Druding's results can be based on the ease of substitution of a silanol group for the water of hydration in the inner coordination sphere of the metal ion. Lithium, with its high charge density, would be the least likely to release its waters of hydration for a silanol group and so should exhibit less retention and consequently a higher R_F value than the other alkali metal ions.

Additional evidence for an ion exchange reaction mechanism, which is very close to the explanation just cited for the alkali metal ions was developed by Burwell *et al.*²⁰ in studies of the adsorption of coordination complexes on silica gel and their reaction with the gel. They found

In the choice of stationary phase and solvent, the separation of nonionic inorganic complexes may be treated in the same manner as the separation of polar organic compounds. Differences in dipole moment or solubility in the developer are the important factors that govern the effectiveness of the separation.

An exciting recent development is the use of gas chromatography for the separation of volatile nonionic metal compounds such as carbonyls, alkoxides, alkyls, and especially chelates with oxygen, sulfur, selenium, phosphorus, and nitrogen as donor atoms¹¹². Since its inception in the late nineteen-fifties the application of gas chromatography to the metal chelates of the β -diketones has developed rapidly. The first such studies resulted in the separation of beryllium, aluminum, and chromium by gas chromatography of their acetylacetonates^{8,54}. The method has since been extended not only to acetylacetonates of other metals but also to trifluoroacetylacetonates, hexafluoroacetylacetonates, and other ligands. By use of the new technique, rare earths⁴³ and geometric isomers have been separated, and even optical isomers have been resolved¹³⁹. Further details and applications can be found in the excellent monograph by Moshier and Sievers¹¹².

that the silanol group could displace chloride ion from the coordination sphere of such inert complexes as *cis*- and *trans*-[Co(en)₂Cl₂]⁺ when these complexes were adsorbed onto the surface of the silica gel. The degree of adsorption was also found to be dependent upon the charge and charge distribution of the complex ion. Their observations were confirmed by the thin-layer chromatographic studies of some cobalt(III) complexes made by Druding and Hagel³⁸. The retention values of Burwell *et al.*²⁰ and the *R_F* values of Druding and Hagel³⁸ for appropriate cobalt(III) ammine complexes are compared in Table 1. The greater retention of the *cis* isomers over the corresponding *trans* isomers agrees with the greater charge polarization of the *cis* complex as compared to the *trans* complex.

TABLE 1

RETENTION OF OCTAHEDRAL COORDINATION COMPLEXES ON SILICA GEL AS A FUNCTION OF CHARGE ON THE COMPLEX

Complex	Retention (Burwell <i>et al.</i>) ²⁰	<i>R_F</i> (Druding and Hagel) ³⁸
[Co(NH ₃) ₆] ³⁺	—	0.20
[Co(NH ₃) ₅ NO ₂] ²⁺	0.036 mole/g.SiO ₂	0.23
<i>cis</i> -[Co(NH ₃) ₄ (NO ₂) ₂] ⁺	—	0.60
<i>trans</i> -[Co(NH ₃) ₄ (NO ₂) ₂] ⁺	—	0.80
[Co(NH ₃) ₃ (NO ₂) ₃]	slight	0.98
[Co(NO ₂) ₆] ³⁻	no retention	1.00

In his study of the paper chromatography of cobalt(III) amines, Yoneda¹⁶⁸ suggests that even on cellulose, an ion exchange mechanism may be the basis for separation. In the absence of an electrolyte in the developer, long tails and diffuse spots resulted. However, with an electrolyte in the developer, tailing decreased; the spots were more intense, but some tailing persisted. Use of a base eliminated the tailing. Yoneda cites this as evidence for an ion exchange mechanism involving the hydroxy sites of the cellulose structure. The electrolyte displaces the complex ion from the cellulose. As the value of the charge on the complex decreases, its mobility increases. Yoneda¹⁶⁸ has even suggested the possibility of determining the ionic charge on a complex by noting its *R_F* value.

We shall now briefly review recent developments in the chromatography of coordination complexes according to the type of separation involved. The coverage is intended to be illustrative rather than exhaustive.

B. SEPARATION AND IDENTIFICATION OF DIFFERENT ELEMENTS BY FORMATION OF COORDINATION COMPLEXES

In this phase of chromatography, *viz.*, the separation and identification of different metal ions by complex formation with the same ligand, the formation of the complex ion is only the means for effecting the separation of the elements⁵³.

In some cases complex formation is mandatory; *e.g.*, amphoteric elements such as arsenic and antimony, which hydrolyze completely and immediately on an alumina column, can be chromatographed as tartrate complexes¹³². The order of elution of species can also be changed by conversion to complexes¹³².

Several months after the first articles^{1,93} on the paper chromatography of cations had appeared, Elbeih *et al.*⁴⁴ pointed out that the formation of complexes plays an important role in the separation of cations. R_F values have been correlated with stability constants of complexes^{73,74}.

As might be expected, ion exchange methods, with their high sensitivities to slight differences in charge densities, have been the most popular of chromatographic separations. Only a few of the more recent separations of unusual complexes will now be cited. Chromatographic separation of elements by complex formation is almost as old as the invention of modern ion exchange resins. The now classical separation of iron, cobalt, and nickel depends on the formation of hexachloro complexes of each ion under various conditions. Kurt A. Kraus and his co-workers have made extensive studies of the separation of many elements as chloro complexes by means of anion exchange resins⁸³. Commercial separation of the lanthanide elements from one another is usually accomplished on cation exchange columns; formation and stability of the lanthanide-EDTA complex are the dominant equilibria here⁷⁰.

In a few instances such as those just cited, where either a preparation is involved or a subsequent quantitative determination of each species is desired, column methods are preferred because of their greater capacity. Usually, however, the purposes of separating individual elements is to identify rapidly and qualitatively the components of a mixture. In such cases, a method more suitable to semimicro or micro quantities is then used (paper, gas, or thin-layer chromatography or ion electrochromatography).

Table 2 lists a few recent separations of different elements as coordination complexes. In most of these separations, the mechanism of separation can be correlated with the charge density of the complex species or with the difference in

TABLE 2

SEPARATION OF ELEMENTS BY FORMATION OF COORDINATION COMPLEXES

<i>Elements separated</i>	<i>Ligand</i>	<i>Chromatographic medium</i>	<i>Ref.</i>
Various elements	chloride ion	column; Dowex 1	83
Cd, Cu, Ni, Zn	glycinate ion	paper	63
Ca, Cu	ammonia	column; alumina	81
Hg, Cu, Cd, Bi, Pb	diethyldithiocarbamate ion	thin-layer; silica gel	136
Zn, Cd	pyridine, thiocyanate ion	paper	146
Lanthanides	bis(2-ethylhexyl)phosphate ion	thin-layer; silica gel	36, 105
Lanthanides	α -hydroxybutyric acid	thin-layer cellulose electrophoresis	3, 17
Cr, Co	ammonia	column; silica gel	61
Cu-Cd, Th-Fe- UO_2^{2+} , Th-Ce-Fe, Al-Zr, Ti-Fe-Al	chloride ion	ion exchange-impregnated paper	94
Transition metals	fluoride ion	ion exchange-impregnated paper	95
Ni, Co, Mn, Zn, Sb, Sn, Al, Cr	ammonia, hydroxide ion	ion electrophoresis	103
Fe, Co, Mn, Ni	nitrosophthalol	thin-layer; silica gel	137
Cu, Ni, Co	thiocyanate ion	paper	145
Ni, Co, Cd, Cu, Zn	thiocyanate ion	thin-layer; silica gel	147
Ag, Zn, Cu, Cd, Co	ethylenediamine	thin-layer; silica gel	151
Co from Cu, Ti, Zn, Ni	ethylenediaminetetraacetate ion	column; silica gel	149
Zn, Cu	citrate ion	column; silica gel	150
Ag, Au, Pd, Pt	dithizone	thin-layer; silica gel	57
Hg, Pb, Cu, Bi, Cd, Zn	dithizone	thin-layer; silica gel	56
Cu, Cd, Hg, Pb, Bi	dithizone	paper	113, 114
Be, Al, Cr	acetylacetone	gas chromatography	8, 54
Lanthanides	2,2,6,6-tetramethyl-3,5-heptanedione	gas chromatography (Teflon, 2% Apiezon H on Gas Pack F)	43

stability of the complex ion itself with respect to substitution or solvolysis. In Lederer's work⁹⁴⁻⁹⁶ with ion exchange-impregnated paper, separation is based on differences in the formation of either chloro or fluoro complexes of the metal ions with the solvent, resulting in desorption from the cation exchanger. Kolthoff and Stenger's separation⁸¹ of copper from calcium depends upon the formation of the copper(II) ammine complex, which has a lower charge density than does the calcium ion. On the other hand, Jursik's separation of some post-transition metal bis-glycinates⁶³ is related entirely to the charge densities of these complexes.

C. PREPARATION AND PURIFICATION OF COORDINATION COMPLEXES

With increasing frequency, the inorganic chemist is turning to the use of chromatography in his attempts to prepare and purify isomerically pure coordina-

tion complexes. Chromatography is usually applied to separate the product from its reactants, but in many cases a separation of the product from possible by-products is also desired. Ion exchange methods have been the almost universal choice for preparative work because of the capacity of columns for preparative quantities of material and because of the ionic nature of the substances to be separated. Usually, the product is of quite different charge density from the reactants (starting materials for the most common preparations of coordination complexes are hydrated metal cations and either neutral or anionic ligands).

Although preparative thin-layer chromatography has been extensively applied to the preparation and isolation of small quantities of valuable organic and biochemical substances and although Kauffman and Benson⁶⁹ have demonstrated the feasibility of preparative separations of inorganic complexes by TLC, no applications of preparative thin-layer chromatography to coordination compounds have been reported. This technique would be of greatest advantage in cases where

TABLE 3

USE OF CHROMATOGRAPHY IN THE PREPARATION AND PURIFICATION OF COORDINATION COMPLEXES

Product	Other components	Stationary phase	Ref.
Substituted acetyl-acetonates of Co^{III} and Cr^{III}	HAcac^a , Co^{3+} , Cr^{3+}	column; alumina	31
Metal trifluoro-acetylacetonates	metal benzoylacetonates	column; alumina	48
$[\text{Co}(\text{Acac})_3]^a$	reaction by-products	column; Florosil	31
$[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$	$[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]^{2+}$	ion electrophoresis	124
$[\text{Co}(\text{H}_2\text{O})_4(\text{Acac})_2\text{NO}_2]$	other ions	column; Dowex 1	35
$[\text{Co}(\text{en})_2\text{am}]^{2+}$	amino acids (Ham)	paper; thin-layer; cellulose	18
$[(\text{H}_2\text{O})_4\text{Cr}(\text{OH})(\text{SO}_4)-\text{Cr}(\text{H}_2\text{O})_4]^{3+}$	$[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$, OH^-	column; Dowex 50	50a
Bis(2,3-diaminopropionate)cobalt(III) chloride	$\text{Na}_3[\text{Co}(\text{CO}_3)_3]$, 2,3-diaminopropionic acid hydrochloride	column; Dowex 50	50b
$[\text{Co}(\text{trien})\text{am}]^{2+}$	amino acids (Ham)	paper; thin-layer; cellulose	19
Nitro-en-pn complexes of Co^{III}	$[\text{Co}(\text{NH}_3)(\text{en})(\text{NO}_2)_3]$, pn	paper	65
$[\text{Co}(\text{gly})_2(\text{L-val})]^d$	amino acids, Co complexes	column; alumina	138
Co^{III} amines	anions	column; Amberlite IRA 140	2
$[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$	$[\text{Cr}(\text{H}_2\text{O})_5(\text{NCS})]^{2+}$, $[\text{Cr}(\text{H}_2\text{O})_4(\text{NCS})_2]^+$	column; Dowex 50	76
$[\text{Cr}(\text{NH}_3)_6]^{3+}$	$[\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})]^{2+}$, $[\text{Cr}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$	column; alumina	62
$[\text{Cr}(\text{en})(\text{H}_2\text{O})_4]^{3+}$	$[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$	column; Dowex 50	27a
$[\text{Cr}(\text{en})_2(\text{H}_2\text{O})\text{Cl}]^{2+}$	$[\text{Cr}(\text{en})_2\text{Cl}_2]^+$	column; Dowex 50	100
$\text{Cr}(\text{en})_2(\text{H}_2\text{O})\text{Cl}]^{2+}$	other ions	column; Dowex 50	101
$[\text{Cr}(\text{en})_2\text{F}_2]^+$	$[\text{Cr}(\text{en})_2(\text{H}_2\text{O})\text{F}]^{2+}$	column; Dowex 50	49, 142a
$[\text{Cr}(\text{en})_2(\text{H}_2\text{O})\text{NCS}]^{2+}$	$[\text{Cr}(\text{en})_2(\text{NCS})\text{Cl}]^+$	column; Dowex 50	144
$[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$	reactants	column; Dowex 50	117
Troponeiron carbonyls	iron carbonyls	column; alumina	14

TABLE 3 (continued)

Product	Other components	Stationary phase	Ref.
Organometallic carbonyl complexes of cobalt	acetylenes and cobalt compounds	column; alumina	86, 87
Organometallic carbonyl complexes of iron	acetylene and iron carbonyls	column; alumina	153
<i>cis</i> -[Rh(C ₅ H ₅ N) ₃ Cl ₂] [Ru(H ₂ O) ₂ Cl ₂]	<i>trans</i> -[Rh(C ₅ H ₅ N) ₄ Cl ₂]Cl other Ru ^{III} complexes	column; alumina column; Dowex 50W, Dowex 1	131 34
[Ru(H ₂ O) ₂ Cl ₂] ⁻ [Pd(Acac) ₂]	other Ru ^{III} complexes reaction products	column; Dowex 1 column; alumina	15 142
Alkyl and aryl derivatives of Pt ^{II} phosphines	starting materials and isomers	column; alumina	26, 27
C ₅ H ₅ Fe(CO)(COCH ₂)L (L = (C ₆ H ₅) ₃ P, (C ₆ H ₅) ₂ P, (C ₆ H ₅ O) ₂ P)	C ₅ H ₅ Fe(CO) ₂ CH ₃ and phosphines or phosphites	column; alumina	7
Rh ₃ [(C ₆ H ₅ O) ₃ P] ₄ (SCN) ₂ [Mo(CO) ₄ -(C ₆ H ₅) ₃ M]SCN] (M = P, As, Sb)	Rh[(C ₆ H ₅ O) ₃ P] ₂ COCl, KSCN <i>cis</i> -[Mo(CO) ₃ [(C ₆ H ₅) ₃ M] ₂ SCN], <i>trans</i> -[Mo(CO) ₃ -(C ₆ H ₅) ₃ M] ₂ NCS]	column; alumina column; alumina	63 155
Thiocyanatocarbonyl Mn ⁺ complexes	Thiocyanatopentacarbonyl-manganese(I), amines, phosphines, arsines, stibines	column; Florosil, alumina	45
[Co(fac) ₃] ⁺ [CH ₃ SFe ₂ (CO) ₄] ₂ S	Co ₂ (CO) ₈ , hfac Fe ₂ (CO) ₉ , [CH ₃ SFe(CO) ₃] ₂ , methyl thiocyanate	column; alumina column; alumina	75 28
[(CH ₃ S) ₂ Fe ₂ (CO) ₄ C ₇ H ₅] Fulveneiron carbonyls [MCl _n Br _{6-n}] ²⁻ (M = Ir, Os, Pt, Re; n = 0, 1, ... 6)	<i>anti</i> - and <i>syn</i> -[CH ₃ SFe(CO) ₃] ₂ fulvenes and iron carbonyls M ^{IV} , Cl ⁻ , Br ⁻	column; alumina column; alumina paper; high voltage electrophoresis	80 152 12, 120-122
Chloroaquo and chlorohydroxo complexes of Rh ^{III} , Ir ^{III} and Pt ^{IV}	Rh ^{III} , Ir ^{III} , Pt ^{IV} , Cl ⁻ , OH ⁻	paper; high voltage electrophoresis	11
L-β-[Co(trien)Ox]Cl ^f	L-β-Co(trien)Cl ₂ Cl, H ₂ Ox, K ₂ Ox	column; Dowex 50	92
[Co(en)(gly) ₂] ⁺ [Co(gly) ₂ Ox] ⁻ [Rh(R)-pn]Cl ₄ ⁻	CoCl ₂ , en, glycine CoCl ₂ , glycine, Ox ⁻ [Rh(pn) ₂ Cl ₂] ⁺ , [Rh(pn) ₃] ²⁺	column; Dowex 50 column; Dowex 1 column; Dowex 1	106 106 51a
Bis[1,3-(2'-pyridyl)-1,2-diaza-2-propenato]-rhodium(III) chloride	RhCl ₃ , ligand	column; Chromedia P-11 (cellulose phosphate)	6
Tris[1,3-(2'-pyridyl)-1,2-diaza-2-propenato]rhodium(III) chloride	corresponding ClO ₄ ⁻	column; Deacidite FF (anion exchange resin)	6

^a HAcac = acetylacetone; Acac = acetylacetonate ion; ^b en = ethylenediamine; Ham = amino acid; am = amino acid anion; ^c trien = triethylenetetramine; ^d gly = glycinate ion; val = valinate ion; ^e hfac = hexafluoroacetylacetone; fac = hexafluoroacetylacetonate ion; ^f Ox = C₃O₄⁴⁻.

the reaction yields are small or where the desired compound is of high value such as in the preparation of complexes of the platinum metals.

The general choice of ion exchange column methods is again a logical one for most preparations and purifications. Whereas separation of product from reactants is usually quite easy, separation of product from by-products is often more difficult because of similarities in charge and structure. Yet, even with ion exchange resins, it is possible to separate geometric isomers^{15,34,100,101}.

Use of silica gel, alumina, or cellulose as column adsorbents has been most frequent in cases where the complexes to be separated are of very similar charge to the reactants or where neutral complexes are involved.

Table 3 summarizes some of the applications of chromatography to preparative coordination chemistry. The work of Connick and Fine³⁴ illustrates the specificity of ion exchange methods. By using both cation and anion resins, they were able to separate other isomers from a solution containing the desired product $[\text{Ru}(\text{H}_2\text{O})_3\text{Cl}_3]$, and then were able, by chromatography, to identify the presence of two isomers of neutral trichloroaquoruthenium(III) complexes.

Another very useful preparative application of chromatography, on cellulose, is that of Buckingham and co-workers^{18,19}. In studying the reaction between some cobalt(III) hydroxo-aqua-tetramines and amino acids, they used cellulose chromatography (paper, thin-layer, and column) to separate the resulting amino acid cobalt complex from unreacted amino acid and other reaction products.

Thus far, preparative chromatography has been limited largely to octahedral complexes; only a few examples^{26,27,142}, involving square planar complexes appear in the literature. The technique is not limited to mononuclear complexes but has also been applied to polynuclear complexes^{50a}.

D. SEPARATION OF DIFFERENT COORDINATION COMPLEXES OF THE SAME ELEMENT

This type of chromatography of coordination complexes is very closely related to the previous type except that here will be included those separations that are used primarily to study reaction kinetics and to determine the charge on a coordination species.

The number of square planar complexes investigated is few compared to the number of octahedral complexes separated. Thus it is difficult at present to report whether the trends observed in the separation of octahedral complexes apply as well to complexes of other coordination numbers.

(i) Complexes with different charges

In several publications^{23,59,100,101,116,144}, chromatographic separation has been used to follow the rate of hydrolysis reactions. In these reactions, an anionic

TABLE 4

SEPARATION OF DIFFERENT COMPLEXES OF THE SAME ELEMENT BY CHROMATOGRAPHY

Complex	Stationary phase	Purpose	Ref.
Co ^{III} amines	thin-layer; silica gel	identification	39
Co ^{III} nitroammines	column; alumina	charge identification	61
Co ^{III} nitroammines	paper	separation	82
Co ^{III} nitroammines	ion electrophoresis	charge identification	102
Co ^{III} nitroammines	paper	separation	157
Co ^{III} amines	paper	charge identification	158
Ethylenediamine- <i>l</i> -propylenediamine complexes of Co ^{III}	cellulose	identification	42
[Co(NH ₃) ₄ X ₂] ⁺	ion electrophoresis	follow kinetics	124
Co ^{III} amines	paper	separation	119
[Co(en) ₂ X ₂] ²⁺ (X = Cl, Br, NCS, NO ₂)	ion electrophoresis	separation	10
[Co(en) ₂ Ac ₂] ²⁺ , [Co(en) ₂ (H ₂ O)Ac] ²⁺	column; Dowex 50	follow kinetics	23
[Co(en) ₂ Ac ₂] ²⁺ , [Co(en) ₂ (H ₂ O)Ac] ²⁺	ion exchange paper; Amberlite SA-2	follow kinetics	24
Cr ³⁺ , [Cr(H ₂ O) ₅ (NCS)] ²⁺ , [Cr(H ₂ O) ₄ (NCS) ₂] ⁺	column; Dowex 50	separation	76
[Cr(H ₂ O) ₄ Ox] ²⁺ , [Cr(H ₂ O) ₂ Ox ₂] ^{-c}	column; Dowex 50	follow kinetics	21
[Cr(H ₂ O) ₅ X] ²⁺ (X = F, Cl, Br, I)	column; Dowex 50	separation, follow kinetics	117
Cr ^{III} amines	column; Dowex 50	follow kinetics	123
[Cr(en)NH ₃ (H ₂ O)Br _{3-x}] ²⁺ (x = 1, 2, 3)	column; Dowex 50	identification	59
[Cr(en) ₂ (OH) ₂] ²⁺ , [Cr(en) ₂ Cl ₂] ²⁺	column; Dowex 50	follow kinetics	116
[Cr(en) ₂ Cl ₂] ²⁺ , [Cr(en) ₂ (H ₂ O)Cl] ²⁺	column; Dowex 50	follow kinetics	100
Cr ^{III} amines	thin-layer; silica, alumina	separation; identification	80
[Cr(en) ₂ Cl ₂] ²⁺ , [Cr(en) ₂ (H ₂ O)Cl] ²⁺ , [Cr(en) ₂ (H ₂ O) ₂] ²⁺	column; Dowex 50	follow kinetics	101
[Cr(en) ₂ F ₂] ²⁺ , [Cr(en) ₂ (H ₂ O)F] ²⁺	column; Dowex 50	follow kinetics	49
[Cr(trien)Ox] ²⁺ ^d	column; Dowex 50	follow kinetics	143
[Cr(en) ₂ Cl(NCS)] ²⁺ , [Cr(en) ₂ (H ₂ O)Cl] ²⁺	column; Dowex 50	follow kinetics	144
[Ir(EnS) ₃ Cl ₃]	ion electrophoresis	charge identification	72
[IrCl ₃ (CO){C ₆ H ₅ PCl ₂ }]	thin layer; silica gel	identification	30
[Mo(CO) ₄ {C ₆ H ₅ PCl ₂ }-{C ₆ H ₅ PO}]	column; alumina	identification	51
Pt ^{II} amines	paper	identification	4
Pt ^{II} amines	thin-layer; silica gel	identification	40
Rh chloro complexes	ion electrophoresis; paper; ion exchange	charge identification	96
Ru chloro complexes	column; Dowex 50	identification	22
Ru chloro complexes	column; Dowex 50, Dowex 1	identification	33
[Ru(NO)(H ₂ O) _{5-x} (NO ₂) _x] ^{3-x}	column; Dowex 50	identification	130

^a en = ethylenediamine; ^b Ac = acetate ion; ^c Ox = C₂O₄²⁻; ^d trien = triethylenetetramine.

ligand was replaced by water, resulting in an increase in the charge on the complex. One important limitation to the use of chromatography in the study of reaction kinetics is the fact that no further reaction should take place either with the eluent or between the resin or stationary phase and the adsorbed complex. This restricts the use of chromatography to inert complexes such as those of cobalt(III), chromium(III), and a few other elements, for which reaction times are long compared to the time required to chromatograph the substance.

One of the commonest types of reaction of coordination complexes that has been investigated is a solvolysis reaction in which a solvent molecule replaces one of the ligands, usually an anionic ligand so that the resulting reaction product has a higher charge than the initial complex.

Here again ion exchange is the most popular method as would be expected for separations based on differences in charge. However, inasmuch as many of the separations were developed for the purpose of identification and characterization of ionic charge on a micro or semimicro scale, there are numerous examples of paper chromatographic, thin-layer chromatographic, and ion electrophoretic methods. In most cases, acidic aqueous solvents have been used with a variety of stationary phases. Although separation of octahedral complexes, especially those of chromium(III) and cobalt(III), predominate, several examples of the separation of square planar coordination complexes have appeared in the literature.

It is interesting to compare the results in the first six references in Table 4 for the separation of the various nitroamminecobalt(III) ions, $[\text{Co}(\text{NH}_3)_5-x(\text{NO}_2)_x]^{(3-x)+}$. Although different techniques (column, thin-layer, and paper chromatography; and ion electrophoresis) were used in these experiments, in every case the ionic mobility could be related to the charge on the complex; the chromatographic ionic mobility decreased with increasing charge on the complex. In the case of ion electrophoresis, ionic mobility toward the electrode of opposite polarity from the charge was enhanced by the increased absolute charge on the ion.

In some cases, the chromatographic method was used specifically to characterize the charge on the complex. Kauffman *et al.*⁷² and Maddock and Todesco¹⁰² both used ion electrophoresis in their studies of complexes of type $[\text{MX}_3\text{Y}_3]$ to show that what were supposedly two geometric isomers of a neutral complex were instead the neutral complex and one "polymerization" isomer.

Yoneda's recent study¹⁵⁸ of the paper chromatography of some cobalt(III) amines, in which a 1M aqueous solution of ethylamine was used as developer, suggests that the R_F values can be correlated with the charge on the ion. Ions of 3+ charge have $R_F = 0.10-0.20$; 2+ ions, $R_F = 0.20-0.30$; 1+ ions, $R_F = 0.40-0.60$; neutral complexes, $R_F = 0.70-0.85$; 1- ions, $R_F = 0.85-0.90$, and 3- ions, $R_F = 0.95-1.00$.

The work of Pollard, Banister, Geary and Nickless¹¹⁹ is interesting for its inconsistency when compared to other similar separations of cobalt(III) amine

complexes. They found little relation between R_F values and charge or geometry of the complexes—features that are present in all other research on the subject. They did report some secondary spots and considerable tailing in the chromatography of some of their complexes, a fact which suggests that the complexes were undergoing some kind of decomposition either in solution or during the development of the chromatogram.

(ii) *Complexes with the same charge*

By the use of aqueous ethanol acidified with hydrochloric acid, Basolo *et al.*⁵ were able to separate on paper the *cis* isomers of $[\text{Pt}(\text{NH}_3)_2\text{X}_2]$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) from each other and from the *trans* isomers, but not the *trans* isomers from each other. They accomplished the latter separations by use of acetone–water mixtures.

Dwyer and Sargeson⁴² employed partition chromatography on cellulose to study the composition of the reaction product obtained from *l*-propylenediamine, and *cis*-dichloro- and *cis*-dinitrobis(ethylenediamine)cobalt(III) chlorides. The separations were repeated with a cellulose pulp column, and the complexes obtained in order of elution were $\text{L}[\text{Co}(\text{l-pn})_3]\text{Cl}_3$, $\text{L}[\text{Co}(\text{l-pn})_2\text{en}]\text{Cl}_3$, $\text{L}[\text{Co}(\text{en})_2\text{l-pn}]\text{Cl}_3$, $\text{D}[\text{Co}(\text{l-pn})_3]\text{Cl}_3$, $\text{D}[\text{Co}(\text{l-pn})_2\text{en}]\text{Cl}_3$, $\text{D}[\text{Co}(\text{en})_2\text{l-pn}]\text{Cl}_3$, and $\text{D,L}[\text{Co}(\text{en})_3]\text{Cl}_3$.

E. SEPARATION AND IDENTIFICATION OF GEOMETRIC ISOMERS

Most published separations of coordination complexes by chromatography involve the separation of geometric isomers of both octahedral and square planar complexes. These separations have been developed both for the preparation of isomerically pure complexes and for the identification of reaction products. In some cases^{69,71}, separations have been carried out quantitatively.

Whereas a number of separations of geometric isomers have been carried out on ion exchange resins, silica gel and alumina seem to offer sharper separations, especially for complexes of low ionic charge. The latter two substrates are also popular in thin-layer chromatography.

Certain limitations to the use of chromatography for the determination of geometric configuration must be noted. As Burwell *et al.*²⁰ have pointed out, prolonged contact of some complexes with silica gel results in substitution of the silica gel for a ligand. For cobalt(III) complexes, this substitution would undoubtedly be enhanced by use of a basic solvent. For other complexes, prolonged contact with a solvent might result in solvolysis reactions which yield a change in configuration. The thin-layer chromatographic separation of some cobalt(III) amines on silica gel by Seiler, Biebricher and Erlenmeyer¹³³ seems to provide evidence both for hydrolysis and for geometric rearrangement of some of the complexes.

Although they supposedly used pure complexes for their chromatograms, on development more than one spot appeared.

Isomerization on a silica gel column was observed with $[\text{Pt}\{(\text{C}_4\text{H}_9)_3\text{P}\}_2\text{Cl}_2]$ by Kauffman *et al.*⁷¹. The melting points showed that the recovered isomers were pure; the ratio of *cis* isomer to *trans* isomer had changed significantly although the total weight of both recovered isomers indicated that the separation was quantitative.

Overwhelming evidence exists that for octahedral complexes developed with an acidic or water-containing solvent, the *trans* isomer is more mobile than the *cis* isomer. This regularity has been used, along with other evidence, to assign to $[\text{Cr}(\text{en})_2(\text{H}_2\text{O})\text{F}]^{2+}$ a *trans* configuration^{142a}. Burwell *et al.*²⁰ suggest that the *cis* isomer can have two points of attachment to the silanol sites of the silica gel or, by analogous reasoning, to the acidic sites of an ion exchange resin, whereas the *trans* groups, being on opposite sides of the complex, can attach at only one point. Even if the mechanism is one of mere adsorption rather than ion exchange, this trend should still prevail since the *cis* isomer would be more polarized than the *trans* isomer. As an interesting case in point we may cite Kyuno's separations^{90,91} of $[\text{Co}(\text{NH}_3)_2(\text{CO}_3)(\text{NO}_2)_2]^-$ and $[\text{Co}(\text{NH}_3)_2(\text{C}_2\text{O}_4)(\text{NO}_2)_2]^-$ on an alumina column. In both cases, the order of elution of the three possible geometric isomers was *trans*- $\text{NO}_2 > \text{trans}$ - $\text{NH}_3 > \text{cis}$ isomer. In terms of charge polarization, the least polarized ion is thus the most mobile.

As a perusal of Table 5 will show, in a few cases involving square planar complexes^{5,64}, exceptions are found to this general rule that the *trans* isomer is more mobile than the *cis* isomer. Exceptions to the *trans* $>$ *cis* mobility rule are also found for octahedral complexes^{65,115,130,140,141,157}. In the two papers on the paper chromatography of cobalt(III) amine complexes by Stefanović and Janjić^{140,141} and the one by Yamamoto, Nakahara, and Tsuchida¹⁵⁷, the *cis* isomer always exhibits a higher R_F value than the corresponding *trans* isomer. Unfortunately, Yoneda¹⁵⁸ did not attempt to separate geometric isomers by paper chromatography. He reported only the R_F values for *cis*- and *trans*- $[\text{Co}(\text{en})_2(\text{NO}_2)_2]\text{NO}_3$, which were identical. In another study of the effect of ionic charge on mobility, Pollard *et al.*¹¹⁹ provide data only for *cis*- and *trans*- $[\text{Co}(\text{NH}_3)_4(\text{NO}_2)_2]^+$ where $R_F(\text{trans}) > R_F(\text{cis})$. Despite Yoneda's claim that even paper has ion exchange properties, solvent properties may also play an important role in paper chromatography of charged complexes.

A problem that has not yet been fully solved is the separation of geometric isomers of 2+ or 3+ charge. Unfortunately, separations on paper or thin-layer adsorbents are not as sensitive for the more highly charged ions as they are for the 1+, or 1-, or neutral complexes. Yet in the study of solvolysis reactions, it is the rapid determination of the geometric configuration of the solvolysis products that is often crucial to the interpretation of the reaction mechanism.

It may be in order to mention here separations of complexes which are iso-

TABLE 5

SEPARATION OF GEOMETRIC ISOMERS OF COORDINATION COMPLEXES BY CHROMATOGRAPHY

Complex	Stationary phase	Relative mobility	Ref.
Trisbenzoylacetates of Co, Cr, and Rh	column; alumina	<i>trans</i> > <i>cis</i>	47
Tris(trifluoroacetyl)-acetates of Co, Cr, and Rh	column; alumina	<i>trans</i> > <i>cis</i>	48
Co ^{III} amines	paper	<i>trans</i> > <i>cis</i>	82
Co ^{III} amines	paper	irregular	119
Co ^{III} amines	paper	<i>cis</i> > <i>trans</i>	140
Co ^{III} amines	paper	<i>cis</i> > <i>trans</i>	141
Co ^{III} amines	paper	no difference	158
[Co(NH ₃) ₄ (NO ₂) ₂] ⁺	paper	<i>cis</i> > <i>trans</i>	157
[Co(NH ₃) ₄ Cl] ²⁺ , [Co(NH ₃) ₄ (H ₂ O)Cl] ²⁺ , [Co(NH ₃) ₄ (NO ₂) ₂] ⁺	ion electrophoresis	—	124
Co ^{III} amines	thin-layer; silica gel	<i>trans</i> > <i>cis</i>	39
[Co(NH ₃) ₄ (N ₃) ₂] ⁺	column; alumina	<i>trans</i> > <i>cis</i>	99
<i>trans</i> (RS meso)- [CoL(H ₂ O)Cl] ²⁺ (L = 1,4,8,11-tetra- azaundecane)	column; Dowex 50	—	52
[Co(NH ₃) ₃ (NO ₂) ₃]	ion electrophoresis	<i>trans</i> > <i>cis</i>	102
[Co(NH ₃) ₄ (NO ₂) ₂] ⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	77
[Co(NH ₃) ₄ (NO ₂) ₂] ⁺ , [Co(en) ₂ (NO ₂) ₂] ²⁺	column; Amberlite IR120	<i>trans</i> > <i>cis</i>	110
[Co(en) ₂ (NO ₂) ₂] ⁺	paper	<i>cis</i> > <i>trans</i>	65
[Co(en) ₂ X ₂] ⁺	column; Amberlite IR 120	<i>trans</i> > <i>cis</i>	111
[Co(en) ₂ X ₂] ⁺	ion electrophoresis	<i>trans</i> > <i>cis</i>	10
[Co(en) ₂ Cl] ⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	37
[Co(en) ₂ XY] ⁺	thin-layer; silica gel	<i>trans</i> > <i>cis</i>	133
[Co(en) ₂ (NO ₂) ₂] ⁺	paper	<i>cis</i> > <i>trans</i>	65
[Co(en)EDDA] ⁺ , [Co(NH ₃) ₂ EDDA] ²⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	97
[Co(en)(gly) ₂] ²⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	106
[Co(gly) ₂ Ox] ^{-d}	column; Dowex 1	<i>trans</i> > <i>cis</i>	106
[Co(en) ₂ (Ac) ₂] ²⁺	ion exchange paper; Amberlite SA-2	<i>trans</i> > <i>cis</i>	24
Bis(2,3-diaminopropionato)- cobalt(III) chloride	column; Dowex 50	—	50b
[Co(en) ₂ (H ₂ O) ₂] ²⁺	column; alumina	<i>trans</i> > <i>cis</i>	61
[Co(en)Ox(NO ₂) ₂] ⁻	column; Dowex 1	<i>trans</i> -NO ₂ > <i>cis</i>	88
[Co(en)Ox(NO ₂)NCS] ⁻	column; Dowex 1	<i>trans</i> > <i>cis</i>	89
[Co(NH ₃) ₂ CO ₃ (NO ₂) ₂] ⁻	column; Dowex 1	<i>trans</i> -NO ₂ > <i>trans</i> -NH ₃ > <i>cis</i>	90
[Co(NH ₃) ₂ Ox(NO ₂) ₂] ⁻	column; Dowex 1	<i>trans</i> -NO ₂ > <i>trans</i> -NH ₃ > <i>cis</i>	91
[Co(alin) ₃] ^f	column; alumina; thin-layer; silica gel	<i>trans</i> > <i>cis</i>	25
[Cr(H ₂ O) ₄ Cl] ⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	78
Tris(acetylacetanilido)- chromium(III)	column; alumina	<i>trans</i> > <i>cis</i>	135
Tris(acetylacetanilido)- chromium(III)	thin-layer; silica gel	<i>trans</i> > <i>cis</i>	134

TABLE 5 (continued)

Complex	Stationary phase	Relative mobility	Ref.
Cr ^{III} amines	column; alumina	<i>trans</i> > <i>cis</i>	9
[Cr(en)(NH ₃)(H ₂ O) ₂ Cl] ⁺	column; Dowex 50	I > II	59
[Cr(en) ₂ Cl ₂] ⁺ , [Cr(en) ₂ (H ₂ O)Cl] ²⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	100
[Cr(en) ₂ (H ₂ O)Cl] ²⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	101
[Cr(en) ₂ (H ₂ O)NCS] ²⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	144
[Cr(en) ₂ (OH) ₂] ⁺	column; alumina	<i>trans</i> > <i>cis</i>	155
[Cr(H ₂ O) ₄ (NCS) ₂] ⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	55
[Cr(H ₂ O) ₄ Cl ₂] ⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	78
Iron carbonyls	column; alumina	—	14
[CH ₃ Fe(CO) ₂] ₂	column; alumina	<i>anti</i> > <i>syn</i>	28, 79, 80
<i>cis</i> -[Ir{(C ₂ H ₅) ₂ S} ₂ Cl ₂]	electrophoresis	electrolyte > nonelectrolyte	72
[Cu(gly) ₂] ^c	paper	<i>cis</i> > <i>trans</i>	64
[Mn(CO) ₃ ((C ₆ H ₅) ₃ P) ₂ SCN]	column; alumina	—	155
Pt ^{II} amines	paper	<i>cis</i> > <i>trans</i>	4, 5
Pt ^{II} amines	thin-layer; silica gel	<i>trans</i> > <i>cis</i>	40
Pt ^{II} amines (mononuclear and binuclear)	thin-layer; silica gel	<i>trans</i> > <i>cis</i>	69
Pt ^{II} amines (mononuclear and binuclear)	column; alumina	<i>trans</i> > <i>cis</i>	71
[Rh(H ₂ O) ₂ Cl ₂] ⁻	column; Dowex 1	<i>trans</i> > <i>cis</i>	15, 50
[Ru(H ₂ O) ₃ Cl ₂]	column; Dowex 50	<i>trans</i> > <i>cis</i>	32, 34, 50
[Ru(H ₂ O) ₄ Cl ₂] ⁺	column; Dowex 50	<i>trans</i> > <i>cis</i>	33, 50
[Ru(H ₂ O) ₃ Cl ₂]	column; Dowex 50, 1	<i>trans</i> > <i>cis</i>	50, 107
[Ru(NO)(H ₂ O) _{5-x} - (NO ₂) _x] ^{3-x} + (x < 5)	paper	—	130
[Ru(NO)(H ₂ O) _{5-x} - (NO ₂) _x] ^{3-x} + (x > 5)	Zeo-Karb 225 cation exchange resin	<i>cis</i> > <i>trans</i>	130
[PtL ₂ Cl ₂] (L = (C ₂ H ₅) ₂ S, (C ₄ H ₉) ₂ P)	column; alumina, silica gel	<i>trans</i> > <i>cis</i>	68
Alkyl and aryl derivatives of Pt ^{II} phosphines	column; alumina	<i>trans</i> > <i>cis</i>	26, 27
[CH ₃ COMn(CO) ₄ - P(C ₄ H ₉) ₃] ⁻	thin-layer and column; silica gel	<i>cis</i> > <i>trans</i>	115
[Pt(NH ₃) ₂ X ₂] (X = Cl, Br, I)	paper	—	5
Ditolytellurium(II)	thin-layer; alumina	<i>ortho</i> > <i>meta</i> > <i>para</i>	148
[Pt{(C ₆ H ₅) ₂ S}Cl ₂]	thin-layer; silica gel	<i>trans</i> > <i>cis</i>	68
[(C ₂ H ₅) ₃ PAuSC(S)OR] (R = n or iso-propyl)	thin-layer; silica gel	<i>n</i> > <i>iso</i>	68
Cu(3,4- or 3,5-lutidine) ₂ - Cl ₂ (x = 2 or 4)	thin-layer; silica gel	3,5 > 3,4	68
[Pd(Et ₄ dien)SeCN] ⁺ ^g , [Pd(Et ₄ dien)NCS] ⁺	thin-layer; silica gel	Se-bonded > N-bonded	68

^a en = ethylenediamine; ^b H₂EDDA = ethylenediamine-*N,N'*-diacetic acid; EDDA = its binate anion; ^c gly = glycinate ion; ^d Ox = C₄O₄²⁻; ^e Ac = acetate ion; ^f alin = β-alaninate ion; ^g Et₄dien = *N,N,N',N'*-tetraethyldiethylenetetramine.

meric by virtue of isomerism within the ligand itself, such as the TLC separations of di-*o*-, *m*-, and *p*-tolyltellurium(II) by Vobecky¹⁴⁸ and of dichlorobis(*n*-, *iso*-, *sec*-, and *t*-butyl sulfide)platinum(II) (geometric isomers also), bis- or tetrakis-(3,4- or

3,5-lutidine)copper(II) chlorides, and *O*-*n*- or iso-propyldithiocarbonato(triethylphosphine)gold(I) by Kauffman⁶⁸. Work on the TLC separation of linkage isomers is currently in progress in Kauffman's laboratory. The separation of Se-bonded from N-bonded $[\text{Pd}(\text{Et}_4\text{dien})\text{SeCN}] [\text{B}(\text{C}_6\text{H}_5)_4]$ has already been achieved⁶⁸.

F. RESOLUTION OF OPTICALLY ACTIVE COORDINATION COMPLEXES

A rapid, efficient, and general method for separating optical isomers of coordination complexes has long been a goal sought by inorganic chemists. Indeed, Alfred Werner, the founder of coordination chemistry, used resolution to establish unequivocally the octahedral configuration for tripositive cobalt^{66,67}. Often today, a structure-proof depends upon proving the existence of optical isomers. Pasteur's

TABLE 6

RESOLUTION OF OPTICAL ISOMERS OF COORDINATION COMPLEXES BY CHROMATOGRAPHY

Complex	Stationary phase	Purpose	Ref.
$[\text{M}(\text{Acac})_3]^a$ (M = Co, Cr)	alumina and D-tartaric acid; D-quartz	partial resolution	118
$[\text{Co}(\text{Acac})_3]$	column and thin-layer; D-quartz	partial resolution	46
Tris(hexafluoroacetyl- acetato)chromium(III)	D-quartz (gas chromatography)	partial resolution	139
Hexakis(2-aminoethane- thiol)tricobalt(III) cation	Ion exchange cellulose	total resolution	16
$[\text{Co}(\text{en})_3]^{3+}$, $[\text{Co}(\text{gly})_3]^b$	starch	partial resolution	84
Co L-methylxanthate	alumina	total resolution	85
<i>trans</i> - $[\text{Co}(\text{gly})\text{EDDA}]^c$	starch	partial resolution	98
1,2,6- $[\text{Co}(\text{alin})_3]^d$	L-quartz	partial resolution	25
$[\text{Co}(\text{en})_2\text{X}_2]^+$	paper	partial resolution	140
$[\text{Co}(\text{Acac})_3]$, $[\text{Rh}(\text{Acac})_3]$	D-lactose	partial resolution	29
$[\text{Co}(\text{Acac})_3]$, $[\text{Cr}(\text{Acac})_3]$	D-lactose	partial resolution	108
Cu, Ni β -ketoimines	D-lactose	partial resolution	58
Vanadyl ketoimine and diketoimine	D-lactose	partial resolution	126
Bisacetylacetonepropyl- enediiminooxovanadium(IV)	D-lactose	partial resolution	104, 125
$[\text{Y}(\text{Acac})_3]$, $[\text{Ga}(\text{Acac})_3]$	D-lactose	partial resolution	109
Diphenylbis(acetylaceton- propylenediimino)tin(IV)	D-lactose	partial resolution	127
Bis(N-R salicylaldimino)MII (M = Zn, Cu, Ni, Co; R = C ₆ H ₅ , isopropyl, cyclohexyl)	D-quartz	partial resolution	46
$[\text{NiL}]_2(\text{L}=\text{N}(\text{—CH}_2\text{—CH}_2\text{——N=CH—C}_6\text{H}_4\text{N}_2))$	cellulose	partial resolution	154

^a Acac = acetylacetonate ion; ^b gly = glycinate ion; ^c EDDA = ethylenediamine-*N,N'*-diacetate dinegative anion; ^d alaninate ion.

classical fractional crystallization of diastereoisomers, which is still the most widely used method of resolution, is tedious, inefficient, and not always reliable. In a recent article on future developments in analytical chemistry, L. B. Rogers¹²⁹ predicts that a rapid quantitative *chromatographic* separation of optical isomers will be a major advance within the next five years.

A few separations, on a small scale basis, have been reported, but most of these have yielded only partial resolution of the enantiomers. All of the methods are based on the presence of an optically active resolving agent in the stationary phase. With neutral complexes, such as acetylacetonates and β -ketoimines^{29,58,104,108,109,125,126}, that are soluble in organic solvents, D-lactose has been used quite successfully. The chromatographic resolution of a pentacoordinate complex is of more than passing interest^{125,126}. In separations where aqueous solvents were used, D-quartz seemed to give the best results.

A very tantalizing sentence appeared in Stefanović and Janjić's note¹⁴⁰ on the paper chromatography of some ethylenediamine complexes of cobalt(III) in which they claimed to have separated, on paper, the optical antipodes of *cis*-[Co(en)₂Cl₂]⁺. However, in their subsequent paper¹⁴¹, which gives a more detailed account of their experimental procedures, no mention is made of the separation of optical isomers. Perhaps their first reports may have been based on impurities in their developed chromatograms.

While all separations to date (See Table 6) utilize a resolving agent in the stationary phase, a recent novel paper by Bosnich¹³ suggests that better results might be achieved by use of an optically active solvent. By use of L-2,3-butanediol, he observed that a solution of *cis*-[Co(en)₂Cl₂]⁺ slowly antiracemized to its active isomer. It is quite possible that such a solvent used in chromatography would preferentially desorb one enantiomer from silica gel, alumina, or cellulose, thereby achieving a rapid resolution.

G. SUMMARY

Recent applications of the various types of chromatography (column, thin-layer, paper, ion exchange, and electrophoresis) to the separation, preparation, identification, purification, and resolution of coordination complexes have been briefly reviewed. The techniques have also been used to study reaction kinetics and to determine the charges of such complexes. Despite the rapid progress in the last two decades, a few problems remain to be solved, such as the separation of geometric isomers of high ionic charge and more efficient resolution of optical isomers.

REFERENCES

- 1 T. V. ARDEN, F. H. BURSTALL, G. R. DAVIES, J. A. LEWIS AND R. P. LINSTEAD, *Nature*, **162** (1948) 691.
- 2 R. G. ASPERGER AND C. F. LIU, *Inorg. Chem.*, **6** (1967) 796.
- 3 K. BÄCHMANN AND H. GÖRISCH, *J. Chromatog.*, **23** (1966) 336.
- 4 F. BASOLO, M. LEDERER, L. OSSICINI AND K. H. STEPHEN, *Ric. Sci. Rend. Sez. A*, **2** (1962) 485.
- 5 F. BASOLO, M. LEDERER, L. OSSICINI AND K. H. STEPHEN, *J. Chromatog.*, **10** (1963) 262.
- 6 C. F. BELL AND D. R. ROSE, *Inorg. Chem.*, **7** (1968) 325.
- 7 J. P. BIBLER AND A. WOJCIK, *Inorg. Chem.*, **5** (1966) 889.
- 8 W. J. BIERMANN AND H. GESSER, *Anal. Chem.*, **32** (1960) 1525.
- 9 J. BJERRUM, A. JENSEN AND F. WOLDbye, *Nord. Kemikermøde*, (1956) 9.
- 10 E. BLASIUS AND B. A. BILAL, *J. Chromatog.*, **18** (1965) 134.
- 11 E. BLASIUS AND W. PREETZ, *Z. anorg. allgem. Chem.*, **335** (1965) 1.
- 12 E. BLASIUS AND W. PREETZ, *Z. anorg. allgem. Chem.*, **335** (1965) 16.
- 13 B. BOSNICH, *J. Am. Chem. Soc.*, **89** (1967) 6143.
- 14 E. H. BRAYE AND W. HÜBEL, *J. Organometal. Chem.*, **3** (1965) 25.
- 15 K. L. BRIDGES AND J. C. CHANG, *Inorg. Chem.*, **6** (1967) 619.
- 16 G. R. BRUBAKER, J. I. LEGG AND B. E. DOUGLAS, *J. Am. Chem. Soc.*, **88** (1966) 3446.
- 17 K. BUCHTELA AND M. LESIGANG-BUCHTELA, *Microchim. Acta*, (1967) 380.
- 18 D. A. BUCKINGHAM AND J. P. COLLMAN, *Inorg. Chem.*, **6** (1967) 1803.
- 19 D. A. BUCKINGHAM, J. P. COLLMAN, D. A. R. HAPPER AND L. G. MARZILLI, *J. Am. Chem. Soc.*, **89** (1967) 1082.
- 20 R. L. BURWELL, R. G. PEARSON, G. L. HALLER, P. B. TJOK AND S. P. CHOCK, *Inorg. Chem.*, **4** (1965) 1123.
- 21 D. I. BUSTIN, J. E. EARLEY AND D. PETRIDIC, *Inorg. Chem.*, **6** (1967) 2066.
- 22 H. H. CADY AND R. E. CONNICK, *J. Am. Chem. Soc.*, **80** (1958) 2646.
- 23 V. CARUNCHIO, G. GRASSINI-STRAZZA, G. ORTAGGI AND C. PADIGLIONE, *J. Inorg. Nucl. Chem.*, **27** (1965) 841.
- 24 V. CARUNCHIO, G. ILLUMINATI AND G. ORTAGGI, *Inorg. Chem.*, **6** (1967) 2168.
- 25 M. B. ČEPEL, S. R. NIKETIĆ, T. J. JANIĆ AND V. N. NIKOLIĆ, *Inorg. Chem.*, **6** (1967) 2063.
- 26 J. CHATT AND B. L. SHAW, *J. Chem. Soc.*, (1959) 705.
- 27 J. CHATT AND B. L. SHAW, *J. Chem. Soc.*, (1959) 4020.
- 27a R. F. CHILDERS, K. G. VANDER ZYL, D. A. HOUSE, R. G. HUGHES AND C. S. GARNER, *Inorg. Chem.*, **7** (1968) 749.
- 28 J. M. COLEMAN, A. WOJCIK, P. J. POLLICK AND L. F. DAHL, *Inorg. Chem.*, **6** (1967) 1236.
- 29 J. P. COLLMAN, R. P. BLAIR, A. L. SLADE AND R. L. MARSHALL, *Chem. Ind. (London)* (1962) 141.
- 30 J. P. COLLMAN AND C. T. SEARS, JR., *Inorg. Chem.*, **7** (1968) 27.
- 31 J. P. COLLMAN AND J. Y. SUN, *Inorg. Chem.*, **4** (1965) 1273.
- 32 R. E. CONNICK, in S. KIRSCHNER, ed., *Advances in the Chemistry of the Coordination Compounds*, Macmillan, New York, 1962, p. 15.
- 33 R. E. CONNICK AND D. A. FINE, *J. Am. Chem. Soc.*, **82** (1960) 4187.
- 34 R. E. CONNICK AND D. A. FINE, *J. Am. Chem. Soc.*, **83** (1961) 3414.
- 35 B. P. COTSORADIS AND R. D. ARCHER, *Inorg. Chem.*, **6** (1967) 800.
- 36 A. DANEELS, D. L. MASSART AND J. HOSTE, *J. Chromatog.*, **18** (1965) 144.
- 37 P. N. DIMOTAKIS AND C. D. MYRAT, *J. Inorg. Nucl. Chem.*, **21** (1961) 184.
- 38 L. F. DRUDING, *Anal. Chem.*, **35** (1963) 1582.
- 39 L. F. DRUDING AND R. B. HAGEL, *Anal. Chem.*, **38** (1966) 478.
- 40 L. F. DRUDING AND S. I. SHUPACK, *J. Chromatog.*, **24** (1966) 491.
- 41 D. L. DUGGER, J. H. STANTON, B. N. IRBY, B. L. MCCONNELL, W. W. CUMMINGS AND P. W. MAATMAN, *J. Phys. Chem.*, **68** (1964) 757.
- 42 F. P. DWYER, in S. KIRSCHNER, ed., *Advances in the Chemistry of the Coordination Compounds*, Macmillan, New York, 1962, p. 21.
- 43 K. J. EISENTRAUT AND R. E. SIEVERS, *J. Am. Chem. Soc.*, **87** (1965) 5254.

- 44 I. I. M. ELBEH, J. F. W. McOMIE AND F. H. POLLARD, *Discussions Faraday Soc.*, 7 (1949) 183.
- 45 M. F. FARONA AND A. WOJCICKI, *Inorg. Chem.*, 4 (1965) 1402.
- 46 O. D. FAUT, *11th Annual Report on Research under Sponsorship of the Petroleum Research Fund, Period Ending August 31, 1966*, American Chemical Society, Washington, 1966, p. 142.
- 47 R. C. FAY AND T. S. PIPER, *J. Am. Chem. Soc.*, 84 (1962) 2303.
- 48 R. C. FAY AND T. S. PIPER, *J. Am. Chem. Soc.*, 85 (1963) 500.
- 49 K. R. A. FEHRMANN AND C. S. GARNER, *J. Am. Chem. Soc.*, 83 (1961) 1276.
- 50 D. A. FINE, *Chloride Complexes of Ruthenium(III)*, Thesis, University of California, UCRL 9059, Office of Technical Services, U.S. Dept. of Commerce, Washington, 1960.
- 50a J. E. FINMOLT, K. CAULTON, K. KIMBALL AND E. UHLENHOPF, *Inorg. Chem.*, 7 (1968) 610.
- 50b W. A. FREEMAN AND C. F. LIU, *Inorg. Chem.*, 7 (1968) 764.
- 51 S. O. GRIM, D. A. WHEATLAND AND P. R. MCALLISTER, *Inorg. Chem.*, 7 (1968) 161.
- 51a S. K. HALL AND B. E. DOUGLAS, *Inorg. Chem.*, 7 (1968) 533.
- 52 H. G. HAMILTON AND M. D. ALEXANDER, *J. Am. Chem. Soc.*, 89 (1967) 5065.
- 53 E. HAYEK, *J. Chromatog.*, 2 (1959) 334 (see pp. 347 and 348 and references therein) (in German); *Chromatographic Rev.*, 2 (1960) 171 (in English).
- 54 R. D. HILL AND H. GESSER, *J. Gas Chromatog.*, 1 (Oct. 1963) 11.
- 55 J. T. HOUGEN, K. SCHUG AND E. L. KING, *J. Am. Chem. Soc.*, 79 (1957) 519.
- 56 M. HRANISAVLJEVIC-JAKOVljeVIC, I. PEJKOVIC-TADIC AND K. JAKOVljeVIC, *Thin-Layer Chromatog., Proc. Symp., Rome* (1963) 221.
- 57 M. HRANISAVLJEVIC-JAKOVljeVIC, I. PEJKOVIC-TADIC AND J. MILJKOVIC-STOJANOVIC, *Mikrochim. Ichnoanal. Acta*, (1965) 141.
- 58 T. M. HSEU, D. F. MARTIN AND T. MOELLER, *Inorg. Chem.*, 2 (1963) 587.
- 59 R. G. HUGHES AND C. S. GARNER, *Inorg. Chem.*, 7 (1968) 74.
- 60 M. A. JENNINGS AND A. WOJCICKI, *Inorg. Chem.*, 6 (1967) 1854.
- 61 A. JENSEN, J. BJERRUM AND F. WOLDBYE, *Acta Chem. Scand.*, 12 (1958) 1202.
- 62 E. JØRGENSEN AND J. BJERRUM, *Acta Chem. Scand.*, 12 (1958) 1047.
- 63 F. JURSIK, *J. Chromatog.*, 19 (1965) 448.
- 64 F. JURSIK, *J. Chromatog.*, 26 (1967) 339.
- 65 L. A. P. KANE-MAGUIRE AND T. E. MACDERMOTT, *Inorg. Chem.*, 7 (1968) 769.
- 66 G. B. KAUFFMAN, *Alfred Werner-Founder of Coordination Chemistry*, Springer-Verlag, Berlin, Heidelberg, New York, 1966.
- 67 G. B. KAUFFMAN, *Classics in Coordination Chemistry, Part I: The Selected Papers of Alfred Werner*, Dover, New York, 1968, pp. 155-184.
- 68 G. B. KAUFFMAN, unpublished work.
- 69 G. B. KAUFFMAN AND B. W. BENSON, *Inorg. Chem.*, 6 (1967) 411.
- 70 G. B. KAUFFMAN AND J. S. BLANK, *J. Chem. Educ.*, 37 (1960) 156 and references therein.
- 71 G. B. KAUFFMAN, R. P. PINNELL AND L. T. TAKAHASHI, *Inorg. Chem.*, 1 (1962) 544.
- 72 G. B. KAUFFMAN, J. H. TSAI, R. C. FAY AND C. K. JØRGENSEN, *Inorg. Chem.*, 2 (1963) 1233.
- 73 A. S. KERTES, *Anal. Chim. Acta*, 17 (1957) 470.
- 74 A. S. KERTES, *Anal. Chim. Acta*, 17 (1957) 570.
- 75 M. KILNER, F. A. HARTMAN AND A. WOJCICKI, *Inorg. Chem.*, 6 (1967) 406.
- 76 E. L. KING AND E. B. DISMUKES, *J. Am. Chem. Soc.*, 74 (1952) 1674.
- 77 E. L. KING AND R. R. WALTERS, *J. Am. Chem. Soc.*, 74 (1952) 4471.
- 78 E. L. KING, M. J. M. WOODS AND H. S. GATES, *J. Am. Chem. Soc.*, 80 (1958) 5015.
- 79 R. B. KING, *J. Am. Chem. Soc.*, 84 (1962) 2460.
- 80 R. B. KING AND M. B. BISNETTE, *Inorg. Chem.*, 4 (1965) 1663.
- 81 I. M. KOLTHOFF AND V. A. STENGER, *J. Phys. Chem.*, 38 (1934) 475.
- 82 T. B. KONUNOVA, *Uchz. Zap. Kishinevsk Gos. Univ.*, 68 (1964) 1.
- 83 K. A. KRAUS AND F. NELSON, in *Symposium on Ion Exchange and Chromatography in Analytical Chemistry*, Special Technical Publication No. 195, American Society for Testing Materials, 1958, p. 27; in W. J. HAMER, ed., *The Structure of Electrolytic Solutions*, Wiley New York, 1959, p. 340. The nearly classical papers by KRAUS *et al.*, appearing in *J. Am. Chem. Soc.* beginning in the early 1950's, are too numerous to be cited.
- 84 H. KREBS AND R. RASCHKE, *Naturwissenschaften*, 41 (1954) 63.
- 85 H. KREBS AND W. SCHUMACHER, *Z. anorg. allgem. Chem.*, 344 (1966) 187.

- 86 U. KRUEERKE, C. HOOZAND AND W. HÜBEL, *Chem. Ber.*, 94 (1961) 2817.
- 87 U. KRUEERKE, W. HÜBEL AND G. VANHEE, *Chem. Ber.*, 94 (1961) 2829.
- 88 E. KYUNO, *Nippon Kagaku Zasshi*, 80 (1959) 981.
- 89 E. KYUNO, *Nippon Kagaku Zasshi*, 80 (1959) 984.
- 90 E. KYUNO, *Nippon Kagaku Zasshi*, 81 (1960) 724.
- 91 E. KYUNO, *Nippon Kagaku Zasshi*, 81 (1960) 728.
- 92 E. KYUNO AND J. C. BAILAR, JR., *J. Am. Chem. Soc.*, 88 (1966) 1120.
- 93 M. LEDERER, *Anal. Chim. Acta*, 2 (1948) 261.
- 94 M. LEDERER, *J. Chromatog.*, 1 (1958) 314.
- 95 M. LEDERER, *J. Chromatog.*, 2 (1959) 209.
- 96 M. LEDERER, *Zhur. Neorg. Khim.*, 3 (1958) 1799.
- 97 J. I. LEGG AND D. W. COOKE, *Inorg. Chem.*, 4 (1965) 1576.
- 98 J. I. LEGG, D. W. COOKE AND B. E. DOUGLAS, *Inorg. Chem.*, 6 (1967) 700.
- 99 M. LINHARD, M. WEIGEL AND H. FLYGARE, *Z. anorg. allgem. Chem.*, 263 (1950) 233.
- 100 D. J. MACDONALD AND C. S. GARNER, *J. Am. Chem. Soc.*, 83 (1961) 4152.
- 101 D. J. MACDONALD AND C. S. GARNER, *Inorg. Chem.*, 1 (1962) 20.
- 102 A. G. MADDOCK AND A. B. J. B. TODESCO, *J. Inorg. Nucl. Chem.*, 26 (1964) 1535.
- 103 M. MAKI, *Japan Analyst*, 3 (1954) 393.
- 104 D. F. MARTIN AND K. RAMAIAH, *J. Inorg. Nucl. Chem.*, 27 (1965) 2027.
- 105 D. L. MASSART, G. SAINTE AND J. HOSTE, *Acta Chim. Acad. Sci. Hung.*, 52 (1967) 229.
- 106 N. MATSUOKA, J. HIDAKA AND Y. SHIMURA, *Bull. Chem. Soc. Japan*, 40 (1967) 1868.
- 107 E. E. MERCER AND W. A. MCALLISTER, *Inorg. Chem.*, 4 (1965) 414.
- 108 T. MOELLER AND E. GULYAS, *J. Inorg. Nucl. Chem.*, 5 (1958) 245.
- 109 T. MOELLER, E. GULYAS AND R. H. MARSHALL, *J. Inorg. Nucl. Chem.*, 9 (1959) 82.
- 110 M. MORI, M. SHIBATA AND J. AZAMI, *Nippon Kagaku Zasshi*, 76 (1955) 1003.
- 111 M. MORI, M. SHIBATA AND M. NANASAWA, *Bull. Chem. Soc. Japan*, 29 (1956) 947.
- 112 R. W. MOSHIER AND R. E. SIEVERS, *Gas Chromatography of Metal Chelates*, Pergamon, Oxford, 1965, and references therein.
- 113 H. NAGAI, *Kumamoto Journal of Science*, Series A, 4 (1960) 265.
- 114 H. NAGAI, *Nippon Kagaku Zasshi*, 80 (1959) 617.
- 115 K. NOACK, M. RUCH AND F. CALDERAZZO, *Inorg. Chem.*, 7 (1968) 345.
- 116 D. C. OLSON AND C. S. GARNER, *Inorg. Chem.*, 2 (1963) 558.
- 117 D. E. PENNINGTON AND A. HAIM, *Inorg. Chem.*, 6 (1967) 2138.
- 118 T. S. PIPER, *J. Am. Chem. Soc.*, 83 (1961) 3908.
- 119 F. H. POLLARD, A. J. BANISTER, W. J. GEARY AND G. NICKLESS, *J. Chromatog.*, 2 (1959) 372.
- 120 W. PREETZ, *Z. anorg. allgem. Chem.*, 348 (1966) 151.
- 121 W. PREETZ AND E. BLASIUS, *Z. anorg. allgem. Chem.*, 332 (1964) 140.
- 122 W. PREETZ AND H. L. PFEIFER, *Anal. Chim. Acta*, 38 (1966) 255.
- 123 H. J. PRICE AND H. TAUBE, *Inorg. Chem.*, 7 (1968) 1.
- 124 R. RALEA, D. GIURGIU AND S. FISEL, *Rev. Roumaine Chim.*, 9 (1964) 709.
- 125 K. RAMAIAH, F. E. ANDERSON AND D. F. MARTIN, *Inorg. Chem.*, 3 (1964) 296.
- 126 K. RAMAIAH AND D. F. MARTIN, *J. Inorg. Nucl. Chem.*, 27 (1965) 1663.
- 127 W. J. RANDALL AND D. F. MARTIN, *Abstracts of Papers Presented at the 147th Meeting, American Chemical Society, Philadelphia, April 6-10, 1964*, p. 10L.
- 128 L. H. REYERSON AND R. E. CLARK, *J. Phys. Chem.*, 40 (1936) 1055.
- 129 L. B. ROGERS, *J. Chem. Educ.*, 45 (1968) 7.
- 130 D. SCARGILL, C. E. LYON, N. R. LARGE AND J. M. FLETCHER, *J. Inorg. Nucl. Chem.*, 27 (1965) 161.
- 131 H.-H. SCHMIDTKE, *Z. physik. Chem.*, 34 (1962) 295.
- 132 G. M. SCHWAB AND K. JOCKERS, *Angew. Chem.*, 50 (1937) 546.
- 133 H. SEILER, C. BIEBRICHER AND H. ERLIENMEYER, *Helv. Chim. Acta*, 46 (1963) 2636.
- 134 D. N. SEN, private communication to GBK, April 26, 1967.
- 135 D. N. SEN AND P. UMAPATHY, *Indian J. Chem.*, 4 (1966) 455.
- 136 H. J. SENE, *J. Chromatog.*, 21 (1966) 363.
- 137 H. J. SENE, *J. Chromatog.*, 27 (1967) 331.
- 138 M. SHIBATA, H. NISHIKAWA AND Y. NISHIDA, *Inorg. Chem.*, 7 (1968) 9.

- 139 R. E. STEVERS, R. W. MOSHIER AND M. L. MORRIS, *Inorg. Chem.*, 1 (1962) 966.
- 140 G. STEFANOVIĆ AND T. JANJIĆ, *Anal. Chim. Acta*, 11 (1954) 550.
- 141 G. STEFANOVIĆ AND T. JANJIĆ, *Anal. Chim. Acta*, 19 (1958) 488.
- 142 L. M. VALLARINO AND E. G. SANTARELLO, *Gazz. chim. ital.*, 94 (1964) 252.
- 142a J. W. VAUGHN, O. J. STVAN AND V. E. MAGNUSON, *Inorg. Chem.*, 7 (1968) 736.
- 143 J. M. VEIGEL, *Inorg. Chem.*, 7 (1968) 69.
- 144 J. M. VEIGEL AND C. S. GARNER, *Inorg. Chem.*, 4 (1965) 1569.
- 145 M. R. VERMA AND P. K. GUPTA, *Current Sci. (India)*, 30 (1961) 10.
- 146 M. R. VERMA AND P. K. GUPTA, *Current Sci. (India)*, 31 (1962) 10.
- 147 M. R. VERMA AND J. RAI, *Current Sci. (India)*, 35 (1966) 478.
- 148 M. VOBECKY, V. D. NEFEDOR AND E. N. SINOTOVA, *Zh. Obshch. Khim.*, 33 (1963) 4023.
- 149 F. VYDRA AND V. MARKOVÁ, *Collection Czech. Chem. Commun.*, 30 (1965) 2382.
- 150 F. VYDRA AND V. MARKOVÁ, *Collection Czech. Chem. Commun.*, 31 (1966) 1398.
- 151 F. VYDRA AND V. MARKOVÁ, *J. Inorg. Nucl. Chem.*, 26 (1964) 1319.
- 152 E. WEISS, W. HÜBEL AND A. GERONDAL, *Chem. Ber.*, 95 (1962) 1186.
- 153 E. WEISS, W. HÜBEL, R. MERENYI, J. NIELSEN, A. GERONDAL AND R. VANNIEUWENHOVEN, *Chem. Ber.*, 95 (1962) 1155.
- 154 L. J. WILSON AND N. J. ROSE, *Abstracts of Papers Presented at the 155th Meeting, American Chemical Society, San Francisco, April 1-5, 1968*, Abstract M 84.
- 155 A. WOJCICKI AND M. F. FARONA, in V. GUTMANN, ed., *Proceedings of the 8th International Conference on Coordination Chemistry*, Springer-Verlag, Vienna, New York, 1964, p. 262.
- 156 F. WOLDBYE, *Acta Chem. Scand.*, 12 (1958) 1079.
- 157 I. Y. YAMAMOTO, A. NAKAHARA AND R. TSUCHIDA, *J. Chem. Soc. Japan*, 75 (1954) 232.
- 158 H. YONEDA, *J. Chem. Soc. Japan*, 40 (1967) 2442.