

COORDINATION COMPOUNDS OF MOLYBDENUM*

P. C. H. MITCHELL

Department of Chemistry, The University, Reading (Great Britain)

(Received 4 April 1966)

CONTENTS

- A. Introduction
- B. General Survey
- C. Oxidation States -2 to +1
- D. Oxidation State +2
- E. Oxidation States +3 to +6:
 - (i). Binary halides and their derivatives
 - (ii). Complex halides
 - (iii). Complexes with inorganic oxygen-donors
 - (iv). Complexes with inorganic nitrogen-donors
 - (v). Cyanide complexes
 - (vi). Complexes with organic oxygen-donors
 - (vii). Complexes with organic nitrogen-, phosphorus- and arsenic-donors and with "oxine" and EDTA
 - (viii). Complexes with sulphur-donors

A. INTRODUCTION

In recent years the chemistry of molybdenum compounds has attracted increasing attention because of their intrinsic chemical interest and their industrial and biological importance. Molybdenum exhibits a wide range of oxidation states and stereochemistries and forms complexes with most donor atoms. Individual molybdenum compounds or groups of compounds (e.g., molybdenum carbonyls and oxomolybdenum(V) complexes) have been important in the development of theories of bonding in co-ordination compounds and in the interpretation of their magnetic

* (Abbreviations: Ph, C_6H_5 ; R, alkyl or aryl; tu, thiourea, $(NH_2)_2CS$; digly, diglyme, diethyleneglycol dimethyl ether, $(CH_2OCH_2CH_2)_2O$; py, pyridine; dien, diethylenetriamine, $NH_2CH_2CH_2NHCH_2CH_2NH_2$; diphos, 1,2-bis(diphenylphosphino)ethane, $Ph_2PCH_2CH_2PPh_2$; diars, diarsine, *o*-phenylenebis(dimethylarsine), *o*- $C_6H_4(AsMe_2)_2$; bipy, 2,2'-bipyridyl.)

and spectroscopic properties. Work on molybdenum compounds has been stimulated by current interest in the chemistry of the earlier-transition metals, by the wider application of techniques for handling highly reactive, air- and moisture-sensitive compounds¹, by recognition that molybdenum is essential for the activity of certain enzymes, e.g., xanthine oxidase² and nitrate reductase³, and by the use of molybdenum compounds (e.g., molybdenum disulphide) as anti-wear additives and catalysts in the petroleum industry⁴.

This article surveys the co-ordination chemistry of molybdenum with particular reference to recent work on well-characterised compounds. Oxo-species of molybdenum(V) and molybdenum(VI) have recently been reviewed and will not be discussed in detail here⁵. The solution chemistry of molybdenum is also excluded. (Of particular interest is the work of Haight *et al.* on the role of molybdenum(IV) in molybdate catalysed reactions⁶.) No attempt has been made to include all molybdenum compounds mentioned in the literature but it is felt that the major areas of current interest in molybdenum co-ordination chemistry have been covered. There are no recent comprehensive accounts of molybdenum chemistry. Useful, but uncritical accounts of earlier work are available⁷.

B. GENERAL SURVEY

Molybdenum is the second member of Group VIA of the Periodic Table and the fourth member of the second transition series. The valency-shell electronic configuration of the free atom in the ground state is $4d^5 5s^1$. Molybdenum differs from chromium, the first member of Group VIA, in two important respects: molybdenum exhibits more stable oxidation states than chromium and the higher oxidation states of molybdenum are more stable with respect to reduction and the lower oxidation states are less stable with respect to oxidation than those of chromium. This trend reflects the smaller ionisation potentials⁸ for *d*-electrons and the larger ionic and covalent radii^{7c} of molybdenum.

(i) Oxidation states

In its compounds molybdenum exhibits all oxidation states from -2 to +6. The lowest oxidation states, -2 to +1, occur in complexes with π -acceptor ligands, mainly carbon monoxide, cyclopentadiene and related compounds, nitric oxide, and unsaturated nitrogen-, phosphorus-, and arsenic-donor ligands. In these oxidation states molybdenum behaves as a class-B acceptor.

The best known compounds of molybdenum(II) are those containing the polynuclear ion $\text{Mo}_6\text{Cl}_8^{4+}$. Recently six- and seven-co-ordinate mononuclear complexes with π -acceptor ligands have been prepared.

In oxidation states +3 to +6 molybdenum forms a large number of com-

plexes with nitrogen- and oxygen-donor ligands and with the lighter halogens. Complexes with sulphur-ligands are fairly common, but there are very few complexes with phosphorus- and arsenic-donors. In its higher oxidation states molybdenum behaves as a class-A acceptor.

The chemistry of the +6 and +5 oxidation states is dominated by oxo-molybdenum species (MoO^{4+} , MoO_2^{2+} , MoO_3 , $\text{Mo}_2\text{O}_5^{2+}$; MoO^{3+} , $\text{Mo}_2\text{O}_3^{4+}$, $\text{Mo}_2\text{O}_4^{2+}$), with one or more oxygen atoms as terminal ligands (i.e., bonded to only one molybdenum atom) or as bridging atoms (i.e., bonded to two molybdenum atoms). An important feature is π -donation from oxygen to the metal giving a multiple bond⁵. This type of donor π -bonding occurs less readily as the formal oxidation state of the metal decreases and only a few, poorly characterised oxo-species of molybdenum(IV) and molybdenum(III) are known.

(ii) Co-ordination numbers and structures

The most-common co-ordination number is six giving octahedral and, in certain sulphur complexes (section E.(viii)), trigonal prismatic stereochemistries. In many compounds the metal attains six co-ordination through dimerisation or polymerisation. Co-ordination number four is not known and the only well-established example of co-ordination number five is for MoCl_5 in the vapour state. Co-ordination numbers greater than six are more common.

Polynuclear complexes are known for most oxidation states and many contain one or more Mo-Mo bonds.

(iii) Magnetic properties

Data for molybdenum complexes are included in a recent review article⁹. It should be noted that because of the large spin-orbit coupling constants (λ) of molybdenum(V) and molybdenum(IV), magnetic moments of their mononuclear octahedral complexes are expected to be much less than the spin-only values and to vary with temperature (ca. 1.0 and 1.8 B.M. respectively at 300 °K). The observed magnetic moments at 300 °K are 1.5–1.8 B.M. for Mo^{V} and 2.0–2.5 B.M. for Mo^{IV} . The deviations from the spin-only moments are less than expected because the spin-orbit coupling constants are reduced from the free-ion values through complex formation and because of the effect of asymmetric ligand fields in destroying orbital degeneracy. For octahedral low-spin molybdenum(II) (t_{2g}^4) and molybdenum(I) (t_{2g}^3) similar considerations apply except that the magnetic moments should be greater than the spin-only values (3.3 B.M. for Mo^{II} and 2.3 B.M. for Mo^{I} at 300 °K.)⁹. Observed and calculated magnetic moments for octahedral molybdenum(III) are close to the spin-only value. Co-ordination numbers seven and eight give ground states which appear to be orbitally non-degenerate and the complexes are either diamagnetic (Mo^{II} , Mo^{IV}) or have moments

close to the spin-only value for one unpaired electron (Mo^I , Mo^{III} , Mo^V). Complexes of molybdenum(O) are diamagnetic. Bi- and polynuclear complexes usually have magnetic moments corresponding to less than 1.0 B.M. per Mo atom. Temperature-independent paramagnetism has been observed for spin-paired seven-co-ordinate molybdenum(II) and for molybdenum(IV). The magnetic properties of molybdenum complexes are much more complicated than is sometimes realised and caution is necessary in making deductions about structures, spin-states and even oxidation states.

(iv) *Electronic spectra*

Crystal-field ($d-d$) transitions are observed with complexes of the more electronegative ligands (halogens, saturated oxygen- and nitrogen-donors); but with ligands of low electronegativity, intense charge-transfer bands may cover the crystal-field transitions in the visible region. For a series of molybdenum(V) complexes with organic ligands the energies of the charge-transfer transitions decrease as the electron-donor character of the ligands increases suggesting ligand-to-metal charge transfer¹⁰, whilst in molybdenum(III) complexes the charge-transfer transitions are thought to be from metal-to-ligand¹¹.

Spectrochemical and nephelauxetic series based on the molybdenum(III) ion, have been obtained; these agree with the series for other metal ions^{11,12}.

(v) *Classification of complexes*

A strict classification of molybdenum complexes on the basis of donor atoms or oxidation states is not possible without extensive cross-referencing. Many complexes contain more than one type of ligand and their chemical properties may involve more than one oxidation state. Because of this, substituted carbonyls are grouped together (section C) and so are complexes derived from the molybdenum halides (section E.i). Other complexes are classified on the basis of ligand donor-atoms.

C. OXIDATION STATES -2 TO +1

Many complexes of molybdenum in these oxidation states are derivatives of molybdenum hexacarbonyl, $\text{Mo}(\text{CO})_6$, but a few are known which do not contain carbon monoxide.

(i) *Molybdenum carbonyls and related compounds*

The chemistry of $\text{Mo}(\text{CO})_6$ has been very extensively studied¹³. Up to three

CO groups may be replaced by monodentate ligands and up to four by certain bidentate ligands, but only rarely are all the CO groups replaced in substitution reactions. Complexes of organic ligands^{6c} which have attracted much attention include π -complexes of cyclopentadiene (Cp), e.g., $\text{Cp}_2\text{Mo}_2(\text{CO})_6$, $[\text{CpMo}(\text{CO})_3]^-$, Cp_2MoCl_2 ; cycloheptatriene (C_7H_8), e.g., $(\text{C}_7\text{H}_8)\text{Mo}(\text{CO})_3$; and benzene, e.g., $(\text{C}_6\text{H}_6)\text{Mo}(\text{CO})_3$, $\text{Mo}(\text{C}_6\text{H}_6)_2$, $[\text{Mo}(\text{C}_6\text{H}_6)_2]^+$. Carbonyl anions are also known, e.g., $[\text{Mo}(\text{CO})_4]^{2-}$ and $[\text{Mo}_2(\text{CO})_{10}]^{2-}$.

Interest in molybdenum carbonyls and organometallic compounds has been concentrated in two areas: (a) structural and theoretical studies with particular reference to π -bonding in the hexacarbonyl and substituted carbonyls; (b) substitution and oxidation reactions giving new types of compound, compounds with unusual structures, and compounds of uncommon oxidation states.

(a) *Structures and bonding.* That $\text{Mo}(\text{CO})_6$ has a perfectly octahedral structure with linear Mo—C—O bonds has been shown by infrared, Raman, electron-diffraction, and X-ray measurements^{13b}. The Mo—CO bond is described in terms of a σ -bond formed from the $d(e_g)$, s , and p orbitals of molybdenum and the (sp) -lone pair of carbon, and π -bonds formed by donation from the $d(t_{2g})$ -orbitals of molybdenum to the anti-bonding π -orbitals of CO. The Mo—C bond has an order of more than one and the C—O bond an order of less than three. The order of the C—O bond decreases with increasing π -donation from the metal¹⁴. A molecular orbital treatment of $\text{Mo}(\text{CO})_6$ has been given and used to interpret its electronic spectrum¹⁵.


Replacement of CO by weaker π -acceptors causes a decrease in C—O stretching frequencies because of increased π -bonding between the metal and the remaining CO groups. The lowering is greatest in complexes of nitrogen- and oxygen-donors which lack π -acceptor orbitals, e.g.,^{16a}, $(\text{diglyme})\text{Mo}(\text{CO})_3$, and^{16b} $(\text{dien})\text{Mo}(\text{CO})_3$ and least in complexes of ligands which are good π -acceptors, e.g., in^{16b} $\text{cis-}(\text{PCl}_3)_3\text{Mo}(\text{CO})_3$. Measurements on a large number of tricarbonyls give the following order of decreasing C—O stretching frequencies (interpreted as decreasing π -acceptor ability of the ligands): $\text{PCl}_3 \sim \text{AsCl}_3 \sim \text{SbCl}_3 > \text{Ph}_3\text{P} \sim \text{Ph}_3\text{As} \sim \text{Ph}_3\text{Sb} \gg \text{R}_2\text{S} > \text{tu} \sim \text{digly} > \text{py} \sim \text{dien}$ ^{13b, 14, 16}.

For some tetracarbonyls, $\text{Mo}(\text{CO})_4\text{A}$, the order is $\text{diphos} > \text{diars} > \text{bipy}$ ¹⁷.

(b) *Some chemical reactions of molybdenum carbonyls.*

Substitution reactions. Substituted molybdenum carbonyls may be obtained directly from $\text{Mo}(\text{CO})_6$ or by displacement of another substituent, e.g., from $\text{C}_7\text{H}_8\text{Mo}(\text{CO})_3$. As CO groups are removed further substitution becomes more difficult. Thus diarsine displaces¹⁸ two carbonyl groups at 150° and a further two at 200°. This effect is attributed to increased π -bonding between molybdenum and carbon monoxide as the number of CO groups decreases. In agreement, bipyridyl will displace only two CO groups from $\text{Mo}(\text{CO})_6$ but diphos and diars, which are

better π -acceptors, will displace four CO groups¹⁷. A further CO group is displaced from $\text{Mo}(\text{CO})_4(\text{bipy})$ by Ph_3P , Ph_2S or pyridine¹⁹.



Displacement of an *axial*-CO group occurs because the bipyridyl causes more enhancement of metal-CO π -bonding in the plane than out of the plane.

Oxidation by halogens. Reactions in which some substituted molybdenum carbonyls are oxidised by halogens are summarised in Table I. Tetra- and tricarbonyls generally lose carbon monoxide to give dicarbonyls; but dicarbonyls react without further loss of carbon monoxide^{16f, 17, 20-23}. Similar reactions occur with substituted chromium and tungsten carbonyls, the ease of displacement of CO increasing in the order^{21, 23} $\text{W} < \text{Mo} < \text{Cr}$.

TABLE I

OXIDATION OF SUBSTITUTED MOLYBDENUM CARBONYLS BY HALOGENS

Starting compound ^a	Reagents	Products	Ref.
$\text{Mo}(\text{CO})_3(\text{bipy})$	Br_2	$\text{Mo}^{\text{II}}(\text{CO})_3(\text{bipy})\text{Br}_2$	17
$\text{Mo}(\text{CO})_4(\text{diphos})$	Br_2 and I_2	$\text{Mo}^{\text{II}}(\text{CO})_3(\text{diphos})\text{X}_2$	20
$\text{Mo}(\text{CO})_4(\text{diars})$	Br_2 , I_2	$\text{Mo}^{\text{II}}(\text{CO})_3(\text{diars})\text{X}_2$	21
	I_2 , hot CCl_4	$\text{Mo}^{\text{II}}(\text{CO})_3(\text{diars})\text{I}_2$	22
	I_2 , hot CHCl_3	$\text{Mo}^{\text{III}}(\text{CO})_2(\text{diars})\text{I}_3$	22
$\text{Mo}(\text{CO})_4(\text{dithia})$	Br_2 and I_2	$\text{Mo}^{\text{III}}(\text{CO})_3(\text{dithia})\text{X}_2$	16f
$\text{Mo}(\text{CO})_3(\nu\text{-triars})^b$	Br_2 and I_2	$[\text{Mo}^{\text{II}}(\text{CO})_3(\nu\text{-triars})\text{X}]\text{X}$ and some $\text{Mo}^{\text{II}}(\text{CO})_2(\nu\text{-triars})\text{X}_2$	23
$[\text{Mo}(\text{CO})_3(\nu\text{-triars})\text{Br}]\text{BPh}_4$	LiCl in <i>n</i> -propanol	$\text{Mo}^{\text{II}}(\text{CO})_2(\nu\text{-triars})\text{Cl}_2$	23
$\text{Mo}(\text{CO})_4(\text{diphos})^c$	I_2	$[\text{Mo}^{\text{I}}(\text{CO})_2(\text{diphos})_2]\text{I}_3$	20
$\text{Mo}(\text{CO})_3(\text{diars})_2$	Br_2 and I_2	$[\text{Mo}^{\text{II}}(\text{CO})_2(\text{diars})_2\text{X}]\text{X}$	21

^a dithia = 2,5-dithiahexane; ν -triars = 1,1,1-tris(dimethylarsinomethyl)ethane. For other ligand abbreviations see p. 315.

^b The proportion of the dicarbonyl compound increases with increasing temperature. Similar reactions occur with $\text{Mo}(\text{CO})_3(1\text{-triars})$ and $\text{Mo}(\text{CO})_2(o\text{-triars})$ (1-triars = methyl-bis(dimethylarsino-3-propyl)arsine; *o*-triars = bis(*o*-dimethylarsinophenyl)methylarsine).






^c The anion is the tri-iodide ion, I_3^- .

An important feature of these reactions is the formation of complexes of molybdenum(II) formulated, on the basis of physical measurements (conductivities in nitrobenzene, magnetic moments, and infrared spectra), with co-ordination number seven. A pentagonal bipyramidal structure with axial halogens has been proposed²¹ for $\text{Mo}(\text{CO})_3(\text{diars})\text{I}_2$. The seven-co-ordinate complexes are

spin-paired but have temperature-independent paramagnetic moments from 0.4 to 1.1 B.M.²³

The existence of complexes of molybdenum(II) and (III) with co-ordination number seven has been rationalised with Nyholm's "nine-orbital rule" according to which transition metals in their complexes with polarisable ligands tend to utilise all nine valency orbitals (i.e., five $(n-1)d$, one ns , three np) for σ -bonding pairs, non-bonding pairs, and single electrons. The co-ordination number is limited by the number of σ -acceptor orbitals available²¹. In the spin-paired complexes of molybdenum(II) there are two non-bonding electron pairs, leaving seven σ -acceptor orbitals (d^3sp^3). Similarly molybdenum(III) (d^3 with one unpaired electron) has seven acceptor orbitals and so forms, e.g., $\text{Mo}(\text{CO})_2(\text{diars})\text{I}_3$ ($\mu = 1.40$ B.M.)²². However, low-spin molybdenum(I) (d^5) and molybdenum(II) (with two unpaired electrons) have only six acceptor orbitals and so form six-co-ordinate complexes, e.g., $[\text{Mo}^{\text{I}}(\text{CO})_2(\text{diphos})_2]\text{I}_3$ ($\mu = 1.66$ B.M.)²⁰ and $\text{Mo}^{\text{II}}(\text{CO})_2(\text{diars})\text{I}_2$ ($\mu = 1.98$ B.M.)²².


Under vigorous conditions (e.g., excess halogen) carbon monoxide may be displaced completely:



The compounds $\text{Mo}(\text{CO})_3(\text{Ph}_3\text{PO})_3$, *cis*- and *trans*- $\text{Mo}(\text{CO})_4(\text{PPh}_3)_2$, $\text{Mo}(\text{CO})_4(\text{diphos})_2$ react similarly; analogous oxomolybdenum(V) complexes are also obtained by chlorine oxidation²⁶.

(c) *Compounds with Mo-Mo bonds.* There is a tendency for early-transition elements in oxidation states with unpaired electrons to dimerise through the formation of metal-metal bonds. This tendency can be rationalised by the inert-gas rule which is a special case of the nine-orbital rule²¹. Examples of low-valent molybdenum compounds with metal-metal bonds are the anion $[\text{Mo}_2(\text{CO})_{10}]^{2-}$ which is diamagnetic²⁷ and isoelectronic with $[\text{Mn}_2(\text{CO})_{10}]$ and the compound $[(\text{Cp})_2\text{Mo}_2(\text{CO})_6]$ obtained by the reaction of dicyclopentadiene²⁸ or cyclopentadienyl sodium²⁹ with $\text{Mo}(\text{CO})_6$. The compound contains molybdenum formally in oxidation state +1, but is diamagnetic. The presence of the Mo-Mo bond has been shown by X-ray analysis, the molecule consisting of two $[\text{CpMo}(\text{CO})_3]$ units


united by a Mo-Mo bond of 3.22 Å³⁰. Compounds containing mixed Mo-metal bonds have also been prepared, e.g., Cp(CO)₃MoW(CO)₃Cp and Cp(CO)₃MoFe(CO)₂Cp from CpMo(CO)₃Na and CpW(CO)₃I and CpFe(CO)₂I respectively^{31,32}. The metal-metal bond in (Cp)₂Mo₂(CO)₆ is fairly readily cleaved giving diamagnetic, bridged binuclear complexes:



(ii) Nitrosyl complexes

Rather few nitrosyl complexes of molybdenum are known; among these are the complexes³⁵ [CpMo(CO)₂NO] and K₄[Mo(CN)₅NO] (section E.(v)). Recently the compounds [Mo(NO)₂Cl₂] and [Mo(NO)₂(MNT)₂] have been prepared^{36,37}. (MNT is the maleonitriledithiolate anion, ⁻S-C(CN)=C(CN)-S⁻).

Mo(NO)₂Cl₂ is obtained as a dark-green solid from the reaction of Mo(CO)₆ with NOCl in dichloromethane. According to the infrared spectrum (N-O stretching frequency) the NO groups are *cis* and co-ordinated as NO⁺. A polymeric structure with kinked chains is suggested:



Octahedral complexes [Mo(NO)₂Cl₂L₂] are formed with L = Ph₃P, Ph₃As, pyridine, *p*-toluidine, cyclohexylamine, and Cl (from Ph₄AsCl) and, according to the infrared spectra, the NO⁺ groups are *cis*³⁶ to one another.

With Na₂MNT in methanol, Mo(NO)₂Cl₂ forms a deep-green complex isolated as (Ph₄P)₂[Mo(NO)₂(MNT)₂]. The infrared spectrum is in accordance with *cis*-NO⁺ groups and also suggests that in this compound MNT²⁻ functions as a π-acceptor (in contrast to its behaviour with high oxidation states with which it is a good donor, see section E.(viii))³⁷.

(iii) *Complexes of molybdenum(O) with nitrogen- and phosphorus-donors.*

Unstable, air-sensitive complexes in which molybdenum(O) is fully co-ordinated with N- and P-donors have been prepared, e.g., $[\text{Mo}(\text{bipy})_3]$ from $[\text{Mo}(\text{bipy})_3]\text{Cl}_3$ and Li_2bipy in THF³⁸, $[\text{Mo}(\text{tripy})_2]$ from $\text{Mo}(\text{CO})_6$ and 2,2',2''-tripyriddy³⁹, $\text{Mo}(\text{PP})_3$ from $[(\text{C}_6\text{H}_5)_2\text{Mo}]$ and $o\text{-C}_6\text{H}_4(\text{PR}_2)_2$ and also by reducing a mixture of MoCl_5 and the diphosphine in THF with lithium aluminium hydride⁴⁰, and $\text{Mo}(\text{PF}_3)_6$ from³⁹ $(\text{C}_6\text{H}_5)_2\text{Mo}$ and PF_3 . The existence of these complexes depends on the ability of the ligands to function as π -acceptors.

D. COMPLEXES OF MOLYBDENUM(II)

Molybdenum(II) may be stabilised in two ways: in mononuclear complexes with π -bonding ligands and through metal-metal bonding in polynuclear complexes with the more electronegative ligands. Unlike chromium(II), molybdenum(II), does not form a mono-nuclear aquo ion or complexes derived from it.

(i) *Mononuclear complexes*

Complexes of molybdenum(II) in which the metal is seven-co-ordinate have been described in Section C in connection with halogen oxidation of substituted molybdenum carbonyls.

Six-co-ordinate, low-spin complexes of molybdenum(II), $\text{Mo}(\text{diars})_2\text{X}_2$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) ($\mu = 2.8\text{--}2.9$ B.M.), are obtained in a slow reaction (24 hours) between diars and an aqueous solution of molybdenum(III) and the appropriate halogen acid. The reaction appears to involve disproportionation⁴¹ of molybdenum(III) to molybdenum(II) and (V).

(ii) *Polynuclear halogen complexes*

The halides of molybdenum(II), MoX_2 ($\text{X} = \text{Cl}, \text{Br}, \text{I}$), are polynuclear. X-ray analysis of the compounds $[(\text{Mo}_6\text{Cl}_8)(\text{OH})_4(\text{H}_2\text{O})_2] \cdot 12\text{H}_2\text{O}$ and $[(\text{Mo}_6\text{Cl}_8)\text{Cl}_4(\text{H}_2\text{O})_2] \cdot 6\text{H}_2\text{O}$ has shown that the basic structural unit is the octa- μ_3 -chlorohexamolybdenum(II) ion, $[(\text{Mo}_6\text{Cl}_8)]^{4+}$, shown in Fig. 1(a). Eight chlorines are at the corners of a cube and six molybdenums at the face centres. Each molybdenum is in approximately square co-ordination with four chlorines and is also bonded to four other molybdenums 2.64 Å distant⁴². The metal atoms form an octahedral cluster⁴³.

The bonding in the $\text{Mo}_6\text{Cl}_8^{4+}$ cluster has been discussed in valence-bond^{44, 45} and in molecular-orbital terms^{43, 46}. The treatment most readily comprehensible in chemical terms is that due to Cotton and Haas⁴³. They assume that




Fig. 1. Structures of (a) $[\text{Mo}_6\text{Cl}_8]^{4+}$ and (b) $[\text{Mo}(\text{O}_2\text{CCH}_3)_6]^{2-}$. Closed circles represent molybdenum atoms. In (a) broken lines define the octahedral cluster of molybdenum atoms; Mo-Cl bonds are not shown. In (b) the oxygen atoms of carboxyl groups, Mo-O bonds, and the Mo-Mo bond are shown.

each molybdenum atom uses a d_{xy} -orbital, an s -orbital and the p_x - and p_y -orbitals to bind the four co-planar chlorines. The metal-metal bonding, using d_{z^2} , d_{xz} , d_{yz} , and $d_{x^2-y^2}$, is described in terms of molecular orbitals for the octahedral cluster. It is found that there are just sufficient strongly-bonding orbitals (A_{1g} , T_{1u} , T_{2g} , E_g , T_{2u}) to accommodate the 24 electrons of the cluster, thereby accounting for the diamagnetism and the stability of the cluster. Their treatment also shows the presence on each molybdenum of an empty, centrifugally-directed σ -orbital and so accounts for complexes of the type $(\text{Mo}_6\text{Cl}_8)\text{L}_6$.

The structures of the dihalides, Mo_6X_{12} ($\text{X} = \text{Cl}, \text{Br}, \text{I}$), have not been determined but they are thought to be polymeric, with $\text{Mo}_6\text{X}_8^{4+}$ clusters joined through halogen bridges⁴⁷.

The $\text{Mo}_6\text{Cl}_8^{4+}$ cluster will co-ordinate up to six ligands, one for each molybdenum, and a large number of complexes of the cluster are known. Examples include MX_4L_2 (where $\text{M} = \text{Mo}_6\text{Cl}_8$; $\text{X} = \text{Cl}, \text{OH}$; $\text{L} = \text{py}, \text{NEt}_3, \text{H}_2\text{O}, \text{EtOH}$), MX_6^{2-} ($\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{NCS}$)^{44, 47}, $[\text{Mo}_6\text{X}_8\text{Y}_6]^{2-}$ ($\text{X} = \text{Br}, \text{I}$; $\text{Y} = \text{Cl}, \text{Br}, \text{I}, \text{OH}$), $\text{Mo}_6\text{X}_8\text{Y}_4\text{py}_2$ ($\text{Y} = \text{Cl}, \text{Br}$)⁴⁷, $\text{MCl}_4(\text{Ph}_3\text{EO})_2$ ($\text{E} = \text{P}, \text{As}$)⁴⁸, and MX_4L_2 ($\text{L} = \text{dimethylsulphoxide}, \text{dimethylformamide}, \text{methylsulphonate}, \text{MeSO}_3^-$). According to infrared studies the ligands in the last complex are bonded to $\text{Mo}_6\text{Cl}_8^{4+}$ through oxygen⁴⁹.

There have been some mechanistic studies of the reactions of chloromolybdenum(II) complexes. The rates of chloride exchange (using ^{36}Cl) and bromide substitution in $[\text{Mo}_6\text{Cl}_8]\text{Cl}_6^{2-}$ in water and moist ethanol are the same for both halides and independent of their concentrations suggesting that the reactions proceed through a common rate-determining step, probably aquation. No more than six chlorides may be replaced and there is no reaction with chlorides of the $\text{Mo}_6\text{Cl}_8^{4+}$ cluster⁴⁴.

Halogen in the ions $[\text{Mo}_6\text{X}_8(\text{OH})_6]^{2+}$ ($\text{X} = \text{Cl}^{50}, \text{Br}^{51}$), may be replaced by hydroxide giving species $[\text{Mo}_6\text{X}_{8-n}(\text{OH})_n](\text{OH})_6^{2-}$ ($n = 1, 2, 3$, etc.). There is evidence for a bimolecular rate-determining step, the rate being dependent on the

molybdenum(II) and OH^- concentrations. A bimolecular process is reasonable since the chloro- and bromo- Mo^{II} structures are rigid, and a unimolecular process, involving distortion of the cluster to an activated complex of lower co-ordination number, would involve a very high activation energy. The rate is much less for the bromo complex than for the chloro complex suggesting the operation of steric factors: there is "more room" for $\text{S}_{\text{N}}2$ attack in the chloro compound than in the bromo compound⁵¹.

(iii) *Binuclear carboxylates*

The compounds $[\text{Mo}(\text{O}_2\text{CR})_2]_2$ have been obtained as yellow, diamagnetic solids by heating $\text{Mo}(\text{CO})_6$ with various carboxylic acids. They are stable in the absence of air but give molybdenum blue in moist air. With pyridine the acetate gives⁵² an adduct $[\text{Mo}(\text{O}_2\text{CMe})_2\text{py}]_2$.

X-ray analysis of molybdenum(II) acetate⁵³ has shown that the structure is similar to that of cupric acetate with a short Mo-Mo bond of 2.11 Å (Fig. 1(b)).

E. OXIDATION STATES +3 TO +6

(i) *Binary halides and their derivatives*

(a) *Preparations.* The binary halides of molybdenum are listed in Table II which also gives preparative reactions and some of their properties⁵⁴⁻⁷⁶. Of particular interest is the reduction of MoCl_5 with benzene or chlorobenzene which has made MoCl_4 readily obtainable⁶⁷. The best procedure for the preparation of MoF_5 is said⁵⁷ to be the reaction between MoF_6 and PF_3 .

(b) *Structures and physical properties.* MoF_6 is octahedral with $\text{Mo-F} = 1.840$ Å, as indicated by an electron diffraction study of the vapour. The infrared and Raman spectra are also consistent with an octahedral structure⁵⁵. MoF_5 is tetrameric in the solid (X-ray diffraction)⁵⁶. Each molybdenum is in approximately octahedral co-ordination with four terminal fluorines (two at 1.85 and two at 1.70 Å) and two bridging fluorines (at 2.06 Å) (Fig. 2(a)). Molybdenum is also in octahedral co-ordination in solid MoF_3 . According to Gutmann and Jack⁶⁰, X-ray analysis of MoF_3 indicates that the crystal has the cubic ReO_3 structure but a re-determination of the structure suggests a rhombohedral VF_3 structure⁵⁸. It is possible that the compound studied by Gutmann and Jack was, in fact, an oxo-fluoride⁵⁸.

An electron diffraction study showed that, in the vapour, MoCl_5 is trigonal bipyramidal with all bond lengths equal (2.27 Å)⁶³. Gillespie⁷⁷ considers that a re-investigation of the structure would be worthwhile since in most trigonal bi-

TABLE II
BINARY HALIDES

Compound	Preparation	Properties and structure ^a (where known)	Ref.
MoF ₆	Mo + F ₂	octahedral ^c	54, 55
MoF ₅	Mo(CO) ₆ + F ₂	tetrameric (octahedral)	25, 56
	Mo(CO) ₆ + MoF ₆		56
	MoF ₆ + Mo		56
	MoF ₆ + PF ₃		57
MoF ₄	Involatile residue in preparation of MoF ₅		25, 56
MoF ₃	MoF ₅ + Mo	octahedral	58, 60
	MoBr ₃ + HF		59
MoCl ₅	Mo + Cl ₂	trig. bipyramid vapour	61, 63
	MoS ₂ + CCl ₄	dimeric (octahedral) solid; $\mu = 1.67$	62, 64, 65
MoCl ₄	MoO ₃ + CCl ₄	$\mu = 0.93$	66, 67
	MoCl ₃ + C ₆ H ₆ or C ₆ H ₅ Cl		67
MoCl ₃	MoCl ₅ + H ₂	octahedral; $\mu = 0.67$	68, 69, 70
MoBr ₄	Mo(CO) ₆ + Br ₂	$\mu = 1.28$	24, 71
	MoBr ₃ + Br ₂		71
MoBr ₃	Mo + Br ₂	octahedral; $\mu = 1.24$	70, 72, 73
MoI ₃	Mo + I ₂	$\mu = 1.4$	74
	MoCl ₅ + HI	or 1.85	75, 76
	Mo(CO) ₆ + I ₂		76

^a Magnetic moments (μ) in Bohr magnetons at ca. 20°.

pyramidal molecules the axial bonds are longer than the equatorial bonds. In the crystal, MoCl₅ is dimeric (X-ray diffraction) (Fig. 2(b))⁶⁴. There are four terminal chlorines at 2.24 Å, two bridging chlorines at 2.53 Å, and a Mo-Mo bond of 3.84 Å. The molybdenum is in approximately octahedral co-ordination, the shape of the MoCl₆ octahedron being similar to that of MoF₆ in MoF₅. As in MoF₅ the bridging bonds are longer than the terminal bonds. MoCl₅ is monomeric in




Fig. 2. Structures (not to scale) of⁵⁴ (a) [MoF₆]₄ and⁶⁴ (b) [MoCl₅]₂. Closed circles represent molybdenum atoms.

benzene⁷⁸ and in cyclohexane and carbon tetrachloride⁷⁹. In MoCl_3 ⁶⁹ and MoBr_3 ⁷³ the molybdenum atom is also in octahedral co-ordination. The halide ions are hexagonal close-packed with the metal atoms in octahedral holes; there is distortion which brings pairs of metal atoms close together: in MoCl_3 , $\text{Mo}-\text{Cl} = 2.40$ to 2.55 \AA , $\text{Mo}-\text{Mo} = 2.77 \text{ \AA}$; in MoBr_3 , $\text{Mo}-\text{Br} = 2.57 \text{ \AA}$, $\text{Mo}-\text{Mo} = 3.30 \text{ \AA}$.

Rather few physical measurements have been reported for the molybdenum halides. The electronic spectrum of MoCl_5 vapour shows rather-broad bands at 1.0, 15.2, 21.3, 27.8, 35.7, and $>42.6 \text{ kK}$. Assignments are difficult because of the broadness of the bands and some dissociation to MoCl_4 and chlorine; the two highest frequency bands are assigned to charge-transfer transitions and the other bands to $d-d$ transitions⁷⁹.

The magnetic moment of carefully-purified MoCl_5 has recently been re-determined over the range 83 to 333 °K. The value of 1.67 B.M.⁶⁵ is somewhat higher than previous values (1.52⁸⁰, 1.64⁷⁰). Magnetic moments (where known) of the other halides are in Table II. The magnetic moments of the binary halides are less than the spin-only values and decrease with decreasing temperature. This is attributed to anti-ferromagnetism, the spin-exchange occurring via halogen bridges or through direct metal-metal bonding. It is noticeable that the deviation from the spin-only moments increases in the order $\text{Mo}^{\text{V}} < \text{Mo}^{\text{IV}} < \text{Mo}^{\text{III}}$ (correlating with the shorter Mo-Mo bonds in MoCl_3 and MoBr_3 compared with MoCl_5) and in the order $\text{I} < \text{Br} < \text{Cl}$.

(c) *Reactions of molybdenum halides.* These are summarised in Table III⁸¹⁻⁹⁸. Reactions giving oxo-species of molybdenum(V) and (VI) have been reviewed and are not included here⁵.

Addition reactions. Reactions of this type are common for the halides of molybdenum(IV) and molybdenum(III) but rare for MoCl_5 . MoF_6 will add fluoride giving⁸¹ MoF_7^- and MoF_8^{2-} . The number of ligands added is generally two for MoCl_4 and three for MoX_3 and the products are monomeric, paramagnetic, six-co-ordinate complexes except that with Ph_3AsO (but not, apparently, with Ph_3PO and Me_2SO), MoCl_4 gives⁶⁶ a diamagnetic, eight-co-ordinate complex, $\text{MoCl}_4(\text{Ph}_3\text{AsO})_4$.

Substitution and solvolytic reactions. Molybdenum pentachloride reacts vigorously with compounds containing an OH or NH group giving hydrogen chloride and substituted chloro complexes. The commonest products are of the type MoCl_3L_2 ($\text{L} = \text{OR}$, NRR') which, according to their low magnetic moments and molecular weights, are dimeric^{88,89}. Whether the carboxylates, $\text{MoCl}_3(\text{O}_2\text{CR})_2$, are dimeric is not known⁹². An investigation of the mechanism of these substitution reactions would be of great interest since it is uncertain whether MoCl_5 reacts as a dimer or as a monomer or whether the substitution reactions are followed by dimerisation of the products. In discussing the reaction between MoCl_5 and carboxylic acids in carbon tetrachloride, Larson⁹² suggests that MoCl_5 pre-


TABLE III

REACTIONS OF MOLYBDENUM HALIDES^a

Halide and reaction	Reagent	Product ^b	Ref.
MoF ₆ ; addition	MF in liq. IF ₅	K ₂ MoF ₈ , Rb, CsMoF ₇	81
	NOF, NO ₂ F	NO, NO ₂ MoF ₇	82
MoF ₆ ; reduction	NO, NOCl	NO[MoF ₆]	82, 83
	PF ₃	MoF ₅	57
MoF ₆ ; addition	NOF	NO[MoF ₆]	83
MoF ₆ ; addition	py, NMe ₃ , Me ₂ SO, Me ₂ NCHO	MoF ₄ L ₂	84
MoCl ₅ ; addition	Et ₂ O	MoCl ₅ (Et ₂ O) ₂	85
	POCl ₃	MoCl ₅ (POCl ₃) ₂	86
	PCl ₅	MoCl ₅ (PCl ₅) ₂	87
	NMe ₃	MoCl ₅ (NMe ₃) ₂	88
MoCl ₅ ; substitution and solvolysis	conc. HCl, liq. SO ₂	MoOCl ₃ , MoOCl ₄ ²⁻	5
	ROH, -78°	[MoCl ₅ (OR) ₂] ₂	
		(R = Me, Et, n-Pr)	89
	MeOH	MoCl ₅ (OMe) ₂ , MoCl ₅ (OMe) ₃ , MoCl(OMe) ₄	90
	PhOH and p-CO ₂ H, -CH ₃	[MoCl ₅ (OPh) ₃] ₂	10, 91
	R · CO ₂ H	MoCl ₅ (RCO ₂) ₂	92
	Ph ₃ PO	MoOCl ₃ (Ph ₃ PO) ₂	5
	NH ₃ (-33°)	MoCl ₅ (NH ₃) ₂	88
	NH ₃ (25°)	MoCl(NH ₃)(NH ₃) ₂	88
	NHMe ₂	MoCl ₅ (NMe ₂) ₂ (NHMe ₂)	88
	NHR ₁	[MoCl ₅ (NR ₂) ₂] ₂ (R = Et, n-Pr)	88
	NH ₂ R	MoCl ₅ (NHR) ₄ (NH ₂ R)	
		(R = n-Bu, n-Pr) and [MoCl ₅ (NHR) ₂] ₂ (R = n-Pr)	88
	NH ₂ R'	[MoCl ₅ (NHR') ₂] ₂	
		(R' = Me, Et)	88
MoCl ₅ ; reduction	NMe ₃	(NHMe ₂) [MoCl ₅ (NMe ₂)]	88
	pyridine	MoCl ₅ (py) _x (x = 3, 4, 5)	85, 93
	bipyridyl	MoCl ₅ (bipy)	94
	RCN	MoCl ₅ (RCN) ₂	
		(R = Me, Et, n-Pr)	95
MoCl ₅ ; addition	pyridine	MoCl ₅ (py) _x (x = 2, 3)	67
	bipyridyl	MoCl ₅ (bipy)	94
	Ph ₃ AsO	MoCl ₅ (Ph ₃ AsO) ₄	66
	thioxane	MoCl ₅ (S-thioxane) ₂	96
MoCl ₅ ; substitution	acetylacetone	MoCl ₅ (acac) ₂	67
MoCl ₅ ; addition	pyridine	MoCl ₅ (py) ₃	97
MoCl ₅ ; substitution	NH ₃	MoCl ₅ (NH ₃) ₂ (NH ₃) ₂	88
	bipyridyl	[MoCl ₅ (bipy) ₂] ⁺ ₂	
		[MoCl(bipy) ₄] ⁻	94
MoBr ₅ ; addition	pyridine, γ-picoline	MoBr ₅ L ₂	10, 96, 97, 98
	RCN	MoBr ₅ (RCN) ₃	96
MoBr ₅ ; substitution	NH ₃	MoBr(NH ₃) ₂ (NH ₃) ₂	88
	NH ₂ Me	MoBr(NHMe) ₂ (NH ₂ Me)	88
	NHMe ₂	MoBr ₅ (NMe ₂)(NHMe ₂)	88
	bipyridyl	[MoBr ₅ (bipy) ₂] ⁺ ₂	
		[MoBr ₅ (bipy)] ⁻	94
MoI ₅ ; addition	pyridine	MoI ₅ (py) ₃	76
MoI ₅ ; substitution	bipyridyl	[Mo(bipy) ₃] ³⁺ 3 I ⁻	94
MoI ₅ ; reduction	diars	Mo(diars) ₂ I ₃	76

^a For ligand abbreviations see p. 315. ^b Reaction products are formulated as dimers where there is definite evidence (e.g., molecular weights, low magnetic moments).

serves its dimeric structure and that the axial chlorines react preferentially. However, the pentachloride appears to be monomeric (by the molecular weight) in benzene⁷⁸ and its electronic spectrum in cyclohexane and in carbon tetrachloride is the same as in the vapour (implying that it is also monomeric in these solvents)⁷⁹. A more plausible interpretation is that replacement of the axial chlorines of a trigonal bipyramidal monomer first occurs and is followed by dimerisation. If this is the mechanism it seems likely that the dimeric reaction products contain nitrogen or oxygen rather than chlorine bridges:



Dimeric compounds $[\text{MoCl}_2\text{L}_3]_2$ are obtained from MoCl_5 and phenol and substituted phenols^{10,91} and some amines⁸⁸. A partial X-ray analysis of $[\text{MoCl}_2(\text{OPh})_3]_2$ has shown the presence of bridging chlorines and a Mo-Mo bond of 2.8 Å, considerably shorter than in $[\text{MoCl}_5]_2$ (3.84 Å)¹⁰. It appears from the trend in the magnetic moments of the phenolate complexes ($p\text{-CO}_2\text{H} > \text{H} > p\text{-CH}_3$) that metal-metal interaction increases as the *para*-substituents become more electron-donating. This is reasonable because increased electron donation will cause a reduction of the residual charge on the metal atoms and so the repulsion between them will be reduced. Also, orbital expansion, leading to better orbital overlap in the metal-metal bond, will result from a decrease in the effective nuclear charge of the molybdenum atoms¹⁰.

A qualitative feature of the substitution reactions is that the reactivity of MoCl_5 towards phenols (measured in terms of the vigour and speed of the reaction and the yield of product) increases as the *para*-substituent is¹⁰ $\text{CO}_2\text{H} < \text{H} < \text{CH}_3$ and increases towards ammonia and amines (in terms of the number of Mo-Cl bonds broken) in the order⁸⁸ $\text{NR}_3 < \text{NHR}_2 < n\text{-BuNH}_2 < n\text{-PrNH}_2 < \text{EtNH}_2, \text{MeNH}_2 < \text{NH}_3$.

The deep colours of the phenolate complexes are due to charge-transfer transitions (ligand-to-metal), the energies decreasing as the electron-donor character of the *para*-substituent increases¹⁰.

Substitution reactions are less common for the halides of molybdenum(IV) and molybdenum(III). Iodide is replaced more readily than chloride or bromide, e.g., with bipyridyl, MoCl_3 and MoBr_3 give complexes $[\text{MoX}_2(\text{bipy})_2]^+ [\text{MoX}_4(\text{bipy})]^-$ but MoI_3 gives⁹⁴ $[\text{Mo}(\text{bipy})_3]^{3+} + 3 \text{I}^-$. This difference is in accordance with the class-A character of molybdenum(III).

(d) *Reduction with complex formation.* Reduction of MoCl_5 followed by complex formation is an important route to complexes of MoCl_4 with alkyl cyanides⁹⁵, pyridine^{85,93,99} and bipyridyl⁹⁴. The oxidation products of the organic ligands are not known except that with pyridine the product is probably the 1-4'-pyridyl-pyridinium ion^{93,99}. Ligand-exchange reactions of the alkyl cyanide complexes of MoCl_4 have yielded many new complexes (see below). MoI_3 is not reduced by pyridine or bipyridyl but with diarsine a molybdenum(II) complex, $\text{Mo}(\text{diars})_2\text{I}_2$, is obtained⁷⁶.

(e) *Ligand-exchange reactions.* Complexes MoCl_4L_2 and MoCl_4B are formed in ligand-exchange reactions of $\text{MoCl}_4(\text{RCN})_2$ (where L = pyridine, α -picoline, tetrahydrofuran, pentamethylene oxide, Ph_3PO , Ph_3P , Ph_3As , pyrazine, 2,6-dimethyl-pyrazine; B = bipy, 1,10-phenanthroline)⁹⁶. Similar reactions occur with $\text{MoBr}_3(\text{RCN})_3$ giving⁹⁶ $\text{MoBr}_3(\text{RCN})\text{L}_2$ (L = Ph_3P , Ph_3As), $\text{MoBr}_3(\text{MeCN})(\text{bipy})$, $\text{MoBr}_3(\text{pyrazine})_3$, and $\text{MoBr}_3(\text{THF})_3$. Alkyl cyanides are not displaced by thioxane. The compound $\text{MoCl}_4(\text{thioxane})_2$ (in which thioxane appears to be S-bonded) has been prepared⁹⁶ directly from MoCl_4 .

These complexes of MoCl_4 and MoBr_3 are monomeric non-electrolytes. According to infrared data the substituents are *cis* in $\text{MoCl}_4(\text{MeCN})_2$ and *trans* in $\text{MoCl}_4(\text{PPh}_3)_2$. The magnetic moments of the molybdenum(III) complexes are close to the spin-only value but the molybdenum(IV) complexes have low magnetic moments (2.2 to 2.7 B.M.). The electronic spectrum of $\text{MoBr}_3(\text{RCN})_3$ is similar to that of MoCl_6^{3-} as expected from the crystal-field order $\text{RCN} > \text{Cl} > \text{Br}$; crystal-field transitions are observed at ca. 15, 21, and 25 kK and charge-transfer transitions at higher frequencies. In the molybdenum(IV) complexes crystal-field transitions occur⁹⁶ at ca. 20 and 25 kK (also observed for MoCl_6^{2-}).

(ii) Halide complexes

(a) *Preparations.* The complexes are listed and preparative reactions outlined in Table IV¹⁰⁰⁻¹⁰⁸. Various features of these reactions are noteworthy. In reactions involving more than one halogen (MoF_6 and NaI ¹⁰⁰, K_3MoCl_6 and Br_2 ¹⁰⁴, KI and MoCl_5 ¹⁰³) the complex of the more electro-negative halogen is invariably obtained. In some reactions the product depends on the cation; MoF_8^{2-} with K^+ but MoF_7^- with⁸¹ Rb^+ and Cs^+ ; salts of MoBr_6^{2-} only with¹⁰³ Rb^+ and Cs^+ , salts of MoCl_6^{2-} from MoCl_5 and $\text{a}^\circ\text{chloride}$ in liquid sulphur dioxide with only Rb^+ and Cs^+ (also formed in this reaction are salts of MoOCl_4^{2-} which are the only products with pyH^+ , quinH^+ , and Et_2NH_2^+)^{105,106}. Presumably lattice energy considerations are important in the formation of these compounds. It is not possible to obtain complexes of molybdenum(IV) by electrolytic reduction of molybdenum(VI) in aqueous solution since the reduction goes via molybdenum(V) to molybdenum(III).

TABLE IV

PREPARATION OF COMPLEX HALIDES

Compound	Preparation	Ref.
<i>Molybdenum(VI)</i>		
K_3MoF_6 and $(Rb, Cs)MoF_6$	See Table 3.	
<i>Molybdenum(V)</i>		
K_3MoF_6	$MoF_6 + KF$ in liq. SO_2	81
M_2MoF_6 ($M = Na, K, Rb, Cs$)	$MoF_6 + MI$ in liq. SO_2	100
$KMoCl_6$	Fuse $MoCl_5$ and KCl	101
<i>Molybdenum(IV)</i>		
Na_2MoF_6	$MoF_6 + \text{excess } NaI$ in liq. SO_2	102
K_3MoCl_6	$MoCl_5 + KCl$ in ICl	103
	$^a K_3MoCl_6 + Br_2$ (45°)	104
$(Rb, Cs)_2MoCl_6$	$MoCl_5 + MCl$ in liq. SO_2	105, 106
$(pyH)_2MoCl_6$	$MoCl_5(n\text{-}PrCN)_2 + (pyH)Cl$ in $CHCl_3$	106
$(Me_4N)_2MoCl_6 \cdot \frac{1}{2} MeCN$	$MoCl_5 + (Me_4N)Cl$ in $MeCN$	101a
$(Rb, Cs)_2MoBr_6$	$MoBr_5 + MBr + IBr$ (300°)	103
<i>Molybdenum(III)</i>		
$K_3MoCl_6, K_2MoCl_6(H_2O)$	Electrolytic reduction of Mo^{VI} in HCl	107
K_3MoCl_6	$KI + MoCl_5$	103
$K_3MoBr_6, K_2MoBr_6(H_2O)$	Electrolytic reduction of Mo^{VI} in HBr	108
$Rb_2Mo_2Br_9$	$Rb_2MoBr_6 + \text{liq. } NH_3$	103
K_3MoI_6 (? not isolated)	$K_2MoBr_6 + \text{conc. } HI$	11

^a That K_3MoCl_6 is formed is not certain as no analytical figures are given. At 450° the product is $MoCl_5$.

(b) *Structures and physical properties.* The complex ions in $NaMoF_6$ ¹⁰⁹ and K_2MoCl_6 ¹⁰³ are octahedral as indicated by X-ray analysis. ($Mo-F = 1.74 \text{ \AA}$; $Mo-Cl = 2.31 \text{ \AA}$). The binuclear ion $Mo_2Cl_9^{3-}$ is diamagnetic and is assumed to have a structure similar to that of $W_2Cl_9^{3-}$, i.e., two octahedrons joined by a common face, giving three chlorine bridges, and a metal-metal bond¹¹⁰.

The magnetic moments of the complexes K_3MoCl_6 and $K_2MoCl_6 \cdot H_2O$ are in the expected range, 3.7–3.8 B.M.^{10,70,111,112}. Salts of the ions $MoCl_6^{2-}$ and $MoBr_6^{2-}$ have magnetic moments in the range 2.1 to 2.4 B.M. depending on the cation^{101a,103,105,106}. The reduction below the spin-only value is attributed to spin-orbit coupling. The magnetic moments of the complexes M_2MoCl_6 ($M = K, Rb, Cs, Tl$) and M_2MoBr_6 ($M = Rb, Cs$) have been measured over the temperature range 80 to 290 °K, and plots of μ against temperature for the rubidium and caesium salts are in fair agreement with Kotani's theory for the t_{2g}^2 configuration¹⁰³. Best values of the spin-orbit coupling constants (ζ_{nd}) are 450 cm^{-1} for the chloro-complexes and 500 cm^{-1} for the bromo-complexes (much less than the free ion value⁹ for Mo^{4+} of 950 cm^{-1}). Salts of the ion MoF_6^- have magnetic moments around 1.7 B.M. The Curie-Weiss law is obeyed with rather large values of θ indicating anti-ferromagnetic interactions between the ions¹¹³. It should be noted that because of the large spin-orbit coupling constant for Mo^{5+} (ca. 1000 cm^{-1}) the t_{2g}^1 configuration should give moments much below the spin-

only value. Evidently the spin-orbit coupling constant is considerably reduced in the complexes⁹.

The electronic spectra of the molybdenum(III) halide complexes have recently been re-determined by Furlani and Piovesana¹¹ who give references to earlier work. The spectrum of K_3MoCl_6 in $HCl \gg 9 M$ is the same as that of the solid; but at lower HCl concentrations the spin-allowed $d-d$ bands are shifted to higher frequencies as a result of aquation and, between 1 and 2 M HCl , a spectrum attributed to $[MoCl_5(H_2O)]^{2-}$ is observed. The behaviour of the bromo-complex is similar. Spurious bands are sometimes observed as a result of slight oxidation of the molybdenum(III) solutions. The spectra show spin-forbidden transitions at ca. 10 kK ($^2E_g, ^2T_{1g} \leftarrow ^4A_{2g}$) and ca. 14 to 15 kK ($^2T_{2g} \leftarrow ^4A_{2g}$) and spin-allowed crystal-field transitions at 17 to 21 kK ($^4T_{2g} \leftarrow ^4A_{2g}$) and 20 to 26 kK ($^4T_{1g} \leftarrow ^4A_{2g}$). The values of the crystal-field splitting parameter, Δ , and the nephelauxetic ratio, β , increase in the order $I < Br < Cl$. The spectra of salts of $MoCl_6^{2-}$ show crystal-field transitions at 21.51 ($^3T_{2g} \leftarrow ^3T_{1g}$), 25.64 kK ($^3T_{1g}(P) \leftarrow ^3T_{1g}$) and 27.78 (possibly $^3A_{2g} \leftarrow ^3T_{1g}$)¹⁰⁶. The value of Δ for $MoCl_6^{2-}$ is 22 kK. The spectrum of $KMoCl_6$ has a single asymmetric crystal-field band at 24.1 kK ($^2E_g \leftarrow ^2T_{2g}$) giving $\Delta = 24.1$ kK^{101a}. As expected Δ decreases in the order $Mo^V > Mo^{VI} > Mo^{III}$.

(c) *Chemical properties. Ligand exchange.* The hexahalide complexes are hydrolysed in water, the degree of hydrolysis decreasing from molybdenum(V) to molybdenum(III). Exchange reactions of K_3MoCl_6 have been useful in the preparation of organic complexes of molybdenum(III) (see below).

Oxidation-reduction. In aqueous solution the molybdenum(III) and molybdenum(IV) complexes oxidise rapidly in air. Molybdenum(III) is oxidised to molybdenum(V) by iron(III) and to molybdenum(VI) by Ce^{IV} . In the potentiometric titration curve of molybdenum(III) with Ce^{IV} there is no break corresponding to molybdenum(IV). It is clear that molybdenum(IV) is inaccessible as the chloro-complex in water¹¹⁴.

Reduction of K_3MoCl_6 with diarsine has already been mentioned (p. 323).

(d) *Oxohalides and complexes.* Compounds of molybdenum(V) and molybdenum(VI) of this type are common⁵. Oxo-complexes of molybdenum(III), derived from MoO^+ , have been described but there is no evidence (e.g., from infrared spectra) that they contain terminal oxygen. Wardlaw and Wormell¹¹⁵ prepared complexes $MoOX \cdot xH_2O$ ($X = F, Cl, Br$). On the basis of chemical evidence they considered that the red and green forms of $MoOCl(H_2O)_4$ are *cis*- and *trans*-isomers respectively. The oxo-complexes of molybdenum(III) clearly need reinvestigation with modern techniques.

(iii) Complexes with inorganic oxygen-donors

A few, poorly characterised compounds with inorganic oxoanions are known,

e.g., MoO_2SO_4 ¹¹⁶, $\text{MoO}_2(\text{NO}_3)_2$ ¹¹⁷, $\text{K}_4\text{Mo}(\text{CO}_3)_4 \cdot \text{K}_2\text{CO}_3 \cdot 2\text{H}_2\text{O}$ ¹¹⁸. The structure of MoOPO_4 has been determined¹¹⁹.

(iv) *Complexes with inorganic nitrogen-donors*

Molybdenum, unlike chromium, has very little affinity for ammonia and the best-known molybdenum complexes with an inorganic nitrogen-donor are those with thiocyanate.

(a) *Complexes with NH_3 , NH_2^- , NH^{2-} , and N^{3-}* . From a solution of $(\text{NH}_4)_2[\text{MoCl}_5(\text{H}_2\text{O})]$ in liquid ammonia Wardlaw *et al.*^{107b} obtained a complex $(\text{NH}_4)_2[\text{MoCl}_5(\text{NH}_3)]$ which rapidly hydrolyses in water. Rosenheim *et al.*¹²⁰ reported complexes $[\text{MoCl}_3(\text{NH}_3)_3]$, $[\text{MoCl}_2(\text{NH}_3)_4]\text{Cl}$, and $[\text{Mo}(\text{NH}_3)_6]\text{Cl}_3$; but, in view of recent work on the ammonolysis of molybdenum halides⁸⁸, their formulations are probably incorrect. Examples of ammonolysis products of the molybdenum halides include $[\text{MoCl}_2(\text{NH}_2)(\text{NH}_3)_2]$, $[\text{MoBr}(\text{NH}_2)_2(\text{NH}_3)]$ (from MoX_3 and liquid ammonia) and $[\text{MoCl}_3(\text{NH}_2)_2]$ and $[\text{MoCl}(\text{NH})(\text{NH}_2)_2]$ (from MoCl_5)⁸⁸.

Various nitrido complexes of molybdenum(V) and molybdenum(VI) have been reported, e.g., MoNCl_5^{3-} ,¹²¹ $\text{MoO}_3\text{N}^{3-}$,¹²² and MoNCl_3 ¹²³. The compound MoNCl_3 is obtained as dark brown crystals from the reaction of MoCl_5 and ClN_3 . It is diamagnetic and the presence of a Mo-N multiple bond is shown by a band at 1045 cm^{-1} in the infrared spectrum. The force constant of this bond (7.86 mdynes/\AA) is less than that of the MoO multiple bonds in MoOCl_3 (8.24) and MoOCl_4 (7.92). The compound is hydrolytically-unstable but gives an adduct with pyridine, $\text{MoNCl}_3(\text{py})_3$ ¹²³.

(b) *Complexes with thiocyanate*. Thiocyanate complexes are formed by molybdenum in oxidation states +3, +4 and +5. Preparative reactions are given in Table V¹²⁴⁻¹²⁶.

TABLE V
PREPARATION OF THIOCYANATE COMPLEXES

Compound	Preparation	Ref.
<i>Molybdenum(V)</i>		
$(\text{pyH})_2\text{MoO}(\text{NCS})_2$	$(\text{pyH})_2\text{MoOCl}_6 + \text{HNCS}$; $(\text{pyH})_4\text{Mo}_2\text{O}_4(\text{NCS})_6 + \text{HNCS}$	124
$(\text{Me}_4\text{N})_2\text{Mo}_2\text{O}_4(\text{NCS})_6$	$(\text{Me}_4\text{N})_2\text{MoOCl}_6 + \text{NH}_4\text{NCS}$	124
$(\text{pyH})_4\text{Mo}_2\text{O}_4(\text{NCS})_6$	$(\text{pyH})_2\text{MoOCl}_6 + \text{NH}_4\text{NCS}$	124
$\text{Mo}(\text{NCS})_5(\text{Me}_2\text{CO})_2$ and $\text{MoCl}(\text{NCS})_4(\text{Me}_2\text{CO})_2$	$\text{MoCl}_6 + \text{KNCS}$ in acetone	125
<i>Molybdenum(IV)</i>		
$(\text{pyH})_2\text{Mo}(\text{NCS})_6$	$(\text{pyH})_3\text{Mo}(\text{NCS})_6 + \text{Fe}(\text{CN})_6^{3-}$	126
<i>Molybdenum(III)</i>		
$\text{M}_2\text{Mo}(\text{NCS})_6 \cdot x\text{H}_2\text{O}$ ($\text{M} = \text{K}, \text{NH}_4, \text{pyH}$)	$\text{Mo}^{\text{III}} + \text{NCS}^- + \text{M}^+$	107b

Preparations and chemical properties. Because of the instability of molybdenum(IV) in water the complex $(\text{pyH})_2\text{Mo}(\text{NCS})_6$ is prepared by triturating a suspension of $(\text{pyH})_2\text{Mo}(\text{NCS})_6$ with a concentrated aqueous solution of potassium ferricyanide¹²⁶. A procedure¹²⁷ in which thiocyanate and pyridine are added to a 2:1 mixture of equimolar-solutions of molybdenum(III) and molybdenum(VI) (i.e., $2 \text{ Mo}^{\text{III}} + \text{Mo}^{\text{VI}} \rightleftharpoons 3 \text{ Mo}^{\text{IV}}$) gives mainly the molybdenum(V) complex¹¹⁴.

In solution (e.g., ethanol, dioxan) the compounds are very susceptible to hydrolysis and the molybdenum(IV) complex disproportionates to molybdenum(III) and molybdenum(V). Its instability relative to molybdenum(III) and molybdenum(V) is similar to that of molybdenum(IV) with chloride and water (but quite different from the behaviour with cyanide ion which strongly stabilises molybdenum(IV))¹¹⁴.

Physical properties. There are no X-ray structural data. Infrared data show that the thiocyanates are bound to molybdenum through nitrogen: the CN stretching frequencies for the molybdenum(V) and (IV) complexes¹²⁸ and the CN and CS^{128,129} frequencies for the molybdenum(III) complexes occur in the appropriate range for N-bonded thiocyanate. Infrared data also show the absence of ligand-pyridine¹³⁰ and the presence in the molybdenum(V) complexes of terminal oxygen¹⁰.

The magnetic moments of the molybdenum(III) complexes are close to the spin-only value^{112b}, but the moment of $(\text{pyH})_2\text{Mo}(\text{NCS})_6$ is low because of spin-orbit coupling¹⁰. The magnetic moment of $(\text{pyH})_2\text{MoO}(\text{NCS})_5$ is normal whilst the other molybdenum(V) complexes have low moments as a result of spin-spin interaction through oxygen bridges or a metal-metal bond⁵.

In the electronic spectra of the oxomolybdenum(V) thiocyanate complexes the first crystal-field transition is observed as a weak broad peak near 12.5 kK, whilst ligand-to-metal charge-transfer transitions occur at higher frequencies⁵. Crystal-field transitions in the thiocyanates of molybdenum(IV) and molybdenum(III) are obscured by intense charge-transfer transitions^{10,129}. The energies of the charge-transfer transitions increase in the order $\text{Mo}^{\text{IV}} < \text{Mo}^{\text{V}} < \text{Mo}^{\text{III}}$. The expected order in terms of the formal charges on the metal is $\text{Mo}^{\text{V}} < \text{Mo}^{\text{IV}} < \text{Mo}^{\text{III}}$; the reversal of molybdenum(IV) and (V) may be attributed to the effect of π -donation from oxygen in the oxomolybdenum(V) complexes⁵.

(v) Cyanide complexes

The complexes are listed and preparative reactions and some of their properties are given in Table VI¹³¹⁻¹⁴³.

(a) *Structures, bonding, and physical properties.* The octacyanomolybdate anion in $\text{K}_4\text{Mo}(\text{CN})_8 \cdot 2 \text{ H}_2\text{O}$ has been shown by X-ray analysis to be dodecahedral. Mo-C distances are 2.04 to 2.24 Å (average 2.15 Å), C-N, 1.07 to 1.27 Å

TABLE VI

CYANIDE COMPLEXES

Compound	Preparation ^a	Properties ^b	Ref.
<i>Molybdenum(V)</i>			
MoO(OH)(CN) ₂ (H ₂ O) ₂	K ₄ [Mo(CN) ₈] + dil. HNO ₃	red	131
K ₃ [Mo(CN) ₈]	K ₄ [Mo(CN) ₈] + Ce ^{IV} or Mn ^{VII}	yellow, $\mu = 1.66$	10, 127, 132
K ₃ [Mo(OH) ₂ (CN) ₄] · 2 H ₂ O ^c	MoO(OH) ₂ + KCN + EtOH	blue, $\mu = 1.91$	133
<i>Molybdenum(IV)</i>			
K ₄ [Mo(CN) ₈] · 2 H ₂ O	Mo ^{VI} + N ₂ H ₄ · 2 HCl, KCN	yellow, diamag.	70, 132
	[Mo ₂ O ₄ (NCS) ₄] ⁴⁻ + KCN		127, 134
	K ₃ [Mo(CN) ₈] + [Fe(CN) ₆] ⁴⁻		135
	K ₄ [Mo(CN) ₈] in air		136
M ₂ [Mo(CN) ₈ R ₂] · 4 H ₂ O	see text	red	137
K[Mo(CN) ₅]	Aq. Mo ^{VI} + KCN	white, diamag.	136b
K ₄ [Mo(OH) ₂ (CN) ₄] · x H ₂ O	photolysis of K ₄ [Mo(CN) ₈]	purple	138
K ₄ [MoO ₂ (CN) ₄]	Aq. Mo ^{VI} + KCN	red-brown, diamag.	136b
K ₄ [Mo(OH) ₂ (CN) ₄] · 4 H ₂ O ^c	MoO(OH) ₂ + KCN + KOH	red, diamag.	131, 133
K ₃ [Mo(OH) ₂ (CN) ₄ (H ₂ O)] · 2 H ₂ O	MoO(OH) ₂ + KCN + EtOH	blue	131
K ₃ [Mo(OH) ₂ (CN) ₄]	photolysis of K ₄ [Mo(CN) ₈]	blue	137
K ₃ [MoS(OH)(CN) ₄ (H ₂ O) ₂] · 4 H ₂ O	K ₃ [MoS(CN) ₄ (H ₂ O) ₂] + air	violet	139
K ₄ [MoO ₂ (CN) ₄ (NO)] · 3 H ₂ O	K ₄ [Mo(CN) ₅ (NO)] + air	yellow, diamag.	140
<i>Molybdenum(III)</i>			
K ₃ [Mo(CN) ₆] · 2 H ₂ O	K ₂ [MoCl ₆] + KCN, N ₂	red, $\mu = 1.74$	10, 129, 136a
K ₃ [Mo(CN) ₆]	K ₄ [Mo(CN) ₈] + molten KCN	red, $\mu = 0.6$	141
K ₃ [MoS(CN) ₄] · 2 H ₂ O	Mo ^{VI} + KCN + H ₂ S	light blue	142
<i>Molybdenum(II) or (0)^d</i>			
K ₄ [Mo(OH) ₂ (CN) ₄ (NO)] or K ₄ [Mo(CN) ₄ (NO)]	Mo ^{VI} + H ₂ NOH + OH ⁻ + KCN	purple, diamag.	133, 143

^a All reactions in aqueous solution. ^b Colour and magnetic moment (μ) in Bohr magnetons at ca. 20°.

^{c, d} The compounds may be identical: see text.

(average 1.15 Å), and N-N separations are 3.64 to 4.82 Å with an average of four longer distances at 4.75 Å and fourteen shorter at 3.98 Å. Strictly, the anion belongs to the point-group C_{2v} , but, for a structure with averaged bond lengths and angles, the point-group is D_{2d} , the eight cyanides occupying the vertices of a dodecahedron with triangular faces (Fig. 3(a))¹⁴⁴. Discussions of the structure and bonding have generally assumed D_{2d} symmetry. An alternative structure for eight-co-ordination is the square anti-prism (D_{4d} symmetry, Fig. 3(a)). The dodecahedral and anti-prismatic structures are both obtained by distortions of a cube (Fig. 3(a)). The structures of the other cyanide complexes have not been determined by X-ray crystallography. Because the infrared spectra of K₄[Mo(CN)₈] and K₃[Mo(CN)₈] are very similar it is assumed that the latter compound is also dodecahedral¹⁴⁵.

The bonding in the octacyanides is described from a valence-bond approach in terms of d^4sp^3 hybrids with one non- σ -bonding d -orbital occupied in molybdenum(V) by one electron and in molybdenum(IV) by two electrons in agreement with diamagnetism for the molybdenum(IV) compound^{70, 132} and one unpaired




Fig. 3. (a) Distortions of a cube to a dodecahedron (top) and a square antiprism (bottom) (adapted from ref. 149). The reference axes pass through the face centres of the cube. (b) One-electron molecular-orbital energy level schemes for $[Mo(CN)_8]^{3-}$ (left) and $[Mo(CN)_8]^{4-}$ (right) (adapted from ref. 146h). Full vertical lines show spin-allowed $d-d$ transitions, broken vertical lines spin-forbidden $d-d$ transitions.

electron for molybdenum(V)^{10,132}. More recent crystal-field and ligand-field descriptions show that one of the d -orbitals is strongly stabilised, $d_{x^2-y^2}$ for a dodecahedron and d_{z^2} for a square anti-prism¹⁴⁶. Neither of these orbitals has the correct symmetry for σ -bonding. The $d_{x^2-y^2}$ orbital of the dodecahedron, but not the d_{x^2} orbital of the anti-prism, is capable of π -bonding with the anti-bonding (π^*) orbitals of the cyanides.

Energetically, there is very little difference between the dodecahedral and antiprismatic structures¹⁴⁷. It is possible that the dodecahedral structure is slightly

favoured by π -bonding involving the $d_{x^2-y^2}$ orbital. According to electron-spin resonance measurements on $[\text{Mo}(\text{CN})_8]^{3-}$ enriched with ^{13}C , there is extensive delocalisation of the unpaired electron of molybdenum(V) onto the carbon atoms¹⁴⁸. Four of the cyanides are better disposed for π -bonding than the others and, in agreement with this, up to four cyanides may be replaced by hydroxide giving^{146b} the ion $[\text{Mo}(\text{CN})_4(\text{OH})_4]^{4-}$. Against this it is argued that the effect of an electron pair in $d_{x^2-y^2}$ could be to cause an inhomogeneous swelling of the complex so that σ -bonding is loosened to an extent for which π -bonding cannot entirely compensate¹⁴⁷. How these two descriptions are related is not clear.

Whether the $[\text{Mo}(\text{CN})_8]^{4-}$ ion is dodecahedral or anti-prismatic in solution is uncertain. Raman¹⁴⁹, infrared^{145,149} and electronic spectra^{146c,f,g,h} have been interpreted with both models. Infrared data on $\text{K}_4\text{Mo}(\text{CN})_8 \cdot 2\text{H}_2\text{O}$ are consistent with a dodecahedral structure for the solid but it appears that infrared and Raman spectroscopy cannot determine uniquely the structure of the ion in solution¹⁴⁹.

The structure of the molybdenum(III) complex $\text{K}_4\text{Mo}(\text{CN})_7 \cdot 2\text{H}_2\text{O}$, is not known. Two possibilities are eight-co-ordination, dodecahedral (i.e., an anion $[\text{Mo}(\text{CN})_7(\text{H}_2\text{O})]^{4-}$) or seven-co-ordination, pentagonal bipyramidal (i.e., $[\text{Mo}(\text{CN})_7]^{4-}$). The latter structure would be expected from the nine-orbital rule. The magnetic moment^{10,129}, corresponding to one unpaired electron, is consistent with both structures (for the pentagonal bipyramid an orbital doublet lies lowest)¹⁵⁰. That the complex is readily and rapidly oxidised (unlike $\text{Mo}(\text{CO})_2(\text{diars})\text{I}_3$ and $\text{K}_4\text{Mo}(\text{CN})_8$) suggests that one of the three d -electrons occupies a relatively unstable orbital and that the structures of the molybdenum(III) and molybdenum(IV) compounds are similar.

The structure of the compound $\text{KMo}(\text{CN})_5$ is also unknown^{136b}. The present writer has been unable to repeat the preparation of this compound — an experience shared by Figgis and Lewis⁹.

The compound $\text{K}_2\text{Mo}(\text{CN})_5$ also presents structural problems¹⁴¹. The oxidation number of the molybdenum, determined titrimetrically, is three and the compound appears to be monomeric in solution. C-N stretching frequencies are in the same range as for $\text{K}_4\text{Mo}(\text{CN})_8$, but the low magnetic moment of the compound both in the solid state and in solution strongly suggest dimerisation with spin-pairing.

The purple pentacyanonitrosyl complex obtained by reducing with hydroxylamine an alkaline solution of molybdate and cyanide has been variously formulated as a complex of molybdenum(O), $\text{K}_4[\text{Mo}(\text{CN})_5\text{NO}]$ (either with or without water of hydration)¹⁴³ and as an eight-co-ordinate complex of molybdenum(II)¹³³, $\text{K}_4[\text{Mo}(\text{OH})_2(\text{CN})_5\text{NO}]$. Co-ordination of nitric oxide as NO^+ is indicated by the N-O stretching frequency. According to Riley and Ho^{143b} (whose work is the most convincing) the compound is very sensitive to air and moisture and the apparent presence of water or hydroxide is due to partial decomposition. The pure compound shows an N-O stretching frequency at 1455

cm^{-1} ; a band at 1580 to 1590 cm^{-1} (assigned by Griffith *et al.*¹³³ as the N-O stretch) develops during exposure of the mulls to the atmosphere. The diamagnetism is consistent with molybdenum(O) and the rather low N-O frequency is understandable in terms of back donation from molybdenum(O) favoured by the zero formal charge on the metal.

Atmospheric oxidation of the purple complex gives¹⁴⁰ a yellow, diamagnetic molybdenum(IV) complex, $\text{K}_4[\text{Mo}(\text{CN})_5\text{O}_2\text{NO}] \cdot 3\text{H}_2\text{O}$. The N-O stretching frequency (1580 cm^{-1}) is higher than in the purple molybdenum(O) compound, which is expected since molybdenum(IV) should be a poorer π -donor than molybdenum(O). (If the band at 1580 cm^{-1} is assigned as the N-O stretch in the purple molybdenum(O) or molybdenum(II) complex then it becomes necessary to explain why the frequency is the same in the molybdenum(IV) complex.)¹⁴⁰ Further work on these complexes seems called for.

Hydroxocyanides. The literature on the hydroxocyanides is confused and a careful reinvestigation of these compounds would be worthwhile. Bucknall and Wardlaw¹³¹ reacted molybdenum(V) hydroxide, $\text{MoO}(\text{OH})_3$, with the theoretical amount of potassium cyanide and obtained a *blue* aqueous solution from which potassium hydroxide precipitated *red* crystals, $\text{K}_4[\text{Mo}(\text{OH})_4(\text{CN})_4] \cdot 4\text{H}_2\text{O}$. The *red* complex gave a *blue* aqueous solution from which ethanol precipitated *blue* crystals, $\text{K}_3[\text{Mo}(\text{OH})_3(\text{CN})_4(\text{H}_2\text{O})] \cdot 2\text{H}_2\text{O}$. With excess KCN both compounds gave $\text{K}_4\text{Mo}(\text{CN})_8$.

Adamson and Perumareddi¹⁵¹ obtained a *blue* diamagnetic complex, $\text{K}_3[\text{Mo}(\text{CN})_4(\text{OH})_3(\text{H}_2\text{O})]$ (apparently the anhydrous form of Wardlaw's blue compound) as the ultimate photolysis product of an aqueous solution of $\text{K}_4[\text{Mo}(\text{CN})_8]$. Jakob *et al.*¹³⁷ considered the *blue* photolysis product to be $\text{K}_2\text{Mo}(\text{CN})_4(\text{OH})_2$, presumably an octahedral, paramagnetic complex of molybdenum(IV), which differs from Adamson's *blue* complex by one KOH and one H_2O . Since, according to Wardlaw, the hydroxo complexes are readily interconverted, it is possible that the nature of the solid obtained as the ultimate photolysis product is dependent on conditions of isolation and drying. Collenberg¹³⁸ exposed a solution of $\text{K}_4\text{Mo}(\text{CN})_8$ in molar KOH to bright sunlight and isolated a violet compound, $\text{K}_4[\text{Mo}(\text{OH})_3(\text{CN})_5] \cdot x\text{H}_2\text{O}$.

Griffith *et al.*¹³³ apparently repeated Wardlaw's experimental work and obtained a *blue* complex with a magnetic moment of 1.91 B.M., slightly greater than the spin-only moment for molybdenum(V) or low-spin molybdenum(III). They formulated the complex with molybdenum(V), $\text{K}_3[\text{Mo}(\text{OH})_4(\text{CN})_4] \cdot 2\text{H}_2\text{O}$, or with molybdenum(III), $\text{K}_3[\text{MoO}_2(\text{CN})_4(\text{H}_2\text{O})_2]$. It is possible that this material is a mixture of Wardlaw's and Jakob's compounds.

Steele^{136b} was unable to repeat the preparation of Wardlaw's red and blue complexes but obtained a *brown* complex, $\text{K}_4[\text{MoO}_2(\text{CN})_4]$, by refluxing a solution of KCN and MoO_3 . However, Golding and Carrington^{146a} were able to prepare Wardlaw's *red* complex and gave it the same formula. They report its

electronic spectrum in aqueous solution (a $d-d$ transition at 20 kK and a charge-transfer transition above 42 kK).

Compounds with "eight(quasi-ten)-co-ordination". Compounds of the general formula $M_2[Mo(CN)_8R_2] \cdot 4 H_2O$ ($M = Cd, Mn$; $R = H_2O, NH_3, N_2H_4$) separate as red crystals from aqueous solution¹³⁷. Their aqueous solutions are stable in the dark but hydrolyse in light to $[Mo(CN)_4(OH)_2]^{2-}$. Hoard and Silverton¹⁴⁷ describe the molybdenum as having "eight(quasi-ten)-co-ordination" and suggest an antiprismatic structure with the two R groups located above and below the two square faces of the antiprism.

Electronic absorption spectra. There have been a number of attempts to interpret the electronic spectra of $K_3Mo(CN)_8$ and $K_4Mo(CN)_8$ using crystal-field and ligand-field theory and the original papers should be consulted for details^{10, 146c, f, g, h}. Adamson *et al.*^{146h} give a useful critical account of earlier work. The splitting of the d -orbitals according to their ligand-field calculation is shown in Fig. 3(b) for D_{2d} (dodecahedral) and C_2 symmetry. The effect of a reduction in symmetry from D_{2d} to C_2 is to split the e -level into two a -levels. Peak positions and assignments are given in Table VII. There is good agreement between observed and calculated peak positions although not all possible transitions are observed.

TABLE VII

ELECTRONIC SPECTRA OF MOLYBDENUM CYANIDE COMPLEXES

$K_3[Mo(CN)_8]^{3-}$ ^a	$K_4[Mo(CN)_8] \cdot 2 H_2O$ ^a	$K_4[Mo(CN)_7] \cdot 2 H_2O$ ^b
25.8 ${}^2B_2 \leftarrow {}^2B_1$	19.6 ${}^3B_1 \leftarrow {}^1A_1$	19.8 $d-d$
32.0 ${}^2A_1 \leftarrow {}^2B_1$	23.2 ${}^1A_2 \leftarrow {}^1A_1$	26.3 $d-d$
36.8 ${}^2E \leftarrow {}^2B_1$	27.2 ${}^1B_1 \leftarrow {}^1A_1$	45.5 C-T
39.8 ${}^2E \leftarrow {}^2B_1$	32.45 ${}^1E \leftarrow {}^1A_1$	
	37.4 ${}^1E' \leftarrow {}^1A_1$	
	41.67 C-T	

^a Peak positions in kK (1 kK = 10^3 cm^{-1}) and assignments from ref. 146h. C-T denotes a charge-transfer transition.

^b From ref. 10.

Good agreement is also obtained by König^{146e} assuming D_{4d} symmetry (i.e., the antiprismatic structure) but his assignments have been criticised^{146h} because the strongest optical absorption is assigned to highly-forbidden two-electron jumps.

The spectrum of $K_4Mo(CN)_7 \cdot 2 H_2O$ has been measured but assignments have not been attempted. There are two $d-d$ transitions and one charge-transfer transition.¹⁰

Magnetic measurements. The magnetic moments of the cyanide complexes are included in Table VII. The complexes are all low spin having one or no unpaired electrons.

The electron-spin resonance spectra of $K_3Mo(CN)_8$ and $Ag_3Mo(CN)_8$ as powders have a sharp single line at $g = 1.993$ ^{146c}, or 2.005 ± 0.005 ^{146g}, the

sharpness suggesting that the g -tensor is isotropic. An electrostatic crystal field calculation gives $g = 1.92$, much less than the observed value, indicating extensive mixing of the d -orbitals and the cyanide orbitals^{146c}.

(b) *Chemical properties.* Some of the more important reactions of the complexes are apparent from Table VI.

Reactions of the cyanide ligands. With dimethyl sulphate two cyanides of $K_4Mo(CN)_8 \cdot 2 H_2O$ are methylated and two are displaced giving $[Mo(CN)_4(CH_3NC)_2(H_2O)_2] \cdot 4 H_2O$. With potassium cyanide this compound reverts to the octacyanide¹⁵².

With BF_3 , $K_4Mo(CN)_8$ behaves as a Lewis base giving $K_4Mo(CN)_8(BF_3)_8$ which contains cyanide bridges between molybdenum and boron atoms (i.e., $Mo-C \equiv N-BF_3$)¹⁵³.

Cyanide exchange and photolysis. Unlike many molybdenum complexes the octacyanides do not undergo rapid hydrolysis in water. $K_4Mo(CN)_8$ is inert with respect to exchange of ^{14}C -labelled cyanide in the dark.¹⁵⁴ Hydrolysis does, however, occur when an aqueous solution is irradiated by ultraviolet light, the colour changing from yellow through red and green to blue (cf. p. 338)^{137,151,154,155}. High-intensity irradiation of aqueous $K_4Mo(CN)_8$ produces a transient red species formulated by Jakob *et al.*¹³⁷ as $[Mo(CN)_8(H_2O)_2]^{4-}$ and by Adamson and Perumareddi¹⁵¹ as $[Mo(CN)_7OH]^{4-}$. They consider that the mechanism of photolysis is



where R is the red intermediate and B the final blue product, $[Mo(CN)_4(OH)_3(H_2O)]^{3-}$.

Oxidation-reduction. Molybdenum(IV) is strongly stabilised in the octacyano complex with respect to both oxidation and reduction (presumably because for molybdenum(IV) the stable $d_{x^2-y^2}$ -orbital is fully occupied). Oxidation to $K_3Mo(CN)_8$ requires powerful oxidising agents (Ce^{IV} and Mn^{VII})¹²⁷ and reduction to $K_2Mo(CN)_5$ requires¹⁴¹ molten KCN. There is some reduction of $K_4Mo(CN)_8$ by potassium in liquid ammonia to an unidentified cyano complex which is extremely air-sensitive¹⁵⁶. In contrast, the molybdenum(III) complex $K_4Mo(CN)_7 \cdot 2 H_2O$ rapidly oxidises in air¹³⁶ and the molybdenum(V) complex is reduced by mild reducing agents (e.g., potassium ferrocyanide)¹³⁵.

$K_3Mo(CN)_8$ is highly photosensitive undergoing reduction to $K_4Mo(CN)_8$, and the solid and its solutions should be stored and handled in the dark¹³⁵. Electron transfer between the ions $[Mo(CN)_8]^{3-}$ and $[Mo(CN)_8]^{4-}$ is very fast indicating that a true electron-transfer process is involved and not radical or ion exchange¹⁵⁷.

(vi) Complexes with organic oxygen-donors

Complexes of the molybdenum halides with oxygen-donors are described in section E.(iii) and the molybdenum(II) carboxylates in section D.(iii). Complexes of molybdenum(III) with urea and *N,N*-dimethylformamide MoX_3L_3 ($\text{X} = \text{Cl}, \text{Br}$) and $[\text{Mo}(\text{urea})_6]\text{Br}_3$, have also been prepared¹². Of the remaining complexes the best known are those with acetylacetone and other β -diketones and with oxalate (Table VIII). With two exceptions these complexes are all derived from oxo-molybdenum species. The complexes of molybdenum(V) and (VI) have been reviewed⁵.

TABLE VIII

COMPLEXES WITH ACETYLACETONE AND WITH OXALATE LIGANDS

Oxidation state	Acetylacetone	Oxalate	Ref.
+6	$\text{MoO}_3(\text{acac})_3$	$\text{NaNH}_4[\text{MoO}_3(\text{ox})(\text{H}_2\text{O})_2]$ $\text{K}_2[\text{Mo}_2\text{O}_7(\text{ox})_2(\text{H}_2\text{O})_2]$ $(\text{quinH})_2[\text{MoO}_2(\text{ox})_2]$	5 5 5
+5	$\text{MoO}(\text{OH})(\text{acac})_2 \cdot 4 \text{H}_2\text{O}$ $\text{MoO}_3(\text{acac})$ $\text{Mo}_2\text{O}_3(\text{acac})_4$	$\text{K}_2[\text{Mo}_2\text{O}_7(\text{ox})_2(\text{H}_2\text{O})_2]$	5 5 5
+4	$\text{MoO}_2(\text{acacH})_2$	$\text{MoO}(\text{ox})(\text{H}_2\text{O})_3$ $(\text{NH}_4)_2[\text{Mo}_2\text{O}_7(\text{ox})_2(\text{H}_2\text{O})_2]$	158, 163a 163b, 164, 165, 166
+3	$\text{Mo}(\text{acac})_3$	$(\text{NH}_4)_4\text{Mo}(\text{ox})_4 \cdot 8 \text{H}_2\text{O}$ $\text{Mo}(\text{OH})(\text{ox})(\text{H}_2\text{O})_3$	118 107b, 107c, 159, 161, 162
	$\text{MoO}(\text{acac})_2 \cdot 2\frac{1}{2} \text{H}_2\text{O}$	$\text{Mo}_2\text{O}(\text{ox})_2(\text{H}_2\text{O})_4$ $\text{Mo}_3\text{O}_3(\text{ox})_3 \cdot 12 \text{H}_2\text{O}$	160, 163a 163a

(a) *Acetylacetonates*. A diamagnetic molybdenum(IV) complex, $\text{MoO}_2(\text{acacH})_2$, was obtained as brown needles by reduction with zinc dust of a hot solution of $\text{MoO}_2(\text{acac})_2$ in acetylacetone¹⁵⁸. The compound was originally formulated $\text{Mo}(\text{OH})_2(\text{acac})_2$ but the infrared spectrum showed the absence of OH and suggested the presence of neutral acetylacetone. On treatment of the compound with aqueous alkali "molybdenum(IV) oxide monohydrate" separates quantitatively. Acetylacetone may be replaced by other β -diketones but not by CO, Ph_3P , Ph_3As and PhNC . Reduction of molybdenum(VI) to molybdenum(IV) in an oxygen environment is rather unusual and the evidence for molybdenum(IV) in this compound is not entirely convincing. The compound is described as having the same properties as the "black form" of molybdenum(VI) acetylacetonate prepared by Larson and Moore and shown later¹⁵⁹ to be a diamagnetic molybdenum(V) complex, $\text{Mo}_4\text{O}_3(\text{acac})_4$.

The compound $\text{MoO}(\text{acac})_2 \cdot 2\frac{1}{2} \text{H}_2\text{O}$ is obtained as brown crystals by heating

acetylacetone and $\text{MoOCl}(\text{H}_2\text{O})_4$ in ethanol, but the presence of terminal oxygen has not been proved. Aerial oxidation gives¹⁶⁰ green $\text{MoO}_2(\text{acac})$.

$\text{Mo}(\text{acac})_3$ has been prepared by heating^{107c} $(\text{NH}_4)_2\text{MoCl}_5(\text{H}_2\text{O})$ or^{159,161} K_3MoCl_6 with acetylacetone and by refluxing acetylacetone with¹⁵⁹ $\text{Mo}(\text{CO})_6$. Complexes with other β -diketones were later obtained from the hexacarbonyl¹⁶². The compound oxidises rapidly and exothermically in air to¹⁵⁹ $\text{Mo}_2\text{O}_3(\text{acac})_4$. It does not react with water, benzophenone, pyridine or triphenylphosphine, but with carbon tetrachloride it gives chloromolybdenum materials of variable composition¹⁵⁹. The magnetic moment (3.8 B.M.) is normal and the electronic spectrum shows charge-transfer peaks at 27 and 23.3 kK¹⁶².

(b) *Oxalato complexes.* Various oxalato complexes of molybdenum(III) were prepared by Wardlaw *et al.* from solutions of^{107b} $(\text{NH}_4)_2\text{MoCl}_5(\text{H}_2\text{O})$ or¹⁶³ $\text{Mo}(\text{OH})_3$ and oxalates (see Table VIII).

Aerial oxidation of solutions of molybdenum(III) oxalates gives molybdenum(IV) complexes of empirical formula^{163a} $\text{MoO}(\text{C}_2\text{O}_4)(\text{H}_2\text{O})_3$. A red-violet salt, $\text{M}_2[\text{Mo}_3\text{O}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_5]$ ($\text{M} = \text{K}, \text{pyH}$), has also been isolated^{163b}. A detailed spectrophotometric and titrimetric investigation¹⁶⁴ of the oxidation has shown that molybdenum(V) is formed initially and that the species $[\text{Mo}_3\text{O}_4(\text{C}_2\text{O}_4)_3]^{2-}$ arises from a slow reaction between molybdenum(V) and molybdenum(III). In the polarographic reduction of a solution of molybdenum(IV) oxalate only two waves are observed ($\text{Mo}^{\text{VI}} \rightarrow \text{Mo}^{\text{V}}$ and $\text{Mo}^{\text{V}} \rightarrow \text{Mo}^{\text{III}}$) but for solutions of molybdenum(III) oxalate which have been oxidised in air a third wave, attributed to $\text{Mo}^{\text{IV}} \rightarrow \text{Mo}^{\text{III}}$, is observed. Aqueous solutions of the molybdenum(IV) complex are stable in air for long periods. Saturation of the solution with HCl gas gives a chlorooxalato complex of molybdenum(IV). This behaviour is quite different from that of other molybdenum(IV) compounds (e.g., $\text{K}_4\text{Mo}(\text{CN})_8$) which disproportionate in concentrated hydrochloric acid to molybdenum(III) and molybdenum(V)¹⁶⁵. Stabilisation of molybdenum(IV) by oxygen-donors is unusual and it is possible that the basic structural unit is a trinuclear cluster $[\text{MoO}(\text{C}_2\text{O}_4)(\text{H}_2\text{O})_3]_3$ with metal-metal bonds and oxygen bridges and with one oxalate and one or more water molecules co-ordinated to each molybdenum. Such a structure is consistent with the diamagnetism¹⁶⁶, stability and slow rate of formation of the complex.

The complex $(\text{NH}_4)_4[\text{Mo}(\text{C}_2\text{O}_4)_4] \cdot 8 \text{H}_2\text{O}$ is said to be formed by electrolytic reduction of ammonium molybdate in the presence of oxalic acid. It is brown and diamagnetic and apparently contains eight-co-ordinate molybdenum(IV)¹¹⁸.

(vii) *Complexes with organic nitrogen-, arsenic- and phosphorus-donors and with "oxine" and EDTA*

Molybdenum forms rather few complexes in which these are the only donor


atoms. Substituted carbonyls are described in section C, complexes of molybdenum halides in section E.(i), and molybdenum(II) complexes of phosphorus and arsenic donors in section D. Complexes of these donors with oxo-species are also known⁵, e.g., $\text{MoOCl}_3(\text{MeCN})_2$, $\text{MoOCl}_3(\text{bipy})$, $\text{Mo}_2\text{O}_3\text{Cl}_4(\text{bipy})_2$, $\text{Mo}_2\text{O}_4\text{Cl}_2(\text{bipy})_2$, $\text{MoO}_3(\text{dien})$, $\text{MoOCl}_3(\text{PPh}_3)_2$.

(a) *Pyridine and substituted pyridines.* No complexes of the type $\text{Mo}(\text{py})_6$ are known; but the bidentate ligand 2-aminomethyl-pyridine (amp) forms a complex $[\text{Mo}(\text{amp})_3]^{3+}$ with aqueous molybdenum(III) which may be crystallised as the yellow perchlorate salt ($\mu = 3.7 \text{ B.M.}$)¹⁶⁷.

(b) *2,2'-bipyridyl and 1,10-phenanthroline.* By reacting 2,2'-bipyridyl (bipy) and 1,10-phenanthroline (phen) with $(\text{NH}_4)_3\text{MoCl}_6$, Steele¹⁶⁸ obtained dark red complexes $[\text{Mo}(\text{bipy})_3]\text{X}_3$ and $[\text{Mo}(\text{phen})_3]\text{X}_3$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) with magnetic moments of 3.7 to 3.8 B.M. From MoI_3 and excess bipyridyl, Carmichael *et al.*⁹⁴ obtained a crimson compound $[\text{Mo}(\text{bipy})_3]\text{I}_3$ ($\mu = 3.93$) but they were unable to prepare the corresponding chloride and bromide salts and, instead, obtained compounds $[\text{MoX}_2(\text{bipy})_2]^+[\text{MoX}_4(\text{bipy})]^-$ ($\text{X} = \text{Cl}, \text{Br}$) (crimson, $\mu = 3.62 \text{ B.M.}$). Furlani and Piovesana¹¹ were also unable to prepare tris(bipyridyl) complexes and obtained compounds $\text{BH}[\text{MoCl}_4\text{B}]$ ($\text{B} = \text{bipy}$ or phen) (brown solids, 1:1 electrolytes in nitromethane). Their formulae are based on analyses for chlorine only. Herzog and Schneider³⁸ appear to have prepared a compound $[\text{Mo}(\text{bipy})_3]\text{Cl}_3$ since they describe its reduction to $\text{Mo}(\text{bipy})_3$. The bipyridyl and phenanthroline complexes of molybdenum(III) are stable to oxidation. Stabilisation of molybdenum(III) by π -acceptor ligands is, of course, expected.

(c) *Phosphorus- and arsenic-donors.* Complexes of the higher-oxidation states with these donors are rare. A molybdenum(IV) complex, $\text{MoBr}_4(\text{diars})$, (orange-brown, $\mu = 1.96 \text{ B.M.}$) is formed by the action of excess bromine on $\text{Mo}(\text{CO})_4(\text{diars})$ ²¹.

(d) *Oxine and EDTA derivatives.* With 8-hydroxyquinoline (oxine) and ethylenediaminetetraacetate (EDTA^{4-}), molybdenum(VI) and molybdenum(V) form complexes⁵:



No complexes of molybdenum(IV) with these ligands are known. A complex of molybdenum(III) with oxine is formed by heating $\text{Li}_3\text{MoCl}_6 \cdot \text{LiCl} \cdot 9 \text{H}_2\text{O}$ with a solution of oxine in ethanol. On the basis of elementary analyses the compound was assigned the rather unusual formula $\text{Mo}_4(\text{oxine})_5\text{Cl}_7(\text{H}_2\text{O})_7$. To explain the low magnetic moment (1.2 B.M.) a bridged structure involving eight-co-ordination is suggested¹⁶⁹.


(viii) Complexes with sulphur-donors

Although molybdenum in its higher oxidation states has class-A character, it does have some affinity for sulphur-donor ligands. The sulphides, Mo_2S_3 , MoS_2 , Mo_2S_5 , and MoS_3 are well known⁷. A recent X-ray analysis of Mo_2S_3 has shown that the sulphur atoms form a distorted close-packed array with the metal atoms displaced 0.34 Å from the centres of octahedral holes and metal-metal bonds of 2.85 Å¹⁷⁰. In MoS_2 the metal atom is surrounded by six sulphurs at the vertices of a trigonal prism. MoS_3 is amorphous¹⁷¹.


Molybdates react with hydrogen sulphide with successive replacement of oxygen by sulphur; deep-red tetrathiomolybdates may be isolated¹⁷² as ammonium or alkylammonium salts, R_2MoS_4 .

Numerous colour reactions of molybdates with organo-sulphur compounds are described in the analytical literature but the compounds formed have generally not been well characterised. The reaction of cysteine, $\text{CH}_2(\text{SH})\text{CH}(\text{NH}_2)\text{COOH}$, with molybdenum(VI) and molybdenum(V) in aqueous solution has been studied spectrophotometrically. With molybdenum(VI) the ligand : metal ratio varies from 3 : 1 to 1 : 1 but with molybdenum(V) only the 1 : 1 complex is formed. It is thought that co-ordination is through the ionised mercapto- and either the amino or carboxyl group¹⁷³. The structure of the complex $\text{Mo}_2\text{O}_3(\text{xanthate})_4$ has recently been determined^{5,174}. Some well-characterised complexes in which molybdenum is fully co-ordinated with sulphur are described below.

Complexes with 1,2-dithiols. Recently there has been widespread interest in tris-complexes of molybdenum and other transition metals with ligands of the type $\text{R}-\text{C}(\text{SH})=\text{C}(\text{SH})-\text{R}$ where $\text{R} = \text{CF}_3$ ¹⁷⁵, C_6H_5 ¹⁷⁶, CN ¹⁷⁷ and with toluene-3,4-dithiol, $(\text{H}_2\text{tdt}, \text{I})$ ¹⁷⁸. Complexes of toluene-3,4-dithiol with molybdenum(VI) and (IV) were first prepared by Gilbert and Sandell^{178a}, and formulated $\text{Mo}(\text{tdt})_3$ and $\text{Mo}(\text{tdt})_2(\text{H}_2\text{tdt})$ or $\text{Mo}(\text{tdt})(\text{Htdt})_2$ respectively. The complex $[\text{Mo}(\text{CF}_3-\text{C}(\text{S})-\text{C}(\text{S})-\text{CF}_3)_3]$ (abbreviated to $[\text{Mo}(\text{SS})_3]^{0-}$) has been obtained as black crystals by refluxing $[\text{Mo}(\text{CO})_6]$ with bis(trifluoromethyl)dithietene(II) in methylcyclohexane^{175a}.



I



II

The complex is reduced by hydrazine in ethanol to a blue complex, $[\text{Mo}(\text{SS})_3]^{2-}$, isolated as the $[\text{Ph}_4\text{As}]^+$ salt. Both compounds are diamagnetic. They react in solution to give a singly-charged complex, $[\text{Mo}(\text{SS})_3]^{1-}$, isolated as the $[\text{Ph}_4\text{As}]^+$ salt which has one unpaired electron. The electron-spin resonance spectrum of this compound suggests that the unpaired electron resides in a molecular orbital which is principally ligand in character. It appears that in all three compounds molyb-

denum is formally in the +4 oxidation state and that the electronic configurations (including a ligand orbital, L) are^{175b}: $[\text{Mo}(\text{SS})_3]^0$, d^2L^0 ; $[\text{Mo}(\text{SS})_3]^{1-}$, d^2L^1 ; $[\text{Mo}(\text{SS})_3]^{2-}$, d^2L^2 . Other tris-dithiolato complexes undergo similar redox reactions^{178b}.

The rhenium complex, $\text{Re}(\text{PhC}(\text{S})=\text{C}(\text{S})\text{Ph})_3$, has been shown by X-ray analysis¹⁷⁹ to have a trigonal prismatic structure (D_{3h} -symmetry Fig. 4) and it seems likely that other tris(dithiolato) complexes also have this structure.







Fig. 4. Trigonal prismatic coordination. In the compound $\text{Re}(\text{PhC}(\text{S})=\text{C}(\text{S})\text{Ph})_3$ the bidentate ligands are bonded to the metal atom through pairs of sulphur atoms aa, bb, and cc¹⁷⁹.

*Complexes of molybdenum(III) with multidentate sulphur-ligands*¹⁶⁹. With a number of sulphur-ligands molybdenum(III) gives complexes which are non-electrolytes and have low magnetic moments (0.3–1.6 B.M.). It is possible that the complexes are dimerised through thiol bridges, and that in most of the compounds the molybdenum is eight-co-ordinate. Examples of proposed structures are shown below (the complex (I) with 3-ethyl-thiopropene-1-thiol, $\text{EtSCH}_2\text{CH}_2\text{CH}_2\text{SH}$; and the complex (II) with *N*-2-mercapto-phenylene-2'-pyridylmethylenimine (III)) but they must be regarded as tentative until X-ray structural determinations have been carried out.



With thiourea and substituted thioureas, molybdenum(III) forms complexes with normal magnetic moments (e.g., $[\text{Mo}(\text{NH}_2\text{CSNH}_2)_3\text{Cl}_3]$, $\mu = 3.71$ B.M.) and with low moments (e.g., $[\text{Mo}_2(\text{NH}_2\text{CSNH}_2)_3\text{Cl}_6]$, $\mu = 0.59$ B.M.). It is suggested that the complexes with low moments are dimeric with three halogen bridges¹⁸⁰.

REFERENCES

- 1 See, for example, G. W. A. FOWLES in W. L. JOLLY (Ed.), *Preparative Inorganic Reactions*, Interscience, 1964, Vol. 1, p. 121.
- 2 R. C. BRAY in *The Enzymes*, Academic Press, Vol. 7, 1963.
- 3 R. C. BRAY AND K. R. HARRAP, *Ann. Repts. on Progr. Chem. Chem. Soc. London*, 55 (1958) 348.
- 4 H. KAY, *Ind. Eng. Chem.*, 49 (1957) no. 6, 44A.
- 5 P. C. H. MITCHELL, *Quart. Rev. (London)*, 20 (1966) 103.
- 6 G. P. HAIGHT AND C. VON FRANKENBERG, *Acta Chem. Scand.*, 15 (1961) 2026; G. P. HAIGHT, P. MOHLNER AND A. KATZ, *Acta Chem. Scand.*, 16 (1962) 221, 659; G. P. HAIGHT AND W. F. SAYER, *J. Am. Chem. Soc.*, 74 (1952) 6056.
- 7 (a) *Gmelin's Handbuch der anorganischen Chemie, Molybdän*, Syst. No. 53, Berlin, 1935; (b) N. V. SIDGWICK, *Chemical Elements and Their Compounds*, Oxford Univ. Press, 1950, Vol. II, p. 998; (c) D. H. KILLEFER AND A. LINZ, *Molybdenum Compounds*, Interscience, New York, 1952; (d) G. SPENGLER AND J. GANNSHEIMER, *Angew. Chem.*, 69 (1957) 523; (e) P. PASCAL (Ed.), *Nouveau Traité de Chimie Minérale*, Masson et Cie., Paris, 1959, Vol. XIV.
- 8 C. E. MOORE, *Atomic Energy Levels*, National Bureau of Standards, Washington, 1958.
- 9 B. N. FIGGIS AND J. LEWIS, *Progr. Inorg. Chem.*, 6 (1964) 37. See also J. LEWIS, R. S. NYHOLM AND P. W. SMITH, *J. Chem. Soc.*, (1962) 2592, and references therein.
- 10 P. C. H. MITCHELL AND R. J. P. WILLIAMS, *J. Chem. Soc.*, (1962) 4570.
- 11 C. FURLANI AND O. PROVASANA, *Mol. Phys.*, 9 (1965) 341.
- 12 T. KOMORITA, S. MIKI AND S. YAMADA, *Bull. Chem. Soc. Japan*, 38 (1965) 123.
- 13 (a) J. C. HILEMAN in ref. 1 p. 77, (b) E. W. ABEL, *Quart. Rev.*, 17 (1963) 133; (c) H. ZEISS, *Organometallic Chemistry*, Reinhold, 1960.
- 14 F. A. COTTON AND R. M. WING, *Inorg. Chem.*, 4 (1965) 314.
- 15 H. B. GRAY AND N. A. BEACH, *J. Am. Chem. Soc.*, 85 (1963) 2922.
- 16 (a) R. P. M. WERNER AND T. H. COFFIELD, *Chem. Ind.*, (1960) 936; (b) E. ABEL, M. BENNETT AND G. WILKINSON, *J. Chem. Soc.* (1959) 2323; (c) C. S. KRAHMANZEL AND F. A. COTTON, *J. Am. Chem. Soc.*, 84 (1962) 4432; *Inorg. Chem.*, 2 (1963) 533; (d) G. W. A. FOWLES AND D. K. JENKINS, *Inorg. Chem.*, 3 (1964) 257; (e) F. A. COTTON AND F. ZINGALES, *Inorg. Chem.*, 1 (1962) 145; (f) H. C. E. MANNERSKANTZ AND G. WILKINSON, *J. Chem. Soc.*, (1962) 4454.
- 17 M. H. B. STIDDARD, *J. Chem. Soc.*, (1962) 4712.
- 18 H. L. NIGAM, R. S. NYHOLM AND M. H. B. STIDDARD, *J. Chem. Soc.*, (1960) 1803.
- 19 M. H. B. STIDDARD, *J. Chem. Soc.*, (1963) 756.
- 20 J. LEWIS AND R. WHYMAN, *J. Chem. Soc.*, (1965) 5486.
- 21 H. L. NIGAM, R. S. NYHOLM AND M. H. B. STIDDARD, *J. Chem. Soc.* (1960) 1806.
- 22 J. LEWIS, R. S. NYHOLM, C. S. PANDE AND M. H. B. STIDDARD, *J. Chem. Soc.*, (1963) 3600.
- 23 (a) R. S. NYHOLM, M. R. SNOW AND M. H. B. STIDDARD, *J. Chem. Soc.*, (1965) 6570; (b) J. LEWIS, R. S. NYHOLM, C. S. PANDE, S. S. SANDHU AND M. H. B. STIDDARD, *J. Chem. Soc.*, (1964) 3009; (c) C. D. COOK, R. S. NYHOLM AND M. L. TOBE, *J. Chem. Soc.*, (1965) 4194.
- 24 W. HIEBER AND E. ROMBERG, *Z. Anorg. Chem.*, 221 (1935) 321.
- 25 R. D. PEACOCK, *Proc. Chem. Soc.*, (1957) 59.
- 26 J. LEWIS AND R. WHYMAN, *J. Chem. Soc.*, (1965) 6027.
- 27 E. O. FISCHER AND V. HRISTIU, *Chem. Ber.*, 95 (1962) 253.
- 28 (a) G. WILKINSON, *J. Am. Chem. Soc.*, 76 (1954) 209; (b) R. G. HAYTER, *Inorg. Chem.*, 2 (1963) 1031.
- 29 (a) T. S. PIPER AND G. WILKINSON, *J. Inorg. Nucl. Chem.*, 3 (1956) 104; (b) R. B. KING AND F. G. A. STONE, *Inorg. Syn.*, 7 (1963) 107; (c) E. O. FISCHER, *Inorg. Syn.*, 7 (1963) 136.
- 30 F. C. WILSON AND D. P. SHOEMAKER, *J. Chem. Phys.*, 27 (1957) 809.
- 31 E. W. ABEL, A. SINGH AND G. WILKINSON, *J. Chem. Soc.*, (1960) 1321.
- 32 R. B. KING, P. M. TREICHEL AND F. G. A. STONE, *Chem. Ind.*, (1961) 747.
- 33 W. R. CULLEN AND R. G. HAYTER, *J. Am. Chem. Soc.*, 86 (1964) 1030.
- 34 R. B. KING, *J. Am. Chem. Soc.*, 85 (1963) 1587.

- 35 E. O. FISCHER, O. BECKERT, W. HAFNER AND H. O. STAHL, *Z. Naturforsch.*, 10b (1955) 598.
36 F. A. COTTON AND B. F. G. JOHNSON, *Inorg. Chem.*, 3 (1964) 1609.
37 J. LOCKE AND J. A. MCCLEVERTY, *Chem. Commun. (London)*, (1965) 102.
38 S. HERZOG AND I. SCHNEIDER, *Z. Chem.*, 2 (1962) 24.
39 H. BEHRENS AND U. ANDERS, *Z. Naturforsch.*, 19b (1964) 767.
40 J. CHATT AND H. R. WATSON, *Proc. Chem. Soc.*, (1960) 243; *J. Chem. Soc.*, (1962) 2545.
41 J. LEWIS, R. S. NYHOLM AND P. W. SMITH, *J. Chem. Soc.*, (1962) 2592.
42 C. BROSSET, *Arkiv Kemi*, A20 (1945) 7; A22 (1946) 11.
43 F. A. COTTON AND T. E. HAAS, *Inorg. Chem.*, 3 (1964) 10. See also R. COLTON AND R. L. MARTIN, *Nature*, 207 (1965) 141.
44 J. C. SHELDON, *J. Chem. Soc.*, (1960) 1007, 3106.
45 R. J. GILLESPIE in S. KIRSCHNER (Ed.), *Advances in the Chemistry of the Coordination Compounds*, Macmillan, New York, 1961, p. 34.
46 L. D. CROSSMAN, D. P. OLSEN AND G. H. DUFFEY, *J. Chem. Phys.*, 38 (1963) 73; S. F. A. KETTLE, *Theor. Chim. Acta.*, 3 (1965) 211.
47 J. C. SHELDON, *Nature*, 184 (1959) 1210; *J. Chem. Soc.*, (1962) 410.
48 J. C. SHELDON, *J. Chem. Soc.*, (1961) 750.
49 F. A. COTTON AND N. F. CURTIS, *Inorg. Chem.*, 4 (1965) 241.
50 J. C. SHELDON, *J. Chem. Soc.*, (1963) 4183.
51 J. C. SHELDON, *J. Chem. Soc.*, (1964) 1287.
52 T. A. STEPHENSON, E. BANNISTER AND G. WILKINSON, *J. Chem. Soc.*, (1964) 2538.
53 D. LAWTON AND R. MASON, *J. Am. Chem. Soc.*, 87 (1965) 921.
54 G. BRAUER, *Handbook of Preparative Inorganic Chemistry*, Academic Press, New York, 2nd Edition, 1963, p. 259.
55 J. GAUNT, *Trans. Faraday Soc.*, 49 (1953) 1122; R. GLAUBER AND V. SCHOMAKER, *Phys. Rev.*, 89 (1953) 667; H. H. CLAASSEN, H. SELIG AND J. G. MALM, *J. Chem. Phys.*, 36 (1962) 2888; G. NAGARAJAN, *Australian J. Chem.*, 16 (1963) 906.
56 A. J. EDWARDS, R. D. PEACOCK AND R. W. H. SMALL, *J. Chem. Soc.*, (1962) 4486.
57 T. A. O'DONNELL AND D. F. STEWART, *J. Inorg. Nucl. Chem.*, 24 (1962) 309.
58 D. E. LAVALLE, R. M. STEELE, M. K. WILKINSON AND H. L. YAKEL, *J. Am. Chem. Soc.*, 82 (1960) 2433.
59 V. GUTMANN AND H. J. EMELEUS, *J. Chem. Soc.*, (1949) 2979, (1950) 2115.
60 V. GUTMANN AND K. H. JACK, *Acta Cryst.*, 4 (1951) 244.
61 Ref. 54, p. 1405.
62 S. Y. TYREE, *Inorg. Chem.*, 3 (1964) 149.
63 R. V. G. EWENS AND M. W. LISTER, *Trans. Faraday Soc.*, 34 (1938) 1358.
64 D. E. SANDS AND A. ZALKIN, *Acta Cryst.*, 12 (1959) 723.
65 R. COLTON AND I. B. TOMKINS, *Australian J. Chem.*, 18 (1965) 447.
66 S. M. HORNER AND S. Y. TYREE, *Inorg. Chem.*, 1 (1962) 947.
67 M. L. LARSON AND F. W. MOORE, *Inorg. Chem.*, 3 (1964) 285.
68 Ref. 54, p. 1404.
69 H. G. SCHNERING AND H. WOHLE, *Naturwissenschaften*, 50 (1963) 91.
70 W. KLEMM AND H. STEINBERG, *Z. Anorg. Chem.*, 227 (1936) 193.
71 P. J. H. CARNELL AND R. E. MCCARLEY, *U.S. At. Energy Comm.*, IS-487 (1962) [C. A., 60 (1964) 15406f].
72 Ref. 54, p. 1407.
73 D. BABEL AND W. RUDORFF, *Naturwissenschaften*, 51 (1964) 85.
74 J. LEWIS, D. MACHIN, R. S. NYHOLM, P. J. PAULING AND P. W. SMITH, *Chem. Ind.*, (1960) 259.
75 F. KLANBERG AND H. W. KOHLSCHÜTTER, *Z. Naturforsch.*, 15b (1960) 616.
76 C. DJORDJEVIC, R. S. NYHOLM, C. S. PANDE AND M. H. B. STIDDARD, *J. Chem. Soc. (A)*, (1966) 16.
77 R. J. GILLESPIE, *J. Chem. Soc.*, (1963) 4672, 4679.
78 D. A. EDWARDS AND G. W. A. FOWLES, *J. Chem. Soc.*, (1961) 24.
79 R. F. W. BADER AND A. D. WESTLAND, *Can. J. Chem.*, 39 (1961) 2306.
80 K. KNOX AND C. E. COFFEY, *J. Am. Chem. Soc.*, 81 (1959) 5.
81 G. B. HARGREAVES AND R. D. PEACOCK, *J. Chem. Soc.*, (1958) 4390.

- 82 J. R. GEICHMAN, E. A. SMITH AND P. R. OGLE, *Inorg. Chem.*, 2 (1963) 1012.
- 83 J. R. GEICHMAN, E. A. SMITH, S. S. TROND AND P. R. OGLE, *Inorg. Chem.*, 1 (1962) 661.
- 84 E. L. MUETTERTIES, *J. Am. Chem. Soc.*, 82 (1960) 1082, 6429.
- 85 W. WARDLAW AND H. W. WEBB, *J. Chem. Soc.*, (1930) 2100.
- 86 V. GUTMANN, *Z. Anorg. Chem.*, 269 (1952) 279.
- 87 W. L. GROENEVELD, *Rec. Trav. Chim.*, 71 (1952) 1152.
- 88 G. W. A. FOWLES, *Progr. Inorg. Chem.*, 6 (1964) 1, and references therein. See also G. W. A. FOWLES AND D. NICHOLLS, *Quart. Rev.*, 16 (1962) 19.
- 89 D. C. BRADLEY, R. K. MULTANI AND W. WARDLAW, *J. Chem. Soc.*, (1958) 4647.
- 90 H. FUNK, F. SCHMEIL AND H. SCHOLZ, *Z. Anorg. Chem.*, 310 (1961) 86; H. FUNK, M. HESSELBARTH AND F. SCHMEIL, *Z. Anorg. Chem.*, 318 (1962) 318.
- 91 A. ROSENHEIM AND C. NERNST, *Z. Anorg. Chem.*, 214 (1933) 209.
- 92 M. L. LARSON, *J. Am. Chem. Soc.*, 82 (1960) 1223.
- 93 R. E. MCCARLEY, B. G. HUGHES, J. C. BOATMAN AND B. A. TORÈ, in *Reactions of Co-ordinated Ligands and Homogeneous Catalysis*, *Advan. Chem. Ser.*, 37 (1963) 243.
- 94 W. M. CARMICHAEL, D. A. EDWARDS AND R. A. WALTON, *J. Chem. Soc.(A)*, (1966) 97.
- 95 E. A. ALLEN, B. J. BRISDON AND G. W. A. FOWLES, *J. Chem. Soc.*, (1964) 4531.
- 96 E. A. ALLEN, K. FEENAN AND G. W. A. FOWLES, *J. Chem. Soc.*, (1965) 1636.
- 97 D. A. EDWARDS AND G. W. A. FOWLES, *J. Less-Common Metals*, 4 (1962) 512.
- 98 Ref. 54, p. 1408.
- 99 R. E. MCCARLEY AND T. M. BROWN, *Inorg. Chem.*, 3 (1964) 1232.
- 100 G. B. HARGREAVES AND R. D. PEACOCK, *J. Chem. Soc.*, (1957) 4212.
- 101 (a) S. M. HORNER AND S. Y. TYREE, *Inorg. Chem.*, 2 (1963) 568; (b) B. J. BRISDON AND R. A. WALTON, *J. Inorg. Nucl. Chem.*, 27 (1965) 1101.
- 102 A. J. EDWARDS AND R. D. PEACOCK, *Chem. Ind.*, (1960) 1441.
- 103 A. J. EDWARDS, R. D. PEACOCK AND A. SAID, *J. Chem. Soc.*, (1962) 4643.
- 104 R. C. YOUNG AND R. A. LAUDISE, *J. Am. Chem. Soc.*, 78 (1956) 4861.
- 105 E. A. ALLEN, D. A. EDWARDS AND G. W. A. FOWLES, *Chem. Ind.*, (1962) 1026.
- 106 E. A. ALLEN, B. J. BRISDON, D. A. EDWARDS, G. W. A. FOWLES AND R. G. WILLIAMS, *J. Chem. Soc.*, (1963) 4649.
- 107 (a) K. H. LOHMANN AND R. C. YOUNG, *Inorg. Syn.*, 4 (1953) 97; (b) W. R. BUCKNALL, S. R. CARTER AND W. WARDLAW, *J. Chem. Soc.*, (1927) 512; (c) J. H. BALTHIS, *J. Inorg. Nucl. Chem.*, 24 (1962) 1017; (d) W. G. PALMER, *Experimental Inorganic Chemistry*, Cambridge University Press, 1959, p. 404.
- 108 W. WARDLAW AND A. J. I. HARDING, *J. Chem. Soc.*, (1926) 1592.
- 109 A. J. EDWARDS AND R. D. PEACOCK, *J. Chem. Soc.*, (1961) 4253.
- 110 C. BROSSET, *Arkiv Kemi*, 1 (1949) 353.
- 111 R. J. IRVING AND M. C. STEELE, *Australian J. Chem.*, 10 (1957) 490.
- 112 (a) N. ELLIOTT AND C. EPSTEIN, *J. Chem. Phys.*, 22 (1954) 634; (b) B. M. FIGGIS, J. LEWIS AND F. E. MABBS, *J. Chem. Soc.*, (1961) 3138.
- 113 G. B. HARGREAVES AND R. D. PEACOCK, *J. Chem. Soc.*, (1958) 3776.
- 114 P. C. H. MITCHELL, *Thesis*, Oxford, (1961).
- 115 W. WARDLAW AND R. L. WORMELL, *J. Chem. Soc.*, (1924) 2370; (1926) 1592; (1927), 130, 1087.
- 116 F. H. NICHOLLS, H. SAENGER AND W. WARDLAW, *J. Chem. Soc.*, (1931) 1443.
- 117 M. SCHMEISSER, *Angew. Chem.*, 67 (1955) 493.
- 118 M. C. STEELE, *Australian J. Chem.*, 10 (1957) 367.
- 119 P. KIRKEGAARD AND M. WESTERLUND, *Acta. Chem. Scand.*, 18 (1964) 2217.
- 120 A. ROSENHEIM, G. ABEL AND R. LEWY, *Z. Anorg. Chem.*, 197 (1931) 189.
- 121 E. M. SZUSTOROWICZ, *J. Struct. Chem.*, 4 (1963) 245, 273.
- 122 C. K. JØRGENSEN, *Inorganic Complexes*, Academic Press, 1963, pp. 60-61.
- 123 K. DEHNICKS AND J. STRAHLE, *Z. Anorg. Chem.*, 339 (1965) 171.
- 124 R. G. JAMES AND W. WARDLAW, *J. Chem. Soc.*, (1928) 2726.
- 125 H. FUNK AND H. BOHLAND, *Z. Anorg. Chem.*, 324 (1963) 168.
- 126 G. A. BARBIERI, *Atti Accad. Lincei*, 12 (1930) 55.
- 127 H. W. WILLARD AND R. C. TRUELKE, *J. Am. Chem. Soc.*, 57 (1935) 2609.

- 128 P. C. H. MITCHELL AND R. J. P. WILLIAMS, *J. Chem. Soc.*, (1960) 1912.
129 J. LEWIS, R. S. NYHOLM AND P. W. SMITH, *J. Chem. Soc.*, (1961) 4590.
130 P. C. H. MITCHELL, *J. Inorg. Nucl. Chem.*, 21 (1961) 382.
131 W. R. BUCKNALL AND W. WARDLAW, *J. Chem. Soc.*, (1927) 2981.
132 W. A. RAWLSON, *Australian Chem. Inst. J. Proc.*, 8 (1941) 42.
133 W. P. GRIFFITH, J. LEWIS AND G. WILKINSON, *J. Chem. Soc.*, (1959) 872.
134 N. H. FURMAN AND C. O. MILLER, *Inorg. Syn.* 3 (1950) 160.
135 I. M. KOLTHOFF AND N. J. TOMSICEK, *J. Phys. Chem.*, 40 (1936) 247.
136 (a) R. C. YOUNG, *J. Am. Chem. Soc.*, 54 (1932) 1402; (b) M. C. STEELE, *Australian J. Chem.*, 10 (1957) 404.
137 W. JAKOB AND Z. JAKOB, *Roczniki Chem.*, 36 (1962) 593, 601; W. JAKOB, A. KOSINSKA-SAMOTUS AND Z. STASICKA, *ibid.*, 611.
138 O. COLLEBERG, *Z. Anorg. Chem.*, 136 (1924) 249.
139 K. M. MIKHAILEVICH AND A. N. SERGEEVA, *Nauk. Zap. L'viv'sk. Politekh. Inst.*, 22 (1956) 11, C. A., 52 (1958) 13509^b.
140 W. P. GRIFFITH, *J. Chem. Soc.*, (1963) 3286.
141 W. L. MAGNUSSON, E. GRISWOLD AND J. KLEINBERG, *Inorg. Chem.*, 3 (1964) 88.
142 E. CREPAZ, *Gazz. Chim. Ital.*, 58 (1928) 391.
143 (a) W. HIEBER, R. NAST AND G. GEHRING, *Z. Anorg. Chem.*, 256 (1948) 169; (b) R. F. RILEY AND L. HO, *J. Inorg. Nuclear Chem.*, 24 (1962) 1121; (c) W. JAKOB, E. HEIMO AND A. KANAS, *Roczniki Chem.*, 37 (1963) 703.
144 J. L. HOARD AND H. H. NORDSIECK, *J. Am. Chem. Soc.*, 61 (1939) 2853.
145 E. G. BRAME, F. A. JOHNSON, E. M. LARSEN AND V. W. MELOCHE, *J. Inorg. Nucl. Chem.*, 6 (1958) 99.
146 (a) J. H. GRIFFITHS, J. OWEN AND I. M. WARD, *Proc. Roy. Soc. (London)*, A219 (1953) 529; (b) L. E. ORGEL, *J. Inorg. Nucl. Chem.*, 14 (1960) 136; (c) R. J. GILLESPIE, *Can. J. Chem.*, 39 (1961) 2336; (d) M. RANDIC, *J. Chem. Phys.*, 36 (1962) 2094; (e) R. M. GOLDING AND A. CARRINGTON, *Mol. Phys.*, 5 (1962) 377; (f) G. GLIEMAN, *Theor. Chim. Acta*, 1 (1962) 14; (g) E. KÖNIG, *Theor. Chim. Acta*, 1 (1962) 23; (h) J. R. PERUMAREDDI, A. D. LIEHR AND A. W. ADAMSON, *J. Am. Chem. Soc.*, 85 (1963) 249.
147 J. L. HOARD AND J. V. SILVERTON, *Inorg. Chem.*, 2 (1963) 235.
148 S. I. WEISSMAN AND M. COHN, *J. Chem. Phys.*, 27 (1957) 1440; see also refs. 146a, 145e.
149 S. F. A. KETTLE AND R. V. PARISH, *Spectrochim. Acta*, 21 (1965) 1087; see also H. STAMMREICH AND O. SALA, *Z. Elektrochem.*, 64 (1960) 741; 65 (1961) 149.
150 F. BASOLO AND R. G. PEARSON, *Mechanisms of Inorganic Reactions*, Wiley, London, 1965, p. 55.
151 A. W. ADAMSON AND J. R. PERUMAREDDI, *Inorg. Chem.*, 4 (1965) 247.
152 F. HOLZL AND G. I. XENAKIS, *Monatsh.*, 48 (1927) 689.
153 D. F. SHRIVER, *J. Am. Chem. Soc.*, 85 (1963) 1405.
154 A. W. ADAMSON, J. P. WALKER AND M. VOLPE, *J. Am. Chem. Soc.*, 72 (1950) 4030.
155 S. AŠPERGER, I. MURATI AND D. PAVLOVIĆ, *J. Chem. Soc.*, (1960) 730.
156 P. C. H. MITCHELL AND K. MACKAY, unpublished work.
157 R. L. WOLFGANG, *J. Am. Chem. Soc.*, 74 (1952) 6144; P. GEORGE AND D. H. IRVINE, *J. Chem. Soc.*, (1954) 587.
158 D. GRDENIĆ AND B. KORPAR-COLIG, *Proc. Chem. Soc. (London)*, (1963) 308.
159 M. L. LARSON AND F. W. MOORE, *Inorg. Chem.*, 1 (1962) 852; 2 (1963) 887.
160 G. T. MORGAN AND R. A. S. CASTELL, *J. Chem. Soc.*, (1928) 3262.
161 K. CHRIST AND H. L. SCHAFER, *Angew. Chem.*, 2 (1963) 97.
162 T. G. DUNNE AND F. A. COTTON, *Inorg. Chem.*, 2 (1963) 263.
163 (a) W. WARDLAW AND W. H. PARKER, *J. Chem. Soc.*, (1925) 1311; (b) H. M. SPITTLE AND W. WARDLAW, *J. Chem. Soc.*, (1929) 792.
164 W. WENDLING AND R. RÖHMER, *Bull. Soc. Chim. France*, (1964) 360.
165 E. WENDLING, *Bull. Soc. Chim. France*, (1965) 437.
166 G. M. FRENCH AND J. H. GARSIDE, *J. Chem. Soc.*, (1962) 2006.
167 G. J. SUTTON, *Australian J. Chem.*, 15 (1962) 232.
168 M. C. STEELE, *Australian J. Chem.*, 10 (1957) 489.

- 169 L. F. LINDOY, S. E. LIVINGSTONE AND T. M. LOCKYER, *Australian J. Chem.*, 18 (1965) 1549.
- 170 F. JELLINEK, *Nature*, 192 (1961) 1065; see also A. V. GOROKH, L. N. RUSAKOV AND A. A. SAVINSKAYA, *Dokl. Akad. Nauk. S.S.S.R.*, 156 (1964) 541.
- 171 J. C. WILDERVANCK AND F. JELLINEK, *Z. Anorg. Chem.*, 328 (1964) 309.
- 172 Ref. 54, p. 1416.
- 173 J. T. SPENCE AND H. M. Y. CHANG, *Inorg. Chem.*, 2 (1963) 319.
- 174 A. B. BLAKE, F. A. COTTON AND J. S. WOOD, *J. Am. Chem. Soc.*, 86 (1964) 3024.
- 175 (a) R. B. KING, *Inorg. Chem.*, 2 (1963) 641; (b) A. DAVISON, N. EDELSTEIN, R. H. HOLM AND A. H. MAXI, *J. Am. Chem. Soc.*, 86 (1964) 2799.
- 176 (a) G. N. SCHRAUZER, V. P. MAYWEG, H. W. FINCK, U. MÜLLER-WESTERHOFF AND W. HEINRICH, *Angew. Chem. Internat. Edit.*, 3 (1964) 381; (b) G. N. SCHRAUZER AND V. P. MAYWEG, *Z. Naturforsch.*, 19b (1964) 192; (c) G. N. SCHRAUZER, V. P. MAYWEG AND W. HEINRICH, *Inorg. Chem.*, 4 (1965) 1615; (d) G. N. SCHRAUZER, H. W. FINCK AND V. P. MAYWEG, *Angew. Chem. Internat. Edit.*, 4 (1965) 146.
- 177 M. GERLOCH, S. F. A. KETTLE, J. LOCKE AND J. A. MCCLEVERTY, *Chem. Commun. (London)*, (1966) 29.
- 178 (a) T. W. GILBERT AND E. B. SANDELL, *J. Am. Chem. Soc.*, 82 (1960) 6087; (b) E. I. STIEFEL AND H. B. GRAY, *J. Am. Chem. Soc.*, 87 (1965) 4012.
- 179 R. EISENBERG AND J. A. IBERS, *J. Am. Chem. Soc.*, 87 (1965) 3776.
- 180 V. B. EVDOKMOV, V. V. ZELENSTOV, I. D. KOLLI, W.-H. T'ANG AND V. I. SPITSYN, *Dokl. Akad. Nauk. S.S.S.R.*, 145 (1962) 1282; V. I. SPITSYN, Z. D. KOLLI AND W.-H. T'ANG, *Zh. Neorg. Khim.*, 9 (1964) 99.