

MAGNETIC RESONANCE METHODS IN THE STUDY OF THE ELECTRONIC STRUCTURE OF TRANSITION METAL COMPLEXES

D.R. EATON and K. ZAW

*Department of Chemistry, McMaster University,
Hamilton, Ontario (Canada)*

(Received January 5th, 1971)

CONTENTS

A Introduction	197
B ESR theory	199
C Some ESR results	203
D NMR theory	210
E Some NMR results	214
F Conclusion	221
References	221

A INTRODUCTION

The description of the metal–ligand bonding in transition metal complexes presents a continuing problem for inorganic chemists. Theoretical work in this area has progressed from the valence bond approach of Pauling through crystal field theory to the current molecular orbital approach. Our qualitative understanding of bonding in these complexes can now be described as quite good. They are, however, complicated, many-electron systems and quantitative calculations are still far from reliable. From the point of view of the chemist, much of the interest in this subject lies in just those finer details of the electronic structure which are most inaccessible to theoretical calculation. Experimental approaches to the problem are therefore still of considerable value. Of the various physical techniques which provide information in this area, magnetic resonance methods are probably the most powerful. The full potentialities of the technique are only realized when the complexes are paramagnetic, in which case electron spin–nuclear spin interactions provide a very detailed probe of the electronic structure. It is this area of the subject which will be described in the present review.

In principle, a paramagnetic metal complex offers a choice of magnetic resonance techniques. Transitions involving a change in electron spin quantum number may be studied by ESR and those involving a change in nuclear spin quantum number by NMR. A variety of double resonance techniques (ENDOR, the Overhauser effect) complement the straightforward experiments and related methods such as Nuclear Quadrupole Resonance and




Fig 1 NMR and ESR transitions for $S = 1/2$, $I = 1/2$ system $\cdots \rightarrow$ NMR, \longrightarrow ESR

Mossbauer spectroscopy where applicable, contribute valuable supplementary data. However, in the present article only simple ESR and NMR experiments will be considered.

There are very few systems amenable to study by both ESR and NMR. The primary reason for this can be seen by reference to Fig 1. In this figure the magnetic energy levels for a simple system comprising one electron and one nucleus of spin $1/2$ are shown. Two ESR transitions differing in energy by A , the hyperfine coupling constant, are expected. There are also two NMR transitions, one at a frequency given by $h\nu = g_N\beta_N H - 1/2(A)$ and one at $g_N\beta_N H + 1/2(A)$. The appearance of the spectra, however, depends critically on the electron spin relaxation time T_{1e} . The necessary condition for observing well-resolved hyperfine structure in the ESR spectrum is that $1/T_{1e} \ll A$, i.e. that the relaxation time be long. For the NMR spectrum the nuclear spin relaxation time determines the spectral properties. For a paramagnetic complex the dominant nuclear relaxation mechanism is likely to arise from the fluctuating magnetic fields produced at the nucleus by changes in the orientation of the electron spin. Thus the nuclear relaxation time is determined by the electron spin relaxation time. Furthermore, the limitation on the relaxation time necessary for the observation of sharp NMR lines is much more rigorous than that for ESR. Typically NMR lines are of the order of Hz broad whereas ESR line widths are measured in MHz. Hence the NMR transitions indicated in Fig 1 in practice are never observed. There are, however, certain conditions under which sharp-line NMR spectra can be observed. These arise in cases where the electron spin relaxation time is very short, i.e. $1/T_{1e} \gg A$. In these circumstances, the rapidly fluctuating magnetic fields produced by the electron spin are no longer effective in producing nuclear relaxation. The fluctuations are of such a high frequency that they have negligible components of the correct frequencies to induce nuclear transitions. In these circumstances, the nucleus, instead of experiencing

either the hyperfine magnetic field corresponding to interaction with an α electron spin or the hyperfine field corresponding to interaction with a β spin, experiences an average local field corresponding to the mean value, $\langle S_z \rangle$, of the electron spin polarization. This mean value is not equal to zero since the population of the α and β states are not equal. Hence there is a chemical shift resulting from the electron spin–nuclear spin interaction. Thus it is apparent that the conditions for the ESR experiment ($1/T_c \ll A$) and for the NMR experiment ($1/T_c \gg A$) are incompatible. The two types of experiments are not, however, completely exclusive. Relaxation times are temperature-dependent and in some instances it is possible to carry out ESR experiments in solids at low temperatures where T_c is long and NMR experiments in solutions at high temperatures where T_c is short. Examples of such cases may be found below.

Not only do the two forms of magnetic resonance differ in their practical applicability, they also provide different, but complementary, information for the inorganic chemist. In ESR, attention is focussed on the metal ion and its immediate environs and the question is asked “how have the properties of the free ion been modified by interaction with the ligands?” In NMR, the focus is on the ligands and the question is “how has the electronic structure of the ligand been modified by interaction with the metal ion?”. There is an obvious symmetry to the two approaches and the objective of the present review is to draw attention to some of the interlocking facets of the two fields. The fundamental physics of the situation could probably be best illustrated by reviewing a number of the very elegant experiments which have been carried out on relatively simple systems such as transition metal halides. We choose, however, to adopt a more chemical criterion and to discuss instead work on transition metal chelates, the appeal of which is more chemical in nature. Initially, some of the relevant theory will be briefly reviewed.

B ESR THEORY

Basically the interpretation of the ESR spectrum of an isolated transition metal ion yields values for two different kinds of parameter. The spectroscopic splitting constants (g values) describe the frequencies (or energies) required to bring about transitions between the different electron spin states. The hyperfine coupling constants (A) describe the electron spin–nuclear spin interactions which give rise to the hyperfine structure observed in the spectrum. Both of these parameters may be anisotropic, i.e. their values may depend on the orientation of the molecule with respect to the applied magnetic field. In addition, ions containing more than one unpaired electron will exhibit zero-field splitting effects due to electron spin–electron spin interactions and to spin orbit coupling. These interactions (which are also usually anisotropic) are described by the zero-field splitting parameters D and E . All of these parameters are affected by the presence of ligands around the central metal atom. The ligand effects can be subdivided into two kinds. Initially, one may adopt the crystal field approximation of regarding the function of the ligands as one of providing an electrostatic field which will perturb the electronic structure of the metal ion. This approach proves inadequate for the type of complexes we wish to discuss. The breakdown occurs because chemically interesting complexes are usually covalent rather than ionic in character. In these circumstances, it is no longer satisfactory to consider only a basis set of atomic orbitals centred on the metal ion. Instead one must

deal with molecular orbitals expressed as appropriate linear combinations of metal and ligand atomic orbitals. Using molecular orbital theory, a set of secondary parameters (usually written α, β etc.), which are the coefficients of the various atomic orbitals in the expansions for the molecular orbitals, can be derived from the primary parameters g and A . It is these secondary parameters which are of prime interest to the inorganic chemist since they provide detailed quantitative data on the nature of the metal-ligand bonding.

The basic theory pertinent to the application of ligand field theory to ESR has been developed in a number of excellent textbooks¹ and review articles². The g values of a complex are determined by the expectation values of the components of the vector operator $L + 2S$ (Allowing for relativistic corrections the 2 in this expression becomes 2.0023). In many cases the orbital angular momentum in the ground state of the molecule is completely quenched and an isotropic g value of 2.0023 is obtained to zero-order approximation. This will always be the case if the complex is of sufficiently low symmetry that the orbitals containing the odd electrons are non-degenerate. This approximation, however, does not usually suffice for transition metal complexes. Spin-orbit coupling leads to a mixing of excited state configurations with the ground state configuration and it is necessary to find the expectation values of $L + 2S$ using these corrected ground-state wave functions. Typically, therefore, perturbation theory leads to g values of the form

$$g_{\parallel} = 2 \left(1 - \frac{x\lambda}{\Delta} \right) \quad g_{\perp} = 2 \left(1 - \frac{y\lambda}{\Delta} \right)$$

for an orbitally non-degenerate ground state, where λ is the spin-orbit coupling parameter, Δ is an energy difference between the ground-state configuration and an excited-state configuration and x and y are simple numerical factors. In principle, λ and Δ are obtainable from optical spectroscopy. In practice, it is found that the g values measured do not agree with those calculated unless smaller values of λ are assumed than those obtained for the free ions. Typically, a 20 or 30% reduction in the spin-orbit coupling constant is involved. This effect is attributed to covalency in the metal-ligand bonding. In order to calculate the g values it is necessary to start with the correct molecular orbitals rather than with simple atomic d orbitals. Since such a basis set will allow the electrons to spend part of their time on the ligand rather than on the metal, it is apparent that the effects of spin-orbit coupling will be reduced. The equations for the g values will now contain the coefficients of the atomic orbitals (α, β) in the initial molecular orbitals as well as the parameters noted above. Analysis of the data, therefore, provides values for the molecular orbital coefficients. A number of individual calculations will be discussed in more detail in the sections below.

The hyperfine coupling between electron spins and nuclear spins arises from two sources. The first of these is a direct dipolar interaction. The magnetic moment of the electron produces a field at the nucleus, the direction of which depends on the orientation of the electron spin. The situation is illustrated in Fig. 2. The most notable feature of this interaction is that it is anisotropic, i.e. the magnitude depends upon the orientation of the molecule with respect to the applied magnetic field. Thus, referring to Fig. 2, let us suppose that the electron is located at a fixed position on the x axis. It is apparent that application of the field parallel to the y axis (Fig. 2(a)) results in a dipole-dipole interaction of opposite sign to that obtained by applying the field in the x direction (Fig. 2(b)). In this




Fig. 2 Dipolar interactions between an electron spin and a nuclear spin fixed in space

In the illustration we have kept the position of the molecule fixed with respect to the x and y axes and changed the direction of the field. In the actual experiment the direction of the field is kept constant and that of the molecule, fixed in a crystalline matrix, is varied. The two are obviously equivalent. A model in which the electrons are kept fixed in space is not a valid one for real molecules. However, if the odd electron occupies an orbital with directional properties, e.g. an atomic p or d orbital, it is apparent that some directions will be preferred over others. Analysis of the problem shows that the dipole-dipole coupling is proportional to the average value $\langle (3 \cos^2 \theta - 1)/r^3 \rangle$ where θ is the angle between the direction of the magnetic field and a line joining the odd electron and the nucleus, and r is the separation of the electron and metal nucleus. Numerical values of this average can be readily calculated for the various d orbitals. In a covalent metal complex the odd electrons will be partly delocalized to the ligands. The value of $\langle 1/r^3 \rangle$ for electrons located on the ligand will be much smaller than for an electron located in a metal d orbital and dipolar contributions from such a source are often neglected. Thus in a covalent complex, the anisotropic contribution to the hyperfine splitting will be reduced by a factor α^2 , where α is the coefficient of the d orbital in the molecular orbital containing the unpaired electron. This effect, therefore, provides a method of estimating α independent of that previously discussed involving the g values.

The second source of hyperfine coupling is the Fermi Contact Interaction. This mechanism demands a finite probability of finding the odd electron at the metal nucleus, i.e. there must be spin density in a metal S orbital. Since S orbitals are involved, it is an isotropic interaction. An important consequence of this is that the isotropic hyperfine splitting is the same in solution, where the molecules are freely tumbling, as it is in the solid state. The dipolar coupling, on the other hand, averages to zero when the molecules are freely tumbling and therefore does not contribute to the solution spectrum. A recent paper by McGarvey³ has considered isotropic hyperfine coupling in metal complexes in some detail. Unpaired electrons in d orbitals do not contribute directly to this effect. The major source of the interaction is thought to be spin polarization of electron pairs in

filled metal orbitals notably the 3s orbital in the first transition series. In this respect the situation is similar to that existing in aromatic radicals for which the well-known McConnell relationship^{A6}

$$a_H = Q\rho_c$$


which states that the amount of spin polarization is proportional to the $p\pi$ spin density, holds. There are, indeed, many complexes for which the metal hyperfine coupling provides a measure of the d orbital spin density.^{2a} In such complexes, increasing covalency results in increasing delocalization of the unpaired electrons and decreasing metal hyperfine splitting. Estimates of the molecular orbital parameters obtained in this way are consistent with those obtained by the methods outlined above. There are, however, some notable exceptions, especially amongst Cu^{II} complexes, where the metal hyperfine coupling appears to be increasing with increasing covalency.⁶ The most plausible interpretation involves interaction with higher unfilled metal S orbitals and a specific case will be considered in more detail below. Obviously a degree of caution is necessary in this area.

Finally, in some complexes, hyperfine splitting additional to that expected from the metal nucleus has been observed. This is attributed to interaction with ligand nuclei. Usually, only atoms directly bonded to the metal are involved although there are occasional exceptions to this generalization. In principle, both isotropic (contact) and anisotropic (dipolar) interactions are involved but since the odd electrons are predominantly on the metal so that $\langle 1/r^3 \rangle$ for ligand atoms is small the isotropic contribution usually predominates. A large isotropic ligand splitting is expected in cases where the spin-containing orbital can mix directly with a ligand S orbital. This is the case, for example, for Cu^{II} complexes where the odd electron occupies a σ antibonding orbital. If it is assumed that the ligand orbitals comprise sp^2 hybrids, or some other suitable combination of ligand atomic orbitals, the ligand hyperfine splitting is dominated by the S component of the hybrid and the extent of metal-ligand orbital mixing can be calculated. Usually, parameters obtained by this method agree well with those obtained from the g values indicating that the above assumptions are correct. Line-width considerations restrict the observation of ligand hyperfine couplings of less than a gauss or so and as a result, unless there is direct delocalization of spin to the ligand S orbital as in the above example, ligand splitting will not be seen. Thus, in d^1 vanadyl complexes, the odd electron is in an orbital orthogonal to the ligand σ orbitals and no ligand hyperfine splitting has been reported.

Finally, before discussing detailed results, it should be made clear that bonding parameters derived from ESR results refer only to ligand interactions with metal orbitals containing unpaired electrons. Energetically, the most important interactions are, of course, with the s , p and d metal orbitals which hybridize to a set with appropriate geometrical properties and give rise to completely filled bonding molecular orbitals. We are dealing here, therefore, with weaker additional interactions which, although they may contribute relatively little to the total bonding energy, nevertheless provide insight into the nature of metal-ligand interactions. Exactly the same considerations apply to the discussions of NMR contact shifts.

C. SOME ESR RESULTS

As indicated in the above paragraphs, the determination of bonding parameters from ESR data is, in principle, quite straightforward. In practice the algebraic complexity of the problem tends to obscure this simplicity. Initially, therefore, we will consider a Cu^{II} complex without ligand hyperfine splitting. Cu^{II} has only a single unpaired electron, and its relaxation properties are such that most complexes give relatively sharp lines at room temperature. As a result, a wide range of complexes have been studied. Maki and McGarvey⁴ have carried out a single-crystal ESR study of Cu^{II} acetylacetonate (structure I). The acetylacetonate ligand is typical of the class which chemically speaking give "co-



valent" compounds and the ease of formation of its complexes has led to many studies of their physical properties. Hence we will be able to compare these ESR results with NMR data on other transition metal acetylacetonates. The Cu^{II} compound has planar geometry and its crystal structure is isomorphous with the corresponding diamagnetic Pd^{II} complex. Maki and McGarvey, therefore, made their measurements on Cu^{II} acetylacetonate diluted to 0.5 mole % in the Pd^{II} compound. The use of such magnetically dilute crystals eliminates dipole-dipole interactions which would otherwise broaden the ESR lines. The point group symmetry of the molecule is D_{2h} but the rhombic distortions are sufficiently small to allow the use of the full D_{4h} square planar symmetry in the theoretical treatment. The relevant anti-bonding molecular orbitals can then be written

$$\begin{aligned}
 B_{1g} & \quad \alpha d_{x^2-y^2} - \frac{\alpha'}{2} (-\sigma_x^a + \sigma_y^b + \sigma_z^c - \sigma_y^d) \\
 B_{2g} & \quad \beta_1 d_{xy} - \frac{(1-\beta_1^2)^{1/2}}{2} (p_y^a + p_x^b - p_y^c - p_x^d) \\
 A_{1g} & \quad \alpha_1 d_{3z^2-r^2} - \frac{\alpha'_1}{2} (\sigma_x^a + \sigma_y^b - \sigma_z^c - \sigma_y^d) \\
 & \quad \beta d_{xz} - \frac{(1-\beta^2)^{1/2}}{2^{1/2}} (p_z^a - p_z^c) \\
 E_g & \quad \beta d_{yz} - \frac{(1-\beta^2)^{1/2}}{2^{1/2}} (p_z^b - p_z^d)
 \end{aligned}$$

The superscripts a, b, c and d label the four ligand oxygens and the ligand σ orbitals are hybrids of the oxygen s and p atomic orbitals of the form

$$\sigma_k^m = n P_k^m + (1-n^2)^{1/2} S^m$$

The coefficients α and α_1 give a measure of the ligand contribution to the σ orbitals and β and β_1 of the out-of-plane and in-plane π mixing respectively. The objective is to determine these coefficients. Overlap has been neglected for the π orbitals. The odd electron is in the B_{1g} orbital. These orbitals are eigenfunctions if spin-orbit coupling is neglected. They can, however, be mixed by the spin-orbit coupling $L \cdot S$. It is, therefore, necessary to determine first a new basis set treating \mathcal{H}_{LS} as a perturbation. This basis set can then be operated on by the appropriate magnetic perturbation terms and the results expressed in the form of the spin Hamiltonian.

$$\mathcal{H} = \beta_0 [g_{\parallel} H_z S_z + g_{\perp} (H_x S_x + H_y S_y)] + A S_z I_z + B (S_x I_x + S_y I_y) + Q [I_z^2 - \frac{1}{3} I(I+1)] - \gamma \beta_N H \cdot I$$

In this Hamiltonian, the first term represents the electron Zeeman effect, the second and third arise from hyperfine interactions, the fourth from nuclear quadrupole interactions and the final term from the nuclear Zeeman effect. Spin-orbit coupling mixes the B_{2g} and E_g orbitals with the spin-containing B_{1g} orbital but not the A_{1g} orbital. The expressions for the spin Hamiltonian parameters therefore contain α , β and β_1 but not α_1 . Solution of the problem gives

$$g_{\parallel} = 2.0023 - 8\rho [\alpha\beta_1 - \frac{1}{2}\alpha'(1-\beta_1^2)^{1/2} T(n)]$$

$$g_{\perp} = 2.0023 - 2\mu [\alpha\beta - \frac{1}{\sqrt{2}}\alpha'(1-\beta^2)^{1/2} T(n)]$$

$$A = P[-\frac{4}{7}\alpha^2 - \kappa + (g_{\parallel} - 2) + \frac{3}{7}(g_{\perp} - 2) - 4\rho\alpha'(1-\beta_1^2)^{1/2} T(n) - \frac{3\sqrt{2}}{7}\mu\alpha'(1-\beta^2)^{1/2} T(n)]$$

$$B = P[\frac{2}{7}\alpha^2 - \kappa + \frac{11}{14}(g_{\perp} - 2) - \frac{11\sqrt{2}}{14}\mu\alpha'(1-\beta^2)^{1/2} T(n)]$$

$$Q' = -\frac{e^2 q \alpha^2 \langle r^{-3} \rangle_0}{7}$$

where

$$\rho = \frac{\lambda_0 \alpha \beta_1}{E_{xy} - E_0} \quad \text{and} \quad \mu = \frac{\lambda_0 \alpha \beta}{E_{xz, yz} - E_0}$$

$T(n)$ is a function of the ligand atomic orbitals which can be calculated if n is known. It is assumed that $n = (2/3)^{1/3}$ corresponding to sp^2 hybridization. λ_0 is the spin-orbit coupling constant for the free Cu^{2+} ion and the quantity P given by

$$P = 2\gamma\beta_0\beta_N \langle r^{-3} \rangle_0$$

also refers to the free Cu^{2+} ion. In the expressions for the hyperfine coupling constants, the first term is the dipolar contribution, the second the Fermi contact contribution and the remaining terms are second-order corrections resulting from mixing in of the excited states. A relationship between α and α' is also required and this is given as

$$\alpha' \approx (1 - \alpha^2)^{1/2} + \alpha S$$

where S , the overlap between $d_{x^2-y^2}$ and the ligand σ orbitals is calculated to be 0.094.

$\langle r^{-3} \rangle_0$ refers to the orbitals of the free divalent ion and is calculated to be 7.25 au. The measured experimental quantities are

$$\begin{aligned} g_{\parallel} &= 2.266_1 & g_{\perp} &= 2.053_5 \\ A &= -1.60 \times 10^{-2} \text{ cm}^{-1} & B &= -0.19_5 \times 10^{-2} \text{ cm}^{-1} \\ Q' &= 7 \times 10^{-4} \text{ cm}^{-1} \end{aligned}$$

From optical spectroscopy, the further information obtained is

$$E_{xy} - E_0 = 15,000 \text{ cm}^{-1} \quad E_{xz, yz} - E_0 = 25,000 \text{ cm}^{-1} \quad \lambda_0 = -828 \text{ cm}^{-1}$$

Sufficient information is now available to calculate the bonding parameters. The results obtained are

$$\alpha^2 = 0.81 \quad \beta^2 = 0.99 \quad \beta_1^2 = 0.85 \quad \kappa = 0.33$$

This value of κ agrees well with that obtained from the isotropic hyperfine splitting of Cu^{2+} in ionic salts and confirms the assumption implicit in the above treatment that, in this case, the isotropic splitting is not sensitive to the degree of covalency. The qualitative deduction from the bonding parameters is that the B_{1g} and B_{2g} orbitals have significant ligand character but the E_g orbital is largely metallic. We will compare these conclusions with those obtained for acetylacetonates of the earlier transition metals below.

It is interesting to ask how these interactions are modified by substitution on the acetylacetonate ligand. This question could be answered by carrying out single-crystal studies on substituted Cu^{II} acetylacetonates analogous to the study described above. Experimentally, this would be a rather formidable task and as yet it has not been undertaken. Measurements on solutions or on polycrystalline solids are much less tedious but still yield a limited amount of information on the bonding. The work of Kuska and Rogers⁵ on substituted Cu^{II} acetylacetonates serves to illustrate the possibilities.

Analysis of solution ESR spectra yield two parameters, namely the average g value ($g = \frac{1}{3}(g_{\parallel} + 2g_{\perp})$) and the isotropic hyperfine coupling constant, A . It is apparent from the expressions given previously for g_{\parallel} and g_{\perp} that bonding parameters cannot be obtained directly from the average g . If a McConnell-type relationship exists for the isotropic hyperfine coupling it should give a measure of the spin density in the Cu atom and hence determine the covalency of the spin-containing orbital. Kuska and Rogers⁵ have investigated this possibility and have obtained the results shown in Table 1. Initially, they assumed that A would be given by an equation

$$A = P[-\alpha^2 \kappa_0 + (g - 2.0023) + \text{second-order terms}]$$

where κ_0 is the isotropic coupling constant for the free Cu^{2+} ion and α has been defined above. The values of α^2 obtained using this equation are listed in the table as "apparent α^2 ". These values imply that substitution with electron-withdrawing groups such as CF_3 leads to an increase in the amount of spin on the ligand, i.e. make the bond less ionic. This is contrary to what would be expected on chemical grounds and is not supported by polarographic data on the same compounds. It was concluded that the initial assumption that the hyperfine coupling gives a measure of the metal spin density was in error. McGarvey³ has carried out a perturbation treatment to investigate the effects of coval-

TABLE I

ESR hyperfine splitting constants and other data for copper acetylacetonates

Substituents			ESR A value ($\text{cm}^{-1} \times 10^4$)	g Value	Apparent α^2	Anisotropy α^2
R'	R''	R'''				
CH_3	CH_3	CH_3	78.5	2.121	0.784	0.121
CH_3	H	CH_3	77.2	2.123	0.779	0.753
CH_3	Cl	CH_3	76			
Phenyl	H	CH_3	76.4	2.124	0.777	0.795
Phenyl	H	Phenyl	75.3	2.124	0.770	
CF_3	H	CF_3	54.7	2.150	0.698	0.828


ency on the hyperfine coupling and has confirmed this conclusion. The essence of the problem is that the participation of the $4s$ orbital as well as that of the $3d$ orbitals must be considered. Thus κ_0 will not be a constant but is made up of two terms, one representing spin density in the $3d$ orbital and one spin density in the $4s$ orbital. Kuska and Rogers suggest that the signs of these contributions are opposite and since both will depend on the covalency it follows that the isotropic coupling does not provide a direct measure of α^2 .

In a later paper, Kuska et al.⁶ have obtained values for the anisotropic hyperfine coupling constants of these substituted Cu^{II} acetylacetonates by examining the spectra obtained in frozen chloroform solutions. Combination of this data with the isotropic splitting constants obtained in solution gives the dipolar contribution to the hyperfine splitting. This quantity should be proportional to α^2 and rearrangement of previous equations gives

$$\alpha^2 = \frac{7}{4} \left[\frac{A_{\parallel}}{P} - \frac{|A|}{P} + \frac{2}{3}g_{\parallel} - \frac{5}{21}g_{\perp} - \frac{6}{7} \right]$$

Thus α^2 can be obtained in a manner which does not depend on the mechanism of the contact interaction. Some values of α^2 obtained from the dipolar interactions are also shown in Table I. It is apparent that the trend is the opposite of that obtained from the isotropic A 's and is thus consistent with the polarographic and other data mentioned above. It is interesting to note that the spectrum of the diphenyl acetylacetonate (compound 4 of Table I) is found to be complicated by additional hyperfine splitting from the phenyl protons. No detailed analysis of this additional splitting was achieved.

An example of the successful analysis of hyperfine splitting from ligand nuclei is provided by the work of Maki and McGarvey⁷ on Cu^{II} bis salicylaldehyde imine. This compound has the structure



It was examined as a single crystal magnetically dilute in the corresponding nickel compound. To simplify the spectrum, isotopically pure ^{63}Cu was used. It is to be anticipated that the four lines due to the copper hyperfine splitting will be further split by the two nitrogen atoms directly attached to the metal. A five-line pattern is predicted. In fact a more complicated eleven-line pattern is observed. The additional splitting can be attributed to a pair of nuclei with spin $1/2$ which in the present case can only be protons. The most likely origin of the splitting might be thought to be the N-H protons since these are closest to the metal atom but deuteration disproved this possibility. However, methylation of the adjacent carbon did eliminate the additional splitting which must therefore be attributed to the proton marked H*. Analysis of the spectrum yielded the parameters

$$\begin{aligned} g_x &= 2.040_2 & g_y &= 2.050_0 & g_z &= 2.200_4 \\ A &= -1.85 \times 10^{-2} \text{ cm}^{-1} & B &= -0.21 \times 10^{-2} \text{ cm}^{-1} \\ A(^{14}\text{N}) &= 11.1 \text{ gauss} & A(^1\text{H}) &= 5.5 \text{ gauss} \end{aligned}$$

The similarity of g_x and g_y shows that the Cu-N and Cu-O bonds are similar in character. These values yield the bonding parameters

$$\alpha^2 = 0.83 \quad \beta^2 \geq 0.91 \quad \beta_1^2 = 0.74 \quad \kappa = 0.34$$

Comparison with the values previously given for Cu^{II} acetylacetonate shows that the only significant difference is a rather higher ligand contribution to the in-plane π orbitals. It appears that the contact contribution is most important for the nitrogen hyperfine splitting—dipolar interactions are estimated to contribute only $\sim 1.5\%$ of the total splitting. The contact part is given by

$$W = \left(\frac{\alpha'^2}{4} \right) (2\gamma\beta_0\beta_N) \frac{-8\pi}{9} \delta(r)$$

where α' is the coefficient of the nitrogen orbital in the spin-containing molecular orbital. A value of $\alpha'^2 = 0.25$ (consistent with $\alpha^2 = 0.83$ given above when allowance is made for overlap) gives a calculated coupling of $10.8 \times 10^{-4} \text{ cm}^{-1}$ compared with an experimental value of $10.4 \times 10^{-4} \text{ cm}^{-1}$. The spin-containing orbital has symmetry B_{1g} (predominantly $d_{x^2-y^2}$). It should be noted that the N-H proton lies near a node of this orbital but that the C-H proton, which shows hyperfine coupling, lies close to a direction of maximum electron density.

The d^9 configuration found in the above copper complexes is particularly favourable for ESR studies of metal-ligand bonding. The d^1 configuration found in vanadyl complexes is also favourable. The reasons are similar. It is an $S = 1/2$ ion and in the relatively low-symmetry vanadyl complexes with orbitally non-degenerate ground states the spectra are easy to observe. Kivelson and Lee⁸ have carried out a study of vanadyl acetylacetonate. They use a set of molecular orbitals similar to those used by Balhausen and Gray⁹ in their calculations on this molecule. These orbitals have the form

$$\begin{aligned} \psi_{b_2} &= \beta_2 d_{xy} + \frac{1}{2} \beta'_2 (P_{y_1} + P_{x_2} - P_{y_3} - P_{x_4}) \\ \psi_{b_1} &= \beta_1 d_{x^2-y^2} + \frac{1}{2} \beta_1 (\sigma_1 - \sigma_2 + \sigma_3 - \sigma_4) \end{aligned}$$

$$\psi_{e_{\pi x}} = \epsilon_{\pi} d_{xz} + \epsilon'_{\pi} P_{x5} + \frac{1}{\sqrt{2}} \epsilon''_{\pi} (P_{z1} - P_{z2})$$

$$\psi_{a_1} = \alpha_1 (d_{z^2} + S_0) + \alpha'_1 \sigma_5$$

$$\psi_{a_{||}} = \alpha_{||} (d_{z^2} - S_0) - \frac{1}{2} \alpha'_{||} (\sigma_1 + \sigma_2 + \sigma_3 + \sigma_4)$$

The symmetry of the complex is strictly C_{2v} but the assumption of C_{4v} symmetry implicit in the above orbitals is justified by the rather small rhombic distortion. The subscripts 1-4 refer to the oxygens of the acetylacetonate ligands and 5 to the vanadyl oxygen. As written above, these are all bonding orbitals, $a_1 a_{||}$ and b_1 being σ bonding, b_2 in-plane π bonding and e out-of-plane π bonding. The coefficients in the corresponding anti-bonding orbitals are denoted by stars. All these anti-bonding orbitals are unoccupied except for b_2^* which is singly occupied. Expressions can be derived for $g_{||}$, g_{\perp} , $A_{||}$ and A_{\perp} involving these molecular orbital coefficients which are quite similar in form to those given above for the d^9 case. Experimentally, $g_{||} = 2.054$, $g_{\perp} = 2.023$, $A_{||} = 166.5 \times 10^{-4} \text{ cm}^{-1}$, $A_{\perp} = 63.5 \times 10^{-4} \text{ cm}^{-1}$. It is assumed that $\beta_2^* = 1$, $\beta_2'^* = 0$, i.e., there is no in-plane π interaction. This seems to be generally a valid assumption for vanadyl complexes which is supported by the failure to observe ligand hyperfine structure with nitrogen ligands. This may be contrasted with the behaviour of Cu^{II} complexes. The other derived parameters are

$$\beta_1'^* = 0.84 \quad \beta_1'^* = 0.44 \quad \epsilon_{\pi}^* = 0.95$$

Thus it is deduced that the σ orbitals are considerably delocalized — it may be noted that the overlap is substantial. The out-of-plane π orbitals are only slightly delocalized and it is stated that this probably involves predominantly the vanadyl oxygen.

The analysis of the ESR spectra of metal complexes with more than one unpaired electron is more complicated than those of $S = \frac{1}{2}$ systems. It is necessary to take account of electron-electron dipolar interactions. An illustrative example is provided by the work on Cr^{III} acetylacetonate. This molecule was examined by Singer¹⁰ and more recently by McGarvey¹¹. The latter used a sample magnetically dilute in the Co^{III} compound. The spectrum can be fitted to the spin Hamiltonian

$$\mathcal{H} = g\beta H S + D(S_z^2 - \frac{5}{4}) + E(S_x^2 - S_y^2)$$

The g values are the main source of information on the electronic structure. D and E which describe the dipolar interactions are very sensitive to distortions of the molecule from octahedral symmetry but are not sensitive to the detailed nature of the molecular orbitals. McGarvey found the following parameters

$$g = 1.9802 \pm 0.0005$$

$$|D| = 0.600 \pm 0.001 \text{ cm}^{-1}$$

$$|E| = 0.0085 \pm 0.0005 \text{ cm}^{-1}$$

In addition, he was able to observe hyperfine coupling from the ^{53}Cr isotope ($I = 3/2$ abundance 9.54%) which gave the additional parameters

$$A = 16.2 \pm 0.1 \times 10^{-4} \text{ cm}^{-1}$$

$$B = 16.9 \pm 0.1 \times 10^{-4} \text{ cm}^{-1}$$

The lack of anisotropy in the g value and the small anisotropy of the hyperfine coupling is expected in view of the orbitally non-degenerate ground state ($^4A_{2g}$). The g value differs from that of the free electron since spin-orbit coupling leads to mixing of the ground state with the $^4T_{2g}$ excited state. Crystal field theory leads to the expression

$$g = 2.0023 - \frac{8\lambda}{\Delta E(^4T_{2g})}$$

The observed g value may be compared with those of the hexa-aquo Cr^{III} complex ($g = 1.976$) and the hexacyano complex ($g = 1.992$). The intermediate value for Cr^{III} acetylacetonate implies that the complex is more covalent than the aquo compound but less covalent than the cyanide. There is a similar trend in the metal hyperfine coupling constants, $A(\text{Cr}(\text{H}_2\text{O})_6^{3+}) = 17.0 \times 10^{-4}$, $A(\text{Cr}(\text{CN})_6^{3-}) = 14.7 \times 10^{-4} \text{ cm}^{-1}$ which supports this conclusion. McGarvey has examined a number of Cr^{III} complexes and carried out a more quantitative analysis. The $^4T_{2g}$ excited state arises from a configuration in which an electron has been promoted from the half-filled t_{2g} orbitals to the e_g orbitals. These latter are anti-bonding σ^* orbitals and as such are mixtures of metal d_{z^2} , $d_{x^2-y^2}$, s and p orbitals and appropriate ligand σ combinations. The expression for the g value becomes

$$g = 2.0023 - \frac{8\lambda P_d}{\Delta E(^4T_{2g})}$$

where P_d is a function of the e_g orbital coefficients. The value of P_d of 0.68 is considered to indicate considerable σ mixing. In this treatment π interactions are neglected. This is justified to some extent by the value of the metal hyperfine coupling constant which is relatively little changed from that of the aquo complex. This latter inference rests on the assumption that the hyperfine coupling constant, being a ground-state property, depends only on the coefficients of the half-filled d_{xy} , d_{xz} and d_{yz} orbitals which in turn can be mixed only with ligand π orbitals.

As a final ESR example, we will consider the work on Co^{II} trispyrazolyl borate carried out by Jesson.¹² Co^{II} has a d^7 configuration and the electron spin relaxation is such that its complexes are usually suitable for NMR studies in solution at room temperature. This is the case for the present complex. It appears that the shifts are predominantly dipolar in character and it is therefore important to establish the ESR parameters to aid the interpretation of the NMR*. At liquid-helium temperature the relaxation time is sufficiently long to allow observation of the ESR spectrum. However, at this temperature only the lowest of the Kramers doublets resulting from the combined effect of the trigonal distortion and spin-orbit coupling on the $^4T_{1g}$ ground state is populated. The spectrum can therefore be analysed using the Spin Hamiltonian

$$\mathcal{H} = \beta[g_x H_x S_x + g_y (H_y S_y + H_z S_z)] + AS_z I_z + B(S_x I_x + S_y I_y)$$

* An alternative which is now considered preferable is to measure the single-crystal magnetic susceptibilities (W D Horrocks and E S Greenberg, W D Horrocks and D W Hall, preprints)

The g value and the hyperfine coupling are both highly anisotropic ($g_{\parallel} = 8.46 \pm 0.01$, $g_{\perp} = 0.97 \pm 0.01$, $A = 0.0362 \pm 0.0003 \text{ cm}^{-1}$, $B < 0.0001 \text{ cm}^{-1}$). The anisotropic g value, of course, favours dipolar NMR shifts. Combination of this ESR data with optical spectroscopy allowed calculation of the g values for other Kramers doublets in the ground-state manifold and hence provides basic data for interpretation of the NMR. Since the ESR parameters are determined predominantly by the geometry and spin-orbit coupling of the complex, they do not provide a great deal of direct evidence on the nature of the bonding. This is generally true for orbitally degenerate ground states. In his calculation of the g values, Jesson used an orbital reduction factor of 0.9 for the matrix elements of the orbital angular momentum and reduced the free ion spin-orbit coupling by 20%. Both of these factors are indicative of covalency. Similarly, the analysis of the hyperfine coupling suggested about 15% charge transfer to the ligand. However, the information on metal-ligand interactions obtained by ESR on this type of compound is, at best, qualitative.

D. NMR THEORY

In principle, the theory applicable to the NMR of paramagnetic complexes is more straightforward than the corresponding ESR theory. In practice, the difficulties of interpretation are such that it may be fairly stated that all results reported thus far should be regarded as having qualitative rather than quantitative significance. Basically, there are two problems. The first is to separate the dipolar effects from the contact effects. Only the contact contribution provides direct information regarding the bonding. The contact shifts, when obtained, give directly the spin densities in the S orbitals of the magnetic nuclei. Usually there is no direct overlap of these orbitals with the spin-containing metal orbitals and they have acquired unpaired electron density indirectly from other ligand orbitals. It is the disentangling of these indirect pathways which provides the information on metal-ligand bonding and which constitutes the second problem.

The separation of dipolar and contact effects has been attempted in a number of ways. McConnell^{A5} first pointed out that contact shifts should be identical in both solution and solid whereas dipolar shifts should differ in the different phases. Comparison of solution and polycrystalline solid spectra should therefore establish the presence of dipolar effects. McConnell and Holm^{K1} applied this principle to vanadocene and concluded the shift was contact in origin but the method has not been further exploited. The experimental problem lies in the excessive line widths of NMR lines of polycrystalline solids. It has been suggested^{A21, R1} that the different dependences of nuclear T_1 's and T_2 's on hyperfine coupling constants could form the basis of a method for identifying contact interactions. This suggestion does not appear to have been followed up, probably because of the experimental difficulties of measuring short T_1 's.

The most generally used arguments to establish or disprove dipolar contributions to isotropic shifts have been based on experimental data or theoretical inferences on the presence or absence of magnetic isotropy in a complex. Dipolar (sometimes called pseudo-contact) effects can only be present if there is magnetic anisotropy. Thus McConnell and Robertson^{A5} derived the expression

$$\Delta H = - \frac{\beta^2 H_0 S(S+1)}{27kT} (g_{\parallel} + 2g_{\perp})(g_{\parallel} - g_{\perp}) \frac{(3 \cos^2 \theta - 1)}{r^3}$$

for the dipolar contribution in solution. In this expression, θ is the angle between the principle magnetic axis and a line joining the metal atom and the magnetic nucleus under consideration and r is the distance of this nucleus from the metal atom. This equation demands at least a three-fold axis. The other symbols have their usual significance. It is apparent that the dipolar shifts will be equal to zero if $g_{\parallel} = g_{\perp}$. Later work by Jesson^{A14} has shown that the exact form of the dipolar expression depends on the relative values of the electron spin relaxation time, the correlation time for molecular tumbling and the Zeeman anisotropy energy. The qualitative dependence on the g value anisotropy and the molecular geometry is, however, maintained. Still more recently Horrocks^{A16} has pointed out that previous treatments have neglected the contribution of the second-order Zeeman effect, which may be substantial in some cases, and has suggested that it would be preferable to evaluate the dipolar contribution to isotropic shifts from measurements of the anisotropy of the magnetic susceptibility rather than of the g values. Thus, although it appears that the theory of dipolar shifts is working towards a definitive form, it must be concluded that most of the quantitative calculations published thus far are unreliable. It is probable, though, that many of the qualitative conclusions which have been made in the past will not be substantially altered by refinements in the theory. Thus it has often been postulated that in octahedral or near-octahedral Ni^{II} complexes, dipolar effects can be neglected since magnetic anisotropy is not expected from the orbitally non-degenerate 3A_2 ground state. Shifts of such complexes have therefore been treated as purely contact in origin and it seems unlikely that this assumption can be seriously in error. Similarly, for other complexes such as Cr^{III} acetylacetonate and Ti^{III} acetylacetonate where the g values are known to be either isotropic or only slightly anisotropic, it is improbable that a more refined theory will alter the conclusion that the shifts are predominantly contact in origin. Thus if ESR measurements have established that the g value anisotropy is small or if theoretical arguments show that the ground state is orbitally non-degenerate and well-separated from excited states which can be mixed in by spin-orbit coupling, it seems fairly safe to assume that isotropic shifts observed will be predominantly contact in origin.

Three other lines of argument have been developed to estimate the extent of the dipolar contribution to NMR shifts in paramagnetic molecules. They may be grouped together in that none of them require any prior knowledge of the magnetic properties of the molecule. The first such method depends on estimating the ratios of the pseudo-contact shifts from their geometrical dependence. Providing the geometry of the molecule is known and there is sufficient symmetry to identify a principal axis without ambiguity, values of $(3 \cos^2 \theta - 1)/r^3$ can be calculated for each proton and the dipolar shifts must be proportional to these values. If the experimental shifts diverge widely from these ratios, and particularly if the signs are wrong, it can be deduced that dipolar effects do not predominate. This argument has been applied to Cr^{II} complexes^{C1}. It is open to the objection that relatively small contact contributions can affect the ratios. Secondly, if the mode of spin delocalization can be identified and the shifts show good proportionality to calculated spin densities, this has been taken as good evidence for a contact mechanism. The argument is most persuasive for π delocalization in a non-alternant system for which the alternation of positive and negative spin densities, and hence of high and low field shifts, provides a rather characteristic pattern. An extension of this argument rests on the observation that a methyl group shifts in the opposite direction to a proton substituted

at the corresponding position. Such an observation is readily interpretable by a π delocalization mechanism (see below) but would require very specific geometry for a dipolar mechanism*. Single inferences along these lines are inconclusive but if a large number of data can be fitted to a single π delocalization mechanism, it must be regarded as rather good evidence for the preference of a contact to a dipolar mechanism. If the spin density calculations are reliable, the residual dipolar contribution can be found by subtraction. A third method depends on the comparison of the ratios of shifts for complexes with the same ligand but different metal atoms. This has been applied most extensively to octahedral Ni^{II} and Co^{II} complexes. It is argued that the Ni^{II} complexes show only contact shifts, that the mechanism of delocalization will be the same for both Ni^{II} and Co^{II} , hence that the ratios of the contact shifts will be the same for both complexes and that discrepancies in the observed ratios must therefore be attributed to dipolar effects in the Co^{II} complex.^{A16} The crucial assumption is that delocalization mechanisms are the same for Ni^{II} and Co^{II} and this has led to some controversy.^{G56} Obviously none of these methods is completely unambiguous.

The overall conclusions reached thus far on the question of dipolar versus contact shifts can be summarized as follows. In the first transition series there seems to be little doubt that the shifts of octahedral Cr^{III} and Ni^{II} complexes arise from the contact effect. It is widely agreed that octahedral Co^{II} complexes show large, and in many cases predominant, dipolar effects. Octahedral V^{III} and Cr^{II} and tetrahedral Ni^{II} have been widely studied and all the interpretations advanced have been based on contact effects in spite of the fact that theory indicates dipolar effects should be present. It appears to the present author that by and large the attribution of the shifts in these complexes is probably correct for the reason that all the complexes studied are distorted from octahedral or tetrahedral geometry sufficiently to give orbitally non-degenerate ground states well separated (several thousand cm^{-1}) from excited states. Detailed measurements of the magnetic and spectroscopic properties of these complexes would do much to clarify the situation. Tetrahedral Co^{II} appears to show dipolar shifts^{F8} in spite of its orbitally non-degenerate ground state, presumably because of mixing of low-lying excited states. Data on the remaining first-row ions are rather scarce but the experimental evidence, such as it is, favours predominantly contact interactions. Data on second- and third-row transition metal complexes and on rare earth complexes are even more sparse but it is to be anticipated that dipolar interactions will be relatively more important if only because of the larger spin-orbit coupling constants.

We may now turn to the origin of the contact shifts. McConnell and Robertson^{A5} related the contact shift to the hyperfine coupling constant A by the expression

$$\frac{\Delta H}{H_0} = -A \frac{\gamma_e}{\gamma_H} g \frac{\beta S(S+1)}{3kT}$$

In this expression, ΔH is the shift at applied field H_0 . A , in turn, is related to the spin density ($[\psi(0)]^2$) at the nucleus in question by

* Recent INDO calculations suggest that σ delocalization can also lead to a reversal of CH_3 shifts (W. D. Horrocks and D. L. Johnson, preprint).

$$A = \frac{8\pi}{3h} g_e g_N \beta_e \beta_N [\psi(0)]^2$$

Thus, measurement of the contact shift immediately gives the spin density at the nucleus. Since only S orbitals have a finite density at the nucleus, in the simplest case a proton contact shift immediately gives the contribution of the H $1s$ orbital to the partly filled molecular orbital. However, this simplification is inadmissible since it neglects spin correlation. This problem was first encountered by ESR spectroscopists in the case of aromatic ion radicals. For these radicals the unpaired electron clearly occupies a π molecular orbital which has a node at the hydrogen atoms. Nevertheless, there is finite hyperfine coupling due to hydrogens and π - σ spin polarization must be invoked. The information which one would like to obtain from contact shift measurements relates to the amount of mixing of specific ligand orbitals with specific metal orbitals and hence a description of the covalency of the various types of metal-ligand bonding. In principle, this requires comparison of the observed shifts with S orbital spin densities obtained by complete, many-electron calculations with full allowance for spin correlation. In practice, such calculations are not feasible for molecules including transition metals.

In spite of the above limitations, contact shift measurements can yield valid information on metal-ligand interactions in certain cases. The main requirement is that it should be possible to identify a single dominant delocalization mechanism. If more than one such mechanism is present, the bonding information is likely to be, at best, very qualitative. It is probably most profitable to illustrate the type of information that can be obtained in this way by reference to some specific examples as will be done in the next section but we summarize, at this point, some of the possible modes of spin delocalization.

It is assumed, initially, that the unpaired electrons occupy predominantly metal d orbitals in a manner determined by simple crystal field theory. Thus a $VIII$ complex (approximately octahedral) will have two unpaired electrons in the $t_{2g}(d_{xy}, d_{xz}, d_{yz})$ orbitals. The Boltzmann distribution will determine that the α spin states of these unpaired electrons are favoured in a magnetic field. The spin-containing orbitals can mix directly with an empty ligand π^* orbital of similar energy giving rise to α spin density in ligand $p\pi$ atomic orbitals. The π spin densities can be related directly to hydrogen atom or methyl group hyperfine coupling constants using the theory developed for the ESR of aromatic radicals. This corresponds to π "back-bonding" with metal-to-ligand charge transfer and the ligand hyperfine coupling constants would be a measure of the extent of this process. There is also the possibility that the metal orbitals could interact with a filled ligand π orbital. This corresponds to π bonding with ligand-to-metal charge transfer. Since the d orbitals are less than half-filled, either an α electron or a β electron could be donated. In the former case there will be excess β spin density remaining on the ligand and in the latter case excess α spin density. Since the wave functions and hence the spin density distributions are usually different for the π and π^* orbitals and since the signs of the contact shifts are different for α and β spin densities, we can, in principle, distinguish these possibilities. However, if the symmetry is less than octahedral, the d_{xy}, d_{xz} and d_{yz} orbitals can also mix with ligand σ^* orbitals. This can lead to direct delocalization of α spin to hydrogen atoms on the ligand. In general, the spin density in σ orbitals will fall off rather rapidly with an increasing number of bonds separating the magnetic nucleus from the metal atom and this argument is often used to identify or refute σ delocalization. It has ob-

vious hazards. Still more subtle delocalization mechanisms may be encountered. Thus unpaired electrons in essentially non-bonding π orbitals may polarize the spins of paired electrons in neighbouring σ bonding orbitals leading to excess β spins in the ligand σ orbital. Again the main delocalization may be to the σ orbitals of the atom attached directly to the metal but this can polarize filled ligand π orbitals. Since incorrect identification of the delocalization mechanism will lead to incorrect inferences with regard to the bonding, it is apparent that results in this area must be assessed with some care.

Finally, we might say a word about the calculation of ligand spin densities. The most desirable approach would obviously be a complete calculation involving all metal and ligand orbitals with full allowance for electron correlation. Needless to say such calculations have not been carried out. In a number of cases (notably metallocene complexes), extended Huckel calculations on the complete molecule have been used to obtain spin densities^{K7}. These lack allowance for correlation effects but the reasonable agreement with experiment suggests they may be fair approximations. This does not imply, however, that the method will necessarily work equally well in other cases especially where both positive and negative spin densities occur. A more common procedure has been to carry out a calculation on a radical derived from the ligand, to compare the relative spin densities at different positions, and hence to deduce the fraction of an electron delocalized which gives a measure of the metal-ligand interaction. The majority of applications have been to π radicals of the non-alternant type showing large negative spin densities. In such cases it is essential to include electron correlation and single electron Huckel calculations are inadmissible. Earlier workers resorted to valence bond calculations (which automatically include correlation) and these were surprisingly successful in a number of cases^{G5}. Later, MacLachlan developed a rather simple molecular orbital technique for dealing with this situation and this has been widely used^{B3}. Calculations on σ radicals are more limited and have been confined to the extended Huckel^{A15} and linear combination of bond orbital methods^{G26} neither of which include correlation. In general, the success of these calculations is such that the conclusions reached with regard to the type of orbital containing the unpaired electrons and qualitative estimates of the amount of delocalization can be regarded as reliable in favourable circumstances. In the present state of the art, this is perhaps the maximum amount of information which can be derived from contact shift studies.

E. SOME NMR RESULTS

The study of the isotropic NMR shifts of paramagnetic metal complexes has been an active area of research for almost ten years. During this period a rather substantial number of papers has been published and an attempt has been made to provide a bibliography. In the present section, only a small number of selected papers will be reviewed. These have been chosen, somewhat arbitrarily, to try and illustrate the utility of this type of experiment in studies of metal-ligand bonding. The contributions of the papers listed to other areas of interest, ranging from isomerism in metal complexes to hyperconjugation in the organic ligands, have been substantial so that the following examples by no means represent a balanced selection.

The first series of complexes to be extensively studied were the Ni^{II} aminotroponimine-





Fig 3 ^1H spectrum of Ni^{II} N,N' -bis(p -1,3-butadienylphenyl)aminotroponeimineate CDCl_3 solvent, TMS reference


ates of structure (II)^{G6} In the light of subsequent work, this was an extremely fortuitous series of compounds with which to start. In contrast to other compounds, the results can be interpreted in a relatively simple manner, the ligands allow wide synthetic flexibil-



ity and the NMR relaxation properties are particularly favourable. There is only one complication, namely that there is an exchange process, fast on the NMR time scale, between a planar diamagnetic isomer and a tetrahedral paramagnetic isomer. As a result, the observed shifts are the average of the paramagnetic and diamagnetic forms. For our present purpose we will neglect this aspect of the studies. It was originally considered that the fast isomer exchange determined the fast electron spin relaxation but this does not appear to be correct. However, for whatever the cause, the NMR lines are sharp enough to resolve spin-spin splitting in many cases and shifts of up to 15,000 Hz at 60 MHz are observed. A typical spectrum is shown in Fig 3. The most notable qualitative feature about this spectrum is the existence of both high and low field shifts. Closer examination of this and related molecules shows that the fundamental pattern is very simple — protons attached to successive carbons shift in opposite directions. This pattern is maintained indefinitely as the ligand π system is extended. This pattern can be interpreted very simply with a model involving contact shifts due to spin delocalization in the π system. The proton hyperfine coupling constants can be used to obtain π spin densities from the relationship

$$A_{\text{H}} = Q\rho_{\text{c}}$$

where Q is a constant of numerical value approximately -22.5 gauss. The theory pertinent to the use of this equation was developed to interpret the ESR of aromatic radicals. High field shifts correspond to positive π spin densities and low field shifts to negative π spin densities. A simple valence bond formulation suffices to determine which positions will experience positive spin densities and which negative spin densities. Thus, if a radical is formed by transferring an electron from the aminotroponeimineate ion to the metal, the principal contributing structures will be



Reasonable structures placing positive spin at the α and γ positions can be written, but there is no such possibility at the β position. Experience with similar radicals, such as the allyl radical, suggests that this position will acquire negative spin density by spin polarization of the filled π orbitals. Similar valence bond structures can easily be written for the more complex ligands obtained by substituting aromatic moieties on the nitrogen or at the γ position. Spin densities on these radicals can be calculated by the valence bond method and compared with the experimental results. Since we are not dealing with "free" radicals but only with ligands with some radical character, the fit of the model must be judged from the relative spin densities at the different positions. If a common scaling factor gives satisfactory agreement at all positions, this scaling factor is a measure of the fraction of an electron delocalized and hence of the amount of π interaction. Table 2 gives some comparisons of calculated and experimental spin densities. The implication is that about 1/10 of an electron is delocalized to the aminotroponeimineate framework and about 1/30 to aromatic fragments attached to the nitrogen. The overall agreement is better than might have been expected from the rather unsophisticated method of calculation.


Thus the immediate conclusion is that π interaction is sufficiently important to give rise to about a 10% contribution from structures involving delocalization of the unpaired electrons to the ligand. It should be noted that tetrahedral Ni^{II} has unpaired electrons in the d_{xy} , d_{xz} and d_{yz} orbitals which are capable of direct overlap with the ligand π system so there is no necessity to postulate any indirect mode of delocalization. Furthermore,

TABLE 2

Spin densities in nickel^{II} N,N' -bis(*p*-1,3-butadienyphenyl) aminotroponeimineate

Proton	ρ_{obs}	ρ_{calc}	Proton	ρ_{obs}	ρ_{calc}
α	+0.0410		a	-0.00586	-0.00465
β	-0.0210		b	+0.00422	+0.00643
γ	+0.0568		c	-0.00281	-0.00219
ortho	+0.00792	+0.01099	α	+0.00342	+0.00288
meta	-0.00869	-0.00629	c	+0.00305	

the interaction must involve ligand-to-metal charge transfer since molecular orbital calculations^{E8} show that the spin density distribution in the π^* orbital is incompatible with the experimental results. Considering the consistency of the results over the substantial number of compounds studied, it seems certain that the shifts arise predominantly from this mechanism. Smaller contributions from other sources are, however, by no means excluded. Thus the observation of shifts in *N*-alkyl substituents and discrepancies in some of the proton shifts of aromatic substituents at positions relatively close to the metal are certainly indicative of some σ contribution. Recently, ^{13}C shifts have been observed for one of these compounds and it is considered that they are also indicative of some σ delocalization^{G55}. Similarly, although the arguments against a substantial dipolar effect are strong, a small contribution from this source cannot be excluded. Thus, even in this well-studied system, one must be content with qualitative conclusions about the bonding. Even with more reliable calculations on the spin density distribution, it is unlikely that one will be able to be more explicit than to estimate 10% for the π delocalization.



The Ni^{II} salicylaldehydes of structure (III) are a similar series of compounds which have also been rather extensively studied^{G17}. Again, the suggested interpretation involves delocalization into the ligand π system by $d\pi-p\pi$ interaction. In this case, the delocalization is smaller, amounting only to about 1/30th of an electron. Probably the interacting π orbital is energetically less favourably situated for interaction with the d orbitals than in the aminotroponimineates. The reservations expressed in the last paragraph apply equally well here but the predominant mechanism seems to be well established.

In both of the above cases the rather large number of chemically distinct ligand protons allows extensive comparison of observed and calculated spin densities which is a valuable aid in establishing the delocalization mechanism. However, in cases with less favourable relaxation conditions, the lines are broad and a complex ligand is likely to give rise to difficulties in the assignment of lines. For comparisons between different metal ions it is therefore better to choose a simple ligand. A series of first-row transition metal acetylacetonates has been examined with this consideration in mind^{B2}. The observed isotropic shifts are given in Table 3. There are only two chemically distinct types of proton, present in the ratio of six to one, so for any reasonable resolved spectrum there is no assignment problem. In spite of this the lines were too broad in the cases of Ti^{III} and Cr^{III} to locate the single C-H proton. It was argued that, in the early transition metals at least (up to Fe^{III}), dipolar effects are not important. This argument is by no means watertight for all the complexes but in the authors' opinion is probably substantially correct because there is a smooth progression of the shifts (as seen from the figures of Table 3) in a series which includes both complexes where dipolar effects can be certainly eliminated and those where it is less certain. The suggested interpretation was based on spin delocalization.

TABLE 3

NMR of paramagnetic acetylacetonates

	Ti ^{III}	V ^{III}	Cr ^{III}	Mn ^{III}	Fe ^{III}
CH ₃	~ -3500	-2744	-2320	-1505	-1243
H		-2404		-1085	+1644

Shifts in Hz at 60 MHz from TMS

TABLE 4

Molecular orbital calculations

Acetylacetonate				
Energy top bonding	+0.9417 β			
Energy bottom anti-bonding	-0.6103 β			
	Spin densities			
	Top bonding		Bottom anti-bonding	
	Huckel	McLachlan	Huckel	McLachlan
O	+0.0890	+0.0710	+0.0799	+0.0852
C ₁ (C-CH ₃)	+0.1260	+0.0640	+0.4201	+0.4979
C ₂ (C-H)	+0.5683	+0.7291	+0.000	-0.1658

to the π system. Table 4 contains calculated π spin densities for both the bottom anti-bonding π orbital and the top bonding π orbital. Both the CH₃ and the single H shifts can be related to these spin densities but in the first case the Q value should be taken as positive ($\sim +27$ gauss) and in the second case negative. These calculations lead to the following predictions for the contact shifts.

(a) For metal-to-ligand charge transfer to the lowest unoccupied π orbital, the CH₃ will have a large low field shift and the single hydrogen will also move to low field shift though somewhat less.

(b) Ligand-to-metal charge transfer will leave unpaired spin density in the highest occupied π orbital. For a less than half-filled d shell, an α electron will be transferred. This gives rise to a large low field shift of the single hydrogen and a small shift of indeterminate sign for the CH₃. For a half-filled or more shell, the signs are reversed, i.e. a large high field shift is predicted for the single hydrogen. The experimental results can be fitted rather nicely to these predictions. At the beginning of the series (Ti^{III}) it is postulated that the d orbitals are relatively close in energy to the π^* orbital and metal-to-ligand transfer predominates. On passing along the series, the increasing nuclear charge lowers the energy of the d orbitals and ligand-to-metal charge transfer must also be considered. Delocalization in the π^* orbital decreases from about 7% to 1% of an electron from Ti^{III} to Fe^{III}. The change in sign of the C-H shift at Fe^{III} is predicted from the ligand-to-metal charge transfer mechanism. Perhaps even more significantly the interpretation is

consistent with the chemistry of these ions. The complexes showing the largest tendency to metal-to-ligand charge transfer involve those ions most easily oxidized and where ligand-to-metal charge transfer is involved the ions are reducible to a lower oxidation state. It should be emphasized, though, that results of this kind do not have more than qualitative significance. It is plausible from the expectation that the d orbitals will be closer in energy to the π orbitals of the ligand than to the σ orbitals that π delocalization will be more important than σ delocalization. There is also some experimental evidence to this effect from the early observation of Forman et al.^{B1} that substitution of CH_3 for the single hydrogen of the VIII complex reversed the sign of shift as expected for a π radical. There could, however, easily be enough σ delocalization to spoil quantitative interpretations in addition to the uncertainties introduced by possible dipole interactions and inaccurate spin density calculations. Chemically, however, results of this kind are of importance in demonstrating trends in the bonding and in giving a qualitative idea of the extent of π interaction.

It is appropriate to discuss next a case where several delocalization mechanisms have been shown to occur simultaneously. In a recent paper, Cramer and Drago^{G46} have discussed contact shifts in complexes of the type NiL_6^{2+} where L is a pyridine or substituted pyridine. There had been previous reports of shifts for nickel pyridine complexes with distorted octahedral or distorted tetrahedral geometry. The use of regular octahedral complexes enabled the authors to dismiss the possibility of dipolar effects with some certainty and also to eliminate direct delocalization into the π system. The unpaired electrons reside in an e_g orbital which can interact with ligand σ orbitals but which is orthogonal to the π orbitals. At room temperature there is fast ligand exchange so that only averaged shifts are observed but the contact shifted resonances can be obtained at -40°C . The observed shifts are shown in Table 5. For the pyridine complex itself, the general pattern is

TABLE 5
 NiL_6^{2+} contact shifts

Ligand L		$\Delta\nu$ (c/sec)
Pyridine	O	-3820
	M	-1420
	P	- 445
4-Methylpyridine	O	-4570
	M	-1500
	P- CH_3	+ 422
3-Methylpyridine	O	-4320
	M	-1575
	P	- 450
	M- CH_3	- 313
4-Vinylpyridine	O	-2631
	M	-1236
	H_3	- 195
	H_2	+ 53
	H_1	- 218

Shifts in Hz at 60 MHz relative to TMS

that expected for σ delocalization. The shifts are to low field, indicating the protons can sense directly the α spin on the metal, and diminish rather rapidly with distance from the nickel. The results on some of the substituted pyridines indicate that this mechanism is not completely satisfactory. In particular, the high field shifts of the CH_3 of 4-methylpyridine and of H2 of 4-vinylpyridine cannot be explained in this manner. The suggestion is made that polarization leads to spin density in the π system. Extended Hückel calculations suggested about 4×10^{-2} spin density in the σ orbitals. β spin density in the π orbitals is about an order of magnitude less than this and it was not possible to identify the specific orbital. These results seem quite plausible*. It might be noted, though, that the observation of both high and low field shifts does not necessarily implicate π orbitals. Such effects have been observed in some substituted Ni^{II} aminotroponimineates in which the substituent has no π system. The only firm conclusion is that one-electron calculations for such systems are not sufficient.

The above papers are indicative of the kind of attempts which have been made to obtain information on metal-ligand bonding directly from contact shift measurements. There are a large number of variations on the themes already introduced. In this final section we would like to indicate some of the more indirect uses of NMR in this area. The work of Jesson et al.^{F23} on cobalt^{II} pyrazolylborates has already been mentioned. The predominant shift mechanism is dipolar and so does not provide direct information on the bonding. There is, however, a significant contact contribution so that bonding parameters are likely to be obtained as a by-product from detailed studies of this kind. A very recent paper by McGarvey^{A18} has re-examined the theory of these d^7 systems and introduced some quantitative changes in the interpretation. We refer the reader to the original paper for details. Work of this kind involving continuing refinement of the theory emphasizes the limitation of most of what has been published so far to qualitative significance. At the other end of the scale a recent paper by Rakshys^{G53} illustrates very nicely the usefulness of the NMR technique in an empirical way. The molecules studied were complexes of the type $\text{Ni}(\text{acac})_2\text{L}_2$ (acac = acetylacetonate, L = substituted anilines) in which the contact shifts were correlated with stability constants. Both electronic and steric effects appear to be important and the NMR measurements provide a sensitive probe for disentangling these effects. Finally we might mention the rather limited number of studies on complexes with mixed ligands^{G13,G19}. The interest here lies in the approach which contact shifts offer to the question of how substituent changes in one ligand affect bonding to other ligands. This is an important problem in studies of substitution mechanisms of complexes and in homogeneous catalysis. It has been demonstrated that rather large effects are observed in the NMR spectra of mixed complexes. The suggested interpretations have been based on differences in the occupation of the spin-containing metal orbitals as the various ligands are changed. This would appear to be a very promising area which awaits more systematic experimental exploitation.

* Recent INDO calculations suggest that it may be possible to interpret these results purely on the basis of σ delocalization (W D Horrocks and D L Johnson, preprint).

F. CONCLUSION

For the reasons outlined in the Introduction, it is not possible to make explicit comparisons between ESR and NMR results on metal-ligand bonding. It is, however, feasible to make some comments on the consistency of the general pictures which emerge from the two types of studies. The results described above have been very largely confined to chelate complexes of the first-row transition metal ions. Both techniques demonstrate that covalency is a factor of major importance in this class of compounds. In both cases interactions allowed by symmetry considerations are most important. Thus, ligand hyperfine splitting is observed in Cu^{II} complexes but not in V^{IV} complexes. π delocalization appears to be dominant from contact shift studies of tetrahedral Ni^{II} complexes but is subsidiary to σ effects in octahedral Ni^{II} complexes where spin can reach the π system only indirectly. ESR results indicate up to 20% contribution of ligand σ orbitals to formally d orbitals but in general less than 10% ligand contribution to π orbitals. NMR results agree with this limitation on π bonding. Perhaps one of the more direct comparisons is that of 5% ligand π contribution in vanadyl acetylacetonate from ESR and about the same percentage in V^{III} acetylacetonate from NMR. The NMR technique is more sensitive but less specific. Thus, a detailed single-crystal ESR study can separate out the different modes of spin delocalization without ambiguity. It will probably not, however, detect 2 or 3% contribution of a given type of ligand orbital. Such a contribution will be sufficient to impart substantial contact shifts to the NMR spectrum but there is no simple experimental criterion for disentangling the mechanisms. This sensitivity of the NMR method means that relatively exotic indirect modes of spin delocalization which can be safely neglected in ESR must be considered in interpreting the NMR spectra. The theory for ESR is rather well developed and allows quantitative interpretations to be put forward with some confidence. That for NMR is more complex and cannot be applied with certainty without sophisticated many-electron calculations on complete complexes. The overall result is that ESR gives us a reliable quantitative description of a few classes of complexes. NMR has given a large amount of essentially qualitative data on a much wider range of compounds. Both techniques are capable of much wider exploitation and jointly are likely to continue to make a substantial contribution to our understanding of the electronic structure of transition metal complexes.

REFERENCES

ESR

- 1 For example, A. Carrington and A. D. McLachlan, *Introduction to Magnetic Resonance*, Harper and Row, 1967.
- 2 Recent reviews (a) B. R. McGarvey, *Transition Metal Chem.*, 3 (1966) 90 (b) H. A. Kuska and M. T. Rogers, in E. T. Keiser and L. Kevan (Eds.), *Radical Ions*, Interscience, 1968, p. 579 (c) E. König, in H. A. O. Hill and P. Day (Eds.), *Physical Methods in Advanced Inorganic Chemistry*, Interscience, 1968, p. 266.
- 3 B. R. McGarvey, *J. Phys. Chem.*, 71 (1967) 51.
- 4 A. H. Maki and B. R. McGarvey, *J. Chem. Phys.*, 29 (1958) 31.

- 5 H A Kuska and M T Rogers, *J Chem Phys*, 43 (1965) 1744
- 6 H A Kuska, M T Rogers and R E. Dillinger, *J Phys Chem* 71 (1967) 109
- 7 A H Maki and B R. McGarvey, *J Chem. Phys*, 29 (1958) 35
- 8 D Kivelson and S K. Lee, *J Chem Phys*, 41 (1964) 1896
- 9 C J Ballhausen and H B. Gray, *Inorg Chem*, 1 (1962) 111
- 10 L S. Singer, *J Chem Phys*, 23 (1955) 379
- 11 B R. McGarvey, *J Chem Phys*, 41 (1964) 3743
- 12 J P. Jessor, *J Chem. Phys*, 45 (1966) 1049

NMR

A Theory

- 1 N Bloembergen, L M. Purcell and R V. Pound *Phys Rev* 73 (1948) 679 Relaxation by paramagnetic ions
- 2 N Bloembergen and W C Dickinson, *Phys Rev*, 79 (1950) 179 NMR shifts with paramagnetic ions
- 3 W C Dickinson, *Phys Rev*, 81 (1951) 717 Shifts with paramagnetic ions
- 4 H M. McConnell and C H. Holm, *J Chem Phys*, 26 (1957) 1517 Knight shifts Na and liquid NH_3
- 5 H M. McConnell and R E. Robertson, *J Chem. Phys* 29 (1958) 1361 Theory of isotropic nuclear resonance shifts
- 6 H M. McConnell and D B. Chesnut, *J Chem. Phys*, 28 (1958) 107 Spin density and hyperfine coupling
- 7 H M. McConnell, *J Chem. Phys*, 28 (1958) 1188 Spin density matrices
- 8 N Bloembergen and L O. Morgan, *J Chem. Phys* 34 (1961) 842 Relaxation in paramagnetic solutions
- 9 W C. Lin and L E. Orgel, *Mol Phys*, 7 (1964) 130 Unpaired spin in unsymmetric aminotropone-imine complexes of Ni^{II}
- 10 R M. Golding, *Mol Phys*, 8 (1964) 561 Temperature dependence of contact shifts
- 11 G N. La Mar, *J Chem Phys*, 43 (1965) 1085 Dipolar shifts in paramagnetic complexes
- 12 G N. La Mar, *J Amer Chem Soc*, 87 (1965) 3567 Relaxation in paramagnetic complexes
- 13 J P M. Bailey and R M. Golding, *Mol Phys* 12 (1967) 49 Spin polarization in aliphatic fragments
- 14 J P. Jessor, *J Chem Phys*, 47 (1967) 579 Isotropic shifts of Co^{II} complexes
- 15 R S. Drago and H. Peterson, Jr, *J Amer Chem Soc*, 89 (1967) 5774 Hyperfine coupling constants of sigma radicals
- 16 W D. Horrocks, Jr, *Inorg Chem*, 9 (1970) 690 Evaluation of dipolar shifts
- 17 R J. Kurland and B R. McGarvey, *J Magnet Res* 2 (1970) 286 Theory of dipolar shifts
- 18 B R. McGarvey, *J Chem Phys*, 53 (1970) 86 Theory of Co^{II} complexes
- 19 Y N. Molin, *Chem Phys Letters* 5 (1970) 511 Spin density distribution in ligands
- 20 M C R. Symons, *J Chem Phys*, 53 (1970) 1620 Theory ^{19}F and ^{13}C shifts in simple complexes
- 21 H. Sternlicht, *J Chem Phys*, 42 (1965) 2250 Dipolar relaxation

B Vanadium

- 1 A. Forman, G N. Murrell and L E. Orgel, *J Chem Phys* 31 (1959) 1129 β -Diketone complexes
- 2 D R. Eaton, *J Amer Chem Soc* 87 (1965) 3097 Bonding in β -diketone complexes
- 3 E. Rohrscheid, R E. Ernst and R H. Holm, *Inorg Chem*, 6 (1967) 1315 β -Diketone complexes
- 4 E. Rohrscheid, R E. Ernst and R H. Holm, *Inorg Chem*, 6 (1967) 1607 β -Ketoamine complexes
- 5 E. Rohrscheid, R E. Ernst and R H. Holm, *J Amer Chem Soc*, 89 (1967) 6472 Salicylaldehyde and salicylaldimine complexes
- 6 R H. Holm, D H. Gerlach, J G. Gordon and M G. McNamee, *J Amer Chem Soc*, 90 (1968) 4184 Tris β -thioketonates
- 7 Y T. Chen and G W. Everett, Jr, *J Amer. Chem Soc*, 90 (1968) 6660 3-Substituted camphor complexes

- 8 J. Reuben and D. Fiat, *Inorg. Chem.*, 8 (1969) 1821 ^{17}O vanadyl complexes
- 9 G. W. Everett and Y. T. Chen, *J. Amer. Chem. Soc.*, 92 (1970) 508 Carvone and pulegone complexes

C Chromium

- 1 G. N. La Mar and G. R. Van Hecke, *J. Amer. Chem. Soc.*, 91 (1969) 3442 *o*-Phenanthroline and α, α' -bipyridine complexes
- 2 G. N. La Mar and G. R. Van Hecke, *J. Amer. Chem. Soc.*, 92 (1970) 3021 Temperature dependence of methyl *o*-phenanthroline complexes

D Manganese

- 1 B. B. Wayland and W. L. Rice, *Inorg. Chem.*, 5 (1966) 54 Contact shifts of Aq compounds
- 2 B. B. Wayland and R. S. Drago, *J. Amer. Chem. Soc.*, 87 (1965) 2372 Hexamethylphosphoramide compounds
- 3 H. Sternlicht, R. G. Shulman and E. W. Anderson, *J. Chem. Phys.*, 43 (1965) 3123 Bonding to adenosine triphosphate
- 4 Z. Luz, B. L. Silver and D. Fiat, *J. Chem. Phys.*, 46 (1967) 469 ^{17}O $\text{Mn}(\text{acac})_3$
- 5 D. G. Davis and R. J. Kurland, *J. Chem. Phys.*, 46 (1967) 388 ^{13}C of cyanides
- 6 R. M. Golding, P. Healey, P. Newman, E. Sinn, W. C. Tennant and A. H. White, *J. Chem. Phys.*, 52 (1970) 3105 $\text{Mn}(\text{dithiocarbamates})$

E Iron

- 1 R. M. Golding, W. C. Tennant, C. R. Kanekar, R. L. Martin and A. H. White, *J. Chem. Phys.*, 45 (1966) 2688 Fe^{III} dithiocarbamates
- 2 V. V. Voevodskii, Y. N. Molin and K. I. Zernarsky, *Zh. Strukt. Khim.*, 8 (1967) 864 Spin densities in Fe complexes
- 3 L. H. Pignolet, D. Forster and W. De W. Horrocks, Jr., *Inorg. Chem.*, 7 (1968) 828 Dihalo-bis(triarylphosphine) Fe^{II}
- 4 K. Wuthrich, R. G. Shulman, B. L. Wyluda and W. S. Caughey, *Proc. Natl. Acad. Sci. U.S.A.*, 62 (1969) 636 Porphyrin Fe^{III} cyanides
- 5 M. L. Nicholas, *J. Amer. Chem. Soc.*, 92 (1970) 4141 Fe^{III} phenanthroline complexes
- 6 R. E. De Simore and R. S. Drago, *J. Amer. Chem. Soc.*, 92 (1970) 2343 Bipyridyl and phenanthroline complexes
- 7 J. P. Jesson, S. Trofimenko and D. R. Eaton, *J. Amer. Chem. Soc.*, 89 (1967) 3158 Polypyrazolylborates
- 8 D. R. Eaton, W. R. McClellan and J. F. Weiher, *Inorg. Chem.*, 7 (1968) 2040 Aminotroponemimines of Fe, V, Cr, Mn and Co

F. Cobalt

- 1 R. S. Milner and L. Pratt, *Discussions Faraday Soc.*, 34 (1962) 88 Amino acids
- 2 C. C. McDonald and W. D. Phillips, *J. Amer. Chem. Soc.*, 85 (1963) 3736 Histidine complexes
- 3 W. D. Horrocks and G. N. La Mar, *J. Amer. Chem. Soc.*, 85 (1963) 3512 Phosphine complexes
- 4 J. A. Happe and R. L. Ward, *J. Chem. Phys.*, 39 (1963) 1211 Pyridine/acac
- 5 W. D. Horrocks, R. C. Taylor and G. N. La Mar, *J. Amer. Chem. Soc.*, 86 (1964) 3031 Adducts of Ni^0 and Co^0 -acac
- 6 Z. Luz and S. Meiboom, *J. Chem. Phys.*, 40 (1964) 2686 Exchange of methanol complexed to Co^{2+} and Ni^{2+}
- 7 Z. Luz and S. Meiboom, *J. Chem. Phys.*, 40 (1964) 1058 Solvation of Co^{II} in MeOH , $\text{MeOH}-\text{H}_2\text{O}$ systems
- 8 G. N. La Mar, W. D. Horrocks and L. C. Allen, *J. Chem. Phys.*, 41 (1964) 2126 $\text{Bis}(\text{triarylphosphine})\text{Co}^{\text{II}}, \text{Ni}^{\text{II}}$ dihalides

- 9 Z Luz and R G Shulman, *J Chem Phys*, 43 (1965) 3750 Aquo complexes
- 10 G N La Mar, *J Phys Chem*, 69 (1965) 3212 Arylphosphine compounds
- 11 G W Everett, Jr and R H Holm, *J Amer Chem Soc*, 87 (1965) 5266 Planar and tetrahedral equilibria
- 12 B B Wayland and R S Drago, *J Amer Chem Soc*, 88 (1966) 4597 $\text{Co}(\text{OP}(\text{N}(\text{CH}_3)_3)_2)_2\text{X}_2$, $\text{Co}(\text{C}_5\text{H}_5\text{N})_2\text{X}_2$
- 13 B B Wayland, R J Fitzgerald and R S Drago, *J Amer Chem Soc*, 88 (1966) 4600 *N,N*-dimethylacetamide, Co^{II}
- 14 N A Matwiyoff, *Inorg Chem*, 5 (1966) 788 Solvation of cations in *N,N*-dimethylformamide
- 15 G W Everett, Jr and R H Holm, *J Amer Chem Soc*, 88 (1966) 2442 β -Ketoamino complexes
- 16 W D Horrocks and L H Pignolet, *J Amer Chem Soc*, 88 (1966) 5929 Kinetics of triphenyl phosphine exchange
- 17 R W Kluber and W D Horrocks, *J Amer Chem Soc*, 88 (1966) 1399 Quinoline oxide and isoquinoline oxide adducts of $\text{Ni}(\text{acac})_2$ and $\text{Co}(\text{acac})_2$
- 18 W D Horrocks and J R Hutchison, *J Chem Phys*, 46 (1967) 1703 H_2O shifts Co^{II} with various anions
- 19 S Thomas and W L Reynolds, *J Chem Phys*, 46 (1967) 4164 $\text{Co}(\text{DMSO})_6^{2+}$ and $\text{Ni}(\text{DMSO})_6^{2+}$
- 20 R W Kluber and W D Horrocks, *Inorg Chem*, 6 (1967) 166 Pyridine complexes of bis(benzoylacetate)-Ni and Co
- 21 W D Horrocks, G R Van Hecke and D De W Hall, *Inorg Chem*, 6 (1967) 694. Diteriary phosphine complexes
- 22 G N La Mar, *Inorg Chem*, 6 (1967) 1939 L_2CoX_2
- 23 J P Jesson, S Trofimenko and D R Eaton, *J Amer Chem Soc*, 89 (1967) 3148 Polypyrazolylborates
- 24 N H Agnew, L F Larkworthy and G A Webb, *Inorg Nucl Chem Letters*, 3 (1967) 303 Vinyl pyridine complexes of Co and Ni
- 25 B B Wayland and W L Rice, *Inorg Chem*, 6 (1967) 2270 Ammine complexes
- 26 B B Wayland and W L Rice, *Inorg Chem*, 6 (1967) 54 Aquo complexes
- 27 J F O'Brien and W L Reynolds, *Inorg Chem*, 6 (1967) 2110 Co and Ni acetonitrile complexes
- 28 H P Fritz, W C Gretner, H J Keller and K E Schwarzhans, *Z Naturforsch*, 23b (1968) 906 Salicylaldehyde/pyridine complexes of Co and Ni
- 29 R W Kluber and S Kopycinski, *J Inorg Nucl Chem*, 30 (1968) 1891 Aldehyde and ketone adducts of $\text{Ni}(\text{acac})_2$ and $\text{Co}(\text{acac})_2$
- 30 R S Drago and B B Wayland, *Inorg Chem*, 7 (1968) 628 Absence of steric effects in $\text{Co}(\text{Py})_2\text{X}_2$
- 31 G W Everett, Jr and R H Holm, *Inorg Chem*, 7 (1968) 776 Structural interconversion of planar and tetrahedral Co and Ni
- 32 B R McGarvey and J Pearlman, *J Magnet Res*, 1 (1969) 178 Pseudo-contact d^5 ion-complexes
- 33 R J Fitzgerald and G R Hubaker, *Inorg Chem*, 8 (1969) 2265 Square planar complexes—dithioacetylacetonate
- 34 L Pratt and B B Smith, *Trans Faraday Soc*, 65 (1969) 1703 Aminopolycarboxylates of Co^{II}
- 35 T Vonezaw, I Morishima, Y Akana and K Fukuta, *Bull Chem Soc Japan* 43 (1970) 379 Substituted anilines complexes with Ni-, Co-acac
- 36 U Sakaguchi, Y Arata and S Fujiwara, *J Mol Spectry*, 34 (1970) 312 Tetra-coordinated dihalobis(picoline) Co^{II}
- 37 R Engel, *J Chem Soc D*, (1970) 133 Decoupling with paramagnetic ions

G Nickel

- 1 W D Phillips and R.E. Benson, *J Chem Phys*, 33 (1960) 607 Observation of shifts in aminotroponemineates
- 2 R E Benson, D R Eaton, A D Josey and W D Phillips, *J Amer Chem Soc*, 83 (1961) 3714 Spin densities in Ni-aminotroponemines
- 3 D R Eaton, A D Josey, R E Benson, W D Phillips and T L Cairns, *J Amer Chem Soc*, 84 (1962) 4100 Electron spin density in π -systems

- 4 E.A. LaLancette, D.R. Eaton, R.E. Benson and W.D. Phillips, *J. Amer. Chem. Soc.*, **84** (1962) 3968 Ni-salicylaldehydes
- 5 D.R. Eaton, A.D. Josey, W.D. Phillips and R.E. Benson, *Discussions Faraday Soc.* **34** (1962) 77 Conjugated aminotroponimineates
- 6 D.R. Eaton, A.D. Josey, W.D. Phillips and R.E. Benson, *J. Chem. Phys.*, **37** (1962) 347 π spin densities in aminotroponimineates
- 7 D.R. Eaton, A.D. Josey, W.D. Phillips and R.E. Benson, *Mol. Phys.*, **5** (1962) 407 ^{19}F contact interactions
- 8 D.R. Eaton, W.D. Phillips and D.J. Caldwell, *J. Amer. Chem. Soc.* **85** (1963) 397 Configuration and magnetic properties of aminotroponimineates
- 9 D.R. Eaton, A.D. Josey, W.D. Phillips and R.E. Benson, *J. Chem. Phys.*, **39** (1963) 3513 Spin-spin coupling constants from paramagnetic complexes
- 10 D.R. Eaton, A.D. Josey and W.A. Sheppard, *J. Amer. Chem. Soc.* **85** (1963) 2689. ^{19}F contact interactions CF_3 , OCF_3 , SCF_3 , SF_5 and SO_2 groups
- 11 R.H. Holm, A. Chakravorty, G.O. Dudek, *J. Amer. Chem. Soc.*, **86** (1964) 379 Conformational equilibrium
- 12 E.A. LaLancette and D.R. Eaton, *J. Amer. Chem. Soc.*, **86** (1964) 5145 Contact shifts in $\text{Bis}(\text{Ph}_3\text{P})\text{NiX}_2$
- 13 A. Chakravorty and R.H. Holm, *J. Amer. Chem. Soc.*, **86** (1964) 3999 Mixed-ligand salicylaldehydes
- 14 A. Chakravorty and R.H. Holm, *Inorg. Chem.* **3** (1964) 1010 α -Hydroxynaphthaldehyde complex
- 15 D.R. Eaton and E.A. LaLancette, *J. Chem. Phys.*, **41** (1964) 3534 Metallo-pyrromethenes
- 16 Z. Luz and S. Meiboom, *J. Chem. Phys.*, **40** (1964) 1066 Solvation of Ni in MeOH
- 17 R.H. Holm, A. Chakravorty and G.O. Dudek, *J. Amer. Soc.* **86** (1964) 379 Salicylaldehyde complexes
- 18 G.W. Everett, Jr. and R.H. Holm, *J. Amer. Chem. Soc.*, **87** (1965) 2117 $\text{Bis}(\beta\text{-ketoamino})\text{Ni}^{\text{II}}$ complexes
- 19 D.R. Eaton and W.D. Phillips, *J. Chem. Phys.* **43** (1965) 392 Mixed-ligand complexes
- 20 R.H. Holm, G.W. Everett, Jr. and W.D. Horrocks, *J. Amer. Chem. Soc.* **88** (1966) 1071 Pyridine and picoline complexes
- 21 R.H. Holm, A. Chakravorty and L.J. Theriot, *Inorg. Chem.*, **5** (1966) 625 $\text{Bis}(\text{pyrrole-2-aldimino})$ complexes
- 22 B.B. Wayland, R.S. Drago and H.F. Henneke, *J. Amer. Chem. Soc.*, **88** (1966) 2455 Amides and lactams of Ni
- 23 G.N. La Mar and L. Sacconi, *J. Amer. Chem. Soc.*, **89** (1967) 2282 High-spin pentadentate Schiff bases
- 24 G.N. La Mar, *Mol. Phys.*, **12** (1967) 427 Schiff base complexes of Ni
- 25 G.N. La Mar, *Inorg. Chem.*, **6** (1967) 1921 Macrocyclic Schiff bases of Ni
- 26 D.R. Eaton, A.D. Josey and R.E. Benson, *J. Amer. Chem. Soc.*, **89** (1967) 4040 Aminotroponimineates $-\sigma$ systems
- 27 D.R. Eaton and W.R. McClellan, *Inorg. Chem.*, **6** (1967) 2134 Conjugation involving Group IV atoms
- 28 N.A. Matwyoff and S.V. Hooker, *Inorg. Chem.* **6** (1967) 1127 Ni and Co perchlorates in CH_3CN
- 29 D.K. Ravage, T.R. Stengle and C.H. Langford, *Inorg. Chem.*, **6** (1967) 1252 Ni^{2+} CH_3CN
- 30 R.E. Ernst, M.J. O'Connor and R.H. Holm, *J. Amer. Chem. Soc.*, **90** (1968) 5735 Distereoisomeric complexes
- 31 R.E. Ernst, M.J. O'Connor and R.H. Holm, *J. Amer. Chem. Soc.*, **90** (1968) 5305 Enantiomeric Ni-complex in optically active solvent.
- 32 M.L. Wicholas and R.S. Drago, *J. Amer. Chem. Soc.* **90** (1968) 6946 Bipyridyl and substituted bipyridyl complexes
- 33 R.J. Fitzgerald and R.S. Drago, *J. Amer. Chem. Soc.* **90** (1968) 2523 Ni^{II} alkylamine complexes.
- 34 K. Fricke and H. Suhr, *Ber. Bunsenges. Phys. Chem.*, **72** (1968) 430 $\text{Ni}(\text{acac})_2$ + amines
- 35 G.N. La Mar and L. Sacconi, *J. Amer. Chem. Soc.*, **90** (1968) 7216 σ -Coordinated complexes
- 36 D.H. Gerlach and R.H. Holm, *J. Amer. Chem. Soc.*, **91** (1969) 3457 β -Aminothiones

- 37 I Bertini, D L Johnston and W D Horrocks, *J Chem. Soc., D*, (1969) 1471. Schiff bases
- 38 G N La Mar and E O Sherman, *J Chem. Soc., D* 14 (1969) 809 Kinetics of planar to tetrahedral isomerization
- 39 I S. Frankel, *J Mol Spectry*, 29 (1969) 273 Decoupling by paramagnetic ions
- 40 G N La Mar, *Inorg Chem.*, 8 (1969) 581 Pyridine adducts of hetero atom substituted (acac)
- 41 J R Hutcheon, G N La Mar and W D Horrocks, *Inorg Chem*, 8 (1969) 126 Pyridylnitrone complexes
- 42 R J Fitzgerald and R S Drago, *Inorg Chem*, 8 (1969) 2254 Diaminobutane complexes
- 43 Z Luz, *J Chem. Phys*, 51 (1969) 1206 $\text{Ni}(\text{MeOH})_5\text{Cl}^+$
- 44 L H Pignolet, W D Horrocks and R H Holm, *J Amer Chem Soc*, 92 (1970) 1855 Thermodynamics and kinetics of planar-tetrahedral equilibrium of Ni-phosphines
- 45 G N La Mar and E O Sherman, *J Am Chem Soc*, 92 (1970) 2691 Kinetics of square planar-tetrahedral isomerization
- 46 R E Cramer and R S Drago, *J Amer Chem Soc*, 92 (1970) 66 Substituted pyridines
- 47 I Bertini, D L Johnston and W D Horrocks, *Inorg Chem*, 9 (1970) 693 Tetrahedral benzaldimine complexes
- 48 F L Ho, L E Erickson, S R Watkins and C N Reilly, *Inorg Chem*, 9 (1970) 1139 Amino acid complexes.
- 49 L E Erickson, F L Ho and C N. Reilly, *Inorg Chem.*, 9 (1970) 1148 Iminodiacetic acid and nitrilotriacetic acid complexes
- 50 I Bertini, D L Johnston and W D Horrocks, *Inorg Chem*, 9 (1970) 698 5-Coordinated benzaldimine complexes.
- 51 C E Strouse and N A Matwiyoff, *J Chem Soc., D*, (1970) 439 ^{13}C contact shifts in amine-Ni compounds
- 52 R W. Kluber, R Kukla and W D Horrocks, *Inorg Chem* 9 (1970) 1319 Ligand exchange kinetics of 4-picoline
- 53 J W Rakshys, Jr, *Inorg Chem*, 9 (1970) 1521. Aniline adducts of $\text{Ni}(\text{acac})_2$
- 54 R W Kluber and W D Horrocks, *Inorg Chem. Acta*, 4 (1970) 183 Imine and nitrone adducts of $\text{Ni}(\text{acac})_2$
- 55 D Dodrell and J D Roberts, *J Amer Chem Soc* 92 (1970) 4484 ^{13}C Ni^{II} aminotroponemimines.
- 56 J I Zink and R S Drago, *J Amer Chem Soc*, 92 (1970) 5339 Contact and dipolar shifts

H Copper

- 1 N C Li, R L Scruggs and E D Becker, *J Amer Chem Soc*, 84 (1962) 4650 Amino-acid and peptide complexes
- 2 B B Wayland and W L Rice, *J Chem Phys*, 45 (1966) 3150 Cu- and Ni-amines
- 3 C C Hinckley, *Inorg Chem*, 7 (1968) 396 Acetonitrile
- 4 A F Garito and B B Wayland, *J Amer Chem Soc*, 91 (1969) 866 Pyridine adducts of $\text{Cu}(\text{acac})_2$
- 5 M K Kim and A E Martell, *J Amer Chem. Soc*, 91 (1969) 872 Cu^{II} and Ni^{II} peptide complexes
- 6 C I Lin and N C Li, *J Inorg Nucl Chem* 32 (1970) 2796 DMF complexes to Co β -diketonates
- 7 C H Ke, R J Kurland, O J Lin and N C Li, *J Phys Chem*, 74 (1970) 1728 Pyridine and pyridine oxide adducts of $\text{Cu}(\text{acac})_2$

I Second period

- 1 E W Randall and D. Shaw, *Mol Phys*, 10 (1965) 41 Paramagnetic Re^{III} and Re^{IV} phosphines
- 2 J Chatt, G J Leigh, D M P Mingos, E W Randall and D Shaw, *Chem Commun*, (1968) 419 Re phosphines and arsines

J Actinides

- 1 J C Sheppard and J L Burdett, *Inorg Chem*, 5 (1966) 921 Np-CH₃OH system
- 2 N Paladino, G Lugi, U Pedretti, M Brunelli and G Gia Cometti, *Chem Phys Letters*, 5 (1970) 15 Pseudocontact shift in tetra-allyl-U compounds

K π -Complexes

- 1 H M McConnell and C H Holm, *J Chem Phys*, 28 (1958) 749 and 27 (1957) 314 Nickelocene, vanadocene and chromocene
- 2 Y S Karimov and V M Chibrikov, *J Phys Chem Solids*, 12 (1963) 1683 Dibenzene and ditoluene Cr
- 3 H P Fritz, H J Keller and K E Schwarzhans, *Z Naturforsch*, 21b (1966) 809 Dicyclopentadienyl compounds of V, Cr, Co, Ni and Fe
- 4 H P Fritz, H J Keller and K E Schwarzhans, *J Organometal Chem*, 7 (1967) 105 Ferricinium cation
- 5 H P Fritz, F H Koehler and K E Schwarzhans, *J Organometal Chem*, 16 (1969) 14. Bis-pi-indenyl-Co^{II}
- 6 S E Anderson and R S Drago, *J Amer Chem Soc*, 91 (1969) 3656 Aromatic complexes
- 7 M F Rettig and R S Drago, *J Amer Chem Soc*, 91 (1969) 1361 Metallocenes
- 8 Y S Karimov, V M Chibrikov and L F Schegolov, *J Organometal Chem* 16 (1969) 14 Dibenzene and ditoluene Cr

L Second sphere and ion pairs

- 1 G N La Mar, *J Chem Phys*, 41 (1964) 2992 Pseudo-contact shifts in ion pairs of Co
- 2 G N La Mar, *J Chem Phys*, 41 (1964) 2992 Isotropic shifts in Co and Ni ion pairs
- 3 T R Stengle and C H Langford, *J Phys Chem*, 69 (1965) 3299 Relaxation Cr³⁺ ion pairs
- 4 G N La Mar, *J Chem Phys*, 43 (1965) 235 Ion pairs and ionic distances in Co, Ni compounds
- 5 W D Horrocks, R H Fischer, J R Hutchison and G N La Mar, *J Amer Chem Soc*, 88 (1966) 2436 Ion pairing in Ni, Co compounds
- 6 G N La Mar, R H Fischer and W D Horrocks, *Inorg Chem*, 6 (1967) 1798 Ion pairs, interionic distances, shifts in Ni, Co compounds
- 7 R H Fischer and W D Horrocks, *Inorg Chem*, 7 (1968) 2659 Ion pairs and interionic distances
- 8 P K Burkart, H P Fritz, W Gretner, H J Keller and K E Schwarzhans, *Inorg Nucl Chem Letters*, 4 (1968) 237 Contact shifts of tetra-alkyl ammonium ions
- 9 I M Walker and R S Drago, *J Amer Chem Soc*, 90 (1968) 6951 Ion pairing in tetrahedral complexes
- 10 J C Fanning and R S Drago, *J Amer Chem Soc*, 90 (1968) 3987 Ion pairing with hexacyanoferrate
- 11 D R Eaton, *Can J Chem*, 47 (1969) 2645 2nd-sphere Co^{II} complexes
- 12 A J Leffner, *J Phys Chem*, 74 (1970) 2810 2nd-sphere solvation
- 13 L S Frankel, *J Phys Chem*, 74 (1970) 1645 Orientation of 2nd-sphere ligands
- 14 D G Brown and R S Drago, *J Amer Chem Soc*, 92 (1970) 1871 Fermi contact shifts in ion pairs
- 15 H P Fritz, W C Gretner, H J Keller and K E Schwarzhans, *Z Naturforsch*, 25b (1970) 174 Ion pairs Co^{II} and Ni^{II}

R Reviews

- 1 D R Eaton and W D Phillips, *Advan Magnet Resonance*, 1 (1966) 103
- 2 E de Boer and H Van Wuijgen, *Progr NMR Spectry*, 2 (1967) 111
- 3 R H Holm, *Accounts Chem Res*, 2 (1969) 307
- 4 H J Keller and K E Schwarzhans, *Angew Chem Intern Ed Engl*, 9 (1970) 196