

## IRON (1984)

Peter Thornton

### CONTENTS

1	INTRODUCTION	132
2	GENERAL	132
3	IRON(IV)	132
4	IRON(III)	133
4.1	Cyanide complexes	133
4.2	Complexes of anionic nitrogen ligands	134
4.3	Amine complexes	134
4.4	Pyridine complexes	135
4.5	Diimine complexes	135
4.6	Schiff base complexes	136
4.7	Oxide and oxo-complexes	137
4.8	Hydroxides	140
4.9	Alkoxides and complexes with OR ligands	141
4.10	Carboxylates	141
4.11	Diketonates	144
4.12	Complexes of oxyanions	145
4.13	Water complexes	146
4.14	Complexes of neutral oxygen donors	146
4.15	Sulfides	147
4.16	Thiolates and other SR ligands	147
4.17	Dithiocarbamates	147
4.18	Complexes of other sulfur ligands	148
4.19	Fluorides and fluoro-complexes	148
4.20	Chlorides, bromides, iodides and halide complexes	149
4.21	Other iron(III) complexes	151
5	IRON(II)	151
5.1	Cyanide complexes	151
5.2	Phthalocyanines and other complexes of anionic nitrogen ligands	152
5.3	Ammonia complexes	153
5.4	Amine complexes	153
5.5	Pyridine complexes	153
5.6	Diimine complexes	153
5.7	Schiff base complexes	155
5.8	Complexes of phosphorus ligands	156
5.9	Oxides	156
5.10	Hydroxide and hydroxo-complexes	156
5.11	Complexes of OR ligands	156
5.12	Carboxylates, amino acid and related complexes	156
5.13	Diketonates	157
5.14	Complexes of oxyanions	157
5.15	Water complexes	158
5.16	Complexes of neutral oxygen donors	159
5.17	Sulfides and sulfido-complexes	159
5.18	Complexes of other sulfur ligands	160
5.19	Fluorides and fluoro-complexes	160
5.20	Other halides and halide complexes	160
5.21	Hydrides and hydrido complexes	162
5.22	Other iron(II) complexes	162
5.23	Mixed valence compounds	162

6	BIOLOGICAL IRON COMPOUNDS	162
6.1	Haem proteins	163
6.2	Peroxidases	168
6.3	Cytochromes	169
6.4	Porphyrin complexes	176
6.5	Iron-sulfur complexes	181
6.6	Iron-molybdenum-sulfur compounds	186
6.7	Haemerythrin	187
6.8	Siderophores	188
6.9	Ferritin	189
7	ORGANOIRON COMPOUNDS	190
7.1	Carbonyls and related compounds	190
7.2	Nitrosyls	197
7.3	Alkyls and aryls	198
7.4	Alkene complexes	203
7.5	Alkyne complexes	204
7.6	Diene complexes	205
7.7	Cyclopentadienyls and related compounds	209
	References	218

### 1. INTRODUCTION

This review covers the developments in the chemistry of compounds of iron reported in 1984. For operational reasons it has been necessary to consider only papers cited in Chemical Abstracts for that year. Therefore a little work from 1983 is included and some papers from late 1984 will be discussed in the survey for 1985.

The order of topics is that of decreasing oxidation state, but there are separate sections for biological and organometallic compounds.

### 2. GENERAL

X-ray photoelectron (XPE) spectra have been obtained [1] for many iron complexes and reveal a clear correlation between spin states and the satellite intensity in the Fe 2p spectra.

### 3. IRON (IV)

Moessbauer spectra show [2] the presence of high spin Fe(IV) in the perovskite type oxides  $\text{Ca}_{1-x}\text{Sr}_x\text{FeO}_3$  ( $x \leq 1$ ) or  $\text{Sr}_{1-x}\text{La}_x\text{FeO}_3$  ( $x < 0.6$ ), but at low Sr concentrations  $\text{Fe}^{4+}$  disproportionates to  $\text{Fe}^{5+}$  and high spin  $\text{Fe}^{3+}$ . Another perovskite material,  $\text{Ca}_x\text{La}_{1-x}\text{FeO}_y$ , also contains  $\text{Fe}^{4+}$ ; this was studied [3] by electron microscopy. The enthalpy of solution of  $\text{SrFeO}_{2.97}$  in 6.32M HCl at 298K is  $-214 \text{ kJ mol}^{-1}$  [4].

Among Fe(IV) coordination compounds, an alternating chain exchange mechanism was used [5] to account for the magnetochemistry of  $[\text{FeO}(\text{hemiporphyrizinate})]$ ,

whose doublet ground state was established by Moessbauer spectra. An Fe(IV) porphyrin complex with  $O^{2-}$  is invoked [6] to account for some electron transfer reactions of the Fe(III) - PPIXDME complex studied by cyclic voltammetry, ring-disc voltammetry and electronic spectra. Another example has been found [7] in which  $H_2O_2$  oxidises an Fe(II) porphyrin to an  $[FeO]^{2+}$  species. In related chemistry  $[Fe(MeCN)_4(ClO_4)_2]$  disproportionates [8]  $H_2O_2$  to form an  $[FeO]^{2+}$  complex and  $^1O_2$ ; this system oxygenates such varied organic molecules as ethanol or  $PPh_3$ . It is also suggested [9] that Fe(IV) is formed when  $Fe^{2+}$  reduces a Ti(IV) peroxo complex in sulphuric acid.

#### 4. IRON (III)

##### 4.1. Cyanide Complexes

There are a number of structural studies on  $[Fe(CN)_6]^{3-}$  salts.  $Li_3[Fe(CN)_6] \cdot 2(\text{hexamethylenetetramine}) \cdot 5H_2O$  has been shown [10] by X-rays to have the expected octahedral anion but Moessbauer spectra show there are two slightly different Fe environments. In  $(Me_4N)_2 Na[Fe(CN)_6] \cdot H_2O$  an  $[Na(H_2O)]^+$  moiety is coordinated [11] to three N atoms, unsymmetrically occupying an  $N_6$  cavity.  $Zn_3[Fe(CN)_6]_2$  has been prepared [12] anhydrous and hydrated, with structural identification [13] of the usual anion for the  $14 H_2O$  compound. In  $(Me_4N)Mn[Fe(CN)_6] \cdot 8H_2O$  the anion is linked [14] by CN bridges to  $[Mn(H_2O)_4]^{2+}$  units. In other work on  $[Fe(CN)_6]^{3-}$  salts, Raman spectra were used [15] to study the phase transitions at  $\sim 130K$  for the K salt and  $RbPb[Fe(CN)_6] \cdot 3H_2O$  has been prepared. [16] Thermodynamic data have been obtained [17] for the formation of  $\{Ti[Fe(CN)_6]\}^{2-}$ . Linkage isomerism occurs [18] in  $MFe(CN)_6$ , ( $M=Al, Ga, In, Tl$ ). Prussian blue materials in which either  $Fe^{2+}$  or  $Fe^{3+}$  are replaced by other metals have lowspin Fe when it is bound to C, but high spin when it is bound to N. [19]

Electrochemical work on  $[Fe(CN)_6]^{4-} - [Fe(CN)_6]^{3-}$  systems have highlighted the use of this system in photoelectrochemical cells. Such cells

are improved [20] by roughening the electrodes and addition of CdSe.

Spectroscopic studies in this field include reflectance spectra on Ni electrodes [21] and interfacial FTIR spectra [22].

There have been many studies of redox reactions of cyanide complexes of iron. A study [23] of reactions with reduced or oxidised plastocyanin led to a warning that tunnelling rates do not necessarily reflect the distance involved and that rearrangement energies must be considered. A kinetic investigation [24] of the reduction of  $[\text{Fe}(\text{CN})_6]^{3-}$  on Nb-doped  $\text{SrTiO}_3$  electrode surfaces formed part of a study of semiconductor electrode - electrolyte interfaces. Other species oxidised by  $[\text{Fe}(\text{CN})_6]^{3-}$  are pyridinium salts (to pyrroles [25] or other more complex heterocycles [26]), quaternised 2-phenylquinolines [27] (to quinones with loss of Ph.), cresols, [28] nitrotoluenes, [29] phenanthrene, [30] glyceraldehyde phosphate, [31] 10-methylacridinium, [32] resorcinol [33] (first order in reactants and  $\text{OH}^-$ ), and acrylic and lactic acids [34] (zero order in  $[\text{Fe}(\text{CN})_6]^{3-}$ , catalysed by Ru (III)).

Compounds of the type  $\text{KFe}[\text{M}(\text{CN})_8]$  (M = Mo, W) have been prepared. [35] A cyclic voltammetry study of  $[\text{Fe}(\text{CN})_5\text{L}]^{n-}$  (L = various ligands, n = 2 or 3) showed [36] the effect on  $E^\ominus$  was greatest when L was a good  $\pi$ -acceptor.

#### 4.2. Complexes of Anionic Nitrogen Ligands

Amidino complexes  $[\text{Fe}\{\text{RNC}(\text{R})^1\text{NR}\}_3]$  (R = H, Me, Ph, *p*-tolyl) have been prepared [37] and are thought to be associated with Fe-N-C-N-Fe bridges like the isoelectronic carboxylates.

Magnetism studies and EPR and Moessbauer spectra showed [38] that the phthalocyanine (P) complex  $[\text{Fe}(\text{P})\text{Cl}]$  exists in a mixture, not an equilibrium, of  $^4A_2$  and  $^6A_1$  states. Another Fe(III) - Pc complex was one of a number used [39] to modify a Pt electrode for the reduction of  $\text{O}_2$  in fuel cells. Porphyrin complexes are treated in section 6.

#### 4.3. Amine Complexes

The solvent extraction of  $\text{Fe}^{3+}$  by long chain amines such as dinonylamine has

been studied.[40] Formation constants for glucosamine complexes have been determined.[41] The Moessbauer spectrum of an inosine -5'- monophosphate complex revealed [42] the effects of differing ribose configuration on the three iron environments.

The guanine (L) complex  $[\text{FeL}_2\text{Cl}_3]$  is believed [43] to be polymeric with bridging guanines. Physical measurements suggested [44] hydrazine bridges made  $[\text{Fe}(\text{N}_2\text{H}_4)_2(\text{NCS})_2]$  an octahedral polymer. Complexes of dihydrobis (imidazolyl) borate have been prepared. [45]

Hydrolysis of the 1,4,7- triazacyclononane (L) complex  $[\text{FeLCl}_3]$  at pH9 gave [46]  $[\text{Fe}_2\text{L}_2(\text{H}_2\text{O})_2\text{O}_2]^{2+}$  as I,  $\text{PF}_6^-$  or  $\text{ClO}_4^-$  salts, but  $\text{Br}^-$  gave the  $[\text{Fe}_8\text{L}_6\text{O}_2(\text{OH})_{12}]^{8+}$  cation in which 4[FeL] moieties are each linked to one of two other [FeL] units by an OH bridge and to one of two more [FeL] units by 2OH bridges; the latter fragments are linked by an O bridge to Fe which does not carry an amine ligand.

#### 4.4. Pyridine Complexes

IR and Moessbauer spectra of  $\text{FeCl}_3$  in pyridine suggest [47] the formation of a polymer based on  $\text{Fe}(\text{py})_3\text{Cl}_3$  units. The X-ray crystal structure of  $[\text{FeL}_3\text{Cl}_3]$  (L = 4-cyanopyridine) shows [48] this has a *mer*-configuration. 2-Hydroxypyridine forms a 1:1 complex with  $\text{Fe}^{3+}$  with a higher formation constant than its 3- or 4- isomers, which suggests [49] the O atom also coordinates. Pyridine -4-aldehyde (L) forms [50]  $[\text{FeL}_2\text{Cl}_3] \cdot 4\text{H}_2\text{O}$  and  $[\text{FeL}_2(\text{H}_2\text{O})_4]^{3+}$  complexes. Formation constants for pyridoxine- $\text{FeCl}_3$  complexes have been determined. [51]

The addition of  $\text{Co}^{2+}$  to  $[\text{Fe}(2\text{-picolylamine})_3]\text{Cl}_2 \cdot \text{EtOH}$  affects [52] the temperature dependance of the highspin-lowspin transition, in line with the authors' model of "elastic" interactions between highspin and lowspin ions.

#### 4.5. Diamine Complexes

There has been much structural work on bipy and phen complexes. Those studied are  $[\text{Fe}(\text{bipy})_3]^{2+}$ ,  $[\text{Fe}_2\text{OC}_6\text{H}_6]^{2-}$ , *cis*- $[\{\text{Fe}(\text{phen})_2\text{Cl}\}_2\text{O}]^{2+}$ ,  $2\text{Cl} \cdot 4.5\text{H}_2\text{O}$  [53],  $[\text{Fe}(\text{phen})\text{Cl}_3\text{L}]$  (L =  $\text{H}_2\text{O}$ , MeOH, 1,2,4-triazole)

(*fac*-isomers) [54],  $[\text{Fe}(\text{phen})\text{Cl}_4]^-$  [54],  $[\text{FeL}_2\text{Cl}_2]^+$  (L = bipy, phen) as  $[\text{FeCl}_4]^-$  or  $[\text{Fe}(\text{LCl}_4)]^-$  salts [55], with a neutron diffraction study of a lattice bipy showing [56] a small distortion in this molecule, and a polarized neutron diffraction study of the  $[\text{FeCl}_4]^-$  salt revealing [57] the spin distribution. In  $[\{\text{Fe}(\text{phen})_2(\text{H}_2\text{O})_2\}_2\text{O}]^{4+} \cdot 4\text{NO}_3 \cdot 5\text{H}_2\text{O}$  a typical structure was found [58] with a bridging FeOFe angle of  $155.1^\circ$ ;  $\nu_s(\text{FeOFe})$  and  $\nu_{as}(\text{FeOFe})$  were located with  $^{18}\text{O}$  and are excited in the Raman spectrum by the  $\text{O}^{2-}$ -Fe(III) CT band. These results were used to claim that the complex is a good model for haemerythrin and ribonucleotide reductase.

The complex  $[\text{Fe}(\text{phen})\text{Cl}_3]$  (dmf) was found [59] as an impurity in the preparation of  $[\text{Fe}(\text{phen})_3]^{2+}[\text{Fe}_2\text{Cl}_6\text{O}]^{2-}$ . Formation constant determinations suggested [60] that bipy forms  $[\text{Fe}(\text{OH})_2(\text{bipy})_n]^+$  ( $n = 1, 2$ ) and  $[\text{Fe}(\text{OH})(\text{bipy})_2]^{2+}$  at pH 3.5 - 5.0.

There have been many kinetic studies of  $[\text{Fe}(\text{bipy})_3]^{3+}$  and  $[\text{Fe}(\text{phen})_3]^{3+}$  reactions. The dissociation and racemization of these in SDS solutions showed [61] micelle formation effects. Their reduction by  $\text{OH}^-$  was reinvestigated [62] with a discussion of the relative probabilities of OH adduct formation and deprotonation of the ligand. The oxidation of bromomalonic acid by  $[\text{Fe}(\text{phen})_3]^{3+}$  has been studied, [63] as has the oscillating reaction [64] of this system with  $\text{BrO}_3^-$  and the cation's oxidation of 4-methoxytoluene by an outer sphere process. [65] There has also been a kinetic study [66] of the  $\text{Fe}^{3+}$  oxidation of  $\text{NH}_2\text{OH}$  to  $\text{N}_2$  in the presence of phen and an unspecified Fe(III) - phen complex was the best catalyst for the decomposition [67] of  $\text{O}=\text{C}(\text{CH}_2)_3\text{C}(\text{Me})\text{OOH}$ . See also ref.147.

#### 4.6. Schiff Base Complexes

This section includes various  $\text{>C} = \text{NR}$  ligands, including semicarbazones, oximes and hydrazones. Time and space restrictions prevent detailed discussion of much work or giving structural diagrams.

Details are given [68] of the preparation of  $[\text{FeL}(\text{MeCN})_2]^{2+}$  where L is a macrocycle produced from  $\text{PhCOCOME}$  and  $\text{NH}_2\text{C}_3\text{H}_6\text{NH}_2$ . In

$[\text{Fe}(\text{tim})(\text{SCH}_2\text{Ph})_2]^+$  there are two cations in each unit cell, one with the axial phenyls tilted away from the macrocycle plane, the other with them tilted towards it. [69] Another crystal structure shows [70] square pyramidal coordination in a salicylidene semicarbazide complex.

There has been some interesting magnetochemistry. An Fe(III) bis complex of the salicylaldehyde -  $\text{N} - (3 - \text{aminopropyl}) - \text{aziridine}$  Schiff base is a spin-crossover compound [71] whose magnetism is affected by grinding and by solvation by  $\text{CH}_2\text{Cl}_2$ ; an unusual result is that the rate of spin change is faster than the Moessbauer time scale if the material is more crystalline. [72] The spin state of complexes of related 14 to 16 membered macrocycles seems to depend [73] on axial ligands as well as ligand ring size, with spin states of  $1/2$ ,  $3/2$  and  $5/2$  all being found. Bis(imidazole) iron (III) complexes of the tetradentate ligand formed from hydroxyacetophenone, ethylenediamine and 2,4-pentanedione have ESR and visible spectra for a highspin - lowspin equilibrium, being thermochromic. [74] The magnetism of  $[\text{Fe}_2\text{L}_2\text{O}]$  (L = tetradentate isothiosemicarbazide) has been studied, [75] and the mixed metal complex  $[(\text{PhOFe})\text{L}(\text{CuCl}_2)]$  (L is a  $\text{sal}_2\text{pn}$  derivative) is also antiferromagnetic. [76]

In solution chemistry some binuclear complexes with a  $\text{sal}_2\text{en}$  type ligand have been studied by differential pulse polarography [77], and a kinetic study has been made [78] on the electrolytic reduction of  $[\text{Fe}(\text{sal}_2\text{en})\text{Cl}]$ . The hydrolysis of an Fe(III)  $\text{sal}_2\text{en}$  carboxy derivative at various pH shows the sixth coordination site is occupied by  $\text{H}_2\text{O}$  or  $\text{OH}^-$ ; an  $\text{H}_3\text{O}_2^-$  bridged complex is also suspected. [79] Many other, mostly preparative, studies require no detailed comment. [80 - 99]

#### 4.7. Oxides

This section omits many reports where only small proportions of Fe are present or where the work is primarily technological in character.

The breakdown of passive films on Fe by  $\text{Cl}^-$  has been described [100] as involving the formation of a polymeric hydrated oxide.

There have been many studies on  $\text{Fe}_2\text{O}_3$ . For  $\alpha$ - $\text{Fe}_2\text{O}_3$  these include the preparation of needles, [101] the DTA of its formation from  $\gamma$ - $\text{FeOOH}$ , [102] the anisotropy of its electrical properties, [103] the effect of particle size on its Moessbauer spectrum, [104] and on the catalysis of the oxidation of 1-butene to butadiene, [105] the kinetic study of its solution in acids using  $\text{Me}_2\text{COH}$  radicals and  $[\text{Fe}(\text{edta H}_2)]$  [106]. For  $\gamma$ - $\text{Fe}_2\text{O}_3$  there have been studies of preparation by sputtering, [107] neutron diffraction, [108] temperature dependence of the Moessbauer spectrum, [109] and the transition to  $\alpha$ - $\text{Fe}_2\text{O}_3$ . [110] It was not specified which form of  $\text{Fe}_2\text{O}_3$  was formed [111] in the pyrolysis of  $[\text{Fe}(\text{urea})_6]^{3+} \cdot 3\text{NO}_3^-$ . Together with  $\text{MgO}$  or  $\text{Al}_2\text{O}_3$ ,  $\text{Fe}_2\text{O}_3$  cyclises [112] 1,3-pentadiene to cyclic hydrocarbons.  $\text{Fe}_2\text{O}_3$  absorbs  $\text{H}_2\text{S}$  strongly, but if  $\text{O}_2$  is present S is formed at low and  $\text{SO}_2$  at high temperatures. [113] The hydrogenation of CO over  $\text{Fe}_2\text{O}_3$  -  $\text{SiO}_2$  is most successful when  $\text{FeCl}_3$  is used to prepare the catalyst. [114] In Ni-doped  $\text{Fe}_2\text{O}_3$  the O vacancies play a key role in the oxidation of CO. [115] The Moessbauer spectrum shows [116] at least two components with different spin are present in the high pressure phase of  $\text{Fe}_2\text{O}_3$ . Unspecified Fe oxides enhance the Raney Ni - Zn catalysed hydrogenation of 2-ethylanthraquinone. [117]

Defect-free crystals of  $\text{Fe}_3\text{O}_4$  of at least 5g weight can be made [118] by a method called "skull-melting". Gas-phase reactions of  $\text{FeO}^+$  with alkanes have been observed [119] using collision induced dissociation mass spectrometry.

There have been many reports of work on mixed oxides which contain  $\text{Fe}^{3+}$ . These include the determination [120] of the free energy of formation of  $\text{LiFeO}_2$ , the preparation of the  $\text{Na}_2\text{O}$ -structured  $\text{Li}_5\text{FeO}_4$  [121] and the X-ray crystal structures [122] of the ordered and disordered forms of  $\text{LiFe}_5\text{O}_8$ . A thermodynamic study [123] of the Fe-Na-O system gave  $\Delta G^\circ$  for



$\text{Na}_2\text{FeO}_2$ ,  $\text{Na}_2\text{Fe}_2\text{O}_4$  and  $\text{Na}_4\text{FeO}_3$ . In work with alkaline earth oxides, the magnetoplumbite structure (based on  $\text{Fe}_3\text{O}_4$ ) was found [124] for  $\text{Ca}_{2.95}\text{Fe}_{14.85}\text{O}_{25}$ , and a solubility study of  $\text{BaFe}_{12}\text{O}_{19}$  in  $\text{BaO-B}_2\text{O}_3$  suggests [125] this is a good medium for growing single crystals. Mixed oxides with transition metal 2+ ions have been studied, including the formation [126] of  $\text{Fe}_2\text{NiO}_4$  from the coprolysis of their Tutton salts, the growth [127] of long crystals of  $\text{Fe}_2\text{MO}_4$  ( $\text{M} = \text{Co}, \text{Ni}$ ). These and other ferrites ( $\text{Fe}_2\text{MO}_4$ ) have been made [128] by the pyrolysis of  $\text{Fe}_2\text{M}(\text{C}_2\text{O}_4)_3(\text{N}_2\text{H}_6)$ , ( $\text{M} = \text{Mn}, \text{Co}, \text{Ni}, \text{Zn}$ ). The magnetic anisotropy of  $\text{Co}_{0.16}\text{Fe}_{2.84}\text{O}_4$  has been reported. [129] Fast electron exchange was found [130] in  $\text{Fe}_2\text{Cu}_x\text{Ge}_{1-x}\text{O}_4$  ( $0.6 < x < 0.8$ ). The Moessbauer spectrum shows [131]  $\text{Fe}^{3+}$  occupies both octahedral and tetrahedral sites in  $\text{FeAlO}_3$ , but most  $\text{M}^{3+} - \text{Fe}^{3+}$  mixed oxides involve lanthanide ions. A hydrothermal synthesis of  $\text{FeLaO}_3$  crystals is reported. [132] CO oxidation by  $\text{FeLa}_{0.8}\text{Sr}_{0.2}\text{O}_3$  rapidly increases [133] above 240 °C. X-ray and IR studies of  $\text{LaFe}_x\text{Ni}_{1-x}\text{O}_3$  are recorded. [134] There has been a kinetic study [135] of garnet formation in sintering  $\text{Fe}_2\text{O}_3$  with  $\text{Y}_2\text{O}_3$ , and IR, DTA, and X-ray diffraction studies of this system showed [136] the formation of  $\text{YFeO}_3$  and  $\text{Y}_3\text{Fe}_5\text{O}_{12}$ . Ordering temperatures have been found [137] for  $\text{Y}_2\text{FeSbO}_7$ ,  $\text{Bi}_2\text{FeSbO}_7$  and  $\text{BiYFeSbO}_7$ . Moessbauer spectra showed [138]  $\text{Sc}^{3+}$  occurs in tetrahedral and dodecahedral sites in  $\text{Eu}_{3-x}\text{Sc}_{2+x}\text{Fe}_3\text{O}_{12}$ . Antiferromagnetic resonance has been studied [139] in  $\text{FeTmO}_3$ . There is an unusually low absorption [140] at 1300 nm in  $\text{Gd}_{0.2}\text{Y}_{2.8}\text{Fe}_5\text{O}_{12}$ . Some mixed oxides formally contain  $\text{M}^{4+}$ . The  $\text{Fe}_2\text{O}_3 - \text{GeO}_2$  system produces [141]  $\text{Fe}_2\text{GeO}_5$  and  $\text{Fe}_8\text{Ge}_3\text{O}_{18}$ , the latter being a new structure type, but if Fe is present,  $\text{Fe}_4\text{Ge}_2\text{O}_9$ , paramagnetic above 100K, is formed. [142] The  $\text{BaM}_2\text{Fe}_4\text{O}_{11}$  ( $\text{M} = \text{Sn}, \text{Ti}$ ) ferrites have  $\text{Fe}^{3+}$  in octahedral and tetrahedral sites, but those in the latter can move between two sites. [143] There have been X-ray powder and electron microscopy studies

[144] of  $\text{Fe}_2\text{TiO}_5$ .

Bridging oxo ligands occur in  $[\{\text{FeCl}_3\}_2\text{O}]^{2-}$ , studied as alkylammonium

[145] or  $[\text{Mg}(\text{dmf})_6]^{2+}$  [146] salts, and in the *fac*-configured

$[\{\text{Fe}(\text{H}_2\text{O})_3(\text{phen})_2\}_2\text{O}]^{4+}$  cation. [147]

#### 4.8. Hydroxides

A thorough review has been published [148] on the hydrolysis of Fe(III) salts.

The conditions are described [149] for preparing the  $\text{Fe}(\text{OH})_3$  catalyst for reducing NO to  $\text{NH}_3$ . A new form of  $\text{Fe}(\text{OH})_3$  has been prepared [150] by the

oxidation of  $\text{FeSO}_4$  aqueous solutions by  $[\{\text{Co}(\text{en})(\text{dien})\}_2\text{O}_2]^{4+}$ ; it has

two Fe sites of about equal abundance, is paramagnetic above 190K,

superparamagnetic from 97.5 to 190K and ordered below 97.5K. Magnetization

measurements near 1K were used to find [151] the proportion of hydroxo

oligomers in solutions of  $\text{Fe}^{3+}$  salts in aqueous  $i\text{-PrOH}$ .

Most work in this section concerns  $\text{FeOOH}$ . To make the  $\alpha$ -form free from

$\text{Fe}_3\text{O}_4$  one must use aqueous  $\text{FeSO}_4$  at pH between 5 or 12 at below room

temperature. [152] Other work has been published [153] on this reaction. An

IR study [154] concentrated on surface behaviour and the transition to

$\text{Fe}_2\text{O}_3$ . Conditions have been described [153, 155] for forming  $\beta$ - $\text{FeOOH}$  by

hydrolysis. The Neel temperature of  $\beta$ - $\text{FeOOH}$  depends on the preparative

conditions, with interstitial  $\text{H}_2\text{O}$  having an effect. [156] A discussion

[157] of  $\beta$ - $\text{FeOOH}$   $\text{Cl}_n$  particles, synthetic or natural, dealt with the

possible role of this material in intercalating aminoacids for prebiotic

polypeptide formation. The dehydration in vacuo of  $\beta$ - $\text{FeOOH}$  was monitored

[158] by magnetization studies. The  $\gamma$ -form of  $\text{FeOOH}$  was formed [159] in the

air oxidation of aqueous  $\text{FeCl}_2$  at pH 2-5. The light induced dissociation of

$\gamma$ - $\text{FeOOH}$  is assisted when citrate is present. [160] Neutron diffraction and

Moessbauer experiments showed [161] ferromagnetic coupling between layers of

$\text{Fe}^{3+}$  and antiferromagnetic coupling between layers. The deposition of

hydrated  $\text{FeOOH}$  on gold electrodes has been studied. [162]


#### 4.9. Alkoxides and Complexes with $OR^-$ Ligands

This section includes phenoxides, catecholates and complexes of polydentate ligands in which an  $OR^-$  entity plays a significant role.

The formation of 1:1 Fe(III) complexes of nitro-, dinitro- and methylphenoxides in dmsO was studied [163] by EPR, which showed reduction to Fe(II) was easy. Two crystal structures of Fe(III) complexes containing bridging OEt ligands have been reported; [164] one has a binucleating ligand, the other is  $[Fe_2(acac)_4(OEt)_2]$ . Comparison of the magnetochemistry of the two compounds and of similar OMe bridged ones showed J values did not follow the usual rules for antiferromagnetism and the authors wondered whether ferromagnetism was also involved. One of many bimetallic isopropoxides studied [165] is  $FeAl_3(O^iPr)_{12}$ .

Among studies of catechol type complexes there have been a kinetic study [166] of complex formation, a study [167] of substituent effects on formation constants with reference to modelling siderophores and determination [168] of the formation constants of 4-nitrocatechol complexes. Formation constants have also been determined [169] in the  $Fe^{3+}$  - pyrogallol - edta system. Fe(III) is photoreduced in solutions containing galloxyanine, which may use both catechol and azole-carboxylate bidentate sites; the system might be a model for bleomycin complexes. [170]

There has been a normal vibrational analysis [171] of a tris complex with N-hydroxyurea. Alkoxo bridges have been found [172] in



(2-hydroxyethyl)amine]; this antiferromagnetic compound shows pentagonal bipyramidal Fe coordination. An Fe(III) complex of catechol-3,5-disulphonate catalyses the conversion of benzenes to phenols. [173] A complex with an azocresol ligand is reported. [174]

#### 4.10. Carboxylates

This section includes monocarboxylates, dicarboxylates, aminoacid complexes, edta complexes and those of other ligands containing  $O_2CR^-$  units.

Studies of allegedly simple  $\text{Fe}(\text{O}_2\text{CR})_3$  compounds include the thermal decomposition [175] of iron (III) benzoate and  $[\text{Fe}_2(\text{fumarate})_3 \cdot 5\text{H}_2\text{O}]$ ,  $\text{Fe}(\text{O}_2\text{CR})_3$  species are claimed [176] as participating in the solvent extraction by  $\text{C}_{m+2n-1}(\text{CH}_2)_n\text{CO}_2\text{H}$  in benzene ( $m = 5, 6$ ;  $n = 0-3$ ). The extraction of  $\text{Fe}^{3+}$  from  $\text{H}_2\text{O}$  by octanoic acid or  $\alpha, \alpha$ -dialkylcarboxylates is assisted by the presence of Al or Cr, which give heteropolynuclear complexes. [177] The reaction of  $[(\text{CpCrS})(\text{SCMe}_3)\text{Fe}(\text{CO})_3]$  with carboxylic acids gives [178]  $[\text{Cr}_2\text{FeS}(\text{O}_2\text{CPh})_6]$  or  $[(\text{CpCr})_3(\text{FeO}_2\text{CCMe}_3)_4\text{S}_4]$ ; the structure of the latter comprises a cubane system with monodentate pivalate. [179] The products of the reaction of  $\text{FeCl}_3$  with  $\text{CF}_3\text{CO}_2\text{H}$  are given [180] as  $[\text{Fe}_2\text{Cl}_2(\text{O}_2\text{CCF}_3)_4]$ ,  $[\text{Fe}_2\text{Cl}(\text{O}_2\text{CCF}_3)_5]$  and  $\text{Fe}(\text{O}_2\text{CCF}_3)_3$ , which are all believed to have polymeric structures with bidentate carboxylates and octahedral Fe.

The pyrazolylborate complexes  $[\text{Fe}_2(\text{O}_2\text{CMe})_2(\text{HBpz}_3)]$  and  $[\text{Fe}_2(\text{OH})(\text{O}_2\text{CMe})_2(\text{HBpz}_3)_2]^+$  have been thoroughly investigated. They both have three bridging ligands, are antiferromagnetic and are claimed to be analogues of haemerythrin. [181, 182]

The magnetochemistry of the familiar  $[\text{Fe}_3\text{L}_3\text{O}(\text{RCO}_2)_6]^+$  structure has been investigated [183] for aqua complexes of unsaturated acids and the (l. = py, R =  $\text{CF}_3$ ) complex has been prepared [184], but

$\text{Fe}_3\text{O}(\text{O}_2\text{CCF}_3)_7 \cdot \text{H}_2\text{O}$  has two different Fe sites. [184] The electrochemistry of compounds erroneously described [185] as

$[\text{Fe}_n\text{Cr}_{(3-n)}(\text{O}_2\text{CMe})_6(\text{OH})_2]^+$  is reported. The Fe(II, III, III) mixed valence complex  $[\text{Fe}_3(\text{H}_2\text{O})_3\text{O}(\text{O}_2\text{CPh})_6]$  gives  $\text{Fe}_2\text{O}_3$  on thermal decomposition [186], possibly through the intermediates

$[\text{Fe}_3(\text{OH})_2\text{O}(\text{O}_2\text{CPh})_4]^+$  and " $\text{Fe}(\text{O}_2\text{CPh})_3$ ". Moessbauer spectra are recorded [187] for  $[\text{Fe}_3\text{O}(\text{O}_2\text{CR})_6]$ , (R =  $\text{CF}_3$ ,  $\text{CCl}_3$ ), at different temperatures. Two  $[\text{Fe}_3\text{O}]$  triangles are fitted together in

$[\text{Fe}_4\text{O}_2(\text{H}_2\text{O})_6(\text{O}_2\text{CCF}_3)_8] \cdot 2\text{H}_2\text{O}$ , which has one carboxylate bridge between each pair of Fe atoms, leaving half the  $\text{O}_2\text{CCF}_3$  ligands

monodentate; the compound is prepared [188] by evaporation in air of solutions of  $[\text{Fe}_3(\text{H}_2\text{O})_3\text{O}(\text{O}_2\text{CCF}_3)_6] \cdot 3.5\text{H}_2\text{O}$ . The complex " $\text{Fe}_8\text{L}_2(\text{L}')_2\text{O}(\text{O}_2\text{CMe})(\text{OH})_{19}$ ", (HL = lactobionic acid, L' = dextran glucopyranose) has been used [189] for parenteral iron therapy.

Research on oxalate complexes includes the use of  $\text{K}_3[\text{Fe}(\text{C}_2\text{O}_4)_3]$  as catalyst for the photochemical dissociation of arenediazonium salts, [190] the thermal decomposition of this salt and  $\text{Fe}_2(\text{C}_2\text{O}_4)_3 \cdot 5\text{H}_2\text{O}$  on various supports [191] and of the ammonium salts of  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$  and the tricitrate complex. [192] The kinetics of the pyrolysis of the mixed metal oxalate complexes  $[\text{Fe}_2\text{Mg}(\text{C}_2\text{O}_4)_8]^{8-}$ ,  $[\text{Fe}_2\text{Mg}(\text{C}_2\text{OH})_4]^{4-}$  and  $[\text{Fe}_2\text{Mg}(\text{C}_2\text{O}_4)_2(\text{OH})_6]^{2-}$  is also reported. [193] A 1:1 complex with maleic acid monoanion is reported. [193a]

The glycine complex  $[\text{FeCl}_3(\text{glycine})_3]$  has been prepared, [194] and the X-ray crystal structure of  $[\text{Fe}_2\text{Cr}(\text{H}_2\text{O})_3\text{O}(\text{glycine})_6]^{7+}$  perchlorate is reported. [195] The  $\text{Fe}^{3+} - \text{H}_2\text{O}_2$  - histidine system catalytically oxidises indigo carmine. [196] Formation constants were measured [197] for  $\text{Fe}^{3+}$  complexes with 3-imidazolylpropenoic acid.

The ever popular complexes of edta continue to appeal. Some readers may be comforted that a repeat [198] of the 1975 determination of the crystal structure of  $\text{Na}[\text{Fe}(\text{H}_2\text{O})\text{edta}]$  gives the same result. The K, Ag and Tl salts are similar. [199] Studies on the adsorption of edta and its Fe complexes on  $\text{Fe}_3\text{O}_4$  clarified [200] the dissolution of the latter, which is faster when  $\text{Fe}^{2+}$  is added; the larger formation constant for the Fe(III) complex over its Fe(II) sister assists electron transfer in dissolution. Another report [201] also discusses this reaction. Two papers [202] describe the use of edta and similar complexes of Fe(III) as mediators in microbial fuel cells because of their fast electrode reactions and reduction by E. coli. Formation constants of hexamethylenediaminetetraacetate complexes are reported. [203] Reports on salicylate-type complexes include two preparations, [204,205] and a kinetic study which showed [206] that edta displaces salicylates from Fe(III)

complexes by an associative mechanism.

Citrate complexes of Fe(III) can be solvent extracted with long chain alkylamines in  $\text{CHCl}_3$ . [207] A kinetic study of complexation of  $\text{Fe}^{3+}$  by citric, diglycolic and tartaric acids showed [208] that 1:1 complex formation involved  $[\text{Fe}(\text{OH})]^{2+}$ . The formation constant of the 2:1 complex with tartrate has been measured; [209] coordination is believed to occur through a carboxyloxygen and two alkoxide oxygens. The  $[\text{Fe}(\text{tartrate})_2(\text{L})]$  complexes, (L = saccharose, sucrose) have been studied [210] spectrophotometrically. Tartrate also forms [211] a mixed metal complex with  $\text{Fe}^{3+}$  and  $\text{Cd}^{2+}$ . Formation constant determinations [212] of complexes with picolinate or other pyridinecarboxylate ligands indicate some are dinuclear. The dipicolinate (X) complexes  $[\text{FeX}_2]^{n-}$ , ( $n = 1, 2$ ) have been studied; [213] kinetic measurements show ascorbate dianion reduces the Fe(III) complex  $10^5$  times faster than the monoanion. The formation of red and blue complexes of  $\text{Fe}^{3+}$  with chromoxane cyanine R, probably a carboxylate ligand, is discussed [214] for use in histological staining.

#### 4.11. Diketonates

Complexes with 2,4-pentanedionate predominate. There has been a kinetic study [215] of the formation of its 1:1 complex with  $\text{Fe}^{3+}$ . For  $[\text{Fe}(\text{acac})_3]$  there have been measurements of  $\Delta H$  of sublimation and fusion, [216] and cyclic voltammetry. [217] This molecule catalyses the cyclodimerization of norbornadiene [218] and the photochemical oxidation of tetralin by  $\text{O}_2$ . [219]  $[\text{Fe}(\text{acac})_3]$  reacts with  $\text{AlEt}_3$  to give  $[\text{Et}_2\text{Al}(\text{acac})]$  and  $[\text{Al}(\text{acac})_3]$ , but the Fe products were not identified. [220]  $[\text{Fe}(\text{acac})_3]$  with  $\text{AlCl}_3$  or  $\text{SnCl}_2$  catalyses the hydrogenation of 1-hexene and 1,3-pentadiene by forming an Fe alkene complex. [221,222]

In work with other diketonates,  $[\text{Fe}(\text{dbzm})_3]$  has been found [223] to catalyse the coupling of  $\text{PhMgBr}$  to  $\text{PhCH:CHBr}$  to give styrene. Formation constants have been measured [224] for complexes with isonitrosoacetylacetone. A kinetic

study [225] of hemiacetal formation in the reaction of  $[\text{Fe}(\text{hfacac})_3]$  with MeOH showed two molecules of the latter are involved in the rate-determining step, one being used to assist proton transfer. The vacuum sublimation of Fe(III) thenoyltrifluorothiodiketonate is described. [226] Fe(III) trifluoro (nicotinoyl) acetate is another spin-crossover compound; 27% of it is high spin at room temperature. [227]  $[\text{Fe}(\text{OH})]^{2+}$  catalyses the ketoenol tautomerisation of 2,4-pentanedione or ethylacetoacetate better than  $\text{Fe}^{3+}$ , which is less effective than some  $\text{M}^{2+}$  ions. [228] Tetracycline may coordinate as a diketonate; there have been kinetic [229] and thermodynamic [230] studies of its complexation with  $\text{Fe}^{3+}$ . Arnebin-I might also coordinate to  $\text{Fe}^{3+}$  as a diketonate. [231]

#### 4.12. Complexes of Oxyanions

This miscellaneous section includes both inorganic and organic ligands. Moessbauer spectra were used [232] to study ordering processes in the Fe(III) layered silicates nontronite and celadonite. Other work on silicates did not include material of special interest to this journal's readers. It was another busy year for phosphates. New compounds were prepared. [233] The crystal structures of synthetic tavorite,  $\text{LiFe}(\text{PO}_4)(\text{OH},\text{F})$  [234] and of  $\text{Na}_3\text{Fe}_2(\text{PO}_4)_2(\text{OH})_2\text{F}$  [235] show  $\text{FeO}_6$  octahedra and  $\text{PO}_4$  tetrahedra, but IR spectra were taken [236] to show  $\text{FeO}_4$  and  $\text{PO}_3^-$  being present in other phosphates. Among polyphosphates there have been more preparations, [233] including  $\text{KFe}(\text{H}_2\text{P}_2\text{O}_7)_2$  [237],  $\text{Fe}_2\text{P}_6\text{O}_{18}$  and its nonahydrate, [238] and  $\text{K}_2\text{Fe}_2\text{P}_8\text{O}_{24}$ . [239] The conditions for precipitation of  $\text{NH}_4\text{Fe}_3(\text{OH})_6(\text{SO}_4)_2$  are described. [240] The Al and Fe structures in  $\text{KM}_3(\text{OH})_6(\text{SO}_4)_2$  have been compared. [241] X-ray determination of the electron distribution in monoclinic  $\text{Fe}_2(\text{SO}_4)_3$  found [242] the electron density bisecting the Fe-O bonds. The pyrolysis of  $\text{Fe}(\text{OH})\text{SO}_4$  and  $\text{Fe}_2\text{O}(\text{SO}_4)_2$  has been studied. [243]  $\text{Fe}_2(\text{SO}_4)_3 \cdot 7\text{H}_2\text{O}$  can catalyse acid esterification. [244] Among other metals, Fe dissolves electrolytically in dmsO

SO<sub>2</sub> to give [245] Fe<sub>2</sub>(dmsc)<sub>10</sub>(S<sub>2</sub>O<sub>7</sub>)<sub>3</sub> (SO<sub>2</sub>. Polymeric octahedral structures with bridging anions are suggested [246] for such selenites as Fe<sub>7</sub>(OH)<sub>3</sub>(SeO<sub>3</sub>)<sub>9</sub>.18H<sub>2</sub>O. The reaction of Fe<sub>2</sub>(MoO<sub>4</sub>)<sub>3</sub> with FeCl<sub>3</sub> at 350° in vacuum gives FeMoO<sub>4</sub> Cl, a layer structure with corner - shared MoO<sub>4</sub> tetrahedra and FeO<sub>4</sub> Cl square pyramids and showing Fe-Fe interactions across layers. [247]

Many organic anionic ligands are of biochemical origin. Formation constants have been determined [248] for arabinose and galactose complexes. Fe(III) complexes of sorbitol, glucuronic acid and glucosamine have been prepared, [249] the last being given a dinuclear formulation. Various complexes of 2' - or 3' - adenosine monophosphates have been prepared, and phosphate oxygen is the probable strongest coordination site. [250] Other complexes reported include those of hydroxyanthraquinones, [251] hydroxamates [252] (of which a special case is one derived from 1-hydroxy-2-indolinone [253]), and carboxylate hydrazides. [254]

#### 4.13. Water Complexes

By studying various Fe(III) chelates whose structures did (or did not) contain coordinated H<sub>2</sub>O it was found [255] that the presence of this was necessary to achieve the Fenton reaction of forming OH in reaction with O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub>. Fe<sup>3+</sup> catalyses the hydroxylation of benzene to phenol by H<sub>2</sub>O<sub>2</sub>. [256] There has been a kinetic study [257] of the oxidation of aqueous SO<sub>2</sub> by Fe<sup>3+</sup> salts.

#### 4.14. Complexes of Neutral Oxygen Donors

New preparations include [FeL<sub>4</sub>Cl<sub>2</sub>]<sup>+</sup>, (L = 2-aminopyridine-N-oxide), [258] and [Fe { "h<sub>2</sub>P(O)CH<sub>2</sub>P(O)Ph<sub>2</sub> }<sub>3</sub>]<sup>3+</sup> [259]. The dmsc ligands are O-bonded in [FeCl(dmsc)<sub>5</sub>]<sup>2+</sup>[Fe<sub>2</sub>OCi<sub>6</sub>]<sup>2-</sup> at 103K but are disordered at higher temperatures. [260]

Vibrational spectra were used [261] to study the conformation change in MeP(E) (OC<sub>6</sub>H<sub>4</sub>OC<sub>2</sub>H<sub>4</sub>)<sub>2</sub>O when it is coordinated to FeCl<sub>3</sub>. [Fe(dmf)<sub>3</sub>Cl<sub>2</sub>]<sup>+</sup> [FeCl<sub>4</sub>]<sup>-</sup> converts ketophenols to benzofurans. [262a] Cryptand (222)



complexes are reported. [262b]

#### 4.15. Sulphides

Material concerning cluster complexes and other biochemical model systems can be found in section 6.

There has been a kinetic study [263] of the effect of  $\text{SO}_4^{2-}$  on the reduction of  $\text{Fe}^{3+}$  by  $\text{S}^{2-}$ . The complex magnetic behaviour of  $\text{KFeS}_2$  was analysed [264] with the Néel temperature assigned as the sharp maximum at 12.5K rather than the broad maximum at 250K; there is exchange between nearest and next-nearest neighbours in chains and between chains.  $\text{FePS}_3$  has a Néel temperature of 120K; the resultant moment is perpendicular to the layer planes, with ferromagnetic nearest neighbour interactions. [265]

#### 4.16. Thiolates and other SR<sup>-</sup> Complexes

Biochemical model compounds are described in section 6. The  $\text{Ph}_4\text{P}$  salts of  $[\text{Fe}(\text{SR})_4]^-$ ,  $[\text{R} = 2,4,6-(^i\text{Pr})_3\text{C}_6\text{H}_2]$  have been prepared and show a large FeSC angle of  $112^\circ$ ; they are stable to self reduction to Fe(II). [266] Complexes of the S-N donors 1-amidino-2-thioureas have been prepared. [267]

#### 4.17. Dithiocarbamates

A new example [268] of this class is  $[\text{Fe}(\text{S}_2\text{NHC}_6\text{H}_4-4\text{-OEt})_3]$ . An extensive study of the structures of many tris (dithiocarbamato) complexes shows that Fe-S distances average 2.30 Å for lowspin and 2.45 Å for highspin compounds; the effect is an apparent increase in the vibrational amplitude of S atoms. [269] For the tris-morpholinedithiocarbamate as  $\text{CH}_2\text{Cl}_2$  solvate highspin was found [270] from 77 to 320K; in solvates of this compound the solvent molecules collect round the morpholine moiety, induce its deformation and bring about a change to lowspin. Notwithstanding this result the unsolvated compound is described [271] as exhibiting a triplet - quintet equilibrium. Other workers [272] concluded that the only factor governing the spin state was steric hindrance which affects the SCS angle, with greater interference giving lowspin compounds.

The structure of  $[\text{FeBr}(\text{S}_2\text{CN}^i\text{Pr}_2)_2]$  is normal. [273] Heat capacity measurements on  $[\text{FeI}(\text{S}_2\text{CNMe}_2)_2]$  at 0.4 - 300K showed [274] 40% monomer and 60% dimer at low temperature and zero field splitting about 100 times greater in magnitude than antiferromagnetic exchange.

#### 4.18. Complexes of Other Sulphur Ligands

New preparations of this type are  $[\text{Fe}(\text{S}_2\text{CC}_6\text{H}_4\text{OH})_3]$  (all three isomers [275]),  $[\text{Fe}(\text{H}_2\text{O})_2\text{X}_2]\text{Cl}$ ,  $[\text{HX} = 3-(4\text{-pyridyl})\text{triazoline-5-thione}]$ , [276]  $\text{FeCl}_3$  complexes of substituted dithiobiurets (some of which may be dimers), [277] and  $[\text{Fe}_2(\text{py})_2(\text{P}_2\text{S}_8)_3]$ . [278] Formation constants have been determined [279] for  $\text{Fe}(\text{III})$  complexes of various  $(\text{RO})_2\text{PS}_2^-$  or  $\text{Me}(\text{RO})\text{PS}_2^-$ .

#### 4.19. Fluorides and Fluoro-Complexes

$\text{FeF}_3$  and  $\text{FeF}_3 \cdot \frac{1}{3} \text{H}_2\text{O}$ , described as hexagonal tungsten bronzes, have been shown [279] by Moessbauer spectroscopy to show magnetic ordering temperatures of  $97 \pm 2$  and  $128.7 \pm 0.5\text{K}$  respectively; the spectrum of amorphous  $\text{FeF}_3$  has also been described. [280] Thin films of  $\text{FeF}_3$  have been obtained by vacuum sublimation and characterised spectroscopically. [281]

A Moessbauer study [282] of  $\text{KFeF}_4$  above  $T_N$  showed spin fluctuations in this two-dimensional antiferromagnet. The same technique showed phase transitions in  $\text{RbFeF}_4$  at 381 and 417K, the first abrupt and the second gradual, as the  $\text{FeF}_6$  octahedra tilt with respect to the  $c$ -axis. [283] Angle-resolved photoemission from  $\text{K}_2\text{FeF}_5$  and SCF  $X\alpha$  calculations for  $[\text{FeF}_6]^{3-}$  are reported. [284] Neutron diffraction, magnetization and Moessbauer studies for  $\text{LiMnFeF}_6$  show [285] that the  $\beta$ -phase is ferrimagnetic below 113K with some  $\text{Fe}^{3+}$  not coupled to others. Revision of the space group for  $\text{NH}_4\text{MnFeF}_6$  gave better R values and a more regular structure. [286] Moessbauer spectra are also reported [287] for  $\text{Ba}_2\text{NiFeF}_9$ ,  $\text{Ba}_2\text{CrFeF}_9$  and  $\text{NaBaFe}_2\text{F}_9$ .

The X-ray crystal structure of  $\text{Fe}_3\text{F}_8 \cdot 2\text{H}_2\text{O}$  shows [288] two  $\text{Fe}^{3+}$ , coordinated by six  $\text{F}^-$ , for each  $\text{Fe}^{2+}$ , coordinated by four  $\text{F}^-$  and two

H<sub>2</sub>O. For this compound and its Mn<sup>2+</sup> analogue there is a Néel temperature near 157K when only the Fe<sup>3+</sup> spins order; this is anthropomorphically described [289] as "idle spin" due to "magnetic frustration".

#### 4.20. Chlorides, Bromides, Iodides and Halide Complexes

The enthalpy of formation of FeCl<sub>3</sub> is reported [290] as - 396.02kJmol<sup>-1</sup>.

Similar data for four hydrates have also been determined. [291]

The biggest topic in this section is the intercalation of FeCl<sub>3</sub> in graphite.

There is a new review on this. [292] Moessbauer and powder X-ray methods indicate [293] that intercalation involves the formation of [FeCl<sub>4</sub>]<sup>-</sup>, but other workers found [294, 295] Fe<sup>2+</sup> sites, with Cl atoms surrounding a vacant site. Neutron diffraction studies [296, 297] showed two-dimensional ordering below 30K for the first and second stages, with the former developing three-dimensional ordering below 3.8K. FeCl<sub>3</sub> was one of the halides used [298] for graphite intercalation for Li batteries. An ammonia complex was also investigated [299] for energy storage. The FeCl<sub>3</sub><sup>-</sup> graphite intercalate when reduced catalyses the oxidation of SO<sub>2</sub> in H<sub>2</sub>O, with probably both FeCl<sub>2</sub> and FeO(OH) present. [300] Thermal analysis studies [301] gave ΔH and ΔS for its decomposition to C<sub>12</sub>FeCl<sub>3</sub> or C + FeCl<sub>3</sub>. Simultaneous intercalation of FeCl<sub>3</sub> and CoCl<sub>2</sub> gives alternate layers of each; the material has magnetic phase transitions at 8.8 and 8.0K. [302]

FeOCl is an effective host for intercalation compounds. With RNH<sub>2</sub>, (R up to C<sub>18</sub>) the basal separations increase from 7.92 Å to 57.74 Å as about 0.55 mol of guest enter. [303] Moessbauer spectra of the 2-Mepy intercalate show [304] some Fe<sup>2+</sup> sites form as well as an "intermediate valence" site. A phase study of NH<sub>3</sub> and RNH<sub>2</sub> intercalates found [305] that HCl was lost above 120°C. Another X-ray and Moessbauer investigation supported [306] a charge transfer model for bonding. Reactions of the aminopyridine (L) intercalate FeOClL<sub>0.25</sub> with methanol and ethylene glycol gave [307] FeO(OMe) and Fe<sub>2</sub>O<sub>2</sub>(C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).

New work on  $[\text{FeCl}_4]^-$  and other halo complexes includes the synthesis and UV and EPR spectra [308] of the  $[\text{Mo}(\text{Et}_2\text{dtc})_4]^+$  salt of  $[\text{FeCl}_4]^-$ , and the synthesis [309] of the  $[\text{Ti}(\text{cp})_2\text{Cl}(\text{MeCN})]^+$  salt. The crystal structure of the bis - 8-hydroxyquinolinium chloride salt is reported. [310]  $\text{FeCl}_3$  reacts with  $\text{MgCl}_2$  in THF to give  $[\text{MgCl}(\text{THF})_5]^+[\text{FeCl}_4]^-$  which converts to  $[\text{Mg}(\text{THF})_4(\mu\text{-Cl})_2\text{FeCl}_2]$ , whose structure is also reported. [311] The range of compounds  $\text{Li}(\text{Fe,Al})\text{Cl}_4$  have been prepared and their conductance and magnetism studied. [312] In the near IR polarized single crystal spectrum of  $\text{Cs}_2\text{FeCl}_5 \cdot \text{H}_2\text{O}$  at 10K overtone and combination water bands were seen, [313] but the K analogue has a more complex spectrum. The self-ionisation of  $\text{FeCl}_3$  in organic solvents has been studied by EPR; [314] in acetic acid dimerisation and trimerisation were suspected from conductance and molecular weight data. [315]  $[\text{FeCl}_4]^-$  is also formed by the reaction of  $\text{Fe}_2\text{O}_3$  with  $\text{PCl}_5$  in  $\text{POCl}_3$  or with  $\text{S}_2\text{Cl}_2$  and chlorine; the former reaction also produces  $2\text{FeCl}_3 \cdot 3\text{POCl}_3$ , but reaction with  $\text{POCl}_3$  alone gives  $\text{FeCl}_3$  and  $\text{Fe}(\text{PO}_2\text{Cl}_2)_3$ . [316] The resonance Raman spectrum of  $[\text{Et}_4\text{N}][\text{FeI}_4]$  is reported. [317] Kinetic studies [318,319] of the reaction of  $[\text{FeX}_4]^-$  with  $\text{Y}^-$ , ( $\text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{NCS}$ ) show ion pairs may be formed in an associative mechanism but the dielectric constant of the solvent is important. Solvent extraction of  $[\text{FeCl}_4]^-$  can occur by the quaternary ammonium salt "Hyamine 1622", [320] or  $\text{Me}(\text{octyl})_3\text{NH}^+$ , which selectively transports  $\text{Fe}^{3+}$  through a liquid membrane. [321]

New reactions of  $\text{FeCl}_3$  include the formation of a complex with the cyanamide derivative  ${}^i\text{Pr}_2\text{NC}\equiv\text{N}$  and the complex's reaction with chloroalkanes. [322]  $\text{FeCl}_3$  catalyses the bromination of anisole, [323] the copolymerization of styrene and methylmethacrylate, [324] initiates the polymerization of S-carboxyalkylcysteine (though other chlorides are better), [325] is used with  $\text{PdCl}_2$  in the hydrochlorination of ethene, [326] in the benzylation of alkyltoluenes, [327] the conversion of azoxybenzenes into 4-chloroazobenzenes,

[328] and the oxidative coupling of isoeugenol, [329] and it reacts with derivatives of benzoic acid. [330]

#### 4.21. Other Iron (III) Compounds

The iron (III) complex of bleomycin  $A_2$  is highspin at pH 4, but when the solution pH is raised to 7 a lowspin complex forms as the  $\alpha$ -amino nitrogen of  $\beta$ -aminoaniline coordinates; lowspin complexes can be formed at pH 4 by adding  $NCS^-$ ,  $N_3^-$  or  $S_2O_3^{2-}$  or DNA, which can displace these ligands, possibly with interaction between DNA and bleomycin. [331]

### 5. IRON (II)

#### 5.1. Cyanide Complexes

A series of papers [332-4] investigates the overtone and combination water IR spectrum in  $K_4[Fe(CN)_6] \cdot 3H_2O$ . Electronic spectra show [335] charge transfer bands between  $[Fe(CN)_6]^{4-}$  and pyridinium type cations.

A number of mixed metal complexes are described. The photodecomposition of  $n$ -CdS in  $[Fe(CN)_6]^{4-/3-}$  cells containing  $K^+$  and  $Cs^+$  leads to the formation [336] of  $(K,Cs)_2 [CdFe(CN)_6]$ , while a thermodynamic study of the interaction of  $[Fe(CN)_6]^{4-}$  with a Tl amalgam electrode showed [337] the formation of  $[TlFe(CN)_6]^{3-}$ . Trivalent lanthanide ions appear from IR spectra to bond to N atoms in forming  $K[MFe(CN)_6] \cdot 4H_2O$ . [338,339] The thermal decomposition of  $UFe(CN)_6$  has been studied. [340]

Reactivity studies of  $[Fe(CN)_6]^{4-}$  include the pyrolysis of  $R_3S^+$  and  $Ph_2I^+$  salts, [341] its use as a commercial  $NH_3$  synthesis catalyst, [342] its reaction with HCHO under light to give aminoacids, [343] its use with  $[Cr(C_2O_4)_3]^{3-}$  to generate  $H_2$  from irradiated  $10^{-2}M$  aqueous solutions ( $FeSO_4$  can also be used), [344] and unspecified products from a reaction with styrene oxide. [345]

In the area of mixed ligand complexes, electronic spectra were used [346] to study solvent effects in  $[Fe(CN)_5py]^{3-}$ ,  $[Fe(CN)_4 bipy]^{2-}$  and  $[Fe(CN)_2(bipy)_2]$ . There has been a careful discussion [347] of solvent and pressure effects on the substitution kinetics of  $[Fe(CN)_5L]^{3-}$ ,

(L = 4-CNpy, 2-Mepyzazine). A bright red material is formed when  $\text{Ni}^{2+}$  or a nickel electrode reacts with  $[\text{Fe}(\text{CN})_4 \text{ bipy}]^{2-}$  and this is believed to have a nitrogen from one  $\text{CN}^-$  bound to Fe [348] The reaction of pyridine with  $[\text{Fe}(\text{CN})_5(\text{Hen})]^{2-}$  is unusually fast in MeCN. [349]  $[\text{Fe}(\text{CN})_6]^{4-}$  reacts with  $[\text{Co}(\text{en})_2(\text{NH}_3)(\text{H}_2\text{O})]^{3+}$  to give  $[(\text{NC})_5\text{Fe}(\mu\text{-CN})\text{Co}(\text{NH}_3)(\text{en})_2]$  which shows an intervalence transition. [349] An ingenious kinetic study of the intramolecular electron transfer in  $[(\text{NC})_5\text{FeLCo}(\text{dien})]^{2-}$ , (L = pyrazine-2,6-dicarboxylate) shows that the outer sphere reaction between  $[\text{Fe}(\text{CN})_5(\text{LH})]^{4-}$  and  $[\text{Co}(\text{dien})\text{L}]^+$  is 5.7 times faster at 25 °C, suggesting that a stacked ion-pair may give a suitable structure. [350]

### 5.2. Phthalocyanines and other Complexes of Anionic Nitrogen Ligands

New preparations include phthalocyanines with 4  $t\text{Bu}$ , 8 Me or MeO or 16 Cl substituents and their bispyridine derivatives. [351] Raman spectra are recorded [352] for Fe(II) tetrasulphonated phthalocyanine, as has the EPR spectrum of [FePc] in a polymer matrix. [353] The question of the electronic groundstate of [FePc] has been re-examined; electron density maps and Moessbauer spectra indicated [354] a  $^3\text{E}_g$  ground state, while magnetochemistry and Moessbauer studies led to a description [355] as mostly  $^3\text{A}_1$ .

[FePc] has been prepared as a stable electrode film on Au, [356] and C electrodes in fuel cells are improved by incorporating Fe tetrachloro-phthalocyanine. [357] Magnetochemistry and dielectric measurements were included in a study of biphthalocyanines, which appear to undergo a structural change near 340K. [358]

Fe(II) polyphthalocyanine catalyses the decomposition of  $\text{H}_2\text{O}_2$  by two mechanisms. [359] Partial oxidation is located on Fe in  $[\text{Fe}(\text{Pc})(\text{pz})\text{I}_{0.26}]$ , which contains  $\text{I}^-$ , and  $[\text{Fe}(\text{Pc})(\text{bipy})\text{I}_{1.06}]$ , which contains both  $\text{I}^-$  and  $\text{I}_3^-$ . [360] Reaction of [FePc] with KCN gives [361]  $[\text{Fe}(\text{PcH})\text{CN}]$  and  $[\text{Fe}(\text{Pc})\text{CN}]$ ; the conductance of both has been measured. [362]

The preparation of  $\text{FeX}_2 \cdot 2\text{H}_2\text{O}$ , ( $\text{HX} = \text{H}_2\text{N.NH.C(S)OEt}$ ) is reported. [363]

### 5.3. Ammonia Complexes

Molecular orbital calculations, [364] of which few particulars are given, purport to show that  $\text{Fe}^{2+}$  (and other metal ions) do not use d orbitals in forming complexes with  $\text{F}^-$ ,  $\text{H}_2\text{CO}$  and  $\text{H}_2\text{O}$ , but they do with  $\text{NH}_3$ !

### 5.4. Amine Complexes

The trinuclear complex of 4-ethyltriazole (L),  $[\text{Fe}_3\text{L}_6(\text{H}_2\text{O})_6]^{6+}$  exhibits a gradual highspin - lowspin transition for the central  $\text{Fe}^{2+}$  ion; its Fe-N bonds contract  $0.14 \text{ \AA}$  on cooling, but no such effect is seen for the outer  $\text{Fe}^{2+}$  and there is no magnetic exchange. [365] In related work  $[\text{Fe}(\text{1-propyltetrazole})_6]^{2+}$  also shows a spin equilibrium, [366] the first such to be convertible by light, with the  $^5\text{T}_{2g}$  state stable below 50K. New preparations include other azole complexes, [367] and  $[\text{FeCl}_2(\text{PPh}_3)(\text{N-sulphinylaniline})]$ . [368] A kinetic study of the reaction of histamine with  $[\text{Fe}(\text{CN})_5(\text{H}_2\text{O})]^{3-}$  and the spectroelectrochemistry of the product indicated [369] that histamine coordinates through the azole nitrogen.

### 5.5. Pyridine Complexes

Two studies involve highspin-lowspin transitions. For the tris complex of 2-(pyridin-2-yl) benzothiazole this transition shows thermochromism. [370] The effect of particle size on the equilibrium is studied when  $[\text{Fe}(\text{NCS})_2(2\text{-pyCH:NHPh})]$  is supported on to C or  $\text{MgO}$ , with a greater fraction of lowspin as loading increases. [371]

In new preparative work, complexes have been prepared with 2-picoly-1-p-nitrophenylketone, [372] bromazepam, [373] 8-hydroxyquinolate-5-(p-tolyl)sulphonamide, [374] bis(antipyril)piperazine, [375] and pyridoxine. [204]

### 5.6. Diamine Complexes

Resonance Raman spectra of fully deuterated  $[\text{Fe}(\text{bipy})_3]^{2+}$  allowed the C-H modes to be identified. [376] The same technique, together with surface-

enhanced spectra, studied the absorption of  $[\text{Fe}(\text{phen})_3]^{2+}$  on Ag and Fe.

[377]  $[\text{Fe}(\text{bipy})_3]^{2+}$  on graphite was used [378] to release  $\text{H}_2$  and  $\text{O}_2$  from  $\text{H}_2\text{O}$ .

The kinetics of racemisation of  $[\text{Fe}(\text{phen})_3]^{2+}$  continues to be studied. In the presence of  $\text{C}_n\text{H}_{2n+1}\text{SO}_3^-$  stronger hydrophobic interactions coming when  $n$  is increased accelerated this reaction and the dissociation of the complex, but the latter reaction is retarded at low sulphonate concentration.

[379] These reactions have also been studied [380] for

$[\text{Fe}(4\text{-MeC}_6\text{H}_4\text{N};\text{C}(\text{Ph})\text{-2-py})_3]^{2+}$  as well as  $[\text{Fe}(\text{phen})_3]^{2+}$ .

Nucleophilic attack on the tris-complexes is still attracting interest. The kinetics of the reaction of  $\text{OH}^-$  with  $[\text{Fe}(4,7\text{-diphenylphen})_3]^{2+}$  are reported. [381] Another study, using the nucleophiles  $\text{OH}^-$  and  $\text{CN}^-$  reached the unsurprising conclusion that this was a very complex reaction with many possible intermediates. [382] The aquation of  $[\text{Fe}(\text{phen})_3]^{2+}$  and the  $\text{Fe}^{2+}$  complex of the hexadentate ligand  $(2\text{-py-CH:NC}_2\text{H}_4\text{NHCH}_2)_2$  in microemulsions shows that in some cases the reacting complex is at the interface. [383]  $[\text{Fe}(\text{bipy})_3]^{2+}$  is said [384] to exist as a covalent hydrate during its role in the oscillating reaction of  $\text{BrO}_3^-$  with malonic acid. An equilibrium constant for the association of  $[\text{Fe}(\text{phen})_3]^{2+}$  with  $\text{ClO}_4^-$  is reported. [385]

Much work concerns spin transitions in  $[\text{Fe}(\text{diimine})_2(\text{NCS})_2]$  compounds.

DSC experiments [386] showed a first order transition for the bipy complex, but two steps were found, suggesting the spin transition and structure modification do not quite coincide.  $^{13}\text{C}$  spin lattice relaxation time studies [387] indicated that there is no spin diffusion in the solid phen complex as the transition occurs. The 4,7- $\text{Me}_2\text{phen}$  complex exhibits a 45K hysteresis, the exact values depending on the crystallinity of the sample, [388] but there is a much smaller hysteresis for the bis complex of

1,10-phenanthroline-2-carbaldehydephenylhydrazone. [389] For the latter complex there is a first order transition, but not for  $[\text{FeL}_2(\text{NCS})_2]$ , ( $\text{L} =$



2,2'-bi-5-methyl-2-thiazoline according to DSC results. [390]

Resonance Raman spectra show special enhancement of some C-C vibrations in tris complexes of bipy substituted at the 4 and 4' positions by C<sub>12</sub> or C<sub>18</sub> ether chains. [391] New preparations include bis complexes with a new phenyl substituted terpyridyl, [392] a pyridine-phenanthroline tridentate, [393] and a pyridine-thiazole-pyridine tridentate ligand. [394]

Studies in solution concern the formation of ternary complexes with polypyridine ligands, [395] a ferrozine (sulphonated bipy) complex, [396] and a ternary bipy-benzohydroxamate complex. [397]

### 5.7. Schiff Base Complexes

In addition to Schiff base complexes, this section includes semicarbazones, thiosemicarbazones, oximes, hydrazones and other complexes in which a  $\text{C}=\text{N}$  unit is significant.

The trigonal bipyramidal complex of the bis-Schiff base of salicylaldehyde with N'-methyldipropylenetriamine has the two Schiff base nitrogens in axial positions. [398] The Fe(II) complex of the pentadentate ligand from diacetylpyridine and two methyl histidines reacts irreversibly with O<sub>2</sub> in py, probably by ligand oxidation. [399] The tris complex of dimethylpyrimidinetetrone monoxime shows coordination through the oxime N and a neighbouring O. [400] The [Fe(dmgl)<sub>2</sub>]-py(or 4-CN py) system is used [401] to illustrate a spectrophotometric method for studying two successive equilibria. The product of the reaction of FeCl<sub>2</sub>·4H<sub>2</sub>O with cyclohexanedioxime and BF<sub>3</sub>·2HO<sub>2</sub>CMe is an oxime complex in which two ligands are linked by two borates in a manner already known for other metals. [402] The most interesting of many semicarbazone complexes is one with one anionic and one bidentate NO<sub>3</sub><sup>-</sup>. [403] A Schiff base from 1,3-diaminobenzene may form a binuclear Fe(II) complex. [404] Other formation constant [405] and preparative work [90,96,406-416] requires no further comment.

### 5.8. Complexes of Phosphorus Ligands

$[\text{Fe}(\text{acac})_3]$  reacts with dmpe and  $\text{AlMe}_3$  to give  $[\text{Fe}(\text{dmpe})(\text{acac})]^+$   
 $[\text{AlMe}_4]^-$ ; the  $\text{BPh}_4$  salt can be converted to  $[\text{Fe}(\text{dmpe})\text{Me}_2]$  with  $\text{LiMe}$ .

[417]  $[\text{Fe}(\text{PMe})_4\text{MeBr}]$  reacts with  $\text{LiCH}_2\text{PMe}_2$  to give  
 $[\text{Fe}(\text{PMe}_3)_3\text{XPMc}_2\text{CH}_2]$ , ( $\text{X} = \text{H}, \text{Me}$ ), but  $\text{LiCH}(\text{PMe}_2)_2$  gives  
 $[\text{Fe}(\text{PMe}_3)_3\text{H}(\text{Me}_2\text{PCHPMe}_2)]$  which reductively eliminates to  
 $[\text{Fe}(\text{PMe}_3)_3(\text{dmpm})]$  [418].

### 5.9. Oxides

The rotational spectrum of the  $\text{FeO}$  molecule has been observed and assigned.

[419] Also for  $\text{FeO}(\text{g})$ , the laser excitation and rotationally resolved laser  
 induced fluorescence spectra have been measured, apparently with unusual  
 results. [420]

Studies of solid oxides include the magnetism [421] of  $\text{Fe}_x\text{O}$ , ( $x = 0.929 -$   
 $0.950$ ), the determination [422] of  $\Delta G$  of formation of  $\text{Fe}_2\text{SnO}_4$ , studies of  
 the  $\text{FeO-V}_2\text{O}_3$  - other oxide systems, [423,424] and the discovery [425] that  
 the rutile structured semiconductor  $\text{FeVNbO}_6$  becomes a "spin glass" below  
 20K.

### 5.10. Hydroxide and Hydroxo-Complexes

The decomposition of  $\text{Fe}(\text{OH})_2$  to  $\text{Fe}_3\text{O}_4$ ,  $\text{H}_2\text{O}$  and  $\text{H}_2$  has been studied  
 [426] by Moessbauer spectra. The hydrolysis of  $\text{Fe}^{2+}$  in aqueous  $\text{NaCl}$  gave  
 formation constants [427] for  $[\text{Fe}(\text{OH})]^+$  and  $[\text{Fe}_2(\text{OH})_2]^{2+}$ .

### 5.11. Complexes of $\text{OR}^-$ Ligands

Polarography studies on  $[\text{FeX}_3]^-$  or  $[\text{Fe}(\text{phen})\text{X}_2]$ , ( $\text{X} = 2\text{-nitrosophenols}$ )  
 show the ligands are reduced before  $\text{Fe}$ . [428] A complex of lapachol ( $\text{HX}$ ), a  
 quinone,  $[\text{Fe}(\text{H}_2\text{O})_2\text{X}_2]$ , has been prepared. [429]

### 5.12. Carboxylates, Aminoacids and Related Complexes

Crystals of  $[\text{FeX}(\text{O}_2\text{CR})]$ , ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ;  $\text{R} = \text{Ph}, \text{PhCH}_2, \text{PhCH}:\text{CH}$ ) have been  
 prepared [430] by the new but simple reaction of  $\text{Fe}$  with  $\text{RCO}_2\text{H}$  and  
 organohalides. Formation constants have been determined [431] for complexes  
 of  $\text{Fe}(\text{II})$  with 5-hydroxy-8-quinolinecarboxylic acid.

Two groups have studied the pyrolysis of  $\text{FeC}_2\text{O}_4$  and its dihydrate in differing conditions and have identified the various products. [432, 433] Equilibrium constants for the system  $\text{FeC}_2\text{O}_4 - \text{N}_2\text{H}_4 - \text{H}_2\text{O}/\text{EtOH}$  have been determined. [434] New aminoacid compounds prepared are  $[\text{Fe}(\text{L} - \beta - \text{phenylalaninate})_2]$  [435] and  $[\text{Fe}(\text{glutamate})(\text{lysinate})]$ . [436] An  $\text{Fe}(\text{II})$  complex with benzylpenicillin is believed to show coordination through a carboxylate O and a thiazolidine N. [437] The oxidation [438] of aminoacids by Fenton's reagent is first order in  $\text{Fe}^{2+}$ .

An interesting dinitrogen complex  $[\text{Fe}(\text{edta})\text{N}_2]^{2-}$  has been prepared, together with its edta analogue; [439] pyrolysis studies are also reported. [440] The reactions of  $[\text{Fe}(\text{edta})]^{2-}$  with  $\text{SO}_3^{2-}$  and NO have been studied with special reference to the N and S compounds which are formed. [441, 442] The  $\text{Fe}(\text{II})$  complex of  $\text{C}_3\text{H}_7\text{NHCOCH}_2\text{N}(\text{CH}_2\text{CO}_2\text{H})\text{C}_2\text{H}_4\text{N}(\text{CH}_2\text{CO}_2\text{H})_2$  has been synthesized; it cleaves DNA if  $\text{O}_2$  is present, especially in the presence of reducing agents like Na ascorbate, possibly by an OH radical attacking the deoxyribose ring. [443]

### 5.13. Diketonates

$[\text{Fe}(\text{N}_2\text{H}_4)_2(\text{tfacac})_2]$  has been prepared. [444]  $[\text{Fe}(\text{hfac})_2]$  forms 1:1 complexes with bidentate or tetradentate N ligands and with  $[\text{M}(\text{acen})]$  and  $[\text{M}(\text{salen})]$ , ( $\text{M} = \text{Co}, \text{Ni}, \text{Cu}$ ). [445]

Formation constants have been determined [224] for the isonitrosoacetylacetone (HX) complexes  $[\text{FeX}_2]_2$  and  $[\text{FeX}_3]^-$ .  $\text{Fe}(\text{II})$  complexes are included among many prepared by arnebin-I [231] and 1-hydroxy-9-fluorenone. [446]

### 5.14. Compounds of Oxyanions

Studies on  $\text{FeSO}_4$  predominate. When the heptahydrate is heated with  $\text{M}_2\text{CO}_3$  the Li system is unique in not reacting until all the water is lost, [447] but for  $\text{Be}_2(\text{OH})_2\text{CO}_3$  there is no mutual reaction. [448]  $\text{FeSO}_4$  is implicated in many organic reactions: the dimerisation of amines by  $\text{H}_2\text{O}_2$ , [449] the ring cleavage of oxaziridines, [450] the oxidation of alkenes to vicinal diacetates by  $\text{S}_2\text{O}_8^{2-}/\text{MeCO}_2\text{H}$ , [451] the dealkylation at the

6-position of 2,6-di-t-butyl-4-methylphenol, [452] and the promotion of graphite oxidation. [453]

The chlorosulphates  $[\text{Fe}(\text{SO}_3\text{Cl})_2]$ ,  $[\text{FeL}_4(\text{SO}_3\text{Cl})_2]$  and

$[\text{FeL}_2'(\text{SO}_3\text{Cl})_2]$ , (L = monodentate ligand, L' = bidentate ligand),

have been prepared, [454] as have  $[\text{FeL}_4(\text{MeSO}_3)_2]$ , (L = py,  $\frac{1}{2}$  bipy),

[455] and  $[\text{Fe}(\text{O}_2\text{SeC}_6\text{H}_4\text{R})_2]$ , (R = H, Me, Cl, Br). [456]

$[\text{Fe}_9(\text{PO}_4)_8\text{O}_8]$  has been prepared and shown to have a structure comprising FeO blocks of NaCl structure and  $\text{FePO}_6$  blocks which are alternating rows of  $\text{FeO}_4$  and  $\text{PO}_4$  tetrahedra; the loss of electron to give the mixed valence formulation occurs in the FeO region. [457] Thermodynamic data were obtained [458] for the formation of this from  $\text{Fe}_3(\text{PO}_4)_3\text{O}_3$  and its reduction to  $\text{Fe}_3(\text{PO}_4)_2$ .

$[\text{Fe}(\text{H}_2\text{O})_2(\text{uracil})_2]$  is believed [459] to show coordination through N(3) and the keto O at position 2. Complexes with 2'-AMP and 3'-AMP are believed to show phosphate coordination, perhaps with adenine coordination also in the 2'-isomer. [250]

Fe(II) complexes with substituted hydroxamic acids have been prepared, [460] and formation kinetics studied for the 1:1 acetohydroxamic acid complex. [461] Complexes of  $[\text{ONC}(\text{CN})\text{C}(\text{O})\text{NH}_2]^-$  have been prepared in which coordination is probably through the amide O and nitroso N. [462]

### 5.15. Water Complexes

Moessbauer spectra of various  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  materials are all, unsurprisingly, explained by the same crystal field parameters; [463] the phase-change in the  $\text{SiF}_6^{2-}$  salt is of the order-disorder type. [464] There is an  $S_N1$  rate-determining step in the solid state reaction of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  with KNCS to give  $[\text{Fe}(\text{H}_2\text{O})_4(\text{NCS})_2]$ . [465] The mechanism of the photooxidation of aqueous solutions of  $\text{Fe}^{2+}$  salts has been studied, [466] as has the air oxidation of these solutions in the presence of  $\text{Cu}^{2+}$  ions. [467]

### 5.16. Complexes of Neutral Oxygen Ligands

A wide range of complexes of dmf and its thio analogue have been prepared.

[468] Heat capacity data on  $[\text{Fe}_x\text{Co}_{(1-x)}(\text{C}_5\text{H}_5\text{NO})_6](\text{ClO}_4)_2$  showed no coupling between Fe and Co; crystal field anisotropies appeared much greater than exchange effects. [469] The 2:1 complex of py-4CHO with  $\text{FeCl}_2$  is thought, implausibly, to show O coordination. [50] Other new complexes are  $[\text{Fe}_2\text{Cl}_4\text{X}]^{4-}$ , (X = tetrasuccinylurea), [470] and  $[\text{Fe}\{\text{PhS}(\text{O})\text{C}_2\text{H}_4\text{S}(\text{O})\text{Ph}\}_3]^{2+}$ , also O bonded. [471]

The use of trioctylphosphine oxide and other organophosphorus compounds in the solvent extraction of  $\text{Fe}^{2+}$  from acetate eutectics into  $\text{C}_{12}\text{H}_{26}$  is described. [472]

### 5.17. Sulphides and Sulphido-Complexes

Studies on the material commonly bearing the formula FeS include its electrochemical reduction in propylenecarbonate, [473] its use as an electrode material in button-sized batteries with Li, [474] and its preparation from  $\text{FeS}_2$  reacting with  $\text{SO}_2$ ,  $\text{N}_2$  and S. [475] For both FeS and  $\text{FeS}_2$  there are reports on the  $\underline{\text{K}}$ -absorption edge structure [476] and solubility in LiCl-KCl eutectics. [477] Various spectra of  $\text{FeS}_2$  have been recorded, including the reflectance spectrum, [478] the PE, electronic and Moessbauer (with MO calculations), [479] and a more complete Raman spectrum, in which mixing of S-Fe-S stretches and lattice bands showed one ought not to think of this as a molecular crystal, which we probably would not have done anyway. [480] A photoelectrochemical study of  $\text{FeS}_2$  showed that  $\text{SO}_4^{2-}$  was formed. [481] Ferrimagnetic  $\text{Fe}_7\text{S}_8$  is formed [482] from hydrated iron (III) sulphate,  $\text{H}_2$  and  $\text{H}_2\text{S}$ .

Among mixed metal sulphides there have been studies [483] on the pressure effect on  $T_c$  for  $\text{FeCr}_2\text{S}_4$ , magnetization studies [484] on  $\text{FeSb}_2\text{S}_4$  and on [485]  $\text{Fe}_x\text{Zn}_{(1-x)}\text{Cr}_2\text{S}_4$ .  $\chi$  calculations on  $[\text{FeS}_4]^{n-}$  ( $n = 4-6$ ) were used to interpret bond lengths and magnetochemical and Moessbauer data.

[486] The new cluster complex  $[\text{Fe}_8\text{S}_6\text{I}_8]^{3-}$  is formed by the reaction

of  $\text{Fe}$ ,  $\text{I}_2$ ,  $\text{Et}_4\text{NI}$  and  $[\text{Fe}_6\text{S}_6\text{I}_6]^{2-}$  in  $\text{CH}_2\text{Cl}_2$ ; its structure comprises an  $\text{Fe}_8$  cube with an  $\text{I}$  on each  $\text{Fe}$  and an  $\text{S}$  atom over each cube face. [487]  $[\text{Fe}(\text{WS}_4)_2]^{3-}$  has been prepared, with the same structure as its  $\text{Mo}$  analogue, two bidentate  $\text{WS}_4$  ligands coordinating tetrahedral  $\text{Fe}$ ; physical data suggest the extra negative charge over the 2-anion is held on a  $\text{S}$  atom, not a metal. [488] An X-ray structure for  $(\text{Ph}_4\text{P})_2 [\text{FeCl}_2(\text{MoOS}_3)]$ . MeCN shows bridging by two  $\text{S}$  atoms. [489] Work on  $\text{Fe-S}$  compounds of avowed biochemical relevance is described in section 6.

#### 5.18. Complexes of Other Sulphur Ligands

The complexes  $[\text{X}_2\text{Fe}(\text{S}^t\text{Bu})_2\text{Mo}(\text{CNR})_4]$ , ( $\text{X} = \text{Cl}, \text{Br}$ ;  $\text{R} = \text{C}_6\text{H}_{11}, ^t\text{Bu}$ ) have been prepared; their electronic spectra vary with temperature according to whether *syn* or *anti* conformers are present. [490] Ethyl *N*-phenyldithiocarbamate coordinates  $\text{Fe}^{2+}$  by its thione  $\text{S}$ . [491] A *cis* configuration is adopted by  $[\text{Fe}(\text{CO})_2(\text{dithioacetylacetonate})_2]$ . [492] All three potential donor atoms coordinate  $\text{Fe}^{2+}$  in its bis complex with 1, 5-dithiacyclooctan-3-ol. [493] Complexes of tetradentate thio-encl-hydrazine ligands have been prepared. [494] The reaction of  $\text{MoCl}_5$ ,  $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ ,  $\text{BzSH}$  and  $\text{Et}_2\text{NCS}_2\text{H}$  gives [495]  $[\text{Mo}(\text{Et}_2\text{dtc})_4]^+ [\text{FeCl}_3(\text{BzSSBz})]^-$ .

#### 5.19. Fluorides and Fluoro-Complexes

The far IR spectrum of  $\text{FeF}_2$  was used to study its antiferromagnetism. [496] Its Raman scattering has also been recorded. [497] Coherent anti-Stokes Raman spectra in  $\text{Fe}_x\text{M}_{(1-x)}\text{F}_2$ , ( $\text{M} = \text{Mn}, \text{Zn}$ ) revealed electronic bands in the  $^5\text{T}_{2g}$  state. [498] Ordering transitions in  $\text{K}_2\text{Fe}_x\text{Co}_{(1-x)}\text{F}_4$  were studied by elastic neutron scattering. [499] Jahn-Teller effects in  $[\text{FeF}_6]^{4-}$  were found [500] in EPR and IR spectra by doping in  $\text{KZnF}_3$ . Powder neutron diffraction in  $\text{FeZrF}_6$  is recorded. [501]

#### 5.20. Other Halides and Halo-Complexes

$\text{FeCl}_2$ , with  $\text{Fe}$  and  $\text{Fe}_3\text{O}_4$ , is found by Moessbauer, EPR and SEM studies of

graphite. [502] Mass spectra gave [503] thermochemical, thermodynamic and volatility data for  $\text{FeI}_2$ ,  $\text{Fe}_2\text{I}_4$  and  $\text{Fe}_2\text{I}_2$ . Laser properties in the visible region have been found [504] when  $\text{FeI}_2$  is photodissociated at 193 nm above 720K. PE spectra of  $\text{FeI}_2$  led to calculation of the energetics of the  $\text{Fe(II)} \text{I}^-$  system, including gaseous and ionic species. [505]

Among mixed metal halides there have been neutron scattering [506] and magnetization [507] studies on the 3-D Ising system  $\text{Fe}_x\text{Mg}_{(1-x)}\text{Cl}_2$ . The magnetic phase diagram of  $\text{Fe}_x\text{Mn}_{(1-x)}\text{Cl}_2 \cdot 2\text{H}_2\text{O}$  shows remarkable complications, with two antiferromagnets, ferromagnetism, intra- and interchain interactions and spin-glass regions. [508] Moessbauer spectra are recorded for  $\text{Fe}_x\text{M}_{(1-x)}\text{Cl}_2 \cdot y\text{H}_2\text{O}$ , ( $\text{M} = \text{Mn}, \text{Co}, \text{Ni}$ ;  $y = 4, 6$ ) [509] and for  $\text{CsFe}_x\text{M}_{(1-x)}\text{Cl}_3 \cdot 2\text{H}_2\text{O}$ . [510] Below 0.7 times  $T_N$   $\text{RbFeCl}_3 \cdot 2\text{H}_2\text{O}$  is a pure one-dimensional Ising antiferromagnet; its spin-cluster resonance intensity was studied. [511]

New preparations of halo complexes include a phenothiazine salt [512] of  $[\text{FeCl}_3]^-$ , and the group  $[\text{Fe}(\text{MeCN})_6][\text{FeI}_4]$ ,  $[\text{Fe}(\text{MeCN})_6](\text{I}_3)_2$  and  $[\text{Fe}(\text{thf})_6][\text{Fe}(\text{thf})\text{I}_3]_2$ . [317] Phase changes in  $(\text{MeNH}_3)_2[\text{FeCl}_4]$  were studied by Brillouin scattering. [513] The salts  $(\text{RC}_6\text{H}_4\text{NH}_3)_2[\text{FeCl}_4]$ , ( $\text{R} = \text{H}, \text{Cl}, \text{OMe}$ ) are weakly ferromagnetic below their Neel temperatures of 80-90K as canting of the antiferromagnetic spins occurs. [514] For the salt with  $\text{R} = \text{Br}$  and for  $[3\text{-ClC}_6\text{H}_4\text{NH}_3][\text{FeBr}_2\text{Cl}_2]$  the transitions to ferromagnetism occur at 94 and 98K respectively and short range two-dimensional antiferromagnetism sets in at 140 and 190K. [515] In other work on  $[\text{RNH}_3]_2[\text{FeCl}_4]$  salts heat capacity measurements were used to investigate phase transitions. [516]

The free energy of formation of  $[\text{Fe}(\text{H}_2\text{O})_4\text{Cl}_2]$  has been calculated [517] as  $-1280 \text{ kJ mol}^{-1}$ . The enthalpy of solution in methanol has also been measured. [518] X-ray diffraction on concentrated aqueous solutions of  $\text{FeBr}_2$  revealed the presence of  $[\text{FeBr}(\text{H}_2\text{O})_5]^+$ . Solvates of  $\text{Fe(II)}$  halides and perchlorate with dmf and its thio analogue were assessed for

charge distributions. [520]

The reaction of  $\text{FeCl}_3$  in  $\text{thf}$  with  $[\text{Re}(\text{cp})_2\text{H}]$  gave  $[\text{Fe}_4\text{Cl}_8(\text{thf})_6]$ , with 4 doubly and 2 triply bridging Cl. [521] With  $\text{FeCl}_2$  or  $\text{FeBr}_2$  this reagent gives  $[\text{Fe}(\text{cp})_2\text{H}_2\text{FeCl}_2]$  or  $[\text{Fe}(\text{cp})_2\text{H}_2\text{FeBr}_2]$ , whose structures are not known. [522]

### 5.21. Hydrides and Hydrido Complexes

Moessbauer spectra showed the existence up to 70K of  $\text{FeH}$  in an  $\text{H}_2$  matrix. [523] At 12K in an  $\text{H}_2$  or rare gas matrix  $\text{FeH}_2$  can be formed. [524] An ion beam study of the reaction of  $\text{Fe}^+$  with  $\text{H}_2$  or  $\text{D}_2$  led to a study of  $[\text{FeH}]^+$ , including its dissociation energy and Lewis acid reactions with organooxygen compounds. [525]

Mg and Fe react together with  $\text{H}_2$  to give  $\text{Mg}_2\text{FeH}_6$ , in which D substitution helped physical studies to show the presence of  $[\text{FeH}_6]^{4-}$  in a  $\text{K}_2\text{PtCl}_6$  structure. [526]

### 5.22. Other Iron (II) Compound

Doping  $\text{Fe}^{2+}$  into InP allowed the study of photoluminescence in the  $^5\text{T}_{2g} \rightarrow ^5\text{E}$  transition. [527]

### 5.23. Mixed Valence Compounds

Pyrazine (pz) has been associated with two reports which describe  $\text{Fe(II)}$  and mixed valence compounds. The compounds  $[\text{Fe}(\text{acac})_2\text{L}]_n$ , ( $\text{L} = \text{pz}$ , 4,4'-bipy or derivatives) react with  $\text{I}_2$  to give partial oxidation; the more conjugated the bridging ligand is the higher the solid state conductance. [528] In  $[(\text{NC})_5\text{FepzFe}(\text{CN})_5]^{5,6-}$  there are localized  $\text{Fe(II)}$  or  $\text{Fe(III)}$  valencies; in  $[(\text{NC})_5\text{FepzRu}(\text{NH}_3)_5]^{0,1-}$  Fe is in oxidation state II in both complexes. [529]

## 6. BIOLOGICAL IRON COMPOUNDS

This section includes iron proteins and other well-defined entities which contain iron and occur in living systems. It also includes surveys of work on model compounds. Work on natural systems which does not bear on the role of iron is excluded.



### 6.1. Haem Proteins

The many spectroscopic studies on deoxy haems will be discussed in order of increasing energy.  $^1\text{H}$  NMR spectra of carp and adult human Hb (and its oxy and carbonyl derivatives) showed differences in the tertiary and quaternary structures. [530] Such spectra have also been used to probe the electronic structure of Fe and haem-protein interactions. [531] A clear and thorough review of EPR work on Hb and Mb includes derivatives in all oxidation states, treatment of spin states and compounds with other metals; there are very clear diagrams and a useful analysis of the bonding question. [532]

The use of isotopic substitution of Fe and N in Mb led to the assignment of the  $243\text{ cm}^{-1}$  band in its resonance Raman (rR) spectrum as a "pyrrole tilting" mode in histidine and the  $220\text{ cm}^{-1}$  band as an Fe-N stretch mixed with histidine and/or porphyrin vibrations; [533] this is supported by another group who emphasise the strong coupling of the latter band to a Soret band, [534] but others find the Fe-N stretch ranges over  $40\text{ cm}^{-1}$  in response to protein effects, though six-coordination renders the structure more rigid. [535] The pH dependence of the rR spectrum of Hb can be accounted for by haem-protein interactions during protonation [536]; a similar study was made for Mb and cyano-Mb. [537] Yet another report showed clear differences between the  $\text{O}_2$  carrying and peroxidase proteins both in protein interactions affecting side chain conformations and in the axial histidines. [538] There are similarities in the rR spectra of a highspin haem protein first reported in 1983 when isolated from *Chromatium vinosum* with that of Hb, but it is thought that the new material may use vinyl groups for the haem-protein link. [539]

The Moessbauer spectrum of Mb shows less conformational variety than Hb, which shows less still than carbonyl Hb; this is related to cooperativity effects. [540]

In other physical studies, magnetic susceptibility studies down to 1.7K on Mb and photodissociated carbonyl Mb used carbonyl Mb for diamagnetic correction

and found both compounds under study had  $S = 2$  and  $g = 2$ , with the only difference being in the zero field splitting. [541]

Many physical studies have investigated the R and T states of haem proteins. Kinetic and thermodynamic data for the noncooperative reactions of dimeric Mb from Busycon canaliculatum suggest that a protein may be in an R state as regards one ligand, but this might be a T state for another. [542] A mathematical treatment of the T - R transition claims to produce results in agreement with earlier experiments. [543] The effect of sucrose, glycerol or ethylene glycol on the T - R transition correlates with slower rates in more viscous media, using CO rebinding experiments from laser photolysed carbonyl-Hb. [544] Studying the oxygenation in aqueous alcohols with changes in pH and KCl concentration showed that the energy difference between the T and R states depended on the polarity of the solvent as well as protein-solvent interactions. [545]

A most interesting crystallography result showed an Hb tetramer with the  $\alpha$  units oxygenated but not the  $\beta$ ones; the authors claimed that in living systems Hb might gain or lose  $O_2$  from only the  $\beta$  haems. [546] However the work was criticised on the grounds that the material studied may have been an equal mixture of oxygenated and deoxygenated Hb. [547] In another lengthy theoretical paper minimal energy structures for R and T states were calculated. [548a] Other workers emphasise the effect of quaternary structural changes. [548b] In sea cucumbers deoxy-Hb is tetrameric but it becomes dimeric on oxygenation. [549]

The  $\nu R$  spectra of some oxy Hb vibrations shows complex pH dependant excitation profiles which show the haem distorting from  $D_{4h}$  symmetry. [550] Redetermination of the magnetic susceptibilities of aqueous solutions of oxy-Hb and carbonyl-Hb found that both were diamagnetic; values greater than zero which are sometimes reported might arise from the presence of some deoxy-Hb. [551]

A kinetic study of  $O_2$  and CO binding by the dimeric Hb from the mollusc

Scapharca inaequalis shows that this has cooperativity and leads to the assertion that one does not need to have different types of chain to induce cooperativity, as had previously been supposed. [552] Quantum yields for the photodissociation of oxy-Hb suggest there are two distinct phases in the  $O_2$  binding process, [553] while another photochemical study of this reaction using either  $\alpha$  or  $\beta$  haems in which Fe was replaced by Co showed that  $O_2$  was more easily lost from  $\beta$  haems. [554] Aniline reacts with haems during oxygenation by  $\pi$ - $\pi$  interaction with a pyrrole ring. [555]

The  $\Delta E_q$  values obtained from Moessbauer spectra of haem complexes of polyvinylimidazoles and their CO and  $O_2$  adducts were used to assess steric effects. [556] Removal of methyl or vinyl groups from haems unsurprisingly affects CO and  $O_2$  binding through size and conformation changes. [557]

The most striking result from much work on carbonyl, alias carboxy, haems comes from rR spectra of carbonyl-Hb and cyanomet-Hb from Chironomus thummi thummi, which indicate that earlier crystallographic results on these must be mistaken and the Fe-C bonds are less than  $2\text{\AA}$  long and the FeCO angle is  $8^\circ$  greater than supposed; the new values are more in line with those from human Hb. [558] There has been a flash photolysis study [559] of CO binding to chains of Hb - Zurich at 20-300K and a theoretical kinetic study of the rebinding of small ligands like CO to haem proteins after flash photolysis. [560] Variations in the reactivity of various myoglobins to CO is attributed to differences in the Fe- $N_h$  plane distance as well as in base dissociation. [561] Time resolved X-ray absorption spectra were used to follow the recombination of carbonyl-Mb after laser photolysis to test the technique of using synchrotron sources for such experiments. [562] A kinetic study of the oxidation of carbonyl-Hb by  $[\text{Fe}(\text{CN})_6]^{3-}$  including the effects of  $D_2O$  and inositol hexaphosphate showed the reaction was a 3 stage process but that deoxy-Hb was not an intermediate. [563]  $[\text{Mn}_2\text{Fe}_2]$  hybrid haemoglobins react with CO only at Fe; placing Mn in either the  $\alpha$  or the  $\beta$  haems and pH variation allowed development of a two state model for cooperativity. [564]

Other small molecules interact with haems. EPR was used to follow the reaction of NO with oxy-Hb and establish that 3 NO molecules bind by 200 min but the remaining 8 haem is not bound until much later. [565] Exciting the Soret bands of nitrosyl haems (and horseradish peroxidase) using  $^{15}\text{N}$  and  $^{18}\text{O}$  allowed location of the Fe-NO stretching ( $600\text{ cm}^{-1}$ ), and Fe-NO bending ( $574\text{ cm}^{-1}$ ) vibrations in the rR spectra, but a band at  $554\text{ cm}^{-1}$  previously thought to be a stretch is now thought to be a bend, which casts doubt on the assignment of a  $567\text{ cm}^{-1}$  band in oxyhaems as an Fe- $\text{O}_2$  stretch; the clear differences between nitrosyl haem and carbonyl haem spectra are attributed to stronger binding by NO, probably using the  $\pi^*$  orbital. [566] Applying pressure of up to 1400 atm on aqueous solutions of nitrosyl -Hb and other derivatives produced changes in the  $^1\text{H}$  NMR spectra which were attributed to quaternary effects rather than R - T state transitions. [567] Applying  $\gamma$ -radiation to solutions of oxy-Mb saturated with  $\text{N}_2\text{O}$  yields deoxy, met and ferryl proteins. [568] The equilibrium constants for the binding of  $\text{MeCO}_2\text{PbEt}_3$  to Hb were redetermined and show that 2 molecules coordinate, not 3 as previously reported. [569] The oxidising agent  $[\text{FeO}_4]^{2-}$  binds Hb and other enzymes at phosphate sites, but then oxidises them powerfully; this reaction is prevented by adding 2,3-diphosphoglycerate, in line with this thinking. [570] There has also been a kinetic study of the oxidation of Hb by diazonium salts. [571] EPR and  $^{13}\text{C}$  NMR spectra were used to follow the oxidation of nitroxide by Hb with  $\text{H}_2\text{O}_2$ . [572]  $\text{O}_2$  bound to Hb, rather than free Hb, promotes the autocatalysis of the  $\text{NO}_2^-$  - Hb reaction. [573] Some miscellaneous results. Reconstitution of Mb using Fe(II) chlorins allowed the formation of  $\text{O}_2$  complexes, but not with good reversibility. [574] A chlorin ring system was found as a tetracarboxylate as the haem d, residue in nitrate reductase. [575]

Turning to met-Hb and related species, the photochemical formation from oxy-Hb has been further studied. [576] Reagents for effecting this oxidation include  $[\text{Fe}(\text{bipy})_3]^{3+}$ , for which the effect of such added ligands as citrate was

studied; [577] ferricytochrome  $b_5$  also performs this oxidation, which in this case is inhibited by CO, which leads the authors to believe the oxidation occurs after loss of  $O_2$ . [578] An X-ray study to  $1.9\text{\AA}$  resolution of met-Mb after saturation with 7 atm of Xe shows 4 sites for these in cavities and suggests molecules entering these cavities may induce conformational change. [579] The heat capacity and dielectric constant of met-Mb crystals below 30K have been determined. [580] There has been a magnetization study [581] of met-Hb and met-Mb from 2 to 0.1K.

The greatest weight of papers on met-haems concerns intricacies in NMR spectra ( $^1\text{H}$  in all cases). Changes in pH for solutions of met-Hb, cyano-Hb and deoxy-Hb change the hydrogen bonding of the histidines and induce conformational changes; Hb seems to have a more flexible pocket than Mb. [582] Similar work involved tryptophan residues also and included carbonyl-Mb. [583] A further assignment of histidine protons in cyano-Mb and some derivatives with Et, Br or  $O_2\text{CMe}$  ring substituents is presented. [584] The met-Mb of Indian elephants has distal glutamine, not histidine, and for this it is suggested from NMR that either amine N or carbonyl O can be used for hydrogen bonding to water; [585] a later paper from this group gave more assignments. [586] Other pH variation work on cyano-Mb and deoxy-Mb suggests that at pH 5-6 there is a structural modification of the haem pocket related to  $\pi$ -interactions between a histidine and the haem ring. [587] Also for cyano-Mb, NOE and relaxation rate experiments centred on the haem pocket isoleucine indicate that the magnetic anisotropy of the Fe is rhombic. [588] For azido-Hb the NMR spectra showed a spin equilibrium between  $S = 1/2$  and  $S = 5/2$  states. [589]

EPR is also applied to haem derivatives, as in the use of spin traps to follow the photoreduction of chlorohaemin; H and  $C_2H_4OH$  radicals form in EtOH, but no radicals are formed if py is absent, suggesting py stabilises Fe(II) here. [590]

There are also many studies of electronic spectra of met-haems. Temperature

variation of these confirms fluoro-Mb as highspin, cyano-Mb as lowspin, and a mixture of each for the  $N_3^-$ ,  $OH^-$  and imidazole derivatives. [591]

Electronic spin-echo envelope spectroscopy confirms earlier experiments in showing that  $H_2O$  is directly bonded to Fe in met-Mb but not in met-Hb. [592] Comparison of the crystal and solution spectra of oxy-Hb, cyano-Hb or azido-Hb using photoacoustic difference spectra showed no differences, but there were for fluoro-Hb and aquomet-Hb, for which the strong hydrogen bonding proclivities of the axial ligands were thought responsible for minor structural changes on crystallisation. [593] Electronic, CD and MCD spectra of mixed valence haemoglobins with specific association of oxidation states with the  $\alpha$  or  $\beta$  haems showed the Soret MCD was sensitive to the spin state of Fe(III) and the CD were not the sum of the component units and must therefore reflect the interaction of haem units. [594]

Electron transfer to met-Hb by cytochrome  $b_5$  was followed kinetically by MCD spectra, with identification of a hydroxy-Hb complex. [595] Another kinetic study followed the aquomet-Hb reaction with Tb complexes. [596]

The electroreduction of the ferrihaem dimer by cyclic voltammetry involves the formation of Fe(II) monomers. [597] A polymer containing haemin acts as a  $CN^-$  ion exchanger. [598] The  $\alpha$  and  $\beta$  haems of Hb were separated and fitted to Fe(III) sulphopc; the  $\alpha$  globin forms a monomer and the  $\beta$  a tetramer with a new CD spectrum. [599]

## 6.2. Peroxidases

The variation of the  $^1H$  NMR spectrum of horseradish peroxidase (HRP) with pH using selective deuteration to assist assignments showed that at high pH the deprotonated distal histidine coordinates Fe with its plane perpendicular to the proximal histidine. [600] The rR spectrum of HRP supports this, showing highspin 5-coordinate Fe at neutral pH and lowspin 6-coordination at higher pH; in these alkaline forms bands were found which innocents might confuse with the "marker" bands of Fe(IV). [601] Nevertheless a short communication on X-ray absorption edge and EXAFS for HRP and related compounds reports HRP-I

as being an Fe(IV) - porphyrin  $\pi$  complex. [602] CD, MCD and EPR spectra of the haem protein indoleamine 2,3-dioxygenase show similarities to HRP and it is concluded histidine coordinates the fifth coordination site and the sixth position is hindered; when the substrate L-tryptophan binds the EPR and MCD spectra are those for lowspin Fe(III), so that some conformational change has occurred. [603]

Hartree-Fock-Slater molecular orbitals were used to show how the bond strength of CO varies when it is coordinated to haem proteins with explanations of IR data in terms of redox potentials of proteins, with peroxidases being able to form better hydrogen bonds to CO than other proteins. [604] Kinetic data for the rebinding of CO to cytochrome c peroxidase and HRP after flash photolysis show similar rates which are different for that with Mb. [605] Another kinetic study, of the reaction of  $\text{CN}^-$  with Fe(III) chloroperoxidase at various pH and in the presence of  $\text{NO}_3^-$ , claims to show that this anion bonds to the protonated enzyme and  $\text{CN}^-$  to the unprotonated form. [606] There has also been a rate study for the decomposition of  $\text{O}_2^{2-}$  by Fe(II) in L- $\alpha$ -phosphatidylcholine. [607]

Investigation of a model complex led to the conclusion that HRP works by an outer sphere mechanism but cytochrome P450 operates by substrate formation. [608]

### 6.3. Cytochromes

Sorting out this big topic is made difficult by the non-systematic nomenclature. Many papers concerned with toxicology or other aspects of cytochromes (cyt) in which serious Fe chemistry does not appear are not cited. Two general papers which require recording concern the use of  $^1\text{H}$  NMR of haems to reveal the electronic structure of Fe and the haem protein interactions in, inter alia cyt c and cyt c oxidase, [531] and the discovery of a cyt reduction type mechanism in the simultaneous electron transfer to Fe and proton transfer to alkoxide in the reduction of a haemin-bispharmacopate complex. [609] Many papers discuss the properties and reactivity of cyt c in

its many manifestations. There has been a more extensive assignment of the haem and histidine  $^1\text{H}$  resonances for ferricyt c. [610] NOE techniques were used to probe the methionine-haem interactions in ferrocyt c, finding 4 conformers with the aid of computer graphics and concluding the solid and solution structures were similar. [611] The  $^1\text{H}$  spectra of the oxidised and reduced forms of cyt c-550 from Paracoccus denitrificans have also been assigned, with successive changes with increasing pH attributed to loss of coordinated histidine, of propionate, the displacement of methionine by lysine and the loss of lysine. [612] In an NMR study of a dihaem cyt c-552 from Pseudomonas perfectomarinus in the oxidised, reduced and intermediate forms no coordination by methionine was found. [613] The combined use of CD and  $^1\text{H}$  (NOE) NMR spectra for cyt c-553 from Desulfovibrio vulgaris and D. desulfuricans showed that coordinated sulphur showed S chirality in the reduced form and R chirality when oxidised. [614] The observation of non-Curie law behaviour for the  $^1\text{H}$  NMR spectrum of oxidised cyt c-554 from Alcaligenes faecalis could not be explained; this cytochrome has the fastest self-exchange rate yet found. [615] For cyt c-555, which has an unusually low redox potential, there are large ring shifts for the methyls in the oxidised form; many other lines in the  $^1\text{H}$  spectrum are assigned. [616]

The EPR spectrum of a cyt in ubiquinolcytochrome c reductase (complex III) showed three signals, showing a 1:1:1 mixture of cyt b-566, cyt b-562 and cyt  $c_1$ , the two former having two histidine ligands and the latter one histidine and one methionine. [617]

The observation of both Stokes and Anti-Stokes lines in the rR spectrum of cyt c is the first time for a molecular system, so the paper includes an extensive theoretical treatment. [618] The excitation profiles for the Soret b and rR spectrum of reduced cyt c are also discussed. [619] Such a spectrum for cyt c-554 gave the allegedly unusual result that the oxidised form appears 5-coordinate. [620]

It is suggested that the CT spectrum of oxidised cyt c is a good measure of



its ease of reduction, and that the axial ligands are important because each process involves electron transfer to a  $d_{xy}$  orbital. [621] Electronic spectra and differential pulse and cyclic voltammetries for cyt c-551 at various pH showed the loss of methionine. [622]

The reactions of cyt c have attracted many studies, including a short review.

[623] The reaction of  $O_2$  with cyt c and cyt c oxidase is faster for the former in low ionic strength buffers but faster for the latter with high ionic strength buffers. [624] Electron transfer reactions with  $[Fe(CN)_5X]^{n-}$ ,

(X = aminoacid) are second order processes with two binding sites. [625]

Reduction by  $[Fe(CN)_6]^{4-}$ , followed by  $^1H$  NMR, is faster in cyt c from

Candida krusli than that from a horse. [626] The reduced and oxidised forms

give comparably strong complexes with  $[Co(CN)_6]^{3-}$ , according to  $^{59}Co$

NMR. [627] There have also been kinetic studies of reactions with Ru and Os

bipy and phen complexes, [628] and reductions of ferricyt c by

$[Ru(NH_3)_5L]^{2+}$ , (L = N donor). [629]

The use of  $RuO_2$  or  $IrO_2$  or other electrodes in CV studies of cyt c showed

the significance of surface charge on the electrodes. [630,631] Adding Cu or

Fe compounds to cyt c enhances the rate of reduction by thioglycolic acid,

probably through the formation of a negatively charged metal-thioglycolate

complex; adding phen to this brew assists catalysis by Cu, destroys that by Fe

and induces pH dependence. [632] Oxidation of cyt c by oxidised azurin or

plastocyanin is not a simple process. [633] The reduction of cyt c by

ascorbate proceeds by electron tunnelling over  $\sim 16\text{\AA}$  below  $26^\circ C$  but at higher

temperatures there is complex formation before reduction occurs. [634] The

reduction of cyt c in the presence of 6-hydroxydopamine and  $O_2$  is an

oscillating system; this was then used to induce amusing oscillations in

reactions of catalase, superoxide dismutase, mannitol and desferrioxamine.

[635] Cyt c does not form a strong complex with cyt  $c_1$ , so the earlier

suggestion that such a complex was formed in the reaction between them is

discredited. [636] There have also been kinetic studies of the reactions of

cyt c with cyt oxidase [637] and bacterial reaction centres. [638].

Now for cyt  $c_3$ . This has been identified, along with some 4Fe-4S ferredoxins, in electron transport proteins in Desulfovibrio desulfuricans.

[639] A most useful X-ray structural analysis ( $R = 0.176$ ) of cyt  $c_3$  from Desulfovibrio vulgaris found the 4 haems to be closely packed with adjacent haem planes perpendicular; histidine acts as axial ligands and all the Fe-N (hist) distances are slightly different in the range 1.88-2.12 Å. The haems are exposed to solvent, are non-equivalent and up to 0.37 Å distorted from planarity. [640] Fuller assignment of the  $^1\text{H}$  NMR spectrum of cyt  $c_3$  has been obtained by 2D methods; the paper includes an analysis of the method.

[641] Studying the  $^1\text{H}$  spectrum at various pH and redox potential shows electron exchange between the haems is faster than  $10^5 \text{ s}^{-1}$ ; the midpoint redox potentials of some haems are pH dependent and it was concluded that the protein has to operate at different redox potentials. [642] EPR was used to follow the dithionite reduction of orthorhombic lowspin Fe(III) in oxidised cyt  $c_3$  to lowspin Fe(II) and its restoration on oxygenation; it was concluded that the oxidised form took on a tetrahedral distortion after many redox cycles. [643] In another EPR study 2 haems showed high and 2 showed low redox potentials; reaction with colloidal sulphur was believed to involve an exposed low potential haem. [644] Another kinetic study of the cyt  $c_3$  reduction was followed by electronic and CD spectra and stopped flow methods; there was special interest in the fast third step which may involve unusual interactions of some kind. [645] Cyt  $c_3$  is one of many systems whose flavin reduction gave correlation of rate with redox potential and the redox centre's accessibility to solvents. [646] Other workers have also compared the redox properties of various cyt  $c_3$  species with the exposure of the haems. [647] An experiment in which the RR spectra of cyt  $c_4$  single crystals were measured is suggested as a prototype for assessing whether an X-ray crystal structure is being carried out on the reduced or oxidised form of a protein. [648] A cyt  $c_6$  from a red alga is thought to contain a c type haem bound to

two cysteine residues. [649] The magnetic properties of cyt c' show this has a mixed spin state. [650] Cyt c' species react with EtNC with a wide range of equilibrium constants which do not correlate with CO reactivity and are attributed to steric effects, showing the distal site to be more accessible than had hitherto been supposed. [651]

The next cytochrome system to be reviewed is cytochrome oxidase and its variants. A new cyt c peroxidase has been isolated from Pseudomonas stutzeri and has two interacting haems with distinct redox potentials; the fully oxidised form contains two lowspin Fe(III), one coordinated by 2 histidines and one by a histidine and a methionine, both being lowspin at low temperatures but one showing a highspin - lowspin transition. The half-reduced state has highspin Fe(III) and lowspin Fe(II); the fully reduced form has one highspin Fe(II) and one lowspin. [652] An extensive EPR study showed that the signal at  $g = 12$  coming from an alternative resting state corresponded to a spin greater than 1, that the signal at  $g = 5$  from the pulsed state had an excited integer spin  $7\text{cm}^{-1}$  above the EPR - silent ground state and that the  $\text{Cu}_A$  site was the same in the resting, pulsed and anaerobically oxidised states. [653]

A pulsed EPR study revealed similar spin-lattice relaxation times for haem a and  $\text{Cu}_A$ , indicating either they had similar environments or that they were near and relaxed one another; the relaxation of nitrosylferrocyt  $a_3$  in mixed valence oxidase showed enhancement by a nearby paramagnetic centre which is probably haem a. [654]

Raman spectral bands at  $210\text{--}220\text{ cm}^{-1}$  for Fe-N(hist) were found in the highspin  $a_3$  haem of  $aa_3$  cyt oxidases, but the bands are lost when the pH is raised or  $\text{CN}^-$  is complexed. [655] The similarity of the MCD spectrum of cyt a in cyt c oxidase with that of a haem a bisimidazole complex shows this structure probably occurs in the enzyme. [656] The X-ray absorption of the "pulsed" (oxygenated) oxidase shows that the S bridge between Fe and Cu is lost, probably being replaced by  $\text{H}_2\text{O}$  or  $\text{O}_2^{2-}$ ; the haem in this form is

like ferric peroxidases and if  $\text{CN}^-$  is present there is a resemblance to cyanohaems. [657]

There have been many reactivity studies in cyt oxidase systems. Four stages were found in the reaction of  $\text{O}_2$  with the half reduced form, with first  $\text{O}_2$  binding to Fe; this then gains an electron in a distinct process. [658] Another kinetic study of this reaction showed that O-O bond breaking is associated with change in the cyt  $a_3$  - Cu cluster. [659] When CO complexes cyta<sub>3</sub> the Fe-C-O unit is linear but not at  $90^\circ$  to the haem and there is strain on the proximal histidine. [660] For the complex with  $\text{N}_3^-$  and NO a  $\Delta m_s = 2$  transition occurs through the interaction of electrons on cyt a(NO) and Cu<sub>B</sub>; at 77K the photolysed complex shows cyt  $a_3$ - $\text{N}_3$ -Cu structure but in the triplet it is thought NO bridges and  $\text{N}_3$  is on Cu. [661] Half-reduced cyt c peroxidase easily forms complexes with  $\text{F}^-$ ,  $\text{CN}^-$  and  $\text{N}_3^-$  at the low-potential haem, but only  $\text{F}^-$  induces highspin; the resting enzyme does not form these complexes. [662] Another report confirms the existence of weak catalase activity. [663] Adding  $\text{H}_2\text{O}_2$  to the pulsed oxidase gave an electronic spectrum which was interpreted as showing a peroxide derivative, [664] but another believes [665] the product is the oxygenated oxidase, which may be a peroxide anyway. Cyt c peroxidase catalyses the oxidation by  $\text{H}_2\text{O}_2$  of  $[\text{Co}(\text{terpy})_2]^{2+}$ ,  $[\text{Fe}(\text{CN})_6]^{4-}$ ,  $[\text{Ru}(\text{RNH}_2)_5\text{py}]^{2+}$ , probably by substrate binding to an  $\text{H}_2\text{O}_2$  - haem complex and with electron transfer within the enzyme as the rate determining step. [666] The use of  $^{51}\text{Cr}^{2+}$  isotope showed subunit II was the electron transfer site for that reagent. [667]

Cytochrome P-450 is another intensely active field. Its activity is well-reviewed, [668-671] including emphasis on camphor-5-exohydroxylase [670] and on the different spin and oxidation states and their effects on the redox processes. [671] The theory of the electronic structure of Fe(III) haem proteins has been reworked to add to the  $^4\text{T}_2$  ground state some  $^6\text{A}$ , or other excited state, with the inevitable better fit to experimental data; cyt

P-450 is said specially to benefit from this treatment. [672] The sensitivity of CD spectra in the Soret region proved useful in replacing the sixth ligand in cyt P-450 to model the interaction of haem proteins with the environment. [673] MCD also provides extra detail and confirms the lowspin character of the oxidised form and the highspin nature of the reduced form, which starts to become lowspin at low temperature. [674] Cyt P-450 complexes with esters, ethers and ketones show spectral effects which have hitherto been attributed to solvent action. [675] The two different processes with different rates which were found for cyt P-450 reduction are attributed to different spin states. [676] Other cyt P-450 reactions studied include the conversion of  $\text{CCl}_4$  to  $\text{COCl}_2$ , [677] of vinylcyclooctane to epoxide, [678] and the demethylation of  $\text{PhNMe}_2$ . [679]

Related reactions to those of cyt P-450 which have been studied include those of a protein containing 7 or 8 haems, one having highspin Fe(II) with a very high  $\Delta E_q$ , which converts  $\text{NH}_2\text{OH}$  to  $\text{NO}_2^-$ ; [680] a rat liver phenylalanine hydroxylase which shows two signals for nonhaem highspin Fe(III); [681] a mechanistic study of the action of protocatechuate 3,4-dioxygenase. [682]

There remain some miscellaneous papers on other cytochromes, and these reports cannot be put in any logical order. Electron and proton transfer through cyt bc complexes is reviewed. [683] rR spectra are used to characterise a, b and c cytochromes with the conclusion that haem-haem interactions may vary within the b class and perhaps also the a class. [684] The nitrate reductase haem has been found to be like cyt  $b_5$ . [685] Six different types of haem occur in hydroxylamine oxidoreductase; 4 haems are  $c_{553}$  type, 2 are  $c_{559}$  and one is a P-460 haem. [686] The CD spectrum of NADH-cyt  $b_5$  reductase is reported. [687] EPR showed the two cyt b-563 in the cyt bf complex from spinach are both lowspin with slightly different g-values. [688] A discussion of the role of  $[\text{Fe}(\text{CN})_6]^{3-}$  in restoring the activity of inhibited cyt bc<sub>1</sub> complex led to the view that this reagent can bind to haem centres. [689] The

polarised single crystal electronic spectrum of cyt c<sub>d</sub>, shows these two haems are perpendicular in reduced and oxidised forms. [690] The formation of a 1:1 complex between cyt f and plastocyanin was established by gel electrophoresis and electronic spectra. [691]

#### 6.4. Porphyrin Complexes

This section comprises the large volume of work on Fe complexes of natural or synthetic porphyrins designed to illuminate biochemical topics involving various haem type proteins.

We begin with Fe(II) complexes of simple porphyrins. X $\alpha$  calculations on these predict a  $^3A_{2g}$  ground state with some  $^3E_g$  excited state mixing if the Fe-N bond is shorter than 2.0Å but a  $^5B_{2g}$  state at longer Fe-N distances. [692] Moessbauer spectra were recorded for various highspin 5-coordinate [Fe PPIX] complexes with axial ligands including improbably phenol and catechol. [693] The role of axial ligands (eg NO, py) was one of the variables assessed in a study of factors affecting the redox properties of Fe porphyrins. [694] A combined electrochemical and spectroscopic study of [Fe(TPP)(CS)] and its adducts with N donors showed it can undergo two electron transfer steps, first to Fe(III) and then to a radical with electron loss on the porphyrin. [695] Fitting TPP with phosphorylcholine groups lets its Fe complexes occupy phospholipid bilayers. [696] [Fe(OEP)(2-Meim)<sub>2</sub>] has been prepared and shows a 1/2 - 5/2 spin equilibrium; it is suggested that the orientation of the imidazoles determines the spin state with steric hindrance in the solid state keeping these highspin but this does not apply in solutions. [697] The rR spectra of other 2-Meim porphyrin complexes have been assigned to show which bands are sensitive to ring substituents. [698] O<sub>2</sub> forms complexes with [Fe(TPP)] and [Fe(OEP)] in Ar matrices at 15K which are isomers with monodentate O<sub>2</sub> ( $\nu$  1190cm<sup>-1</sup>) or bidentate O<sub>2</sub> ( $\nu$ 1105cm<sup>-1</sup>); these are S = 1 or S = 0 spin states respectively. [699] Comparison of CO and O<sub>2</sub> binding to [Fe(TPP)] and its 2,4,6-triphenylphenyl analogue in the presence of imidazoles showed that CO binding was not much affected by steric

effects or by H bonding of phen to imidazole; more polar solvents enhance  $O_2$  binding but hinder CO binding, perhaps because a polar solvent will stabilise the charge separation involved in  $O_2$  complexation. [700]

Other reactions of Fe(II) porphyrins include the synthesis of the  $[Na(crown\ ether)(thf)_2]^+$  salt of  $[Fe(TPP)]^-$  which has a trinegative porphyrin; this group also prepared the diamagnetic  $[Na(thf)_3]^+$  salt of  $[Fe(TPP)]^{2-}$ , in which the porphyrin has 2N atoms out of the ring plane. [701]  $[Fe(TPP)]$  catalyses the reduction of allyl bromide to give a porphyrin complex where the  $\alpha$  and  $\sigma$  positions have  $CH_2$ , C:CH or CH:C:CH<sub>2</sub> groups. [702]

Among Fe(III) porphyrins it has been shown that  $[Fe(TPP)Ph]$  is lowspin with the Fe only 0.17Å out of the ring plane. [703] Other X-ray structures reported are those of  $[Fe(TPP)(SC_6HF_4)]$  [704] and  $[Fe_2(TPP)_2SO_4]$  in which  $SO_4^{2-}$  bridges by one O atom bonding to each Fe. [705] Various porphyrin (por) complexes  $[Fe(por)Cl]$  have been synthesised using haems with selective deuteration of Me. [706] The extensively studied  $[FeL(ClO_4)]$  complex, L = tetrakis(trimethoxyphenyl)porphyrin, has a 3/2 spin ground state but the bis-thf complex has some 5/2 state mixed in. [707] The Fe(III) - PPIX complex gives 1:1 complexes with  $\alpha$  and  $\beta$  cyclodextrins and their methyl and acetyl derivatives. [708]  $[Fe(por)(3-Clpy)_2]^+$  complexes also show interesting magnetochemistry, with basic porphyrins like OEP giving highspin, less basic ones like tetra-3-NO<sub>2</sub>PP giving lowspin and intermediate ligands such as PPIXDME giving intermediate results. [709]  $E_2$  values for reduction of  $[Fe(TPP)(imidH)_2]^+$  complexes depend on the H bonding power of the imidazole; [710] others studied similar systems in an unsatisfactory attempt to correlate EPR data with various cytochrome systems. [711]

We have already met a sulphate-bridged dimer, and many others were studied this year. The <sup>13</sup>C NMR spectra of  $[Fe_2(por)_2O]$ , (por = OEP, TPP) give similar antiferromagnetism to the earlier <sup>1</sup>H results. [712] Using carbethoxytetramethyl-porphyrins allowed <sup>1</sup>H spectra to reveal conformational

isomers as well as antiferromagnetism. [713] Tetrasulphophenylporphine complexes can be monomers or dimers in aqueous solutions depending on the pH.

[714]  $[\text{Fe}_2(\text{TTP})_2\text{O}]$  reacts with  $\text{HN}_3$  to give  $[\text{Fe}(\text{TTP})\text{N}_3]$ ; if this is exposed to UV light  $[\text{Fe}_2(\text{TTP})_2\text{N}]$  is formed, but if py is present this photolysis gives  $[\text{Fe}(\text{TTP})(\text{py})_2]$  and  $[\text{Fe}_2(\text{TTP})_2\text{O}]$ . [715] CV experiments showed 2 reductions and 4 oxidations for  $[\text{Fe}_2(\text{TPP})_2\text{C}]$ , with the formation [716] also of such py complexes as  $[\text{Fe}_2(\text{TPP})_2\text{C}(\text{py})_2]^{2+}$ .

Titration of  $\text{OH}^-$  with  $[\text{Fe}(\text{PPIX})\text{Cl}]$  or  $[\text{Fe}(\text{PPIXDME})\text{Cl}]$  in the presence of imidazoles at 77K gave all sorts of azole derivatives including protonation and deprotonation, but at 298K all these revert to  $[\text{Fe}(\text{por})\text{OH}]$ . [717] The reaction of alkalis with iron (III)  $[\text{Fe}(\text{PPIX})]$  appears to involve the interaction of the alkali metal with carboxylate before giving  $[\text{Fe}_2(\text{por})_2\text{O}]^{4-}$  species. [718]

Iron (III) porphyrin complexes also form an active area. Methoxide reacts with  $[\text{Fe}(\text{TPP})\text{Cl}]$  to give successively  $[\text{Fe}(\text{TPP})(\text{OMe})]$  and  $[\text{Fe}(\text{TPP})(\text{OMe})_2]$ , the latter being the first lowspin Fe(III) porphyrin complex with two O axial ligands. [719] In a spectroelectrochemical and  $^1\text{H}$  NMR study of  $[\text{Fe}(\text{por})\text{Ph}]$ , (por = OEP, TPP) and pyridine complexes thereof it was found that redox changes gave delocalisation of charge between cation and porphyrin; [720] however, when slow scan rates were used, the first oxidation step is followed by an irreversible reaction to an intermediate which can lose one electron to form an Fe(III) -  $\text{NPh}$  complex which in turn can be reduced to an Fe(II) -  $\text{NPh}$  one which can then be irreversibly reduced back to the starting material. [721] Determining the equilibrium constant for forming a complex between  $\text{NCO}^-$  and Fe(III) haem octapeptide in 50% aqueous glycol showed this was a weaker complex than that with met Hb or Met Mb. [722] In the latest of a series of papers MCD spectra were used to conclude that a highspin 5-coordinate Fe(II) complex was an intermediate in the catalysis of the electroreduction of  $\text{O}_2$  by an Fe(III) TCPP complex (TCPP = 4-carboxyphenyl analogue of TPP). [723] An Fe(III) deuteroporphyrin IX complex acts as a



peroxidase model in catalysing the chlorination of monochlorodimedone by  $\text{NaClO}_2$  with the formation of hypochlorite. [724] This complex also oxidises dithionite in a diffusion controlled process which is greatly retarded when human serum albumin is added and which is also affected by different anions and the pH. [725] The solvent dependence of the electronic spectra of  $[\text{Fe}(\text{PPIXDME})\text{NO}]$  and its complexes with N ligands is reported. [726]  $[\text{Fe}(\text{por})\text{NO}]$ , (por = PPIXDME, TPP) reacts with NO to give  $\text{N}_2\text{O}$  and  $[\text{Fe}(\text{por})(\text{NO})(\text{NO}_2)]$ . [727]

An SCF calculation for  $[\text{Fe}(\text{porphine})\text{O}]$  shows the  $\text{Fe} = \text{O}$  configuration is less stable by  $66\text{kJ mol}^{-1}$  than inserting the O between Fe and N and suggests such may be the situation in catalase. [728]

Various studies describe reactions which might model cytochromes. These include a study of redox catalysis at an octane-water interface by a coproporphyrin complex, [729] the demetallation of  $\text{Fe(III) TPP}$  (or its 4-OMe derivative) complexes by reactions with  $\text{B(OH)}_3$  under high pressure and transmetallation reactions with  $\text{Cu}_2\text{O}$ , [730] and the partial dechlorination of  $\text{CCl}_4$ ,  $\text{CHCl}_3$  and  $\text{CH}_3\text{CCl}_3$  but not some other chlorohydrocarbons by  $\text{Fe(II)}$  porphyrins. [731] The self-exchange electron transfer rate for  $[\text{Fe}(\text{TPP})(\text{imid})_2]^{2+/3+}$  is slightly higher than for short cytochromes. [732] The reduction of  $\text{Fe(II)}$  deuterioporphyrin by  $\text{Me}_2\text{CO}$  radicals produced by the irradiation of alkaline aqueous isopropanol is followed by slow reoxidation, probably by water, but this process is much faster if  $\text{S}_2\text{O}_4^{2-}$  is the reductant. [733]

The tetra(4-methoxyphenyl)porphyrin complex  $[\text{Fe}(\text{por})\text{SH}]$  is prepared from the chloride analogue by reaction with S and  $\text{LiBHET}_3$ ; its spectral and magnetic data are similar to cyt P-450 and chloroperoxidase. [734] The X-ray structure gives the expected result for  $[\text{Fe}(\text{OEP})\text{SPh}]$ . [735] The carbanion porphyrin complexes  $[\text{Fe}(\text{TPP})(\text{CF}_3\text{CHCl})\text{SR}]$  have uv-visible and EPR spectra like halothane - cyt P-450 products. [736] Moessbauer spectra of  $[\text{Fe}(\text{PPIX})]$  complexes with S ligands show unusually strong Fe-S bonds. [737]  $[\text{Fe}(\text{TPP})]$

reduces many cyt P-450 substrates, including  $R_3N \rightarrow O$  and arene oxides. [738] Of Fe porphyrins only  $[Fe(TpivPP)(O_2CMe)]$  catalyses alkene oxidation by hypochlorite. [739] The  $[Fe(TPP)]$  complex with thioglycolate ethyl ester forms a 1:1 adduct with  $O_2$  which is a superoxide complex and which has spectral similarities with cyt P-450 intermediates. [740]  $[Fe(TPP)Cl]$  cleaves selectively some C-C bonds in 1,3-diols as a model of the enzymic degradation of lignins. [741]

Cyt-c models include compounds of haemin with haemoctapeptide, wherein a histidine from the peptide takes the 5th position and modifies the effects of added 6th ligands such as picoline or azide. [742] Crystal field calculations on histidine peptide Fe(III) complexes are claimed to show these model cyt c. [743]  $[Fe(4-MeO-TPP)Cl]$  is converted by  $tBuOOH$  to an isoporphyrin with a t-butylperoxy group entering at the meso-position; this may be an alternative process to that usually considered for reactions between metalloporphyrins and organoperoxides. [744]

There has been more useful progress on picket fence and other porphyrins with elaborate structures. The field has been given a readable review. [745] The EPR of some "basket handle" porphyrins  $[Fe(por)Cl]$  shows Fe(III) in a mixed  $3/2 - 5/2$  spinstates. [746] New cyclophane-capped porphyrins include Fe(III) complexes where the spin state is affected by the length of the bridging chains or the size of the axial ligand, which also controls whether 5- or 6-coordination is achieved. [747] Using  $2-C_6H_4NHCOCMe_2(CH_2)_nCO_2R$ , ( $n = 1,2,3,10$ ;  $R = H, Me, CH_2Ph$ ), "pickets" gave Fe(III) complexes with Cl or Br as 5th ligands. [748] Another capped porphyrin links  $C_6H_6$  to  $[Fe(TPP)]$  by 4  $[O(CH_2)_4OC(O)]$  units, with less steric hindrance for CO binding than for analogues with 2 or 3  $CH_2$  groups. [749] Oxidation of an Fe(II) - tetra(2,5- $tBu_2$ -4-OH-phenyl) porphyrin complex with peracids or  $O_2$  and base gives a stable complex which oxidises phenols, while its Fe(III) complex with 1-Meimid is reduced to Fe(II); comparisons were drawn with peroxidase or cyt c peroxidase. [750] Another picket fence porphyrin complex with axial 1-lauryl-

-2-methylimidazole, when dissolved in a phospholipid bilayer, binds  $O_2$  reversibly at pH 7 at  $37^\circ C$ ; electric dipole measurements indicate the porphyrin and bilayer lie parallel. [751] The X-ray structure of a  $t$ Bu picket fence porphyrin- $O_2$ - $SC_6F_5$  complex shows  $O_2$  is monodentate inside the porphyrin cup; the O-O distance is apparently surprisingly short ( $1.14 \text{ \AA}$ ) due to disorder and decomposition by X-rays and the Moessbauer spectrum resembles that of oxygenated cyt P-450. [752] Four isomers were found for another  $[Fe(por)OH]$  picket fence complex, which is converted to an O-bridged dimer by 8hr refluxing in  $CHCl_3$ . [753] The enthalpy and entropy for  $O_2$  and CO binding to the Fe(II) tetrapivalamidophenylporphyrin in phosphatidylcholine lipids are similar to those for Hb. [754] In an extension of earlier work it was found that  $O_2$  and NO have similar trends in binding constants with various Fe(II) capped porphyrin azole complexes and it was concluded that the diatomic ligands adopted the bent monodentate configuration but the enforced linearity of CO coordination would generate different steric constraints for the different regions of the ligand. [755] The ingenious model for cyt c oxidase,  $[Fe(tetranicotylporphyrin)Cu]^{2+}$ , has been prepared and shows weak ferromagnetism between two spins of  $\frac{1}{2}$ , but this coupling is lost in some solvents, though not in  $MeOH/CHCl_3$  with or without 1-Meimid; a diamagnetic product or products are formed in dmf. [756]

### 6.5. Iron-Sulphur Compounds

This section concerns proteins containing iron and sulphur and synthetic compounds specifically designed to mimic them. Compounds containing molybdenum and other metals are deferred to the next section.

General papers include a wide ranging review on S ligands in metalloproteins and enzymes [757] and two reviews on core extrusion analysis, [758,759] one of which highlights the use of detergents and chromatography. [758] EPR is used to determine spin-lattice relaxation times; adding the Dy-edta complex as paramagnetic relaxer allows the determination of the distance of a paramagnetic centre from the protein surface. [760]

Too many papers refer to ferredoxins without specifying which of the various ferredoxin structures is present. Examples of these are an rR study of Se replacement of S, [761] the discovery of an electron transport protein in nitrite reduction, [762] electron transfer with gold electrodes using "rotating disc hydrodynamic voltammetry", [763] and the discovery of [Fe-S] proteins in spinach glutamate synthase [764] and mitochondrial NADH dehydrogenase. [765]

A preliminary X-ray study on a rubredoxin from Desulfovibrio desulfuricans shows this is the smallest so far, with only 45-48 residues, but including the unusual histidine. [766] Rubredoxin and a [4Fe-4S] ferredoxin have been found in Clostridium formicoaceticum. [767] The role of surface charge is important in CV studies of rubredoxin and other proteins using  $\text{RuO}_2$  or  $\text{IrO}_2$  electrodes. [630]

Turning to [2Fe-2S] proteins, a new ferredoxin (Fd) has been isolated from Trichomonas vaginalis. [768] Benzene dioxygenase from Pseudomonas putida consists of a flavoprotein, a [2Fe-2S] Fd and two other [2Fe-2S] units whose EPR resembles Rieske proteins. [769] An admirable group of papers [770-4] uses modern NMR methods to obtain structural information.  $^{13}\text{C}$  enrichment of Fd from the bacterium Anabaena variabilis allowed full assignments with the aid of 2-D and decoupling techniques; relaxation times were shorter in the more paramagnetic reduced form. [770] Similar work on Fd from another bacterium or from spinach gave improved assignments of the cys protons and studies of the hyperfine shifts. [771] The  $^1\text{H}$  and  $^{13}\text{C}$  spectra for the histidine units gave conformational information in reduced and oxidised forms. [772]  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  and Fd-NADP reductase bind in a similar area containing at least 3 glutamate residues. The primary sequence of aminoacids was obtained using 360 or 470 MHz spectra, 2-D methods,  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  binding and chemical modification with carboxypeptidase A and trinitrobenzenesulphonic acid. [774] Another  $^1\text{H}$  NMR study assigned all the protons near Fe and distinguished  $\text{CH}_2$  units near Fe(II) from those near

Fe(III) by the different temperature dependence of their chemical shifts.

[775] The rR spectrum of the Fd from Clostridium pasteurianum which is known as the "red paramagnetic protein" (RPP) shows this has the usual structure but a slight weakening of all the Fe-S bonds and a change in the conformation occur. [776] The kinetics of reductions by  $[\text{Cr}(\text{15-aneN}_4)]^{2+}$  of [2Fe-2S] and [4Fe-4S] proteins show both are inner sphere processes and that binding  $\text{Cr}^{3+}$  to the reduced protein does not affect oxidation by  $[\text{Co}(\text{NH}_3)_6]^{3+}$ ; addition of  $[\text{Cr}(\text{en})_3]^{3+}$  does not inhibit reduction by the Cr(II) complex but actually accelerates it for the [2Fe-2S] protein. [777] The Rieske [2Fe-2S] protein from Thermus thermophilus was isolated and the iron atom which is reduced was shown to have at least one non-cys ligand, not identified at this stage, there being just 2 cys residues between the 2Fe. [778]

The behaviour of ferredoxins containing 3Fe atoms is curiouser and curiouser. There was a report of the crystallisation and preliminary structural work on a 3Fe Fd from Desulfovibrio gigas, but the authors were unable at the time to determine whether they had encountered a planar or 3/4 cubic configuration. [779] ESR spectra on [3Fe-3S] and [3Fe-4S] Fd proteins suggest these are more similar than the X-ray work claimed but this inconsistency was thought possibly to arise from modifications when crystals became frozen aqueous solutions. [780] Unusually fast relaxation occurs in the 3Fe proteins from D. gigas or Azotobacter vinelandii, confirming earlier ideas that these have low lying excited states arising from spin coupling. [281] An EPR study of the conversion of the [3Fe-4S] cluster of Fd II from D. gigas to [4Fe-4S] shows this could occur in in vivo conditions. [782] The [3Fe-3S] and [3Fe-4S] cores can be distinguished by rR spectra. [783] Fd proteins from Pseudomonas ovalis and Mycobacterium smegmatis are shown by rR spectra to contain both 3Fe and 4Fe clusters. [784] A. vinelandii FdI is normally isolated as a combination of [4Fe-4S] and [3Fe-3S] cores, but it can be reconstituted as an 8Fe core, so it was suggested that one Fe is lost in the isolation procedure. [785] The use of  $^{35}\text{S}^{2-}$  to study the exchange of  $\text{S}^{2-}$  in aconitase showed

3S exchanged easily, so it was supposed that the [3Fe-4S] cluster was accessible to solvent; Fe is also taken up easily, suggesting formation of a [4Fe-4S] cluster to convert the inactive enzyme to an active one. [786] The oxidation of Fd I from A. vinelandii by  $[\text{Fe}(\text{CN})_6]^{3-}$  produced an intermediate corresponding to a 3-electron oxidation of the [4Fe-4S] unit; this was believed to be a cysteinyl-disulphide radical formed by the loss of S and Cys from the [4Fe-4S] cluster. [787] In another study of this reaction, using  $^1\text{H}$  NMR spectra, excess of  $[\text{Fe}(\text{CN})_6]^{3-}$  converted the [4Fe-4S] core to a [3Fe-4S] and then a [3Fe-3S] unit. [788]

In work on [4Fe-4S] proteins it was shown that this structure is found in FdI from D. africanus, [789] and in a hydrogenase from D. vulgaris. [790] EXAFS work on a hydrogenase from Methanobacterium thermoautotrophicum shows this is similar to  $[\text{Fe}_4\text{S}_4(\text{SPh})_4]^{2-}$ , but there was uncertainty on the S/Fe ratio. [791] Substituting Se for the core S in oxidised Fd from Clostridium pasteurianum gave little change in the Moessbauer spectrum, but reduction gave a much more complex spectrum and a triple EPR spectrum, attributed to the usual spin state of  $\frac{1}{2}$  being supplemented by  $\frac{3}{2}$  and  $\frac{7}{2}$  states. [792] The [4Fe-4S] proteins exchange D for H more readily than HP or [2Fe-2S] proteins, especially in the oxidised state; electron spin echo decay measurements show this occurs close to the metals. [793]

A [4Fe-4S] cluster in a 7Fe Fd from P. ovalis was found to be both oxidisable and reversible; it is asserted that this is the first such Fd to show this in its native form. [794] A kinetic study of the reaction of the oxidised forms of plastocyanin and azurin with HP proteins showed this was not a simple process. [633]

A new class of Fd containing 5Fe atoms has been found in a lichen Lobaria pulmonaria. [795]

Next, model compounds, beginning with mononuclear ones. Unstable  $[\text{Fe}(\text{SR})_4]^-$  compounds ( $\text{R} = \text{Ph}$ ,  $\text{C}_6\text{H}_4\text{Me}$ ) have been prepared [796] from  $[\text{Fe}\{\text{o}-(\text{SCH}_2)_2\text{C}_6\text{H}_4\}_2]^-$ .  $[\text{Fe}(\text{H}_2\text{O})_4\text{Cl}_2]$  reacts with

$\text{o-C}_6\text{H}_4\text{S}_2^{2-}$  to give planar  $[\text{Fe}(\text{S}_2\text{C}_6\text{H}_4)_2]^{2-}$ , but if  $\text{PMe}_3$  is present square pyramidal  $[\text{Fe}(\text{S}_2\text{C}_6\text{H}_4)(\text{PMe}_3)_3]$  is formed, which in turn reacts with CO to form  $[\text{Fe}(\text{S}_2\text{C}_6\text{H}_4)(\text{PMe}_3)_2(\text{CO})_2]$  and  $[\text{Fe}(\text{S}_2\text{C}_6\text{H}_4)(\text{PMe}_3)_2(\text{CO})]_2$ . [797] Some Fe(III) complexes of short peptides containing two cys units in the sequence found in rubredoxins had similar electronic, CD and MCD spectra to the protein. [798] Fe(II) complexes of similar type showed similar likenesses to the reduced form but only the complex of  $\text{PhCH}_2\text{O}$  cysproleucysOMe had reversible redox behaviour; it could transfer electrons from NADP oxidoreductase to ferricytochrome c. [799]

Some dinuclear Fe-S complexes have been studied. A variety of routes may produce  $[\text{Fe}_2\text{S}_2(\text{S}_5)_2]^{2-}$ , the simplest method being the reaction of  $\text{FeCl}_2$ , S and Na; using Se for S in this synthesis gave the Se analogue whose X-ray structure shows the expected 2 Se bridges and 2 bidentate  $\text{Se}_5^{2-}$  ligands, resulting in tetrahedral Fe. [800]  $[\text{Fe}_2\text{S}_2\text{Cl}_4]^{2-}$  was used to prepare compounds with dicysteinepeptides; an Fe(II) complex with the peptide can be blended with S to give the  $[2\text{Fe}-2\text{S}]$  model system. [801]

Another series of derivatives from  $[\text{Fe}_2\text{S}_2\text{Cl}_4]^{2-}$  is prepared with 4 molecules of pyrrole or 2 of 2,2'-biphenyldiol. [802] Ferrous ammonium sulphate reacts with  $(\text{NH}_4)_2\text{WS}_4$  and  $\text{Et}_4\text{NI}$  in the presence of saccharoses in water to give  $(\text{Et}_4\text{N})_2[\text{Fe}(\text{H}_2\text{O})_2(\text{WS}_4)_2]$ , and  $[\text{FeCl}_2(\text{WS}_4)]^{2-}$  reacts [803] with NaSH to give  $[\text{Fe}_2\text{S}_2(\text{WS}_4)_2]^{2-}$ .

A thorough investigation of linear  $[\text{Fe}_3\text{S}_4(\text{SR})_4]^{3-}$ , (R = Et, Ph) includes magnetization studies and Moessbauer spectra in dmf to show that in both solid and solution phases there is strongly antiferromagnetic Fe(III), though the coupling between the end Fe atoms is described as being between  $-100$  and  $+100 \text{ cm}^{-1}$ ! These results contrast with the lowspin character of ferredoxins; the authors comment that the linear structure has been found in aconitase. [804]

Improved preparations of  $[\text{Fe}_4\text{S}_4(\text{SR})_4]^{3-}$ , (R = alkyl), from  $\text{FeCl}_2$ , NaSR and NaSH in dmf allow these to be formed without previously isolating the

dianions. The Et and <sup>t</sup>Bu compounds have compressed tetragonal structures which are about 2.5% bigger in volume than the dianions. [805] Reacting  $[\text{Fe}_4\text{S}_4\text{Cl}_4]^{2-}$  with  $2\text{SPh}^-$ , (E = O, S), gives  $[\text{Fe}_4\text{S}_4\text{Cl}_2(\text{EPh})_2]^{2-}$ , which are said to be the first  $\text{Fe}_4\text{S}_4$  cores which do not have the compressed  $\text{D}_{2d}$  structure; further reaction of the SPh complex with  $2\text{OR}^-$ , (R = Ph, 4-MePh) gives replacement of the remaining chlorine. [806] As models for the  $[\text{4Fe-4S}]$  clusters in nitrogenase various  $[\text{Fe}_4\text{S}_4\text{L}_4]^{2-}$  species were studied with the expected result that for L = SPh, OPh or Cl there was charge delocalisation, but not for L =  $\text{Et}_2\text{NCS}_2$ ; it was suggested that the unusual spin state of the iron-sulphur cube in nitrogenase was due to unusual coordination. [807]  $[\text{Fe}_4\text{S}_4(\text{S}^t\text{Bu})_4]^{2-}$  exchanges its thiolate ligands with N - Phenylacetylcysteinyl-glycylcysteinamide. [808] Reacting  $[\text{Fe}_4(\text{cp})_4\text{S}_6]$  with  $[\text{MoOCl}_3(\text{thf})_2]$  gives the unusual cation  $[\text{Fe}_4(\text{cp})_4\text{S}_5]^+$  with a modified  $\text{Fe}_4\text{S}_4$  cube in which one  $\text{S}^{2-}$  is replaced by an  $\text{S}_2^{2-}$  which is  $\eta^2$  to one Fe and  $\eta^1$  to two others. [809] Two new hexanuclear clusters are described.  $[\text{Fe}_6\text{S}_6\text{I}_6]^{2-}$ , consisting of 2  $\text{Fe}_3\text{S}_3$  "chairs" linked by Fe-S bonds, is prepared [810] by the reaction of specific proportions of Fe,  $\text{I}_2$ , S and  $\text{I}^-$ .  $[\text{Fe}_6\text{S}_6\text{Cl}_6]^{3-}$ , a hexagonal prismatic structure, is produced from  $\text{FeCl}_2$ ,  $\text{SPh}^-$ , S and  $\text{Cl}^-$  in MeCN. [811] The new clusters  $[\text{Fe}_6\text{S}_9(\text{SR})_2]^{4-}$ , (R = Et,  $\text{CH}_2\text{Ph}$ ) and  $[\text{Fe}_6\text{S}_9(\text{SMe})_2\text{Na}_2]^{6-}$  are described; the latter has a condensation of  $4[\text{Fe}_3\text{NaS}_4(\text{SR})_6]^{5-}$  units. [812]

#### 6.6. Iron-Molybdenum-Sulphur Compounds

This section describes nitrogenases, compounds which are designed to mimic this protein and other Fe-Mo species whose connection with nitrogenase is more tenuous.

The unlikely reagent bipy gives a  $[\text{2Fe-2S}]$  compound on reaction with oxidised Fe protein from Azotobacter vinelandii nitrogenase. [813] Reactions of  $\text{Fe}_3\text{MoS}_4$  clusters include that of  $[\text{Fe}_7\text{Mo}_2\text{S}_8(\text{SEt})_{12}]^{3-}$  with



tetrachlorocatechol ( $\text{H}_2\text{cat}$ ) to give  $[\text{Fe}_6\text{Mo}_2(\text{SEt})_6(\text{cat})_2]^{4-}$  which in MeCN forms  $[\text{Fe}_3\text{MoS}_4(\text{SR})_3(\text{cat})(\text{MeCN})]^{2-}$ ; the MeCN can be replaced by  $\text{CN}^-$ ,  $\text{N}_3^-$ ,  $\text{N}_2\text{H}_4$ ,  $\text{NH}_3$ ,  $\text{NH}_2\text{NHPH}$  or pip and other  $\text{SR}^-$  derivatives were prepared. [814] The crystal structure of  $(\text{Et}_4\text{N})_3[\text{FeMo}_2\text{S}_8\text{O}_2]\cdot\text{EtCN}$  confirms that O is bound only to Mo. [815] Another structure report describes [816]  $[\text{Fe MoS}_3(\text{SC}_2\text{H}_4\text{S})_2]^{3-}$  as "S-shaped". New preparations [817] include  $[\text{Fe}_2\text{MoS}_4(\text{PhSH})_2(\text{OMe})_4]^-$  and  $[\text{Fe}_8\text{MoS}_9(\text{PhCH}_2\text{SH})(\text{OMe})_6]$ , which both reduce  $\text{C}_2\text{H}_2$  and  $\text{N}_2$ ,  $[\text{Fe}_2\text{MoS}_4\text{Cl}_4]^{2-}$ ,  $[\text{Fe}_2\text{MoS}_4\text{Cl}_4(\text{SR})]^{2-}$ , ( $\text{R} = \text{Ph}, \text{CH}_2\text{Ph}$ ), and  $[\text{S}_2\text{MoS}_2\text{FeS}_2\text{MoOS}_2]^{3-}$ . The formation of an FeMo complex was indicated by CD studies of the interaction of  $[\text{Fe}_4\text{S}_4(\text{SR})_4]^{2-}$  with Mo(IV) peptide complexes, ( $\text{R} = {}^i\text{Pr}, \text{Ac-cys-OH}$ ). [818]  $\text{Fe}(\text{O}_2\text{CMe})_2$  is used to prepare  $[(\text{AcO})_2\text{FeS}_2\text{MS}_2]^{2-}$ , ( $\text{M} = \text{Mo}, \text{W}$ ), from which  $[(\text{PhS})_2\text{FeS}_2\text{MS}_2]^{2-}$  can be prepared, and  $[\text{Fe}(\text{MoO}_2\text{S}_2)_2]^{2-}$ , which can also be prepared from  $\text{Fe}(\text{S}_2\text{CNR}_2)_2$  and which is thought to have Fe-O-Mo-O-Fe bridging. [819]

#### 6.7. *Haemerythrin*

There have been many valuable developments on this interesting protein. Two readable reviews have been published. [820, 821] The structure of metaquo-haemerythrin from *Thermiste dyscritum* shows one 6- and one 5-coordinate (sort of trigonal bipyramidal) Fe, but unlike the metazido- form there is no small ligand on the sixth site of the 5-coordinate Fe so the designation "metaquo" is a misnomer. [822] There has been further work on the metazido structure. [823]

A thorough  ${}^1\text{H}$  NMR study of methaemerythrin and semimethaemerythrin concentrated on the bridging histidine resonances, which are not shifted as much as normal for highspin Fe(III) because of the antiferromagnetism. In the semimet-form the his bonded to Fe(II) could be distinguished from those bonded to Fe(III); sulphatosemimet- is more antiferromagnetic than the azidosemimet-form [824] The rR spectra of oxyhaemerythrin and the azidomet-form gave identification of many important vibrations;  $\text{D}_2\text{O}$  substitution indicated

[825] that  $O_2$  binds as  $HO_2^-$ .

Oxidation of deoxyhaemerythrin by  $NO_2^-$  is unusual in producing the semimet-form, probably by an inner-sphere process, whereas outer-sphere oxidants like  $[Fe(CN)_6]^{3-}$  give methaemerythrin; the  $NO_2^-$  oxidation proceeds via an EPR - silent nitrosyl intermediate. [826] In another kinetic study this deoxy- to semimet- oxidation was achieved [827] by various  $[Fe(CN)_4XY]^{n-}$  species ( $X = CN^-$ ,  $Y = CN^-$ ,  $NH_3$ ,  $PPh_3$ , 4- $NH_2$ py;  $XY = bipy$ ) and of semimet- to met- by these reagents and  $[Co(terpy)_3]^{3+}$ . There have been further studies on the related protein rusticyanin, including the kinetics of  $Fe^{2+}$  reduction and EPR of reaction with the Dy(III) - edta complex. [828]

#### 6.8. Iron Transport Proteins

The subject of siderophores has been reviewed again, with emphasis on ligand synthesis, complex formation energetics and complex configuration. [829]  $Fe^{3+}$  is stereospecifically complexed by the  $\Delta$ -isomer of rhodotorulic acid. [830] Various  $RN(OH)C(O)(CH_2)_n C(O)N(OH)R$ , ( $R = Ph, Pr$ ;  $n = 3-10$ ) ligands form non-antiferromagnetic  $Fe_2X_3$  complexes like rhodotorulic acid and these can be converted to  $[Fe_2X_2(OR)_2]$ , ( $R = Me, Et$ ) in alcohols. [831, 832] Mono-, di- and trihydroxamate complexes of Fe can be distinguished by their response to  $CN^-$ ; the first forms  $[Fe(CN)_5(NO)]^{3-}$ , the second loses its blue colour and the third does not react. [833] The reactions of ferrioxamine B with edta and hydroxamates were studied as models for the release of Fe from siderophores. [834] Retrohydroxamate, a polypeptide with 3 N-methylhydroxamate units, has some biological activity as a synthetic siderophore despite being a weaker ligand. [835]

Many new siderophores have been characterized. The bacterial siderophore pyoverdine Pa was found by mass and NMR spectra to be an octapeptide bound to 2,3-diamino-6,7-dihydroxyquinodine to give two hydroxamate and one catechol functional groups. [836] Vibrio cholerae has a siderophore, vibriobactin, which contains three 2,3-dihydroxybenzoate and two threonine residues. [837]

Rhizobactin comes from the  $N_2$ -fixing bacterium Rhizobium meliloti, but has neither catechol nor hydroxamate ligands. [838] A siderophore of ferrichrome type appears in the bacterium Aspergillus ochraceus. [839] That from the microbe Proteus mirabilis contains a unit of isovaleric acid,  $Me_2CHCH(OH)CO_2H$ . [840] A strain of smut fungus has a ferrichrome siderophore of glycol units in a heptapeptide instead of the usual hexapeptide. [841]

The rR spectrum shows 2,3-dihydroxybenzoate and EXAFS shows thioglycolate bond to Fe(III) in ovotransferrin. [842] Pyrophosphate can transport Fe liberated from transferrin into rat liver mitochondria and might fulfil this role in vivo. [843] ESR spectra of an  $Fe^{3+}$ -hist-citrate model system for transferrin confirm Fe is bound to 2 his, 3 citrate and possibly 1  $HCO_3^-$  but the natural system probably has coordination by ascorbate. [844]

#### 6.9. Ferritin

Two reviews [845,846] have appeared, one of them being a good easy general survey suitable for novices. [845] The structure of horse spleen ferritin has been described afresh. [847] The X-band EPR spectra of ferritin and haemosiderin show two features from  $Fe(OH)_3$  cores which are similar; [848] electron microscopy and Moessbauer evidence reinforce the conclusion that these are similar materials but that haemosiderin particles are smaller. [849] The molecular weight of pigeon ferritin is given, with incredible precision, as 354,813, including 5810 Fe atoms per mol; chicken ferritin is smaller. [850] The UV difference spectra of the  $Fe^{2+}$  - apoferritin reaction, together with the effects of  $Zn^{2+}$  and  $Tb^{3+}$ , shows there is more than one site both for  $Fe^{2+}$  binding and for  $Fe^{3+}$  binding after the reorganization that follows oxidation. [851] The release of Fe from ferritin is promoted by  $O_2^-$  and suppressed by adding superoxide dismutase. [852] The commercial preparation Niferex, a neutralised solution of an Fe(III) carbohydrate complex, has a Moessbauer spectrum and X-ray diffraction like ferrihydrite and is therefore like ferritin. [853]

## 7. ORGANOMETALLIC COMPOUNDS

Organising material in this vast topic is very difficult. Papers are described under the heading for the dominant ligand in the chemistry which is described, but such a classification is occasionally both arbitrary and inconsistent. Readers are advised to consult more than one section when looking for a particular topic.

### 7.1. Carbonyls

We begin with  $[\text{Fe}(\text{CO})_5]$ . Ab initio calculations on its photodissociation lead to the conclusion that excitation to a singlet state occurs before crossing to  $^3\text{E}'$  and dissociating. [854] The laser induced fluorescence of  $[\text{Fe}(\text{CO})_5]$  is described. [855]

Numerous examples have been described of the use of  $[\text{Fe}(\text{CO})_5]$  in the isomerization of alkenes. Substrates reported in 1984 include 1- and trans-2-pentene, [856,857] 1-hexene, [858]  $\text{CH}_2=\text{CH}(\text{CH}_2)_n\text{N}(\text{SiMe}_3)_2$ , [859] and this general reaction has undergone kinetic study. [860] Laser irradiation of  $[\text{Fe}(\text{CO})_5]$  produces a thermally active catalyst for the gas phase hydrogenation of  $\text{C}_2\text{H}_4$ . [861]  $\text{PhN}=\text{CHPh}$  has also been hydrogenated with  $[\text{Fe}(\text{CO})_5]$  catalysis, [862] and the role of  $[\text{HFe}(\text{CO})_4]$  in this reaction has been described. [863]  $[\text{Fe}(\text{CO})_5]$  with  $\text{CO}$ ,  $\text{H}_2\text{O}$ , base and a crown ether catalyses the regiospecific hydrogenation of anthracene, quinoline and other N-heterocycles. [864] Two reports describe its use in the carbonylation of benzyl halides to phenylacetic acid derivatives. [865,866] It can, in dmf, induce the replacement of one Cl in  $\text{Cl}(\text{CH}_2)_n\text{CCl}_2$  by H. [867]

Decomposition studies of  $[\text{Fe}(\text{CO})_5]$  include experiments on the process carried out after adsorption on zeolites, [868] or on polythene, where  $[\text{Fe}(\text{CO})_4]$  and  $[\text{Fe}(\text{CO})_3]$  were spectroscopically detected. [869]

Photooxidation in  $\text{CCl}_4$  gave  $\text{FeCl}_2$  and organics formed from a dichlorocarbene resulting from the oxidative addition of  $\text{CCl}_4$  to  $[\text{Fe}(\text{CO})_4]$ . [870]  $\gamma$ -irradiation of  $[\text{Fe}(\text{CO})_5]$  in a Kr matrix gave, [871]

according to EPR,  $[\text{Fe}(\text{CO})_5\text{Kr}]^+$ . The carbonyl reacts with 2-EtimidH (L) to give  $[\text{FeL}_6]^{2+}[\text{FeH}(\text{CO})_5]^-$ , whose anion is fluxional. [872]

Turning to  $[\text{Fe}(\text{CO})_4]$  species,  $[\text{Fe}(\text{CO})_4]^-$  has been shown to abstract halogen from  $\text{CCl}_4$ ,  $\text{CF}_3\text{Br}$  and  $\text{CF}_3\text{I}$ . [873]  $[\text{Fe}(\text{CO})_4]^{2-}$  is a catalyst in the formation of hydrocarbons from CO and  $\text{H}_2$  at 350–450 °C; the role of  $\text{K}^+$  is discussed. [874] Most work in this section concerns  $[\text{Fe}(\text{CO})_4\text{L}]$  complexes. Their XPE spectra show changes in d orbital energies in line with expectations on whether L is  $\pi$ -bonding. [875] The ligand RP:PR, [R = 2,4,6-( $t$ Bu) $_3\text{C}_6\text{H}_2$ ], is monodentate in  $[\text{Fe}(\text{CO})_4(\text{RP:PR})]$ . [876]  $\text{HC}(\text{PPh}_2)_3$  is more versatile, forming monodentate complexes  $[\text{Fe}(\text{CO})_4\text{L}]$ , bidentate complexes  $[\text{Fe}(\text{CO})_3\text{L}]$  and polymetallic complexes  $[(\text{CO})_4\text{FeLFe}(\text{CO})_3]$  and  $[(\text{CO})_4\text{FeLMo}(\text{CO})_4]$ . [877] There seems to be a craze for forming phosphine complexes with molybdenum moieties tacked on. Other examples for 1984 include  $[\text{Fe}(\text{CO})_4\{\text{P}(\text{NMe}_2)_2\text{Mo}(\text{cp})(\text{CO})_2\}]$  with an FeMoP ring, [878]  $[\text{Fe}(\text{CO})_4-\{\text{P}(\text{NPhPF}_2)\text{Mo}(\text{CO})_3\}]$ , [879] and  $[\text{Fe}(\text{CO})_4\{\text{PPh}_2\text{OPPh}_2\text{Mo}(\text{CO})_5\}]$ . [880] There is an unusual equatorial phosphine in  $[\text{Fe}(\text{CO})_4-\{\text{Me}_3\text{Si}\}_2\text{C:PN}(\text{SiMe}_3)_2]$ , whose C=P bond of 1.657 Å is described as short and this ligand has a 30° twist. [881] A tetrameric phosphazene with oxazole rings on alternate P–N bonds forms a complex with 4  $[\text{Fe}(\text{CO})_4]$  groups. [882] The phosphorus ylide complexes  $[\text{Fe}(\text{CO})_4\text{CH}(\text{R})\text{P}^+\text{Ar}_3]$ , (Ar = Ph, *p*-tolyl) are formed when the parent phosphine is added to a mixture of  $[\text{Fe}(\text{CO})_4(\text{SiMe}_2\text{CH}_2)_2]$  and an aldehyde; alkylidene intermediates are invoked for the reaction and the  $\text{Ph}_3$  derivative reacts with  $\text{CF}_3\text{SO}_3\text{H}$  to give  $(\text{PhCH}_2\text{PPh}_3)^+(\text{CF}_3\text{SO}_3)^-$ . [883] New arsine complexes of  $[\text{Fe}(\text{CO})_4]$  include those with  $\text{AsMe}(\text{NMe}_2)_2$ ,  $\text{AsMeX}_2$  or  $\text{AsMe}_2\text{X}$ , (X = Cl, OEt, SEt), [884] and antimony provides  $[\text{Fe}(\text{CO})_4-\{\text{Me}_3\text{Si}\}_2\text{CHSbSbCH}(\text{SiMe}_3)_2]$  and  $[\{\text{Fe}(\text{CO})_4\}_2\text{SbCH}(\text{SiMe}_3)_2]$  which have  $\text{FeSb}_2$  and  $\text{Fe}_2\text{Sb}$  triangles respectively. [885] The reaction of  $(\text{Me}_3\text{E})_3\text{Sb}$ , (E = Si, Ge) with  $[\text{Fe}_2(\text{CO})_9]$  gives  $[\text{Fe}(\text{CO})_4\text{Sb}(\text{EMe}_3)_2]$  but if  $(\text{Me}_3\text{Sn})_3\text{Sb}$  is used the product is  $[\text{Fe}(\text{CO})_4(\text{SnMe}_3)_2]$ . [886]

Bismuth is not so varied and is content with  $[\text{Fe}(\text{CO})_4(\text{BiR}_3)]$ , ( $\text{R} = \text{alkyl}$ ).

[887]

$[\text{Fe}(\text{CO})_4]$  bonds to group IV ligands in forming  $[\text{Fe}(\text{CO})_4(\text{CO}_2)]^-$  when

$[\text{Fe}(\text{CO})_5]$  is adsorbed on  $\text{MgO}$ . [888] Kinetic studies and the observation of

CIDNP showed H atom transfer occurs in the reaction between

$[\text{Fe}(\text{CO})_4(\text{SiCl}_3)\text{H}]$  and dienes. [889]  $[\text{Fe}_2(\text{CO})_9]$  reacts with

$[\{\text{MeC}_5\text{H}_4\}(\text{CO})_2\text{Mn}^+\text{Ge}]$  to give  $[\{\text{Fe}(\text{CO})_4\}_4\text{Ge}]$  and  $[\{\text{Fe}(\text{CO})_4\}_3 - \text{Ge}\{\text{Mn}(\text{CO})_2(\text{MeC}_5\text{H}_4)\}_2]$ . [890] Sn or Ge porphyrins form

$[\text{Fe}(\text{CO})_4\{\text{M}(\text{por})\}]$ ; the X-ray structure shows the Sn-OEP complex has square

pyramidal Sn with Fe axial and trigonal bipyramidal Fe. [891] One of the most

exotic  $[\text{Fe}(\text{CO})_4]$  complexes is formed when  $[\text{Fe}(\text{CO})_5]$  reacts with

$[\text{Cr}_2(\text{cp})_2(\text{O}^t\text{Bu})_2]$  to produce  $[\text{Cr}(\text{cp})(\mu\text{-O}^t\text{Bu})_2\text{Cr}(\text{cp})\text{Fe}(\text{CO})_4]$  which

has an equilateral  $\text{Cr}_2\text{Fe}$  triangle with no bridging ligands between Cr and

Fe. [892]

There is also a good selection of  $[\text{Fe}(\text{CO})_3]$  species.  $\text{Me}_2\text{S}_2$  reacts with

$[\text{Fe}_3(\text{CO})_{12}]$  to give  $[\{\text{Fe}(\text{CO})_3\}_2\text{SMe}]_2$  and  $[\text{Fe}_2(\text{CO})_6(\text{SMe})_2]$ .

[893] Dithiocarbamate, dithiophosphate and xanthate (X) complexes which have

been prepared are  $[\text{Fe}(\text{CO})_3\text{IX}]$  and  $[\text{Fe}(\text{CO})_2\text{X}_2]$ . [894] Photochemistry in

organic glasses at 100K converts  $[\text{Fe}(\text{CO})_4\text{PPh}_3]$  into  $[\text{Fe}(\text{CO})_3\text{PPh}_3]$  or

$[\text{Fe}(\text{CO})_3(\text{PPh}_3)\text{L}]$ , ( $\text{L} = \text{glass medium}$ ); in  $\text{SiEt}_3\text{H}$  the product is

$[\text{Fe}(\text{CO})_3\text{H}(\text{PPh}_3)(\text{SiEt}_3)]$  whose fac - isomer can be converted to the mer -

form by further irradiation. [895] Bipy and phen (L) complexes  $[\text{Fe}(\text{CO})_3\text{L}]$

can be prepared by carbonylation of  $[\text{Fe}_2(\text{CO})_7\text{L}]$  or by reacting the diimine

with  $[\text{Fe}(\text{CO})_3(\text{PhCH}=\text{CHCMe}=\text{CH}_2)]$ . [896] The silicon ligand

$\text{Me}_2\text{SiCH}=\text{CMe}=\text{CH}$  also forms an  $[\text{Fe}(\text{CO})_3]$  complex, [897] but this might be

a diene complex rather than with bonding through Si. Gaseous  $[\text{Fe}(\text{CO})_4]$

radicals react with H in  $\text{H}_2$  to give  $[\text{Fe}(\text{CO})_3\text{H}_n]^-$  ( $n = 0-3$ ). [898]

There are a few complexes with fewer than 3 carbonyl groups per iron atom.

These include  $[\text{Fe}(\text{CO})_2(\text{dtc})_2]$  and  $[\text{Fe}(\text{CO})(\text{xanthate})_2\text{L}]$ , for which the

$\text{PPh}_3$  complex was more stable than those with N donors. [899] There has been

a thorough study of the complexes  $[\text{Fe}(\text{CO})_2(\text{PPh}_3)_2(\text{RNCS})]$ , ( $\text{R} = \text{Me}, \text{Ph}$ ) prepared by the reaction of RNCS with the tris-phosphine complex; RNCS acts as an  $\eta^2$ -ligand with back-bonding to both the  $\text{CN}-\pi^*$  and S-d orbitals, and further reactions are described substituting one  $\text{PPh}_3$  by  $\text{P}(\text{OPh})_3$  and the RNCS by  $4\text{F}-\text{C}_6\text{H}_4\text{N}_2^+$ , while various electrophiles E convert RNCS to RENCS. [900] Reacting  $[\text{Fe}(\text{CO})_2(\text{PPh}_3)_2(\text{CS}_2)]$  with  $\text{Ph}_2\text{PCC}^t\text{Bu}$  gave  $[\text{Fe}(\text{CO})(\text{PPh}_3)(\text{SCOMe})(\text{SC}^t\text{Bu:CHPPh}_2)]$  which could in turn be converted to a wide range of other phosphinoalkenethiolate complexes. [901]

There is the usual abundance of dinuclear complexes. Reacting  $[\text{Fe}_3(\text{CO})_{12}]$  with 2-nitropropane in PhMe gives  $[\text{Fe}_2(\text{CO})_6(\mu\text{-OCCH}_2\text{Ph})(\mu\text{-NCMe}_2)]$ . [902] EHMO calculations on  $[\text{Fe}_2(\text{CO})_6\text{S}_2]$  and related compounds were compared to their PE spectra. [903] The reaction of  $[\text{Fe}(\text{CO})_5]$  with  $[\text{Cr}(\text{CO})_5(\text{SCSC}_2\text{H}_4\text{S})]$  gives  $[\text{Fe}_2(\text{CO})_6(\mu\text{-SC}_2\text{H}_4\text{S})]$ ,  $[\text{Fe}_2(\text{CO})_6(\mu\text{-SCSC}_2\text{H}_4\text{S})]$ ,  $[\text{Fe}_3(\text{CO})_9(\mu_3\text{-S})_2]$  and  $[\text{Fe}_3(\text{CO})_9(\mu_3\text{-S})(\mu_3\text{-CSC}_2\text{H}_4\text{S})]$ ; in the last of these a C atom bridges 2Fe and an S links across to the other one. [904] A dithiolate complex of  $[\text{Fe}_2(\text{CO})_6]$  is said to be produced in the reaction of  $[\text{Fe}_3(\text{CO})_{12}]$  with sulphur. [905] This carbonyl reacts with  $(\text{Me}_3\text{SiN})_2\text{S}$  to form  $[\text{Fe}_2(\text{CO})_6(\text{SNSiMe}_3)]$ , from which  $[\text{Fe}_2(\text{CO})_6(\text{SNH})]$  and  $[\text{Fe}_2(\text{CO})_6(\text{SNMe})]$  were prepared. [906] The dithioformate esters  $\text{HCS}_2\text{R}$ , ( $\text{R} = \text{Me}, \text{Et}, \text{allyl}$ ) form  $[\text{Fe}_2(\text{CO})_6\text{L}]$  with  $[\text{Fe}_2(\text{CO})_9]$  with both S and CS bridging; a similar product is formed with  $\text{RCSOEt}$ , but  $\text{HCSOEt}$  gives  $[\text{Fe}_2(\text{CO})_6\text{L}_2]$  in which only S bridges. [907] Among the products of the reaction of  $[\text{Fe}_3(\text{CO})_{12}]$  with benzalazine [?] are  $[\text{Fe}_2(\text{CO})_6]$  and  $[\text{Fe}_3(\text{CO})_8]$  complexes of  $\text{C}_6\text{H}_4\text{CH}_2\text{NN:CHPh}$ . [908] The structure of  $[\text{Fe}_2(\text{CO})_6(\text{PR})_4]$  complexes are reported. [909] The reaction of 1,2,3-triphenyl-1,2,3-triposphaindane with  $\text{Fe}_2(\text{CO})_9$  gives many products including complexes with  $[\text{Fe}_2 \text{ or } 3(\text{CO})_{8,9,10}]$  units and an  $[\text{Fe}_2(\text{CO})_6]$  complex with 1,2-bisphenylphosphidobenzene. [910]  $(\text{PNCl}_2)_n$  react with  $[\text{Fe}_2(\text{CO})_8]^{2-}$  to give  $[\text{Fe}_2(\text{CO})_8(\text{P}_n\text{N}_n\text{Cl}_{2n-2})]$ , ( $n = 3, 4$ ), with P

bridging 2Fe or an  $[\text{Fe}_3(\text{CO})_{10}]$  complex with bridging by P and CO. [911]

The reaction of  $[\text{Fe}(\text{CO})_5]$  or  $[\text{Fe}(\text{CO})_4(\text{CS})]$  with  $\text{HgCl}_2$  in various solvents gives the binuclear complexes  $[\text{Fe}(\text{CO})_5 \text{HgCl}_2]$ ,  $[\text{Fe}(\text{CO})_4(\text{CS})\text{HgCl}_2]$  and  $[\text{Fe}(\text{CO})_4\{\text{C}(\text{S})\text{OEt}\}(\text{HgCl})]$  as well as trinuclear  $[\text{Fe}(\text{CO})_4(\text{HgCl})_2]$  and polymeric  $[\text{Fe}(\text{CO})_3(\text{CS})\text{Hg}]$ . [912] The versatile triphosphine  $\text{HC}(\text{PPh}_2)_3$ , (L), mentioned earlier [877] also forms  $[\text{Fe}(\text{CO})_5\text{RhL}]^+[\text{Rh}(\text{CO})_2\text{Cl}_2]^-$  in which 2P bond the Fe, one bonds Rh and there is an Fe-Rh bond. [913]

$[\text{Fe}_2(\text{CO})_9]$  has been used to prepare tetrahydrobenzotriazenes by reaction with adiponitriles and RCN. [914]

Trinuclear complexes make up a very big section.  $[\text{Fe}_3(\text{CO})_{12}]$  catalyses the reaction of perfluoroalkyl halides with allylsilanes to give polyfluoroalkenes. [915] The conditions for the conversion of  $[\text{Fe}(\text{CO})_4\text{H}]^-$  into  $[\text{Fe}_3(\text{CO})_{11}\text{H}]^-$  have been reinvestigated. [916] This anion adsorbed on NaY zeolite is a good catalyst for the water gas shift reaction, comparable to  $[\text{Fe}(\text{CO})_5]$  in the homogeneous mode. [917] Reactions of  $[\text{Fe}_3(\text{CO})_9(\text{OCMe})]^-$  have been reported with mechanistic discussion about C-O cleavage;  $\text{HBF}_4$  gives  $[\text{Fe}_3(\text{CO})_9\text{H}(\text{OCMe})]$ ,  $[\text{Fe}_3(\text{CO})_{10}\text{H}]$  and  $[\text{Fe}_3(\text{CO})_9(\text{HOCMe})]$ , but  $\text{MeSO}_2\text{F}$  gives  $[\text{Fe}_3(\text{CO})_9(\text{CMe})(\text{OMe})]$  and  $[\text{Fe}_3(\text{CO})_9(\text{CMe})(\text{OCMe})]$ . [918]

There has been some interesting work on trinuclear complexes with sulphur ligands.  $\text{Na}_2\text{S}$  and  $\text{Na}[\text{HFe}(\text{CO})_4]$  give  $[\text{Fe}_3(\text{CO})_9(\text{S})\text{H}_2]$ , which easily loses two protons to give  $[\text{Fe}_3(\text{CO})_9\text{S}]^{2-}$ , isolated as the  $\text{Et}_4\text{N}^+$  salt.

[919]  $[\text{Fe}_3(\text{CO})_{12}]$  reacts with  $(4\text{-MeOC}_6\text{H}_4\text{PS}_2)_2$  to give  $[\text{Fe}_3(\text{CO})_9\text{S}(\text{PC}_6\text{H}_4\text{OMe-4})]$ , with  $\mu_3\text{-S}$  and PR groups; [920] this and related compounds can also be prepared by the reaction of  $[\text{Fe}(\text{CO})_4]^{2-}$  with  $\text{RPSCl}_2$ , (R = Me, Ph, 4-MeOC<sub>6</sub>H<sub>4</sub>). [921] 3-Chloroperbenzoic acid oxidises  $[\text{Fe}_3(\text{CO})_9\text{S}_2]$  to  $[\text{Fe}_3(\text{CO})_9\text{S}(\text{SO})]$ . [922]

There are a number of phosphorus ligands which form  $\text{Fe}_3$  trinuclear cluster complexes.  $[\text{Fe}_3(\text{CO})_8(\text{PPh}_2)_2\text{H}_2]$  is prepared from



$[\text{Fe}_4(\text{CO})_{13}]^{2-}$ ,  $\text{CF}_3\text{CO}_2\text{H}$  and  $\text{Ph}_2\text{PH}$ ; it has an isosceles  $\text{Fe}_3$  triangular structure and reaction with CO converts it to  $[\text{Fe}_3(\text{CO})_9(\mu_3\text{-PPh})_2]$ . [923] Reactions of  $\text{RHPCH}_2\text{PRR}'$ , ( $\text{R}$  = alkyl;  $\text{R}'$  = R, H) with  $[\text{Fe}_2(\text{CO})_9]$  gave  $[\text{Fe}_3(\text{CO})_9]$  cluster compounds with or without P-C cleavage. [924]

The isocyanide  $\text{F}_3\text{CNC}$  can be stabilised by acting as a bridge in the complex  $[\text{Fe}_3(\text{CO})_{11}(\text{CNCF}_3)]$ ; [925] the  $t\text{BuNC}$  analogue rearranges to  $[\text{Fe}_3(\text{CO})_9(\text{CN}^t\text{Bu})]$ , which can be reduced by  $\text{KHB}^t\text{Pr}_3$  to  $[\text{Fe}_3(\text{CO})_9\text{H}(\text{CN}^t\text{Bu})]$  which is in turn converted by  $\text{H}_3\text{PO}_4$  into  $[\text{Fe}_3(\text{CO})_9\text{H}(\text{HC}(\text{CN}^t\text{Bu}))]$ . [926] The enthalpy change for the reaction of  $[\text{Fe}_3(\text{CO})_{10}\text{H}(\text{CMe})]$  with  $\text{H}_2$  to form CO and  $[\text{Fe}_3(\text{CO})_9\text{H}_3(\text{CMe})]$  was used to show that the  $\text{FeHFe}$  bond energy is greater than that for  $\text{Fe}(\text{CO})\text{Fe}$ . [927]  $[\text{Fe}_3(\text{CO})_9\text{H}(\text{BH}_4)]$  has been prepared from  $[\text{Fe}(\text{CO})_4(\text{COMe})]^-$  and  $[\text{BH}_3.\text{thf}]$ ; its structure was solved using the H positions of the isoelectronic  $[\text{Fe}_3(\text{CO})_9\text{H}_3(\text{CH})]$  and it was seen to have  $\text{BH}_4^-$  capping an isosceles  $\text{Fe}_3$  triangle, but the  $\text{BH}_4$  hydrogens can exchange with the hydride one at  $60^\circ\text{C}$ . [928]  $[\text{Fe}_3(\text{CO})_{10}\text{H}(\text{BH}_2)]$ , having a  $\text{BH}_2$  cap, is prepared from  $[\text{Fe}(\text{CO})_5]$ ,  $[\text{Fe}(\text{CO})_4(\text{COMe})]^-$  and  $[\text{BH}_3.\text{thf}]$  and deprotonates [929] to  $[\text{Fe}_3(\text{CO})_{10}(\text{BH}_2)]^-$ .

There are also a number of trinuclear heteropolymetallic clusters. The reaction of  $[\text{Fe}_3(\text{CO})_{12}]$  with  $t\text{BuSH}$  and  $[\text{MnL}(\text{CO})_3]$ , ( $\text{L}$  = cp, Mecp), gives  $[\text{Fe}_2(\text{CO})_6(\mu\text{-CO})_2(\mu_3\text{-S})(\text{MnL})]$ . [930] The reactions of  $[\text{Fe}_2\text{Ru}(\text{CO})_{12}]$  and  $[\text{FeRu}_2(\text{CO})_{12}]$  with  $\text{PPh}_3$  or  $\text{P}(\text{OMe})_3$  are compared; if one carbonyl is replaced the structure expands but for double reaction some CO bridges form, leading to the conclusion that the di-iron carbonyl has the  $[\text{Fe}_3(\text{CO})_{12}]$  structure but the diruthenium one does not have that of  $[\text{Ru}_3(\text{CO})_{12}]$ . [931] Isonitrile derivatives of  $[\text{FeCo}_2(\text{CO})_9\text{S}]$  have been prepared and their redox behaviour compared with many  $[\text{Fe}_n\text{Co}_{(3-n)}\text{E}]$  derivatives, ( $\text{E}$  = S, Se, PR, CR). [932] Other examples of this type are  $[\text{FeCo}_2(\text{CO})_9\text{PR}]$ , ( $\text{R}$  = Me,  $t\text{Bu}$ , Ph,  $\text{PMe}_2\text{Ph}$ ), the last of these giving chiral

or meso complexes when a CO or one or both Co are replaced respectively; this paper also describes the preparation of  $[\text{FeCoM}(\text{CO})_9\text{H}(\text{PR})]$ , ( $\text{M} = \text{Fe}, \text{Ru}$ ), which is also chiral and has a hydride bridge between Fe and M. [933] Also in this area there are descriptions of reactions of  $[\text{FeCo}(\text{CO})_6 \text{ S M}(\text{cp})(\text{CO})_2]$ , ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ) with  $\text{M}'\text{AsMe}_2$ , [ $\text{M} = \text{M}(\text{cp})(\text{CO})_3$ ,  $\text{Fe}(\text{cp})(\text{CO})_2$ ] leading to the preparation of  $[\text{Fe}(\text{CO})_3\text{S}\{\text{M}(\text{cp})(\text{CO})_2\}_2]$ , ( $\text{M} = \text{Mo}, \text{W}$ ); [934]

On to larger clusters. Low temperature NMR ( $^1\text{H}$ ,  $^{13}\text{C}$ ) of  $[\text{Fe}_4(\text{CO})_{13}\text{H}]^-$  shows the butterfly and tetrahedral structures are in equilibrium in  $\text{CD}_2\text{Cl}_2$ . [935] This area is a favourite with molecular orbitalers. One group sought to account for the difference between the structure of  $[\text{Fe}_4(\text{CO})_{13}\text{H}]^-$  and that of  $[\text{Fe}_4(\text{CO})_{12}\text{H}(\text{CH})]$ ; [936] another established the metal-domination of frontier orbitals in  $[\text{Fe}_4(\text{CO})_{13}\text{C}]$ ,  $[\text{Fe}_4(\text{CO})_{12}\text{C}]^{2-}$ ,  $[\text{Fe}_4(\text{CO})_{12}(\text{H})\text{C}]^-$  and  $[\text{Fe}_4(\text{CO})_{12}(\text{CCO}_2\text{Me})]^-$ ; [937] a third was interested in the unreactivity of the exposed C atoms in  $[\text{Fe}_4(\text{CO})_{12}\text{C}]^{2-}$  and  $[\text{Fe}_5(\text{CO})_{15}\text{C}]$ . [938]

The structure of  $[\text{Fe}_4(\text{CO})_{12}\text{S}_4]^{2-}$ , formed from  $[\text{Fe}_2(\text{CO})_6\text{S}_2]$  and  $\text{LiEt}_3\text{BH}$  in THF, consists of two  $[\text{Fe}_2(\text{CO})_6\text{S}_2]^-$  ions linked by a S-S bond 2.154 Å long. [939]

The tetrahedral cluster  $[\text{Fe}_3\text{M}(\text{CO})_{14}]^{2-}$ , ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ), prepared by the reaction of  $[\text{Fe}_3(\text{CO})_{11}]^{2-}$  with  $[\text{M}(\text{CO})_3(\text{MeCN})_3]$  can also be described as a hexacapped rectangular antiprism of carbonyls. [940] The adventitious preparation of  $[\text{FeRu}_3(\text{CO})_{13}\text{H}_2]$  allowed a more precise determination of the structure of the second form of this compound. [941] Use of  $^2\text{D}$  NMR led to finding the broad hydride resonance in  $[\text{FeCo}_3(\text{CO})_{12}\text{H}]$ . [942] Reacting  $[\text{Ni}(\text{CO})_4]$  with  $[\text{Fe}_3(\text{CO})_{11}]^{2-}$  gives  $[\text{Fe}_3\text{Ni}(\text{CO})_{12}]^{2-}$ , which can be protonated to  $[\text{Fe}_3\text{Ni}(\text{CO})_{12}\text{H}]^-$ , which can also be prepared from  $[\text{Fe}_3(\text{CO})_{11}]^{2-}$  and  $\text{NiCl}_2$  and whose X-ray structure shows an  $\text{Fe}_3\text{Ni}$  tetrahedron with 8 terminal and 4 edge-bridging CO and a hydrogen over an  $\text{Fe}_2\text{Ni}$  face. [943]  $[\text{Fe}_2(\text{CO})_9]$  and  $[\text{Mn}(\text{cp})(\text{CO})_2\text{PBr}_3]$  give the linear cluster  $[\{\text{Mn}(\text{cp})(\text{CO})_2\}_2\text{Fe}_2(\text{CO})_6]$ , with each P bridging 2Fe as well as

forming a double bond to Mn. [944] The most complicated piece of chemistry I have read this year describes reactions of CO with  $[MM^1M^2M^3(\mu_3-E)(\mu_2-AsMe_2)]$ , ( $M-M^3 = Fe, Ru, Co, Mo, W$ , all with various cp and CO ligands), to give tetranuclear complexes with loss of a Co-As fragment; among the species which are produced are  $[FeCo_2Mo(cp)(CO)_{10}S(AsMe_2)]$  and  $[FeCo_2Mo(cp)(CO)_8S(AsMe_2)]$  and they all revert to the starting materials in a vacuum. [945]  $[FeCoPdPt(CO)_6(NO)(dppm)_2]$  is prepared from  $[Fe(CO)_3(NO)]^-$  and  $[CoPdPt(CO)_3I(dppm)_2]$ . [946]  $[FeCo_3(CO)_{12}]^-$  and  $MLCl$ , ( $M = Cu, Ag$ ;  $L = PPh_3, PhPCH:CH:CH, MeCN$ ), react to give  $[FeCo_3(CO)_{12}ML]$ . [947] The seemingly innocuous combination of  $[Fe(CO)_5]$ ,  $[Mo(CO)_6]$  and Na/Hg in diglyme gives  $[Fe_4Mo_2(CO)_{18}C]^{2-}$  which is said to react with  $FeCl_3$  to give  $[Fe_4Mo_2(CO)_{19}C]$ . [948] Many Fe-Rh clusters are reported, e.g.  $[Fe_4Rh(CO)_{14}C]$  and  $[Fe_4Rh_2(CO)_{16}C]$ . [949] Moessbauer spectra of iron cluster compounds and mixed metal clusters allow correlations of Moessbauer parameters with whether the Fe atom has all its carbonyls terminal, some bridging or with H bridges. [950]

## 7.2. Nitrosyls

Preparations are reported for  $[Fe(NO)(PR_3)_4]^+$ , ( $R = OMe, OEt$ ), and  $[Fe(NO)(PR_3)_2(CNR')_2]$ , ( $R' = aryl$ ); all of these have trigonal bipyramidal geometry with equatorial linear NO. [951] The nitrosyls  $[Fe(NO)_2(PPh_3)_2]$ ,  $[Fe(NO)_2(CO)(PPh_3)]$  and  $[Fe(NO)(CO)(PPh_3)L]$ , ( $L = 4-Ac-C_6H_4N_2$ ), are prepared [952] by the action of  $NO_2^-$  on  $[Fe(NO)(CO)_2(PPh_3)_2]^+$  and  $[Fe(CO)_2(PPh_3)_2L]^+$ . In an elaborate series of reactions  $[Fe_2(NO)_4(PPh_2)_2]$  is reduced by Na,  $LiAlH_4$  or  $Na[AlH_2(OCH_2OMe)_2]$  to  $[Fe_2(NO)_4(PPh_2)_2]^{2-}$  with bridging  $PPh_2^-$  ligands. This is reduced by  $LiBHET_3$  to  $[Fe_2(NO)_4(PPh_2)(PPh_2H)]^-$ , with bridging by NO and  $PPh_2^-$ . Reaction of this with BuLi gives the dianion again, but with mixed bridges; the other isomer forms on storing at  $-70^\circ C$ , but reaction with  $CH_2I_2$

gives an isomerising mixture of  $[\text{Fe}_2(\text{NO})_4(\text{CH}_2)(\text{Ph}_2\text{PPPh}_2)]$  and  $[\text{Fe}_2(\text{NO})_4(\text{CH}_2\text{PPh}_2)]$ . [953]  $[\text{Fe}(\text{cp})(\text{NO})]_2$  is reduced by Na to the dinuclear monoanion;  $\text{CH}_3\text{I}$  converts this to  $[\text{Fe}(\text{cp})(\text{NO})\text{Me}]_2^-$ , which is in rapid equilibrium with the monomer, and thermally unstable  $[\text{Fe}(\text{cp})(\text{NO})(\text{Me})_2]$ , which reacts with  $\text{PMe}_3$  to form  $[\text{Fe}(\text{cp})(\text{PMe}_3)(\text{NOMe})\text{Me}]$  wherein bonding of NOME is through N. [954]

### 2.3. Alkyls and Aryls

This section comprises compounds whose behaviour centres on the formation, reactivity or theory of a sigma bond between Fe and C, but this classification has been interpreted generously. We begin with alkyls and acyls. Ab initio SCF calculations on CO insertion into Fe-C or Fe-H bonds in  $[\text{Fe}(\text{CO})_4(\text{CH}_3)\text{H}]$  indicate that the hydrido-acetyl complex is more stable than the methyl-formyl one. [955] The  $[\text{FeMe}]^+$  ion cleaves  $\text{C}_3\text{H}_8$  to form  $[\text{FeEt}]^+$  in equilibrium with  $[\text{Fe}(\text{C}_2\text{H}_5)\text{H}]^+$ , but a stable  $\text{C}_2\text{H}_4$  complex is formed from  $\text{C}_4\text{H}_8$ . [956] Mass spectra have been used to study the gas phase reactions of  $\text{Fe}^+$  with cyclic ketones to give metallacycles and their subsequent decomposition. [957] Ion-molecule reactions of  $\text{Fe}(\text{CO})_n^+$  with hydrocarbons and ketones gave a range of products, including  $[\text{FeC}_2\text{H}_4]^+$ ,  $[\text{Fe}(\text{CO})\text{H}_2]^+$ ,  $[\text{Fe}(\text{CO})\text{C}_2\text{H}_6]^+$ ,  $[\text{FeC}_3\text{H}_8]^+$  and four varieties of  $[\text{FeC}_4\text{H}_8]^{+}$ . [958] A mass spectrometric study of various  $[\text{FeC}_n\text{H}_{2n}]^+$  found that for  $n > 4$  the hydrocarbon broke down; mechanisms are discussed. [959] Sparking halomethanes on an Fe surface gave  $[\text{FeCXY}]^+$ ,  $[\text{FeCX}]^+$  and  $[\text{FeX}_2]^+$ , (X,Y = H,Cl,Br). [960]

Turning to conventional preparative chemistry,  $[\text{FeBr}(\text{mesityl})(\text{thf})_{0.5}]$  has been prepared from  $\text{FeBr}_2$  and  $[\text{Mg}(\text{mesityl})_2(\text{thf})_2]$ . [961] ESR spectra of  $^{13}\text{C}$ -labelled  $[\text{Fe}(\text{CO})_4(\text{RCO})]$ , (R = long chain alkyl) shows these have axial acyl groups with an unpaired electron in the equatorial plane; this result is said to explain the radicals' failure to dimerise. [962]

$[\text{Fe}(\text{CO})_4(\text{CH}_2\text{CO}_2\text{Me})]^-$  is formed from  $[\text{Fe}_2(\text{CO})_8(\text{CH}_2)]$  by reaction with NaOMe in dmsO; other reactions of the dinuclear precursor are reported.

[963] In the presence of various ligands L,  $[\text{Fe}(\text{CO})_3(\text{PMe}_3)_2\text{Me}]^+$  and  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)(\text{Me})\text{X}]$ , ( $\text{X} = \text{I}, \text{CN}$ ) are converted to  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2(\text{OAc})\text{L}]$ ,  $[\text{Fe}(\text{CO})(\text{PMe}_3)_2(\text{OAc})\text{LX}]$  and  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)(\text{Me})\text{L}]^+$ . [964]  $[\text{Fe}(\text{CO})_4(\text{C}_3\text{F}_7)\text{I}]$  reacts with  $\text{SPHET}_2$  to give  $[\text{Fe}(\text{CO})_3(\text{C}_3\text{F}_7)(\text{SPHET}_2)]$ ; reaction of this with  $\text{NH}_3$  gives a cyclic FePS compound which inserts into alkynes to give a  $\text{C}_2\text{PSFe}$  ring which reacts with more alkyne to give a bicyclic structure. [965] At  $-30^\circ\text{C}$ ,  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2(\text{Me})\text{X}]$ , ( $\text{X} = \text{I}, \text{CN}, \text{Me}, \text{CNBPh}_3$ ) react with  $\text{CO}$  to give unstable  $[\text{Fe}(\text{CO})(\text{PMe}_3)_2(\text{Me})(\text{COME})\text{X}]$  which revert to the starting material even under argon. [966] The Fe-C bond in  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2\text{MeI}]$  is subject to insertion with  $\text{C}_6\text{H}_{11}\text{NC}$ ; subsequent oxidative addition and rearrangement yield  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2\{\text{C}(\text{:CH}_2)\text{N}(\text{C}_6\text{H}_{11})\text{C}(\text{:N}^+\text{HC}_6\text{H}_{11})\}]^+$ . [967] The conversion of acetyl to methyl was observed in the FAB mass spectra of  $[\text{Fe}(\text{CO})_2(\text{diars})(\text{COME})\text{L}]^+$ , ( $\text{L} = \text{monodentate ligand}$ ). [968] The photodecarbonylation in a PVC matrix at 12-200K of  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{COME})]$  is thermally reversible; the photolysis of  $[\text{Fe}(\text{cp})(\text{CO})_2\text{Me}]$  gives reversible loss of one CO. [969] The latter compound reacts with  $[\text{Mo}(\text{cp})(\text{CO})_3]^+$  to give the acetyl-bridging  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{COME})\text{Mo}(\text{cp})(\text{CO})_3]^+$ . [970] Carbonyl insertion into  $[\text{Fe}(\text{cp})(\text{CO})_2\text{CH}_2\text{CH}_2\text{R}]$  is retarded by adding  $\text{Ph}_3\text{C}$ , which then leads to elimination. [971] The acetoxy compound  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CO}_2\text{Me})]$ , which can be converted to  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CH}_2\text{OMe})]$ , is prepared from  $[\text{Fe}(\text{cp})(\text{CO})_2]_2\text{Mg}$  by the reaction of  $\text{CO}_2$  to give  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CO}_2)]_2\text{Mg}$  and the reaction of that with  $\text{CF}_3\text{SO}_3\text{Me}$ . [972]  $[\text{Fe}(\text{cp})(\text{CO})_2\text{C}(\text{S})\text{SR}]$ , [ $\text{R} = \text{Fe}(\text{cp})(\text{CO})_2, \text{Re}(\text{CO})_5$ ] react with  $[\text{M}(\text{CO})_5(\text{thf})]$ , ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ) or  $[\text{Mn}(\text{cp})(\text{CO})_2(\text{thf})]$  with the thf being replaced by the uncoordinated S; the R component can be lost to give a  $\text{CS}_2$  moiety bidentate to Cr, Mo, W or Mn. [973] Another mixed metal complex,  $[\text{Fe}(\text{cp})(\text{CO})(\mu\text{-PPh}_2)(\mu\text{-COME})\text{Mn}(\text{CO})_4]$ , is prepared from  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{PPh}_2)]$  and  $[\text{Mn}(\text{CO})_5\text{Me}]$ , but  $[\text{Mn}(\text{CO})_5\text{H}]$  gives the analogue with bridging hydride, not formyl. [974] Condensation of imines with

$[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)(\text{CO}_2\text{AlEt}_2)]$  gives  $[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)\text{COCH}_2\text{CH}(\text{Ph})\text{NHPh}]$  stereoselectively; iodine in  $\text{CH}_2\text{Cl}_2$  converts these to  $\beta$ -lactams. [975]

$\text{O}^- - \text{C}_6\text{H}_4(\text{CH}_2\text{Br})_2$  converts  $[\text{Fe}(\text{CO})_4]^{2-}$  to

$[\text{Fe}(\text{CO})_3(\text{CH}_2\text{C}_6\text{H}_4\text{CH}_2\text{Br})_2]$ ; if  $\text{PPh}_3$  is present

$[\text{Fe}(\text{CO})_2(\text{PPh}_3)(\text{COCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{Br})(\text{CH}_2\text{C}_6\text{H}_4\text{CH}_2\text{Br})]$  is formed.

[976] In another reaction involving dibromoxylene  $[\text{Fe}_2(\text{CO})_6(\text{PPh}_2)_2]$

is converted to  $[\text{Fe}_2(\text{CO})_6\{\text{PC}_6\text{H}_4(\text{CH}_2\text{PPh})_2\}]$ ; LiBu converts this to

a complex in which one Fe is bonded to a xylyl carbon next to an alkylated P.

[977] The porphyrin complexes  $[\text{FeLR}(\text{NO})]$ , ( $\text{L} = \text{OEP}, \text{TPP}$ ;  $\text{R} = \text{Me}$ , aryl) have been prepared. [978]

Aryls are outnumbered by alkyls, but there is some interesting work.

Interactions between the 2-hydrogen and Fe in  $[\text{Fe}(\text{cp})(\text{Ar})(\text{CO})\text{L}]$  complexes,

( $\text{L} = \text{CO}, \text{PPh}_3$ ) have been detected by IR,  $^1\text{H}$  and  $^{13}\text{C}$  NMR and Moessbauer

spectra. [979] The mass spectra of  $[\text{Fe}_2(\text{CO})_6(\text{C}_6\text{H}_4\text{CH}_2\text{NPh})]$  and

similar complexes are reported. [980] A similar complex to this, but with  $^i\text{Pr}$

instead of Ph on N is prepared from  $[\text{Fe}_2(\text{CO})_9]$  and  $[(\text{CO})_5\text{W}:\text{C}(\text{Ph})(\text{NH}^i\text{Pr})]$

using  $[(\text{CO})_5\text{M}:\text{C}(\text{COR})(\text{NHC}_6\text{H}_{11})]$ , ( $\text{M} = \text{Cr}, \text{W}$ ;  $\text{R} = \text{Me}, \text{Ph}$ ) gives

a complex with  $2\text{Fe}(\text{CO})_3$  moieties bridged by  $(\text{RCO})$  and  $(\text{NHC}_6\text{H}_{11})$  groups.

[981]  $[\text{Me}_2\text{Si}\{(\text{C}_5\text{H}_4)\text{Fe}(\text{CO})_2\}_2]$  reacts with  $\text{Ph}_n\text{P}(\text{CH}_2)_n\text{PPh}_2(\text{L})$

to give either  $[\text{Me}_2\text{Si}\{(\text{C}_5\text{H}_4)\text{Fe}(\text{CO})\}_2\text{L}]$  or

$[\text{Me}_2\text{Si}\{(\text{C}_5\text{H}_4)_2\text{Fe}_2(\text{CO})_3\}_2\text{L}]$ ; if  $n$  is small the dimer (CV study)

is preferred but if  $n$  is large or more reagent is used the tetramer is

favoured. [982]

Four papers describe metallocyclic compounds. A ferracyclopentane is prepared

by reacting  $[\text{Fe}(\text{CO})_3\text{L}]^{2-}$ , ( $\text{L} = \text{CO}, \text{PPh}_3$ ) with

$\text{F}_3\text{CSO}_3(\text{CH}_2)_4\text{O}_3\text{SCF}_3$ ; using the dimethylene reagent gave

$[\text{Fe}(\text{CO})_3\text{L}(\text{C}_2\text{H}_4)]$ . [983] Aziridine, oxirane and thiirane react with

$[\text{Fe}(\text{cp})(\text{CO})_2\text{L}]^+$ , ( $\text{L} = \text{CO}, \text{CS}, \text{PPh}_3$ ) to give five-membered cyclic carbene

complexes. [984] Fe atoms react with  $\text{B}_5\text{H}_9$ , toluene and 2-butyne to give

$[\text{Fe}(\text{C}_6\text{Me}_6)(2,3\text{-Me}_2\text{C}_2\text{B}_4\text{H}_4)]$  and three similar products; the

structures of all were described. [985] The reaction of  $[\text{Fe}_2(\text{CO})_9]$  with  $\text{ROC(S)SMe}$ , ( $\text{R}$  = adamantyl) gives  $[\text{Fe}_2(\text{CO})_6 \text{ ROC(S)SMe}]$ , which contains an  $\text{FeSCS}$  ring with the other  $\text{Fe}$  bonded to  $\text{C}$ , the thione  $\text{S}$  and  $\text{Fe}$ . [986]

The next section is about alkylidene and related compounds, taken in order of progressively fewer carbonyl groups and increasing numbers of  $\text{Fe}$  atoms.

$[\text{Fe}(\text{cp})(\text{CO})_2(\text{CH}_2\text{CH:CHOMe})]$  is prepared from  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CHOMe})]^+$  via the intermediate  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CROMe})]^+$ , ( $\text{R}$  = alkyl, aryl). [987]

Rearrangement of  $\text{N}(\text{C}_2\text{H}_4\text{O})_2\text{PCH}_2\text{CH:CH}_2$  in an  $[\text{Fe}(\text{cp})\text{CO}]$  complex gave a methylvinyl complex with a  $\text{P}$ -bonded residue. [988] Alkenyl complexes such as  $[\text{Fe}(\text{cp})(\text{CO})\{\text{P}(\text{OPh})_3\}(\text{CMe:CMePh})]$  can be converted into acylalkenyl ones by reaction with  $\text{CO}$  catalysed by  $\text{Ag}^+$ ,  $\text{Ce}^{4+}$  or ferricinium ions. [989]

$[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)(\text{C:CMe}_2)]^+$  reacts with  $\text{Li}_2\text{Cu}(\text{CN})\text{R}_2$ , ( $\text{R}$  =  $\text{Ph}$ ,  $\text{CH:CH}_2$ ) to give  $[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)(\text{CR:CMe}_2)]$  and with  $\text{PhS}^-$  to give a complex with  $\text{R} = \text{SPh}$ ; the vinyl derivative rearranges to an  $\eta^3 - \text{CH}_2\text{CHC:CMe}_2$  complex. [990] Complicated bridged complexes are formed when alkynes react with  $[\text{Fe}_2(\text{CO})_7(\text{CPh:CHPh})]$ . [991] The synthesis of  $[\text{Fe}(\text{TPP})\text{C:CR}_2]$ , ( $\text{R} = 4\text{-ClC}_6\text{H}_4$ ) has been achieved starting from  $[\text{Fe}_3(\text{CO})_{12}]$  [992].

Ferracarbene complexes, containing  $(\text{Fe:C})$  moieties, will be discussed in a similar order to the alkylidene complexes.  $[\text{Fe}(\text{CO})_4(\text{CROEt})]$ ,  $\text{R} = \text{Bu}$ ,  $\text{Ph}$  react with alkynes and  $\text{CO}$  to give pyrone complexes of  $[\text{Fe}(\text{CO})_3]$  which rearrange to isomers through a ferracyclobutene intermediate which was isolated. [993] The action of  $\text{LiN}^i\text{Pr}_2$  followed by  $\text{Et}_3\text{O}^+$  on  $[\text{Fe}(\text{CO})_5]$  produced  $[\text{Fe}(\text{CO})_4\{\text{C}(\text{OEt})\text{N}^i\text{Pr}_2\}]$  and  $[\text{Fe}(\text{CO})_3\{\text{C}(\text{N}^i\text{Pr}_2)\text{OC}(\text{O})\text{Et}\}]$ , in which the ketonic  $\text{O}$  atom is also coordinated. [994] The complex  $[(\text{cp})(\text{CO})_2\text{Fe}(\text{CH:CH:CMe}_2)]^+$  can be used to form cyclopropanes. [995] The preparations and reactions of  $[\text{Fe}(\text{cp})(\text{CO})\text{L Fe}(\text{CH:CH:CRCH}_2\text{R}')]^+$ , [ $\text{L} = \text{CO}$ ,  $\text{P}(\text{OMe})_3$ ;  $\text{R} = \text{H}$ ,  $\text{R}' = \text{Me}$ ;  $\text{R} = \text{Me}$ ,  $\text{R}' = \text{H}$ ] are reported. [996] The alkenyl complex  $[\text{Fe}(\text{cp})(\text{CO})(\text{PMe}_3)\{\text{C}(\text{OMe}):CH_2\}]$  has been converted to

various carbene complexes  $[\text{Fe}(\text{cp})(\text{CO})(\text{PMe}_3)\{\text{:C}(\text{OMe})\text{R}\}]$ ,

(R = Me,  $\text{CH}_2\text{CS}_2$ ,  $\text{CH}_2\text{C}(\text{S})\text{SMe}$ ). [997] The reaction of

$[\text{Fe}_2(\text{CO})_6(\mu\text{-SMe})(\mu\text{-SC}(\text{OR}))]$  with  $\text{P}(\text{OMe})_3$  shows a CO on the carbene-bonded Fe is the most labile; [998] there are X-ray and  $^{13}\text{C}$  NMR

studies on the mono- $\text{P}(\text{OMe})_3$  complexes for the R = mesityl and

adamantylmethyl. [999] The cyclopropylidene bridging complex

$[\text{Fe}_2(\text{cp})_2(\text{CO})_5(\text{C}_3\text{H}_4)]$  is prepared by the reaction of  $\text{H}_2\text{CN}_2$  with a bridging vinylidene complex. [1000]

$[\text{Fe}(\text{cp})(\text{CO})(\text{MeCN})\text{C}(\text{SMe})_2]^+$  reacts with  $[\text{Co}(\text{CO})_4]^-$  to form

$[\text{Fe}(\text{cp})(\text{CO})(\mu\text{-CO})\mu\text{-C}(\text{SMe})_2\text{Co}(\text{CO})_2]$  which has an S-Co bond. [1001]

The simplest carbene complex, with  $\text{CH}_2$ , is given when  $[\text{Fe}(\text{cp})(\text{CO})_2]^-$

reacts with  $\text{Me}_3\text{CCO}_2\text{CH}_2\text{Cl}$  to form  $[\text{Fe}_2(\text{cp})_2(\mu\text{-CO})(\mu\text{-CH}_2)]$  as the

cis-isomer, but  $\text{C}_5\text{Me}_5$  gives the trans-isomer. [1002] A  $^{31}\text{P}$  NMR study of

the rotation barriers in  $[\text{Fe}(\text{cp})(\text{diphosphine})(\text{vinylidene})]$  complexes showed

steric effects were important from both non-cp ligands. [1003] The porphyrin

carbene complexes  $[\text{Fe}(\text{por})\text{CRR}']$ , (por = TPP, TTP; R = Me,  $i\text{Pr}$ ; R' = MeO, EtO,

$\text{PhCH}_2\text{S}$  can be made from  $[\text{Fe}(\text{por})\text{Cl}]$  and  $\text{Cl}_3\text{CR}$ , which gives a  $(\text{CClR})$

chlorocarbene complex, which is then treated with R'H. [1004]

The ketenylidene complex  $[\text{Fe}_3(\text{CO})_9(\text{CCO})]^{2-}$  reacts with

$[\text{M}(\text{CO})_3(\text{NCMe})_3]$ , (M = Cr, W) to give carbide complexes

$[\text{Fe}_3\text{MC}(\text{CO})_{13}]^{2-}$  which are similar to  $[\text{Fe}_4\text{C}(\text{CO})_{12}]^{2-}$ , but

$[\text{Co}_2(\text{CO})_8]$  gives  $[\text{Fe}_2\text{Co}(\text{CO})_9(\text{CCO})]^-$  in which the CCO is closer to

perpendicular than in the  $\text{Fe}_3$  analogue. [1005] Successive reaction of the

carbene complex  $[\text{Fe}(\text{CO})_4\{\text{C}(\text{OEt})\text{N}^i\text{Pr}_2\}]$  with  $\text{PPh}_3$  and  $\text{BCl}_3$  gives the

carbyne complex  $[\text{Fe}(\text{CO})_3(\text{PPh}_3)\{\text{CN}^i\text{Pr}_2\}]$  in which the carbyne is

equatorial, the phosphine axial and with the shortest Fe-carbyne bond of 1.734

Å. [1006]

$[\text{Fe}_4(\text{CO})_{12}\text{H}(\text{CH})]$  in  $\text{CF}_3\text{SO}_3\text{H}$  gives  $\text{CH}_4$  only after a long induction

period, but this reaction is accelerated by adding [1007a] such "reducing

agents" as  $\text{TiCl}_3$ ,  $\text{MoBr}_2$ , Fe or  $[\text{Fe}_2(\text{CO})_8]^{2-}$ . The use of deuterium



substitution in the reaction of alkenes with  $[\text{Fe}_2(\text{cp})_2(\text{CO})_3(\text{CD})]^+$  allowed the mechanism to be established. [1007b]

#### 7.4. Alkene Complexes

This section describes complexes where the major point of interest lies in the behaviour of a C=C bond linked to an Fe atom. Allyl complexes are included and the ordering of topics is in decreasing numbers of carbonyl ligands and increasing nuclearity.

PRDDO molecular orbital calculations on  $[\text{Fe}(\text{CO})_4(\text{C}_2\text{R}_4)]$ , ( $\text{R} = \text{H}, \text{F}, \text{Cl}, \text{CN}$ ) show the alkene should bond in the equatorial position for each one. [1008] NMR spectra of the vinyl protons in  $[\text{Fe}(\text{CO})_4]$  complexes of chiral alkenes were used to analyse their stereochemistry. [1009] Bromostyrene complexes were formed by reaction with  $\text{Na}_2[\text{Fe}(\text{CO})_4]$ . [1010]  $[\text{Fe}(\text{CO})_4(\text{SiMe}_3)\text{R}]$ , produced by the alkylation of  $[\text{Fe}(\text{CO})_4(\text{SiMe}_3)]^-$ , rearranges to give a siloxycarbene complex and  $[\text{Fe}(\text{CO})_4(\text{RCH}:\text{CHOSiMe}_3)]$ . [1011] The hexaalkene hemicene forms  $[\text{Fe}(\text{CO})_4]$  complexes with up to 3 of these units on each alkene molecule. [1012] The tautomerism of  $[\text{Fe}(\text{CO})_4 \text{ Ph.CH}:\text{CHC}(\text{Me}):\text{NR}]$  and its  $\alpha$ -allyl- $\sigma$ -carbamoyl  $[\text{Fe}(\text{CO})_3]$  analogue has been studied. [1013] The complex  $[\text{Fe}(\text{CO})_3\text{L}_2]$ , ( $\text{L} = \text{cis-cyclooctene}$ ) is recommended as a source of  $[\text{Fe}(\text{CO})_3]$ . [1014] Reactions of exo- and endo- complexes of  $[\text{Fe}(\text{CO})_3]$  with 5,6-dimethylidenebicyclo[2.2.2]oct-2-ene leading to hydroxy and keto ligands are described and lead to the revision of earlier configurational assignments. [1015] Various allyl-alkyl complexes of  $[\text{Fe}(\text{CO})_3]$  have been used in lactam synthesis. [1016] The reaction of  $[\text{FeH}(\text{CO})_4]^-$  with the diazabutadiene ( $\text{L}$ ) complexes  $[\text{M}(\text{CO})_3\text{BrL}]$ , ( $\text{M} = \text{Mn}, \text{Re}$ ) gave  $[\text{Fe}(\text{CO})_3]$  complexes with Fe-M and Fe-allyl bonds. [1017]

Use of the chiral phosphines  $\text{Ph}_2\text{PNRCHMePh}, (\text{L})$ , gave chiral complexes  $[\text{Fe}(\text{CO})_2(\text{NO})(\text{allyl})\text{L}]$  for allyl or 2-methylallyl, [1018] but alkyl phosphines  $\text{PR}_3$  gave reaction at the allyl ligand to form  $[\text{Fe}(\text{CO})_2(\text{NO})(\eta^2\text{-CH}_2:\text{CH}.\text{CH}_2\text{PR}_3)]$ . [1019] A bicyclo [3.2.1] octene complex of  $[\text{Fe}(\text{cp})(\text{CO})_2]^+$  was formed by a new ring expansion reaction of a norbornyl

complex. [1020] A number of complexes of the type  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{ROCH:CHOR})]^+$  are reported. [1021] The reaction of  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2\text{MeI}]$  with  $^t\text{BuNC}$  gives  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2(\text{MeC:N}^t\text{Bu})]^+$ , in which the  $\text{C}\equiv\text{N}$  bond coordinates like an alkene. [1022]

An improved preparation of allyltricarbonyliron lactone complexes involves the reaction of  $[\text{Fe}_2(\text{CO})_9]$  with alkenyl epoxides. [1023] The methylidyne-bridged complex  $[\text{Fe}_2(\text{cp})_2(\text{CO})_3\text{CH}]^+$  reacts with alkenes to give alkylidyne and  $\sigma, \pi$ -bridging vinyl complexes. [1024a] Benzylideneacetone,  $\text{PhCH:CH.CMe:O}$ , acts as an allyl ligand in various CO and  $\text{PR}_3$  complexes. [1024b]

$[\text{FeCl}_2(\text{PMe}_3)_2]$  and potassium pentadienide give  $[\text{Fe}(\text{PMe}_3)_2(\text{pentadienyl})_2]$ . [1025]

### 7.5. Alkyne Complexes

Alkynes react with  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2(\text{MeSCHS})]$  to give as final product the complex  $[\text{Fe}(\text{CO})_2(\text{PMe}_3)_2(\text{MeSCHCOCR:CRS})]$ , ( $\text{R} = \text{e.g. MeO}_2\text{C}$ ); an alkyne complex might be an intermediate. [1026] Alkyne (L) complexes  $[\text{Fe}(\text{cp})(\text{CO})\{\text{P}(\text{OPh})_3\}\text{L}]^+$  react with nucleophiles like  $\text{CH}_2\text{:CHCu}(\text{CN})\text{Li}_2$  to give alkenyl or alkynyl complexes. [1027] With most alkynes  $\text{RC:CR}$ , ( $\text{R} = \text{Me, Ph, CF}_3, p\text{-MeC}_6\text{H}_4$ ),  $[\text{Fe}_2\text{W}(\text{cp})(\text{CO})_9(p\text{-MeC}_6\text{H}_4)]$  gives  $[\text{FeW}(\text{cp})(\text{CO})_5\{\{p\text{-MeC}_6\text{H}_4\}\text{CCR:CR}\}]$ , but  $\text{RC:CSiMe}_3$ , ( $\text{R} = ^t\text{Bu, Me}_3\text{Si}$ ) gives  $[\text{Fe}_2\text{W}(\text{cp})(\text{CO})_7(p\text{-MeC}_6\text{H}_4)(\text{RCCSiMe}_3)]$ . [1028] The alkynyl complex  $[\text{Fe}_3(\text{CO})_9(\text{CCMe})]$ , in which the organo ligand is  $\sigma$ -bonded to one Fe and  $\pi$ -bonded to the others, can be prepared either by the reaction of methanol with electrochemically generated  $[\text{Fe}_3(\text{CO})_{10}(\text{CMe})]^{2-}$  or by the reaction of  $[\text{Fe}_3(\text{CO})_9(\text{CMe})(\text{COEt})]$  with  $[\text{Mn}(\text{CO})_5]^-$ . [1029] The three-way reaction of  $[\text{Fe}_3(\text{CO})_{12}]$ ,  $[\text{Ru}_3(\text{CO})_{12}]$  and  $\text{Ph:CPh}$  gives  $[\text{Fe}_3(\text{CO})_9(\text{PhC:CPh})]$  and ruthenacyclopentane complexes containing 1 or 2 Fe atoms; reaction of  $\text{Ru}_3(\text{CO})_{12}$  with  $[\text{Fe}_3(\text{CO})_9(\text{RCCR})]$ , ( $\text{R} = \text{Et, Ph}$ ) gives replacement of an Fe by Ru but CNDO calculations rationalise the non-formation of a diruthenium complex by maintaining that it is necessary to

maintain alkyne to  $\text{Fe}_2$  back donation. [1030] Benzonitrile is found to show both  $\sigma$  and  $\pi$  coordination in both  $[\text{Fe}_3(\text{CO})_9(\text{NCPH})]$  and  $[\text{Fe}_4(\text{CO})_{12}(\text{NCPH})]$ , which are prepared when this ligand and hydrogen react with  $[\text{Fe}_3(\text{CO})_{12}]$ . [1031] A complex formulated as  $[\text{Fe}_3(\text{CO})_9(\text{MeC}:\text{CO})]^{2-}$  is prepared by the electrochemical reduction of  $[\text{Fe}_3(\text{CO})_{10}(\text{CMe})]^-$ . [1032]

#### 7.6. Diene and Polyene Complexes

This section considers complexes of ligands containing more than one C=C bond. Monomeric complexes are treated first and the order is of increasing complexity of the diene after any general papers. Cyclopentadienyl complexes make up section 7.7. Allyls have been included, possibly unwisely, in section 7.4.

A number of general papers concern various  $[\text{Fe}(\text{CO})_3(\text{diene})]$  complexes. A short review discusses their use in natural product synthesis. [1033] They have been used for the regio- and stereospecific synthesis of olefinic triphenylphosphonium salts through reactions with  $\text{PPh}_3$ . [1034] There has been a CD study of complexes  $[\text{Fe}(\text{CO})_3]$  with chiral dienes. [1035] The natural chiral diene aminoacid gabaculine has been synthesised through these intermediates. [1036] Iron carbonyls react with chiral dienones to give chiral complexes which preserve their chirality when reacting with prochiral dienes to give  $[\text{Fe}(\text{CO})_3(\text{diene})]$ . [1037] Cyclopentanones are formed by reacting nucleophiles like  $\text{LiCMe}_2\text{CN}$  with these complexes. [1038]

The X-ray crystal structure of the butadiene complex  $[\text{Fe}(\text{PMe}_3)(\text{diene})_2]$  is reported. [1039] Butadiene also forms  $[\text{Fe}(\text{PPhMe}_2)_3(\text{diene})]$ , by reaction with  $[\text{Fe}(\text{PPhMe}_2)_3(\text{C}_2\text{H}_4)_2]$ , which is in turn prepared from  $[\text{Fe}(\text{acac})_3]$ ,  $\text{Et}_2\text{Al}(\text{OEt})$  and the phosphine. [1040] Various butadienes containing an  $\text{SiMe}_3$  group form  $[\text{Fe}(\text{CO})_3(\text{diene})]$  complexes on reaction with iron carbonyls. [1041]  $[\text{Fe}_2(\text{CO})_9]$  and *cis*- $[\text{PtCl}_2(\text{Ph}_2\text{P}:\text{C}^t\text{Bu})_2]$  react to give an  $[\text{Fe}(\text{CO})_3]$  complex of cyclobutadiene, but other Pt complexes produced cyclopentadienone complexes. [1042] Various reactions of  $\text{C}_4\text{Ph}_4$

complexes are reported:  $[\text{Fe}(\text{CO})(\text{NO})(\text{PPh}_3)(\text{diene})]^+$  and

$(\text{Ph}_3\text{P})_2\text{N}^+\cdot\text{Cl}^-$  give  $[\text{Fe}(\text{NO})(\text{PPh}_3)\text{Cl}(\text{diene})]$ ;

$[\text{Fe}(\text{CO})_2(\text{NO})(\text{diene})]^+$  and  $(\text{Ph}_3\text{P})_2\text{N}^+\cdot\text{X}^-$  give  $[\text{Fe}(\text{NO})\text{X}(\text{diene})]^2$ ,

( $\text{X} = \text{Cl}, \text{NCO}, \text{NCS}, \text{NO}_2, \text{N}_3$ ); the dicarbonyl cation gives

$[\text{Fe}(\text{NO})(\text{X})(\text{diene})]$  monomer with  $\text{X} = \text{NO}_3, \text{Et}_2\text{dte}$  or  $\text{acac}$ , the last being prepared with  $\text{Tlacac}$ . [1043]

Reactions of  $[\text{Fe}(\text{CO})_3]$  complexes of  $i\text{-PrN:CH:CH:N}^i\text{-Pr}$  and other diazadienes with dimethylacetylenedicarboxylate are reported with crystallographic confirmation of the extensive ligand rearrangement. [1044]

The Moessbauer spectrum of bis(pentadienyl) iron (II) complexes show more electron density on the Fe nucleus than for ferrocene. [1045] The reaction of nucleophiles like  $\text{LiCHPh}_2$  with such diene complexes as  $[\text{Fe}(\text{CO})_3(\text{isoprene})]$  gives allyls at high temperatures and homoallyls at low temperatures; a mechanistic discussion is given. [1046] 1,1-Dimethylsilole,

$\text{Me}_2\text{Si:CH:CH}_2\text{:CH:CH}_2$ , reacts with  $\text{Fe}_2(\text{CO})_9$  to give

$[\text{Fe}(\text{CO})_3(\text{diene})]$ ; one CO ligand can be replaced by  $\text{PPh}_3$ . [1047] Ab initio

molecular orbital calculations on  $[\text{Fe}(\text{CO})_3(\text{diene})]$  complexes show cyclopentadienone coordinates more strongly than cyclobutadiene. [1048]

The stereochemistry of reactions of substituted cyclohexadienes when coordinated to  $[\text{Fe}(\text{CO})_3]$  is discussed; [1049] there is similar work on these and cycloheptatrienyl complexes. [1050] The cyclohexadiene complex reacts with  $\text{LiCMe}_2\text{CN}$  to give an allyl complex but if CO is present an acyl-alkene complex is given. [1051] Rotational barriers in cyclohexadiene complexes have been assessed by  $^{13}\text{C}$  NMR, with methyl being considered more bulky than methoxy. [1052]

$[\text{Fe}(\text{CO})_2(\text{PR}_3)(\eta^4\text{-PhCH:CHCOR}')] complexes have the phosphine in axial or equatorial positions depending on the steric properties of it and the enone; [1053] there has been a kinetic study of the exchange of cyclohexadiene with the enone in these complexes. [1054] New$

$[\text{Fe}(\text{CO})_3(\text{diene})]$  complexes using 3-methoxy-6,6-disubstituted cyclohexa-1, 3-dienes is reported. [1055] The acetylation of the

cyclohexadiene complex  $[\text{Fe}(\text{CO})_2(\text{PPh}_3)(\text{C}_6\text{H}_8)]$  gave a 5-endo-acetyl complex. [1056]  $\text{Ph}_3\text{C}^+$  converts  $[\text{Fe}(\text{CO})_3\text{L}]$ , ( $\text{L} = 5,5\text{-dimethyl-2-trimethylsilyl-1, 3-cyclohexadiene}$ ), to  $\eta^5$ -cations with  $\text{SiMe}_3$  at the centre of delocalisation; this and similar reactions were rationalised with PE spectra and MO calculations. [1057]  $[\text{Fe}(\text{CO})_3(\text{diene})]$  complexes have been prepared with N-alkoxycarbonyl-1,2-dihydropyridine. [1058] EHMO calculations on  $[\text{Fe}(\text{CO})_3(\text{cyclohexadienyl})]$  complexes showed how these could be formed from the diene complex and how nucleophilic addition gave regioselectivity. [1059] There has been a  $^{13}\text{C}$  NMR and Moessbauer study on solid state motions in this cyclohexadienyl complex. [1060]

In cycloheptadienyl complexes  $[\text{Fe}(\text{CO})_3(\text{diene})]^+$  nucleophilic addition is more inhibited by steric effects than in cyclohexadienyl ones. [1061] The action of "soft" nucleophiles like  $\text{NaCH}(\text{CO}_2\text{CH}_3)_2$  on  $[\text{Fe}(\text{CO})_2(\text{PR}_3)(\text{cycloheptatrienyl})]^+$ , ( $\text{R} = \text{Ph}, \text{OPh}$ ) gave cycloheptadiene complexes but "hard" nucleophiles such as  $\text{NaCN}$  produced allyl complexes [1062]. Wittig reactions on a  $[\text{Fe}(\text{CO})_3(\text{diene})]$  complex gave heptatriene complexes which act as diene ligands and may then undergo isomerization [1063]. Addition of the carbonylcarbenes  $\text{N}_2\text{CHCO}_2\text{Et}$  or  $\text{N}_2\text{CPhCOPh}$  to  $[\text{Fe}(\text{CO})_3(\text{C}_7\text{H}_8)]$  gave reactions on the  $\text{C}_7\text{H}_8$  ligand including formation of a cycloheptafuran complex [1064].  $[\text{Fe}(\text{CO})_3(\text{cot})]$  and  $\text{C}_7\text{H}_7^+$  react together to form  $[\text{Fe}(\text{CO})_3(\eta^4\text{-styrylcycloheptatriene})]$  but the use of methylcyclooctatetraene gives a product with a methyl group on the  $\text{C}_7$  ring, implying extensive rearrangement; the new complexes react with  $\text{HBF}_4$  and  $\text{tCne}$  [1065].

Dimethylcyclooctatetraene-1,8-dicarboxylate forms complexes with 1 or 2  $[\text{Fe}(\text{CO})_3]$  units; one of the latter has an Fe-Fe bond [1066]. The reaction of  $[\text{Fe}(\text{CO})_4]^{2-}$  and  $\alpha, \alpha'$ -dihaloxylenes gives 1,2-bismethylenebenzene complexes which easily rearrange to  $\pi$ -xylylene complexes [1067]. Organic halides react with  $[\text{Fe}(\text{C}_6\text{Me}_6)_2]$  to give  $[\text{Fe}(\text{C}_6\text{H}_6)(\text{diene})]$  complexes which react with oxygen to give  $\sigma$ -xylylene complexes coordinated by the

exocyclic double bonds [1068]. Bis  $\eta^5$  complexes of bicyclooctadiene with  $2[\text{Fe}(\text{CO})_2\text{R}]$  moieties,  $[\text{R} = \text{CO}, \text{P}(\text{OPh})_3]$ , give ring expansion with  $\text{PPh}_3$  to 7-membered triene - diene complexes [1069]. Another triene - diene complex is given when Fe atoms react with bicyclononatriene; when this reacts with  $\text{P}(\text{OMe})_3$ ,  $[\text{Fe}\{\text{P}(\text{OMe})_3\}_3(\text{diene})]$  is formed [1070].

Turning next to bigger ring systems, octalene can be converted to bicyclodecapentaene with  $[\text{Fe}(\text{CO})_5]$  by the intermediate formation of allyl complexes [1071]. Heating many  $[\text{Fe}(\text{CO})_3(\text{C}_7\text{H}_7\text{CH:CHR})]$  complexes gave duron complexes but  $[\text{Fe}(\text{CO})_3(\text{C}_7\text{H}_7\text{CH:CHPh})]$  gave migration of the  $[\text{Fe}(\text{CO})_3]$  unit from the arene position to a diene configuration using one endo- and one exo-cyclic double bond [1072]. An  $[\text{Fe}(\text{CO})_3]$  complex of a bicyclododecadiene has been prepared [1073]. Cyclododecatriene and  $[\text{Fe}(\text{CO})_5]$  react to form a complex which reacts with  $\text{Ph}_3\text{C}^+$  to give a trienyl salt [1074]. The reaction of diaminodihydraceptalenes with  $[\text{Fe}_2(\text{CO})_9]$  gives diene complexes which can be reduced electrochemically or with Na to anionic monoene complexes [1075].

IR spectra have been used to study the tautomerism of the newly prepared  $[\text{Fe}(\text{CO})_4(\eta^2\text{-PhCH:CH.CR:NR}')]_n$ , with a  $\sigma$ -N donor, with an  $[\text{Fe}(\text{CO})_3]$  configuration involving  $\eta^3$ -carbarmoylallyl or  $\eta^4$ -azadiene coordination [1076].

The remaining complexes in this section are dinuclear. The chemistry of the  $[\text{Fe}_2(\text{CO})_7]$  complex of the phosphole 2,3-Me<sub>2</sub>-1-Ph-C<sub>4</sub>H<sub>2</sub>P, which shows both diene and phosphine coordination is reported;  $\text{AlCl}_3$  and  $\text{NH}_4\text{OH}$  remove the iron unit, reactions at P are achieved with  $\text{H}_2\text{O}_2$ ,  $\text{S}_8$ , MeI and  $\text{PhCH}_2\text{Br}$ , reaction with  $\text{PhCOCl}$ ,  $\text{Et}_3\text{N}$  and  $\text{H}_2\text{O}$  give the six-membered ring  $\text{PhP}(\text{O}).\text{CH:CMc.CMc:CH.C}(\text{Ph})\text{OH}$  coordinated as a diene to Fe;  $[\text{PdCl}_2(\text{NMe})_2]$  converts it to a bimetallic complex [1077]. Deprotonation of  $[\text{Fe}_2(\text{CO})_6(\text{PPh}_2\text{H})]^-$  to  $[\text{Fe}_2(\text{CO})_6(\text{PPh}_2)_2]^{2-}$  followed by reaction with allyliodide gives (X-ray)  $[\text{Fe}_2(\text{CO})_5(\text{PPh}_2)(\text{PPh}_2\text{C}_3\text{H}_5)(\eta^3\text{-C}_3\text{H}_5)]$  [1078]. An  $[\text{Fe}_2(\text{CO})_6]$  complex of the diene

$\text{MeC}:\text{C}(\text{NEt}_2)_2\text{C}(\text{NEt}_2)\text{CMe}$  is formed when  $[\text{Fe}_3(\text{CO})_{12}]$  reacts with  $\text{Et}_2\text{NC}:\text{CMe}$  [1079]. An allene-type complex is formed by refluxing in hexane  $[\text{Fe}_2(\text{CO})_6(\mu\text{-COEt})(\mu\text{-CPhCHPh})]$ , prepared from  $[\text{Fe}_2(\text{CO})_4(\mu\text{-CPhCHPh})]$  by reaction with  $\text{Et}_3\text{O}^+\text{BF}_4^-$  [1079]. The alkyne  $\text{HOCPhRC}:\text{C}:\text{CPhR}'\text{OH}$  reacts with  $[\text{Fe}_3(\text{CO})_{12}]$  to give  $[\text{Fe}_2(\text{CO})_6(\mu\text{-PhRC}:\text{C}:\text{C}:\text{CPhR})]$  with each double bond binding different Fe's [1080].

### 7.7. Cyclopentadienyl and Related Compounds

This very large survey begins with ferrocene and other compounds containing two  $\pi$ -bonded cyclic ligands. A new preparation of ferrocene uses controlled potential electrolysis of  $\text{TiCp}$  with an iron anode [1081]. The new unstable 19 - electron complex  $[\text{Fe}(\text{cp})(\eta^6\text{-fluorene})]$ , prepared from the  $\text{Fe}(\text{II})$  cation, is a good intermediate for reactions of fluorene [1082]. The preparation and X-ray structure of the ferrocene analogue of isodicyclopentadienide are reported [1083]. New complexes  $[\text{Fe}(\text{cp})(\text{C}_5\text{H}_4\text{R})]$ , ( $\text{R} = \text{NO}, \text{NO}_2$ ) are prepared by the action of  $\text{H}_2\text{O}_2$  on the product,  $[\text{Cr}(\text{cp})(\text{NO})(\text{CN}^t\text{Bu})\{\text{Fe}(\text{cp})(\text{C}_5\text{H}_4\text{NO})\}]$ , from the reaction of  $[\text{Fe}(\text{cp})(\text{C}_5\text{H}_4\text{Li})]$  with  $[\text{Cr}(\text{cp})(\text{NO})_2\text{Cl}]$  and  $^t\text{BuNC}$ ; the complexes react with  $\text{CH}_2(\text{CN})_2$  or  $\text{PEt}_3$  to give complexes with  $\text{R} = \text{N}:\text{C}(\text{CN})_2$  or  $\text{N}:\text{PEt}_3$  [1084]. New tropone ( $\text{L}$ ) complexes  $[\text{Fe}(\text{cp})\text{L}]^+$  are prepared by ring expansion using  $\text{CH}_2\text{N}_2$  on the products of e.g.  $[\text{Fe}(\text{cp})(\text{C}_6\text{H}_5\text{Cl})]^+$  with  $\text{NaOH}$  or  $\text{NaSH}$  [1085].

There have been many other reports of new arene - cyclopentadienyl complexes, including examples with di- or tetrahydroanthracene which have been prepared by the reaction of anthracene with  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\text{X}]$ , ( $\text{X} = \text{Br}, \text{Cl}$ ) [1086]. Xanthene or thioxanthene react with ferrocene to give new  $\eta^6$  - complexes with the loss of one cp ligand; subsequent stereospecific reactions occur on the new complexed ligand [1087].

Many new compounds contain hetero-atoms in the ligand. Using an Fe anode in the controlled potential electrolysis of carbaborane anions gave salts of  $[\text{Fe}(\text{C}_2\text{B}_9\text{H}_{11})_2]^-$  [1088]. Iron atoms react with  $\text{B}_{10}\text{H}_{14}$  and

toluene or mesitylene to form arene ferraboranes with 8, 9 or 10 boron atoms; one of the products, arising from an impurity reaction, was the first polyhedral boron compound to contain oxygen at a cluster vertex [1089].

[Fe(cp)(1-phenyl-3,4-dimethylphosphole)] is prepared by the reaction of the phosphole with  $[\text{Fe}_2(\text{cp})_2(\text{CO})_4]$  [1090].

Now for crystal structures and other physical studies. 1,1',2,2'-

Tetrachloroferrocene has three crystal phases; the low temperature phase has the rings eclipsed but the others are more disordered [1091]. Ab initio MO calculations show anomalously large quadrupole splittings in polychlorinated ferrocene due to less electron donation to iron as the ring becomes more distant from the metal [1092]. The PE spectra of ferrocene and bis(pentadienyl) iron are compared [1093]. PE spectra are combined with EHT and X $\alpha$  calculations for  $[\text{Fe}(\text{C}_4\text{H}_4\text{P}_2)]$  [1094]. Intervalence transitions are described for many polyferrocene complexes and correlated with the presence of  $\pi$ - and  $\sigma$ -bonding [1095].

Studies of Moessbauer spectra include the observation of changes in the quadrupole splitting of substituted phosphoferrocenes when a phenyl substituent is protonated by  $\text{CF}_3\text{SO}_3\text{H}$  to give a  $\sigma$ -bonded species [1096].

Vibrational assignments and the splitting of the two lowest Kramers doublets in  $[\text{Fe}(\text{C}_5\text{D}_5)]^+$  were established by inelastic neutron scattering spectroscopy at 10K; the analysis includes non-axial ligand fields, spin-orbit coupling and the dynamic Jahn-Teller effect [1097]. For

$[\text{Fe}(\text{cp})(\text{C}_6\text{Me}_6)]^{n+}$ , ( $n=0,1$ ), the extra electron in the reduced compound was shown to spend 83% of its time on the iron atom [1098]. Moessbauer and NMR ( $^1\text{H}$ ) spectra of  $[\text{Fe}(\text{cp})\{\text{C}_5\text{H}_4\text{CH}(\text{Ar})\}]^+$  salts [1099]. Spectra of

$[\text{XC}_5\text{H}_4\text{FeC}_5\text{H}_4\text{C}_5\text{H}_4\text{FeC}_5\text{H}_4\text{X}]^+$  show delocalisation for  $\text{X} = \text{Br}$  or

$\text{I}$  but localisation for  $\text{Cl}$ , despite similar redox potentials [1100]. For triiodide salts of mixed valence biferrocenes spectra were taken in magnetic fields to assign the unpaired electron's orbital; the problem is complicated by spin-lattice relaxation effects [1101]. There is a linear relationship



between quadrupole splitting and redox potentials in some substituted ferrocenes [1102].

Gas phase thermal decomposition gives  $381 \pm 12 \text{ kJ mol}^{-1}$  for the first bond dissociation energy of ferrocene [1103]. The ionisation of  $[\text{Fe}(\text{cp})\{\text{C}_5\text{H}_4\text{CH}(\text{OH})\text{C}_6\text{H}_4\text{R}\}]^+$  has been studied [1104]. A mechanistic study of the photochemical loss of arene from  $[\text{Fe}(\text{cp})(\text{arene})]^+$  includes the effects of steric interactions on the arene which may prevent the solvent or anions participating in the transition state [1105]. The reduction of ferrocene and related compounds has been reviewed [1106].

Cyclic voltammetry has been widely applied to ferrocene derivatives; subjects this year have included  $[\text{Fe}(\text{RCOC}_5\text{H}_4)_2]$  [1107] and  $[\text{Fe}(\text{cp})(\text{C}_5\text{H}_4\text{-C}_6\text{H}_4\text{R})]$  [1108]. Ferrocenes including  $\text{C}_4\text{H}_4\text{P}$  and  $\text{Me}_2\text{C}_4\text{H}_2\text{P}$  rings can lose an electron from Fe, but loss of a second electron gives irreversible oxidation of the ligand [1109].  $[\text{Fe}(\text{cp})(\text{cyclophane})]^+$  and  $[\text{Fe}_2(\text{cp})_2(\text{cyclophane})]^{2+}$  are compared with corresponding arene complexes; two distinct steps are found in the reduction of the diiron complexes [1110].

In chemical reactions which are not purely organic in character, the oxidation of ferrocene to  $[\text{Fe}(\text{C}_5\text{H}_4\text{R})(\text{C}_5\text{H}_4\text{R}')]^+ \text{I}_3^-$  is described [1111].

Ferrocene or ferricenium sulphate react with  $\text{Cl}_3\text{CCO}_2\text{H}$  to give that ferricenium salt with solvated acid [1112]. Ferrocenes carrying one or two crown ether ligands linked to cp by  $\text{CH}_2\text{SCH}_2$  units have been synthesised and found to complex alkali metals [1113].  $[\text{Fe}(\text{cp})(\text{arene})]^+$  salts of  $\text{B}_{10}\text{H}_{10}^{2-}$ ,  $\text{B}_{11}\text{H}_{14}^{2-}$  and  $\text{B}_{12}\text{H}_{12}^{2-}$  have been prepared [1114].

Ferrocene reacts with  $\text{FeCl}_3$  to give  $2[\text{Fe}(\text{cp})_2]^+ [\text{Fe}_2\text{OCl}_6]^{2-}$  and  $[\text{Fe}(\text{cp})_2]^+ [\text{FeCl}_4]^-$ , but not the previously reported  $[\text{FeCl}_3]^{2-}$  [1115]. Measurement of  $J(^{57}\text{Fe}-^{13}\text{C})$  in " $[\text{Fe}(\text{cp})_2(\text{CRR}')]$ ",

$[\text{R} = \text{H}, \text{Me}; \text{R}' = \text{H}, \text{Me}, \text{Ph}, \text{Mn}(\text{CO})_3\text{cp}, \text{Ru}(\text{cp})_3, \text{Fe}(\text{cp})(\text{C}_5\text{H}_4)]$ , show these really have bulvalene type bonding [1116].

The next section concerns complexes of ferrocenes which contain other metals

in addition to iron. We work from left to right in the usual representation of the periodic table to order the heterometal. Yb displaces Hg from  $[\text{Hg}(\text{C}_5\text{H}_4\text{Fecp})_2]$  [1117]. Reacting  $[\text{Fe}(\text{C}_5\text{H}_4\text{SeOH})_2]$  with  $[\text{M}(\text{C}_5\text{H}_4\text{R})_2\text{Cl}_2]$ , ( $\text{M} = \text{Zr}, \text{Hf}$ ;  $\text{R} = \text{H}, {}^t\text{Bu}$ ) gives  $[\text{Fe}(\text{C}_5\text{H}_4\text{Se})_2\text{M}(\text{C}_5\text{H}_4\text{R})_2]$  [1118]. A Zr-H moiety adds across CO when  $[\text{Fe}(\text{cp})(\text{CO})\text{H}(\text{PMe}_3)]$  reacts with  $[\text{Zr}(\text{C}_5\text{Me}_5)_2\text{H}_2]$  and  $\text{PMe}_3$  to give  $[\text{Fe}(\text{cp})(\text{PMe}_3)_2\text{CH}_2\text{OZrH}(\text{C}_5\text{Me}_5)_2]$  [1119]. In a new vanadium complex two  $[\text{V}(\text{C}_5\text{H}_4\text{R})]$  units are linked to one ferrocene component by V-C bonds so that the three sandwich fragments are perpendicular; the result is that the cp rings distort towards  $\eta^4$ -geometry [1120]. Chiral  $[\text{Fe}(\text{cp})(\text{CO})(\text{PMe}_3)(\text{EMe}_2)]$ , ( $\text{E} = \text{As}, \text{Sb}, \text{Bi}$ ) react with  $[\text{Mo}(\text{CO})_4(\text{norbornadiene})]$  to give diastereomeric  $[\text{Fe}(\text{cp})(\text{CO})(\text{PMe}_3)(\text{EMe}_2)]_2\text{Mo}(\text{CO})_4$  [1121]. The preparation and X-ray structure of  $[\text{Fe}(\text{cp})\text{C}_5\text{H}_4\text{CH}:\text{CHSi}(\text{Me})(\text{Ph})\text{C}_5\text{H}_4\text{Mn}(\text{CO})_3]$  are reported [1122]. Many complexes with  $\text{Ru}(\text{II})$  and various zerovalent metals are described with  $[\text{Fe}(\text{C}_4\text{H}_2\text{Me}_2\text{P}_2)_2]$ , ( $\text{L}$ ), in which one or both P atoms coordinate; in  $[\text{Ru}_2\text{Cl}_4(\text{PPh}_3)_2\text{L}_2]$  the  $\text{C}_4\text{P}$  rings are still aromatic as the ligands bridge but the P atoms are eclipsed, which is not the case for uncomplexed  $\text{L}$  [1123]. The  $\text{C}_7\text{H}_7$  ring in  $[\text{Fe}(\text{CO})_3(\text{C}_7\text{H}_7)\text{Rh}(\text{CO})_2]$  is  $\eta^3$  to Fe and  $\eta^4$  to Rh, but if the carbonyls are replaced by dppe the bonding becomes  $\eta^3$  to Rh and  $\eta^4$  to Fe [1124].  $[\text{NiCl}_2(\text{Ph}_2\text{PC}_5\text{H}_4)_2\text{Fe}]$  catalyses the stereoselective silylation of crotylmagnesium bromide [1125]. Ferrocenes with a Pd-cp  $\sigma$ -bond and a cyclometallated Pd- $\pi$ -cp linkage are described [1126].  $[\text{Fe}(\text{cp})\{\text{C}_5\text{H}_4(\text{AuPPh}_3)_3\}]^+$  is described [1127]. Moessbauer and  $^{199}\text{Hg}$  NMR spectra of  $[\text{Fe}(\text{cp})(\text{CO})_2\text{HgX}]$ , ( $\text{X} = \text{halide or bidentate S ligand}$ ), show the S ligand forces more electron density on to the Fe atom; the effect is even greater in  $[\text{Hg}\{\text{Fe}(\text{cp})(\text{CO})_2\}_2]$  [1128]. Moessbauer spectra suggest an Hg-Fe bond is present in the compounds formed when  $\text{Hg}(\text{II})$  compounds react with ferrocenophanes [1129]. Ferricinium ions oxidise  $[\text{Fe}_2(\text{C}_5\text{H}_4\text{HgC}_5\text{H}_4\text{CH}_2\text{NMe}_2)_2]$  to the mono- and dications [1130].

$[\text{Fe}(\text{C}_5\text{H}_4\text{COCH}_2\text{COMe})_2]$  forms diketonate complexes with other metals [1131].

Now we come to cyclopentadienyl compounds containing carbonyl ligands. The order is monomers before polymers and in decreasing number of carbonyls.

$[\text{Fe}(\text{cp})(\text{CO})_3]^+$  reacts with diamines to give

$[\{\text{Fe}(\text{cp})(\text{CO})_2\text{CONR}\}_2(\text{CH}_2)_n]$ ,  $[\text{Fe}(\text{cp})(\text{CO})_2\text{CONHCH}_2\text{CH}_2\text{NMe}_2]$  and

$[\text{Fe}(\text{cp})(\text{CO})_2(\text{CONHCH}_2\text{CMe}_2\text{NH}_3)]^+$ ; the latter can be deprotonated and react with the starting material to give

$[\text{Fe}(\text{cp})(\text{CO})_2\text{CONHCH}_2\text{CMe}_2\text{NHCOFe}(\text{cp})(\text{CO})_2]$  [1132]. Products formed by reactions which start with  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_3]^+$  include

$[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2(\text{PR}_3)]$ , ( $\text{R} = \text{Me}, \text{Ph}$ ),  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})(\text{dppe})]$ ,

$[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})\text{H}_2\text{dppe}]$ ,  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\text{H}]$  and

$[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})(\text{PMe}_3)\text{Me}]$  [1133].

An interesting transformation occurs when  $\text{SO}_2$  reacts with  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{PbPh}_3)]$  to give  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{SO}_2\text{Ph})]$  [1134]. Various  $[\text{Fe}(\text{cp})(\text{CO})_2\text{L}]$

complexes, ( $\text{L} = \text{alkene}, \text{alkyne}, \text{allene}$ ) react with  $[(\text{cyclopentadienyl})\text{Fe}(\text{CO})_3]^+$  salts to give oxohydroazulene complexes, with the

$[\text{Fe}(\text{cp})(\text{CO})_2]$  moiety connected to the five-membered ring [1135]. A short

review on  $[\text{Fe}(\text{cp})(\text{CO})_2\text{H}]$  has appeared [1136].  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{EPh}_3)]$ ,

( $\text{E} = \text{Si}, \text{Ge}, \text{Sn}$ ) can be reversibly reduced electrochemically to its monoanion,

in which the unpaired electron is located in a phenyl  $\pi$ -system; further

electrochemical reduction gives decomposition to  $[\text{Fe}(\text{cp})(\text{CO})_2]^-$  and

$\text{Ph}_3\text{EH}$  or  $\text{SnPh}_3^-$  [1137].  $[\text{Fe}(\text{cp})(\text{CO})_2\text{SiMe}_2\text{SiPh}_3]$  is prepared by

the reaction of  $\text{ClSiMe}_2\text{SiPh}_3$  with  $\text{Na}[\text{Fe}(\text{cp})(\text{CO})_2]$  [1138]. Acetic acid

converts  $\text{Na}[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2]$  to  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\text{H}]$ ; one of these

carbonyl groups can be substituted by  $\text{PMe}_3$  and further reaction with

$\text{Me}_3\text{P}:\text{CH}_2$  gives  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})(\text{PMe}_3)\text{Me}]$  [1139].  $[\text{Fe}(\text{cp})(\text{CO})_2\text{Cl}]$  is

formed from  $\text{SO}_2\text{Cl}_2$  and  $[\text{Fe}(\text{cp})(\text{CO})_2]_2$  in  $\text{CH}_2\text{Cl}_2$ , but in  $\text{C}_6\text{H}_6$

this reaction also gives  $[\text{Fe}(\text{cp})(\text{CO})_3]^+[\text{FeCl}_4]^-$  [1140].

$\text{Na}[\text{Fe}(\text{cp})(\text{CO})_2]$  reacts with  $\text{ClC}(\text{O})\text{CCHB}_{10}\text{H}_{10}$  to give a carborane

complex which loses CO to give  $[\text{Fe}(\text{CCHB}_{10}\text{H}_{10})(\text{cp})(\text{CO})_2]$  [1141].  $\text{H}_2\text{C}:\text{C}(\text{OEt})\text{CO}_2\text{Et}$  acts as an alkene ligand (L) in  $[\text{Fe}(\text{cp})(\text{CO})_2\text{L}]^+$  for reactions to prepare lactones [1142].  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\text{CH}_2\text{OH}]$  is formed from  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_3]^+$  and  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2]$  or  $[\text{Mo}(\text{dppe})_2\text{H}_4]$  [1143].  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\text{Br}]$  can be converted to  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_3]^+$  with CO and  $\text{AlCl}_3$  or with  $[\text{Fe}(\text{cp})_2]^+$ ; the latter reacts with  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2]_2$  in THF to give  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2(\text{THF})]^+$ , from which the THF can be substituted to give many cationic or neutral complexes [1144, 1145]. Various  $[\text{Fe}(\text{C}_5\text{H}_4\text{CO}_2\text{H})(\text{CO})_2\text{R}]$ ,  $[\text{R} = \text{Ph}, \text{CH}_2\text{Ph}, \text{C}_5\text{H}_4\text{Mn}(\text{CO})_3]$ , have been prepared; all are stronger acids than ferrocenecarboxylic acid [1146]. The complexes  $[\text{Fe}(\text{cp})(\text{CO})_2\text{AsR}_2]$ ,  $(\text{R} = \text{Me}, ^t\text{Bu})$  and their  $\text{C}_5\text{Me}_5$  analogues are prepared from  $[\text{Fe}(\text{cp})(\text{CO})_2]^-$ ; the products of their reactions with MeI,  $\text{P}(\text{OMe})_3$  and S are described [1147]. (Mesityl) $\text{PCl}_2$  converts  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2]^-$  to  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\text{I}]_2$ , but  $^i\text{Pr}_2\text{NPCl}_2$  gives  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2(\text{R}_2\text{NPCl})]$ , whose reduction by  $\text{LiEt}_3\text{BH}$  gives  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2(\text{RNPH})]$ , which in turn reacts with  $\text{Ph}_3\text{C}^+$  or  $\text{AlCl}_3$  to give  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})_2\{\text{PHNRCMeCH}_2\}]^+$ , which may be formed via an N-P-Fe heteroallyl intermediate [1148]. Thiophosphinite complexes  $[\text{Fe}(\text{cp})(\text{CO})_2\{\text{P}(\text{S})\text{R}_2\}]$ ,  $(\text{R} = \text{alkyl, aryl})$  are prepared from  $[\text{Fe}(\text{cp})(\text{CO})_2\text{Br}]$ ,  $\text{R}_2\text{HPS}$  and  $\text{NEt}_3$  [1149]. Electrochemical reduction of  $[\text{Fe}(\text{cp})(\text{CO})_2\text{SnR}_3]$  and other tin complexes gave new distannanes carrying  $[\text{Fe}(\text{cp})(\text{CO})_2]$  groups [1150]. Norbornyl (X) complexes  $[\text{Fe}(\text{cp})(\text{CO})_2\text{X}]$  have been prepared [1151].  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CS}_2)]^-$  replaces Cl in  $\text{Me}_3\text{ECI}$ ,  $(\text{E} = \text{Si, Sn})$  or I in  $[\text{Ru}(\text{cp})(\text{CO})_2\text{I}]$ ; the Ru complex and related  $[\text{Re}(\text{CO})_5]$  and  $[\text{Fe}(\text{cp})(\text{CO})_2]$  species are alkylated at the thione S and the reactions of those products are also described [1152].

There are not so many articles about complexes which contain an  $[\text{Fe}(\text{cp})(\text{CO})]$  unit.  $[\text{Fe}(\text{cp})(\text{CO})(\text{Et})\text{L}]$ ,  $\{\text{L} = \text{PPh}_3, \text{P}(\text{OCH}_2)_3\text{CMe}_3\}$ , react with CO with or without  $\text{C}_6\text{H}_{11}\text{NC}$  to give  $[\text{Fe}(\text{cp})\text{LL}'(\text{COEt})]$ ,  $(\text{L}' = \text{CO}, \text{C}_6\text{H}_{11}\text{NC})$ ; the

reaction is stereospecific in nitroalkanes [1153]. BuLi reacts with  $[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)(\text{COMe})]$  to form  $[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)\text{COCH}_2]^-$ , which can be alkylated to give  $[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)\text{COR}]$  or  $[\text{Fe}(\text{cp})(\text{CO})(\text{PPh}_3)\text{COCH}_2\text{CRR'OH}]$  [1154].  $[\text{Fe}(\text{cp})(\text{dppe})(\text{CO})]^+$  is reduced by  $\text{LiAlH}_4$  to  $[\text{Fe}(\text{cp})(\text{dppe})(\text{CO})\text{H}]$  (A) at  $-78^\circ\text{C}$  via a formyl intermediate but refluxing in THF gives  $[\text{Fe}(\eta^5\text{-C}_5\text{H}_6)(\text{dppe})(\text{CO})]$  and  $[\text{Fe}(\text{cp})(\text{dppe})\text{Me}]$  (B); (A) decomposes to  $[\text{Fe}(\text{cp})(\text{CO})\text{H}]_2(\mu\text{-dppe})$  in toluene but to (B) in THF [1155]. A similar starting material containing dppp instead of dppe gives  $[\text{Fe}(\text{cp})(\text{dppp})(\text{CO})\text{H}]$ , with monodentate dppp, but dmpe gives  $[\text{Fe}(\text{cp})(\text{dmpe})_2\text{Me}]$  [1156]. Similar chemistry is reported for the formation of  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{dppe})(\text{CO})\text{H}]$  [1157]. A more complex new compound of this type is  $[\text{FeC}_5\text{H}_4\text{PPh}_2\{\text{Fe}(\text{cp})(\text{CO})\}\{\text{C}(\text{O})\text{C}_5\text{H}_4\}]$  [1158]. Reacting  $\text{Me}_3\text{P:CH}_2$  with  $[\text{Fe}(\text{cp})(\text{CO})_n(\text{RNC})_{3-n}]^+$  gives ylide complexes whose reactivity was investigated, with mechanistic discussions [1159]. Electronic spectra were used to study solution equilibria between monomeric  $[\text{Fe}(\text{cp})(\text{CO})(\text{SPh})\{\text{P}(\text{OR})_3\}]^+$ , ( $\text{R} = \text{Et}, \text{Ph}$ ) or  $[\text{Fe}(\text{cp})(\text{CNC}_6\text{H}_4\text{R}')_2(\text{SPh})]^+$ , ( $\text{R}' = \text{OCH}_3, \text{CF}_3$ ) and dimers with  $\text{PhSSPh}$  ligands; electron withdrawing groups favour dimers [1160]. The X-ray structures of  $[\text{Fe}(\text{cp})(\text{CO})\{\text{C}_5\text{H}_5\text{Mn}(\text{CO})_3\}\text{L}]$ , ( $\text{L} = \text{CO}, \text{PPh}_3$ ), confirm that iron has one  $\sigma$ - and one  $\pi$ -bonded cp ligand [1161]. Paramagnetic  $[(\text{C}_{10}\text{H}_8)_2\text{Fe}(\text{CO})]^+$  salts have been prepared [1162]. Other new compounds reported are  $[\text{Fe}(\text{cp})(\text{CO})(\text{CRPPH}_3)]$ , ( $\text{R} = \text{H}, \text{Me}$ ) [1163],  $[\text{Fe}(\text{cp})(\text{CO})\text{L}(\text{CO}^i\text{Bu})]$ , ( $\text{L} = \text{PPh}_3, \text{PPh}_2\text{NET}_2$ ) [1164], various phosphine complexes  $[\text{Fe}(\text{cp})(\text{CO})\text{L}(\text{PhSEt})]^+$  [1165] and  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{PMe}_3)_1(\text{CO})_m(\text{NCMe})_n]$ , ( $1+m+n=3$ ) [1166].

Turning now to polynuclear compounds,  $\text{GeF}_2$  reacts with  $[\text{Fe}(\text{cp})(\text{CO})_2\text{R}]$ ,  $\{\text{R} = \text{Mn}(\text{CO})_5, \text{Mo}(\text{cp})(\text{CO})_3\}$ , to give  $\text{F}_2\text{Ge}[\text{Fe}(\text{cp})(\text{CO})_2]_2$ , but reaction with  $\text{R} = \text{SiPh}_3$  gives polymers [1167].  $[\text{Fe}(\text{cp})(\text{CO})_2]_2$  and its  $\text{C}_5\text{Me}_5$  and  $\text{C}_5\text{H}_4\text{CH}_2\text{Ph}$  analogues, when photolysed at 77K in organic matrices, give  $[\text{Fe}_2(\text{cp})_2(\text{CO})_3]$  which have a triple CO bridge and will be analogues of  $[\text{Fe}_2(\text{cp})(\text{NO})_2]$  with an Fe-Fe bond. [1168, 1169]. The new compound

$[\text{Fe}_2(\text{CO})_4(\text{C}_5\text{H}_4\text{SiMe}_2\text{C}_2\text{H}_4\text{SiMe}_2\text{C}_5\text{H}_4)]$  has two bridging carbonyls as well as a ferrocenophane bridge [1170]. New silicon containing complexes include  $[\text{Fe}(\text{cp})\text{L}_2(\text{SiR}_3)]$ , ( $\text{L} = \text{CO}$ ,  $\text{PMe}_3$ ;  $\text{R} = \text{H}$ ,  $\text{NMe}_2$ ), some of which form polynuclear complexes on reaction with  $[\text{Co}_2(\text{CO})_8]$  [1171, 1172], and polymeric  $[\text{OSi}(\text{Me})_2\text{Fe}(\text{cp})(\text{CO})_2]$  [1173]. Studies on  $[\text{Fe}(\text{cp})(\text{CO})_2]_2$  and related compounds include cyclic voltammetry [1174] and Moessbauer and ir spectra [1175]. Alkenes and cycloalkanes can be prepared from such as  $[\text{Fe}(\text{cp})(\text{CO})_2(\text{CH}_2)_n\text{Fe}(\text{cp})(\text{CO})_2]$  [1176]. When  $[\text{Fe}(\text{cp})(\text{CO})_2]_2$  is used in the hydroformylation of propene or 1-pentene,  $[\text{Fe}(\text{cp})(\text{CO})_2\text{H}]$  was detected [1177]. Interesting fluxional behaviour is found in the structures of  $[\text{Fe}_2(\text{cp})_2(\text{CO})_2(\text{CS})(\text{CNR})]$ ; ( $\text{R} = \text{Me}$ ,  $\text{Pr}$ ) [1178].

The structure of  $[\{\text{Fe}(\text{cp})(\text{CO})_2(\text{SEt})_2\}^+\text{ClO}_3^-]$  has been determined [1179].  $\text{FeCl}_3$  only oxidises one Fe in  $[\{\text{Fe}(\text{cp})(\text{C}_5\text{H}_4)\}_2\text{CO}]$ ; the paper includes a discussion of Moessbauer quadrupole splittings in diferrocenyl monocations [1180].

Many complexes have been prepared which contain other ligands instead of CO on cyclopentadienyl iron units. Poly (1,1'-ferrocenylene) adducts with  $\text{I}_2$ , TCNQ or TCNE show partial oxidation with very high conductance; the TCNQ complex is the best characterised [1181].  $[\text{Fe}(\text{cp})\{\text{P}(\text{OMe})_3\}_3]^+$  has been prepared by controlled potential electrolysis of  $[\text{Fe}(\text{cp})(\text{C}_6\text{H}_n\text{Me}_{6-n})]^+$  with added  $\text{P}(\text{OMe})_3$  [1182].  $\text{OP}(\text{Ph})\text{CHC}(\text{Me})\text{C}(\text{Me})\text{CHC}(\text{Ph})$  behaves as an  $\eta^4$ -ligand to an  $[\text{Fe}(\text{cp})]$  moiety with the P atom folded away from Fe [1183]. Crown ethers containing two ferrocene units have been prepared and found to coordinate  $\text{Ag}^+$  with possibly some electron transfer from Fe to Ag [1184].  $[\text{Fe}(\text{cp})(\text{dppe})(\text{CN})]$  has been prepared from the Cl analogue [1185]. Acetylferrocene has been used to prepare  $[\text{Fe}(\text{cp})\{\text{C}_5\text{H}_4\text{C}(\text{Cl})\text{CHCHO}\}]$ , whose reactions were studied [1186]. The carbene complex  $[\text{Fe}(\text{cp})\{\text{C}(\text{Et})\text{OMe}\}\text{CO}(\text{PPh}_3)]^+$  and its stereoselective addition reactions have been described [1187]. Reaction of  $\text{N}_2\text{CHCHO}$  with  $[\text{Fe}(\text{cp})(\text{NO})]_2$  gives addition across the

Fe=Fe bond to form  $[\{\text{Fe}(\text{cp})(\text{NO})\}_2\text{CHCHO}]$  [1188].

There have been very many studies on cyclopentadienyl complexes, usually ferrocenyl, containing substituents on the cp rings. These include X-ray structures of ferrocenyl pyrazoles [1189] and of 1,1',2,2'-tetrachloroferrocene [1190],  $^{13}\text{C}$  NMR of ferrocenyl ketones [1191], ir studies on these [1192], vibronic coupling in biferrocene and bis(fulvalene) diiron cations [1193] and Moessbauer spectra of ferrocenyl dimers [1194]. However, most work on substituted ferrocene complexes can be described as organic synthesis. Work of this type has been reviewed at length [1195]. It is impossible to do justice to this field in the available time and space; interested readers should consult the bibliography [1196-1229].

Ferrocenophanes form a special class. Most of the work on these has also been synthetic but the work has been supported by X-ray structures [1230, 1231], NMR spectra [1232, 1233], Moessbauer spectra [1234, 1235], electrochemistry [1234, 1236]. Much other work, though often ingenious, is essentially synthetic in character [1237-1247]. There remain one kinetic study [1248] and some INDO-SCF calculations on ferrocenophane reactivity [1249].

- [1] P. Brant and R.D. Feltham, J. Electron Spectrosc. Relat. Phenom., 32(1983)205
- [2] M. Takano and Y. Takeda, Bull. Inst. Chem. Res. Kyoto Univ., 61(1983)406
- [3] J.C. Grenier, M. Pouchard, P. Hagenmuller, M.J.R. Henche, M. Vallet, J.M.G. Calbet and M.A. Alario - Franco, Rev. Chim. Miner., 20(1983)726
- [4] S. Tagaki and T. Inoue, Kinki Daigaku Rikogakubu Kenkyu Hokoku, (1984)71; CA, 101(1984)29295q
- [5] W. Miller, J. Sraehle, A. Datz, L.W. ter Haar, W.E. Hatfield and P. Guetlich, Mol. Cryst. Liq. Cryst., 107(1984)151
- [6] K.A. Macor and T.G. Spiro, J. Electroanal. Chem. Interfacial Electrochem., 163(1984)223
- [7] W.H. Koppenol and J.F. Liebman, J. Phys. Chem., 88(1984)99
- [8] H. Sugimoto and D.T. Sawyer, J. Am. Chem. Soc., 106(1984)4283
- [9] L.S. Vol'dman and V.M. Berdnikov, Koord. Khim., 9(1983)1480
- [10] J. Pickardt, J. Kaehler, N. Rautenberg and E. Riedel, Z. Naturforsch. B, 39(1984)1162
- [11] M. Witzel and D. Babel, Z. Naturforsch. B, 39(1984)201
- [12] E. Garnier, P. Graverau, K. Ahmadi and A. Hardy, Rev. Chim. Miner., 21(1984)144
- [13] P. Graverau and E. Garnier, Acta Crystallogr. C, 40(1984)1306
- [14] H. Henkel and D. Babel, Z. Naturforsch. B, 39(1984)880
- [15] A. Saito, Y. Motioka and I. Nakagawa, J. Phys. Chem., 88(1984)480
- [16] T.A. Slivko, S.B. Stepina, I.F. Poletaev and T.V. Kamyhnikova, Russ. J. Inorg. Chem., 29(1984)556
- [17] T.M. Finogenko, I.D. Isaev, V.A. Fedorov and V.E. Mironov, Izv. Vyssh. Uchebn. Zaved. Khim. Khim. Tekhnol., 26(1983)1288
- [18] D. Kishore, H.S. Singh, G.P. Gupta and K.C. Lal, Proc. Nucl. Phys. Solid State Phys. Symp., 23C(1980)764



- [19] P.G. Rasmussen and E.A. Meyers, Polyhedron, 3(1984)183
- [20] R. Tenne, J. Electrochem. Soc., 130(1983)2163
- [21] B.D. Humphrey, S. Sinha and A.B. Bocarsly, J. Phys. Chem., 88(1984)736
- [22] S. Pons, M. Datta, J.F. McAleer and A. Shinman, J. Electroanal. Chem. Interfacial Electrochem., 160(1984)369
- [23] K.C. Cho and R.H. Austin, J. Chem. Phys., 80(1984)1131
- [24] P. Salvador and C. Gutierrez, J. Electrochem. Soc., 131(1984)326
- [25] P. Nesvadba and J. Kuthan, Collect. Czech. Chem. Commun., 49(1984)543
- [26] P. Nesvadba, P. Strop and J. Kuthan, Collect. Czech. Chem. Commun., 48(1983)3307
- [27] P. Nesvadba and J. Kuthan, Collect. Czech. Chem. Commun., 48(1983)2965
- [28] P.L. Majumder and A. Kundu, J. Indian Chem. Soc., 61(1984)142
- [29] A.K. Bhattacharjee and M.K. Mahanti, Bull. Korean Chem. Soc., 4(1983)120
- [30] A.K. Bhattacharjee and M.K. Mahanti, React. Kinet. Catal. Lett., 22(1983)227
- [31] T.K. Nikolaenko, I.V. Mel'Nichenko and A.A. Yasnikov, Dopov. Akad. Nauk Ukr. RSR, Ser. B: Geol. Khim. Biol. Nauki, (1984)47
- [32] J.W. Bunting and G.M. Kauffman, Can. J. Chem., 62(1984)729
- [33] M. Bhattacharjee and M.K. Mahanti, Bull. Soc. Chim. Fr., (1983), 225
- [34] B. Singh, S. Saxena and S.P. Singh, J. Indian Chem. Soc., 60(1983)795
- [35] S. Ali and Z. Murtaza, Indian J. Chem. A, 23(1984)258
- [36] C.R. Johnson and R.E. Shepherd, Synth. React. Inorg. Met. - Org. Chem., 14(1984)339
- [37] J.A. Clark, M. Kilner and A. Pietrzykowski, Inorg. Chim. Acta,

- 82(1984)85
- [38] B.J. Kennedy, G. Brain and K.S. Murray, Inorg. Chim. Acta, 81(1984)L29
  - [39] O. Hirabaru, T. Nakase, K. Hanabuse, H. Shirai, K. Takemoto and N. Hojo, Angew. Makromol. Chem., 121(1984)59
  - [40] M.Z. Ilic and R.W. Cattrall, Aust. J. Chem., 37(1984)489
  - [41] M. Genchev, S. Manolov and S. Zhekov, Koord. Khim., 10(1984)168
  - [42] D.M.L. Goodgame, P.B. Hayman and I. Sayer, Inorg. Chim. Acta, 92(1984)L13
  - [43] C.M. Mikulski, L. Mattucci, L. Weiss and N.M. Karayannis, Inorg. Chim. Acta, 92(1984)275
  - [44] K.C. Patil, J.P. Vittal and C.C. Patel, Proc. Indian Acad. Sci., Chem. Sci., 92(1983)83
  - [45] S.A.A. Zaidi, T.A. Khan and Z.A. Siddiqi, Synth. React. Inorg. Met. - Org. Chem., 14(1984)717
  - [46] K. Wieghardt, K. Pohl, I. Jibril and G. Huttner, Angew. Chem. Internat. Ed., 23(1984)77
  - [47] N. Marques and A. Pires de Matos, Inorg. Chim. Acta, 84(1984)L29
  - [48] J.C. Daran, Y. Jeannin and L. Martin, Acta Crystallogr. C, 40(1984)406
  - [49] J.E. Teggin and R.M. Milburn, J. Coord. Chem., 13(1984)245
  - [50] B.Ya. Kuvavskaya and N.V. Gerbeleu, Russ. J. Inorg. Chem., 29(1984)854
  - [51] A.M. Agranovich, A.P. Borisova, N.A. Dobrynina and S.P. Gladkikh, Russ. J. Inorg. Chem., 29(1984)1012
  - [52] I. Sanner, E. Meissner, H. Koeppen, H. Spiering and P. Guetlich, Chem. Phys., 86(1984)227
  - [53] P.C. Healy, B.W. Skelton and A.H. White, Aust. J. Chem., 36(1983)2057
  - [54] P.C. Healy, J.M. Patrick, B.W. Skelton and A.H. White, Aust. J. Chem., 36(1983)2031; W.L. Driessen, R.A.G. de Graaff and J.G.

- Vos, Acta Crystallogr. C, 39(1983)1635
- [55] B.N. Figgis, J.M. Patrick, P.A. Reynolds, B.W. Skelton, A.H. White and P.C. Healy, Aust. J. Chem., 36(1983)2043
- [56] B.N. Figgis, P.A. Reynolds and N. Lehner, Acta Cryst. B, 39(1983)711
- [57] B.N. Figgis, P. A. Reynolds and R. Mason, Inorg. Chem., 23(1984)1149
- [58] J.E. Plowman, T.M. Loehr, C.K. Schauer and O.P. Anderson, Inorg. Chem., 23(1984)3553
- [59] W.M. Reiff and E.H. Witten, Polyhedron, 3(1984)443
- [60] I.A. Brikun, I.A. Belenko, A.V. Storozhev and E.A. Mambetkaziev, Izv. Akad. Nauk Kaz. SSR, Ser. Khim., 3(1984)66
- [61] S. Tachiyashiki and H. Yamatera, Bull. Chem. Soc. Jpn., 57(1984)1070
- [62] N. Serpone and F. Bolletta, Inorg. Chim. Acta, 75(1983)189
- [63] A.B. Rovinsky, J. Phys. Chem., 88(1984)4
- [64] A.B. Rovinskii, Teor. Eksp. Khim., 20(1984)120
- [65] C.J. Schlesener and J.K. Kochi, J. Org. Chem., 49(1984)3142
- [66] A.S.P. Vittal, P.V. Rao and K.J.M. Rao, React. Kinet. Catal. Lett., 23(1983)175
- [67] V.G. Glukhovtsev, Yu.V. Il'in and G.I. Nikishin, Izv. Akad. Nauk SSR, Ser. Khim., (1983)1631
- [68] S.C. Jackels and L.J. Harris, Inorg. Synth., 22(1983)107
- [69] A.A. Aruffo, B.D. Santarsiero, V. Schomaker and E.C. Lingafelter, Acta Crystallogr. C, 40(1984)1693
- [70] S.G. Shova, Yu. A. Simonov, M.A. Yampol'skaya and N.V. Gerbeleu, Izv. Akad. Nauk. Mold. SSR, Ser. Fiz. - Tekh. Mat. Nauk, (1984)21; CA, 101(1984)161573h
- [71] W.D. Federer and D.N. Hendrickson, Inorg. Chem., 23(1984)3870
- [72] W.D. Federer and D.N. Hendrickson, Inorg. Chem., 23(1984)3861

- [73] D.P. Riley and D.H. Busch, Inorg. Chem., 23(1984)3235
- [74] N. Matsumoto, K. Kimoto, K. Nishida, A. Ohyoshi and Y. Maeda, Chem. Lett., (1984)479
- [75] N.V. Gerbeleu, S.G. Shova, B.Ya. Kuyavskaya, K.I. Turte, M.A. Yampol'skaya and M.Kh. Veksel'man, Russ. J. Inorg. Chem., 29(1984)157
- [76] W. Kanda, H. Okawa and S. Kida, Bull. Chem. Soc. Jpn., 57(1984)1159
- [77] M. Nakamura, H. Okawa and S. Kida, Inorg. Chim. Acta, 75(1983)9
- [78] A. Kapturkiewicz and B. Behr, J. Electroanal. Chem. Interfacial Electrochem., 163(1984)189
- [79] C.J. Carrano and K. Spartalian, Inorg. Chem., 23(1984)1993
- [80] F. Akhtar, G.C. De, M. Emran and M.T. Hussain, Nucl. Sci. Appl. B, 14 - 15(1983)67
- [81] A.A. Aruffo, T.B. Murphy, D.K. Johnson, N.J. Rose and V. Schomaker, Acta Crystallogr. C, 40(1984)1164
- [82] V. Atre, V.J.T. Raju, K.J. Ratnam and M.C. Ganorkar, Indian J. Chem., 23(1984)691
- [83] P. Chakravarty and P. Khanna, J. Indian Chem. Soc., 61(1984)112
- [84] U.G. Deshpande and J.R. Shah, Angew. Makromol. Chem., 122(1984)113
- [85] S. Dilli and E. Patsalides, J. Chromatogr., 270(1983)354
- [86] A.M. Hundekar and D.N. Sen, Indian J. Chem. A, 23(1984)477
- [87] M. Kinoshita and T. Sugano, Kotai Butsuri, 18(1983)267; CA, 101(1984)7905m
- [88] M. Mohan and M. Kumar, Synth. React. Inorg. Met. - Org. Chem., 14(1984)615
- [89] R.B. Patel and J.K. Verma, J. Indian Chem. Soc., 60(1983)1094
- [90] M.S. Patil, H.O. Deore, M.M. Kukarni and J.R. Shah, J. Indian Chem. Soc., 60(1983)817
- [91] G.R. Pokhariyal, J. Indian Chem. Soc., 60(1983)1089

- [92] R. Raina and T.S. Srivastava, Inorg. Chim. Acta, 91(1984)137
- [93] V.J. Raju, V. Ranabaore, B.B. Kumar and M.C. Ganorkar, J. Indian Chem. Soc., 60(1983)724
- [94] N.R. Shah and J.R. Shah, J. Indian Chem. Soc., 60(1983)1092
- [95] C.L. Sharma, S.S. Narvi and R.S. Arya, Acta Chim. Hung., 114(1983)349
- [96] H. Tanaka and S. Yamada, Synth. React. Inorg. Met. - Org. Chem., 13(1983)915
- [97] Y.M. Temerk, S.A. Ibrahim and M.M. Kamal, Z. Naturforsch. B., 39(1984)812
- [98] J.L. Vats, N. Garg, S.C. Sharma, H.S. Yadav and R.C. Saxena, Synth. React. Inorg. Met. - Org. Chem., 14(1984)69
- [99] A. Wasey, R.K. Bansal, B.K. Puri, F. Kamil and S. Chandra, Transition Met. Chem., 8(1983)341
- [99a] W. Kanda, H. Okawa and S. Kida, Bull. Chem. Soc. Jpn., 56(1983)3268
- [100] T.E. Pou, O.J. Murphy, V. Young, J.O. Bockris and L.L. Tongson, J. Electrochem. Soc., 131(1984)1243
- [101] Y. Bando, Y. Ikeda and K. Okuda, Kenkyu Hokoku - Asahi Garasu Kogyo Gijutsu Shoreikai, 43(1983)199; CA, 100(1984)149957h
- [102] E. Mendelovici, R. Villalba and A. Sagarazu, Mater. Chem. Phys., 10(1984)579
- [103] D. Benjelloun, J.P. Bonnet, J.P. Doumerc, J.C. Launay, M. Onillon and P. Hagenmuller, Mater. Chem. Phys., 10(1984)503
- [104] T. Tsuji, K. Naito and K. Ishigure, Phys. Status Solidi A, 82(1984)K57
- [105] B.L. Yang, F. Hong and H.R. Kung, J. Phys. Chem., 88(1984)2531
- [106] G.V. Buxton, T. Rhodes and R.M. Sellers, J. Chem. Soc. Faraday Trans. I, 79(1983)2961
- [107] H. Ouchi and M. Umesaki, IEEE Trans. Magn., MAG-19(1983)1980
- [108] C. Greaves, J. Solid State Chem., 49(1983)325

- [109] D.A. Khramov and A.V. Polosin, Fiz. Tverd. Tela (Leningrad), 25(1983)2769
- [110] E. Karamazsin, P. Satre and P. Vergnon, J. Therm. Anal., 28(1983)279
- [111] M.S. Lupin and G.E. Peters, Thermochim. Acta, 73(1984)79
- [112] G.V. Isagulyants, K.M. Gitis, M.I. Rozengart, V.N. Kornyshev and G.L. Markaryan, Geterog. Katal., (1983)511; CA, 100(1984)191200f
- [113] T.G. Alkhasov and N.S. Amirgulyan, React. Kinet. Catal. Lett., 24(1984)55
- [114] A.P. Rudenko, I.I. Kulakova and V.S. Savel'ev, Geterog. Katal., (1983)157; CA, 100(1984)159300
- [115] K.H. Kim, J.H. Jun and J.S. Choi, Bull. Korean Chem. Soc., 5(1984)41
- [116] Y. Syono, A. Ito, S. Morimoto, T. Suzuki, T. Yagi and S. Akimoto, Solid State Commun., 50(1984)97
- [117] A.B. Fasman, P.I. Zabolotin, N.A. Maksimova, S.V. Druz, A.M. Zagor'ev and S.D. Mikhailenko, Zh. Prikl. Khim. (Leningrad), 56(1983)2569
- [118] H.R. Harrison, R. Aragon, J.E. Keem and J.M. Honig, Inorg. Synth., 22(1983)43
- [119] T.C. Jackson, D.B. Jacobson and B.S. Feiser, J. Am. Chem. Soc., 106(1984)1252
- [120] N.A. Godshall, I.D. Raistrick and R.A. Huggins, J. Electrochem. Soc., 131(1984)543
- [121] R. Luge and R. Hoppe, Z. Anorg. Allg. Chem., 513(1984)141
- [122] A. Tomas, P. Laruelle, J.L. Dormann and M. Nogues, Acta Cryst. C, 39(1983)1615
- [123] W. Dai, S. Seetharaman and L.I. Staffansson, Scand. J. Metall., 13(1984)32
- [124] A.V. Arakcheeva and O.G. Karpinskii, Dokl. Akad. Nauk SSSR, 273(1983)1127

- [125] A.V. Abramov, V.P. Cheparin and A.P. Cherkasov, Tr. Mosk. Energ. Inst., 586(1982)66
- [126] T. Peev, L. Bozadzhiev, T. Stpilova and S. Nikolov, J. Radioanal. Nucl. Chem., 85(1984)151
- [127] J. Liu and H. Zhang, Nanjing Daxue Xuebao, Ziran Kexue, (1983)588
- [128] D. Gajapathy and K. C. Patil, Mater. Chem. Phys., 9(1983)423
- [129] T.M. Perekalina, E.M. Smirnovskaya, S.A. Cherkezyan and O.I. Lyamina, Kristallografiya, 29(1984)169
- [130] G.J. Baldha and R.G. Kulkarni, Solid State Commun., 49(1984)169
- [131] K.J.D. Mackenzie and I.W.M. Brown, J. Mater. Sci. Lett., 3(1984)159
- [132] M. Yoshimura, K. Yamasawa and S. Somiya, Anal. Chem. Symp. Ser., 17(1983)198
- [133] H. Taguchi and Y. Takahashi, J. Mater. Sci. Lett., 3(1984)251
- [134] V.F. Savchenko, Izv. Akad. Nauk SSSR, Neorg. Mater., 19(1983)1525
- [135] A. Sztaniszlav, E. Sterk, L. Fetter, M. Farkas - Jahnke and J. Labar, J. Magn. Magn. Mater., 41(1984)75
- [136] E. Beregi, E. Hild, A. Szaniszlav, G. Rudnay and J. Sztatisz, J. Magn. Magn. Mater., 41(1984)73
- [137] M.D. Sundararajan, A. Narayanasamy, T. Nagarajan, G.V. Subba Rao, K.A. Singh and L. Haeggstroem, Solid State Commun., 48(1983)657
- [138] Z.M. Stadnik and E. de Boer, J. Phys. Chem. Solids, 45(1984)113
- [139] A.M. Balbashov, G.V. Kozlov, S.P. Lebedev, A.M. Prokhorov and A.S. Prokhorov, Pis'ma Zh. Eksp. Teor. Fiz., 39(1984)461
- [140] T. Hibiya and J. Nakajima, J. Appl. Phys., 54(1983)7110
- [141] V. Agafonov, D. Michel, M. Perez y Jorba and M. Fedoroff, Bull. Mater. Res. Bull., 19(1984)233

- [142] A. Modaressi, R. Gerardin, B. Malaman and C. Gleitzer, J. Solid State Chem., 53(1984)22
- [143] M.C. Cadée and D.J.W. Ijdo, J. Solid State Chem., 52(1984)302
- [144] M. Shiojiri, S. Sekimoto, T. Maeda, Y. Ikeda and K. Iwauchi, Phys. Status Solidi A, 84(1984)55
- [145] E.W. Neuse and M.G. Meirim, Transition Met. Chem., 9(1984)205
- [146] L. Huang, F. Jiang and JH. Lu, Huaxue Tongbao, (1984)14; CA, 101(1984)63979a
- [147] P.C. Healy, J.M. Patrick and A.H. White, Aust. J. Chem., 37(1984)1405
- [148] C.M. Flynn, Chem. Rev., 84(1984)31
- [149] Y. Kanno, M. Shimizu and S. Takeoka, Kogai to Taisaku, 19(1983)1062
- [150] S.C.F. Au - Yeung, G. Denes, J.E. Greedan, D.R. Eaton and T.R. Birchall, Inorg. Chem., 23(1984)1513
- [151] F. Rodriguez, M. Moreno and J.C. Gomez Sal, Z. Naturforsch. A, 39(1984)617
- [152] T. Leskela, Thermochim. Acta, 77(1984)177
- [153] D. Hanzel and H. Bilinski, Mater. Chem. Phys., 10(1984)277
- [154] C. Morterra, A. Chlorino and E. Borello, Mater. Chem. Phys., 10(1984)119
- [155] M. Lorenz, F. Stiebert, N. Zopf and G. Kempe, J. Signalaufzeichnungsmater., 11(1983)313
- [156] D. Chambaere and E. de Grave, J. Magn. Magn. Mater., 42(1984)263
- [157] N.G. Holm, Origins Life, 14(1984)343
- [158] K. Inoue, K. Shibata, S. Ozeki and K. Kaneko, J. Electrochem. Soc., 131(1984)2435
- [159] M. Kiyama, T. Akita and T. Takada, Bull. Inst. Chem. Res. Kyoto Univ., 61(1983)335
- [160] T.D. Waute and F.M.N. Morel, J. Colloid Interface Sci.,



- 102(1984)121
- [161] M. Pernet, X. Obradors, J. Fontcuberta, J.C. Joubert and J. Tejada, IEEE Trans. Magn., MAG-20(1984)1524
  - [162] J.W. Schultze, S. Mohr and M.M. Lohrengel, J. Electroanal. Chem. Interfacial Electrochem., 154(1983)57
  - [163] J. Faus, J.M. Moratal and M. Castello, Transition Met. Chem., 9(1984)294
  - [164] B. Chiari, O. Piovesana, T. Tarantelli and P.F. Zanazzi, Inorg. Chem., 23(1984)3398
  - [165] R.C. Mehrotra and J. Singh, Transition Met. Chem., 9(1984)148
  - [166] A.Ya. Sychev, V.G. Isak, G.G. Duka and M.V. Gontsa, Russ. J. Phys. Chem., 58(1984)147
  - [167] B. Howlin, A.R. Mohd - Nor, J. Silver and P.W.C. Barnard, Inorg. Chim. Acta, 91(1984)153
  - [168] P. Hakkinen, Finn. Chem. Lett., (1984)59
  - [169] S. Koch and G. Ackermann, Z. Chem., 24(1984)265
  - [170] K.T. Douglas, B. Howlin and J. Silver, Inorg. Chim. Acta, 92(1984)135
  - [171] Ya.Ya. Kharitonov, M.A. Sarukhanov and S.A. Slivko, Russ. J. Inorg. Chem., 29(1984)864
  - [172] Y. Nishida, H. Shimo, K. Takahashi and S. Kida, Mem. Fac. Sci. Kyushu Univ. C, 14(1984)301; CA, 101(1984)203213y
  - [173] S. Tamagaki, K. Suzuki and W. Tagaki, Mem. Fac. Eng. Osaka City Univ., 24(1983)145; CA, 101(1984)191237u
  - [174] M.S. Masoud and M.M. El - Essawi, J. Chem. Eng. Data, 29(1984)363
  - [175] P.S. Bassi, B.S. Randhawa and H.S. Jamwal, Thermochim. Acta, 69(1983)367
  - [176] Z. Brzozka and C. Rozycki, Chem. Anal. (Warsaw), 28(1983)585
  - [177] A.I. Khol'kin, K. Gloe, K.S. Luboshnikova, P. Muehl, L.M. Gindin and N.V. Fedjuk, Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk, (1984)75; CA, 100(1984)145845x

- [178] I.L. Eremenko, A.A. Pasynskii, B. Orazsakgatov, O.G. Ellert,
- [179] V.M. Novotortsev, V.T. Kalinnikov, M.A. Porai - Koshits, A.S. Antsyshkina V.N. Ostrikova, L.M. Dikareva, Yu.T. Struchkov and R.G. Derr, Inorg. Chim. Acta, 73(1983)225
- [180] M. Puri and R.D. Verma, Indian J. Chem., 23(1984)612
- [181] W.H. Armstrong, A. Spool, G.C. Papaefthymiou, R.B. Frankel and S.J. Lippard, J. Am. Chem. Soc., 106(1984)3653
- [182] W.H. Armstrong and S.J. Lippard, J. Am. Chem. Soc., 106(1984)4632
- [183] L. Pajdowski and H.M. Ratajczak, Pol. J. Chem., 56(1982)49
- [184] M. Puri and R.D. Verma, Monatsh. Chem., 115(1984)533
- [185] S.V. Kukharensko, V.V. Strelets and O.N. Efimov, Elektrokhimiya, 19(1983)1597
- [186] A. Abras, M.M. Braga and J.C. Machado, J. Radioanal. Nucl. Chem., 86(1984)111
- [187] K. Turte, S.A. Bobkova and R.A. Stukan, Russ. J. Inorg. Chem., 29(1984)422
- [188] V.I. Ponomarev, L.O. Atovmyan, S.A. Bobkova and K.I. Turte, Dokl. Akad. Nauk SSSR, 274(1984)368
- [189] K. Burger, I. Zay and G. Takacsi Nagy, Inorg. Chim. Acta, 80(1983)231
- [190] H. Baumann, B. Strehmel, H.J. Timpe and D. Rehorek, J. Prakt. Chem., 325(1983)753
- [191] P. Wu, X. Xin, A. Dai and Y. Zhang, Cuihua Xuebao, 5(1984)30,38; CA, 100(1984)216403q, 216404r
- [192] B.S. Randhawa and P.S. Bassi, Radiochem. Radioanal. Lett., 59(1983)171
- [193] M. Brezeanu, E. Tatu, S. Bocai, O. Brezeanu, E. Segal and L. Patron, Thermochem. Acta, 78(1984)351
- [193a] M. Vital and R. Jagannathan, Transition Met. Chem., 9(1984)73
- [194] M. Castillo and E. Ramirez, Transition Met. Chem., 9(1984)273

- [195] W. Clegg, O.M. Lam and B.P. Straughan, Angew. Chem. Int. Ed. Engl., 23(1984)434
- [196] A.Ya. Sychev, N.V. Suen and V.G. Isak, Khim. Fiz., (1983)1565
- [197] V.V. Pal'chevskii, N.G. Firsin and V.I. Shcherbakova, Zh. Prikl. Khim. (Leningrad), 57(1984)1378; CA, 101(1984)117730h
- [198] J.M. Lopez - Alcala, M.C. Puerta - Vizcaino, F. Gonzalez - Vilchez, E.N. Duesler and R.E. Tapscott, Acta Crystallogr. C, 40(1984)939
- [199] X. Solans, M. Font Altaba and J. Garcia - Oricain, Acta Crystallogr. C, 40(1984)635
- [200] M.A. Blesa, E.B. Borghi, A.J.G. Maroto and A.E. Regazzoni, J. Colloid Interface Sci., 98(1984)295
- [201] V.S. Dukhanin, I.G. Gorichev and L.V. Malov, Russ. J. Phys. Chem., 58(1984)1468
- [202] K. Tanaka, C.A. Vega and R. Tamamushi, Bioelectrochem. Bioenerg., 11(1983)135,289
- [203] V.V. Pal'chevskii, T.I. L'vova, I.A. Seliverstova, O.M. Susareva and K.D. Shirko, Vestn. Leningr. Univ. Fiz., Khim., (1984)102; CA, 101(1984)236345w
- [204] E.Kh. Khudaiberdiev and M.K. Alyaviya, Koord. Khim., 10(1984)1072
- [205] M.N. Patel and M.R. Chaudhari, Proc. Indian Acad. Sci., Chem. Sci., 93(1984)65
- [206] V.T. Novikov, N.M. Novikova and L.A. Chernova, Koord. Khim., 10(1984)833
- [207] S.K. Gogia, O.V. Singh and S.N. Tandon, Indian J. Chem.A, 22(1983)965
- [208] F.P. Cavaiano, E. di Dio and C. Sbriziolo, Int. J. Chem. Kinet., 16(1984)221
- [209] Yu.A. Komkov, M.A. Ivanov, G.A. Shagisultanova and Yu.B. Yakovlev, Zh. Neorg. Khim., 28(1983)2834
- [210] Yu.A. Komkov, M.A. Ivanov, G.A. Shagisultanova and Yu.B. Yakovlev, Russ. J. Inorg. Chem., 29(1984)862

- [211] Yu. I. Sal'nikov, N.L. Kuz'mina and V.A. Zakharova, Russ. J. Inorg. Chem., 29(1984)1014
- [212] V.V. Pal'chevskii, V.V. Khorunzhii and V.I. Shcherbakova, Koord. Khim., 10(1984)1076
- [213] N.H. Williams and J.K. Yandell, Aust. J. Chem., 36(1983)2377
- [214] J.A. Kiernan, J. Microsc. (Oxford), 134(1984)13
- [215] T.V. Petrenko and V.T. Novikov, Koord. Khim., 10(1984)1203
- [216] J.P. Murray and J.O. Hill, Thermochim. Acta, 72(1984)341
- [217] C.W. Anderson, K.R. Lung and T.A. Nile, Inorg. Chim. Acta, 85(1984)33
- [218] U.M. Dzhemilev, R.I. Khusnutdinov, V.A. Dokichev, G.A. Tolstikov, S.R. Rafikov and O.M. Nefedov, Dokl. Akad. Nauk SSSR, 273(1983)887
- [219] S. Lunak, M. Vaskova and J. Veprek - Siska, Z. Naturforsch. B, 38(1983)1293
- [220] F.K. Schmidt, G.V. Ratovskii, T.V. Dmitrieva, I.N. Ivleva and Yu.G. Borodko, J. Organomet. Chem., 256(1983)309
- [221] D.A. Begum, Chem. Eng. Res. Bull. (Dacca), 6(1982)21
- [222] P.S. Belov, N.V. Borunova, V.M. Ignatov, B.Dilafrasa, V.A. Petukhov, I.L. Mironova and L.Kh. Freidlin, Neftekhimiya, 24(1984)185; CA, 101(1984)72236h
- [223] G.A. Molander, B.J. Rahn, D.C. Shubert and S.E. Bonde, Tetrahedron Lett., 24(1983)5449
- [224] A.M. Robov, V.S. Cherkashina, A.V. Fedorova, E.Yu. Belyaev and V.A. Fedorov, Koord. Khim., 10(1984)190
- [225] V.A. Ivanchenko, V.M. Nekipelov and K.I. Zamaraev, React. Kinet. Catal. Lett., 24(1984)347
- [226] T. Honjo and S. Shima, Bull. Chem. Soc. Jpn, 57(1984)293
- [227] A.A. Adimado, Polyhedron, 2(1983)1059
- [228] T.V. Petrenko and V.T. Novikov, Zh. Obshch. Khim., 53(1983)2516

- [229] A. Aguilar Navarro, M. Centrich Sureda and J.V. Vinade, An. Quim., Ser. A, 79(1983)354
- [230] D.S. Veselinovic and M.D. Jelakic, Glas. Hem. Drus. Beograd, 48(1983)739; CA, 100(1984)180981k
- [231] P. Painuly, S.V. Katti, S.K. Bajpai and J.S. Tandon, Indian J. Chem. A, 23(1984)166
- [232] L.G. Dainyak, A.S. Bukin and V.A. Drits, Kristallografiya, 29(1984)304, 312
- [233] I. Grunze and H. Grunze, Z. Anorg. Allg. Chem., 512(1984)39
- [234] E.A. Genkina, Yu.K. Kabalov, B.A. Maksimov and O.K. Mel'nikov, Kristallografiya, 29(1984)50
- [235] O.V. Yakubovich, M.A. Simonov and O.K. Mel'nikov, Kristallografiya, 29(1984)484
- [236] A. Osaka, M. Takahashi and M.I. Keda, J. Mater. Sci. Lett., 3(1984)36
- [237] J. Gruntze, W. Hilmer, N.N. Chudinova and H. Gruntze, Izv. Akad. Nauk SSSR, Neorg. Mater., 20(1984)287
- [238] R.A. Abduragimova and Kh. G. Gasanova, Zh. Obshch. Khim., 53(1983)2417
- [239] J. Gruntze, N.N. Chudinova and K.K. Palkina, Izv. Akad. Nauk SSSR, Neorg. Mater., 19(1983)1943
- [240] A.K. Molodkin, V.I. Emel'yanov, R. Murty, G.E. Ermolina and G.I. Dorofeeva, Zh. Neorg. Khim., 28(1983)3078
- [241] C. Haertig, P. Brand and K. Bohmhammel, Z. Anorg. Allg. Chem., 508(1984)159
- [242] P.C. Christidis, P.J. Rentzeperis, A. Kirfel and G. Will, Z.Kristallogr., 164(1983)219
- [243] S.M. Swamy, R. Nayak and P.T. Prasad, J. Chem. Technol. Biotechnol. Chem. Technol. A, 33(1983)171
- [244] Z. Xu, F. Liu, M. Wang and H. Huang, Jilin Daxue Ziran Kexue Xuebao, (1983)63; CA, 100(1984)85006n
- [245] N.K. Graham, J.B. Gill and D.C. Goodall, Aust. J. Chem., 36(1983)1991

- [246] K.A.R. Salib and R.A. Bucher, Transition Met. Chem., 8(1983)360
- [247] C.C. Torardi, J.C. Calabrese, K. Lazar and W.M. Reiff, J. Solid State Chem., 51(1984)376
- [248] V.I. Krupenskii, Izv. Vyssh. Uchebn. Zaved., Lesn. Zh., (1984)122; CA, 100(1984)145875g
- [249] M. Tonkovic, O. Hadzija and I. Nagy - Czako, Inorg. Chim. Acta, 80(1983)251
- [250] A. Terron and V. Moreno, Inorg. Chim. Acta, 80(1983)213
- [251] M.S. Masoud, M.S. Tawfik and S.E. Zayan, Synth. React. Inorg. Met. - Org. Chem., 14(1984)1
- [252] M. Birus, N. Kujundzic, M. Pribanic and Z. Tabor, Croat. Chem. Acta, 57(1984)313
- [253] B. Hutchinson, S. Sample, L. Thompson, S. Olbricht, J. Crowder, D. Hurley, D. Eversdyk, D. Jett and J. Bostik, Inorg. Chim. Acta, 74(1983)29
- [254] P.N. Buev, N.I. Pechurova and E.G. Afonin, Vestn. Mosk. Univ., Ser. 2: Khim., 24(1983)377
- [255] E. Graf, J.R. Mahoney, R.G. Bryant and J.W. Eaton, J. Biol. Chem., 259(1984)3620
- [256] S. Tamagaki, K. Suzuki, H. Okamoto and W. Tagaki, Tetrahedron Lett., 24(1983)4847
- [257] B.I. Khorunzhii, S.M. Mot'ko and V.V. Denisov, Izv. Sev.-Kavk. Nauchn. Tsentra Vyssh. Shk., Estestv. Nauki, (1983)77
- [258] A.E. Landers and D.J. Phillips, Inorg. Chim. Acta, 74(1983)43
- [259] T.S. Lobana, H.S. Cheema and S.S. Sandhu, Transition Met. Chem., 9(1984)330
- [260] V.I. Ponomarev, L.D. Arutyunyan and L.O. Atovmyan, Kristallografiya, 29(1984)910; CA, 101(1984)283420r
- [261] O.A. Raevskii, A.F. Solotnov, A.S. Shtepanek and T.N. Kudrya, Izv. Akad. Nauk SSSR, Ser. Khim., (1984)797; CA, 101(1984)47609z

- [262] [a] M. Varma, R.S. Varma and M.R. Parthasarathy, J. Prakt. Chem., 325(1983)382; [b] M. Perdicakis and J. Bessiere, C.R. Acad. Sci., Ser. 2, 298(1984)199
- [263] M.A. Salam, Pak. J. Sci. Ind. Res., 26(1983)20
- [264] A. Mauger, M. Escorne, C.A. Taft, N.C. Furtado, Z.P. Arguello and T. Parsenio, Phys. Rev. B, 30(1984)5300
- [265] K. Kurosawa, S. Saito and Y. Yamaguchi, J. Phys. Soc. Jpn, 52(1983)3919
- [266] M. Millar, S.A. Koch and R. Fikar, Inorg. Chim. Acta, 88(1984)L15
- [267] N.K. Roy and C.R. Saha, Indian J. Chem. A, 23(1984)484
- [268] N.K. Kaushik, G.R. Chhatwal and A.K. Sharma, J. Thermal Anal., 26(1983)309
- [269] K. Chandrasekar and H. B. Buerger, Acta Cryst. B, 40(1984)387
- [270] K. Staahl, Inorg. Chim. Acta, 75(1983)85
- [271] S. Wajda, K. Drabent and A. Ozarowski, Inorg. Chim. Acta, 45(1980)L201
- [272] K. Staahl and I. Ymen, Acta Chem. Scand. A, 37(1983)729
- [273] P.C. Christidis, P.J. Rentzeperis, F. Vakoulis and C.A. Tsipis, Inorg. Chim. Acta, 83(1984)87
- [274] M. Yoshikawa, N. Sorai, H. Suga and S. Seki, J. Phys. Chem. Solids, 44(1983)975
- [275] C.A. Tsipis, M.P. Sigalas and C.C. Hadjikostas, Z. Anorg. Allg. Chem., 505(1983)53
- [276] B. Singh and H.N. Tiwari, J. Indian Chem. Soc., 60(1983)526
- [277] K.L. Madhok, Polyhedron, 3(1984)39
- [278] R. Micu - Semeniciu, S. Popse and I. Haiduc, Rev. Roum. Chim., 28(1983)605
- [279] M.G. Gabdullin, A.R. Garifzyanov and V.F. Toropova, Russ. J. Inorg. Chem., 29(1984)1384; Y. Calage, M. Leblanc, G. Ferey and F. Varret, J. Magn. Magn. Mater., 43(1984)195

- [280] M. Eibschuetz, M.E. Lines and L.G. van Ultert, Phys. Rev. B, 29(1984)3843
- [281] G. Gevers, A.S. Barriere, J. Grannec, L. Lozano and B. Blanchard, Phys. Status Solidi A, 81(1984)105
- [282] J. Slivka, H. Keller, W. Kuendig and B.M. Wanklyn, Phys. Rev. B, 30(1984)3649
- [283] H. Baumeler, H. Keller, W. Kuendig, I.M. Savic and B.M. Wanklyn, Helv. Phys. Acta, 57(1984)101
- [284] K. Broomfield and P.M.A. Sherwood, J. Chem. Soc. Faraday Trans. 2, 79(1983)1321
- [285] G. Courbion, R. de Pape, J. Teillet, F. Varret and J. Pannetier, J. Magn. Magn. Mater., 42(1984)217
- [286] R.E. Marsh, J. Solid State Chem., 51(1984)405
- [287] P. Liu, F. Varret, A. de Kozak, M. Samouel, M. Leblanc and G. Ferey, Solid State Commun., 48(1983)875
- [288] E. Hardtweck, Z. Anorg. Allg. Chem., 501(1983)131
- [289] M. Leblanc, G. Ferey, Y. Calage and R. de Pape, J. Solid State Chem., 53(1984)360
- [290] E.G. Lavut, B.I. Timofeyev and V.M. Yuldasheva, J. Chem. Thermodyn., 16(1984)101
- [291] G.I. Novikov, I.A. Belov, P.K. Rud'ko and G.D. Poleshko, Vesti Akad. Nauk BSSR, Ser. Khim. Nauk, (1984)113; CA, 100(1984)198628k
- [292] R. Schloegl, W. Jones and H.P. Boehm, Synth. Met., 7(1983)133
- [293] R. Schloegl and W. Jones, J. Chem. Soc. Dalton Trans., (1984)1283
- [294] S.E. Millman and G. Kirczenow, Phys. Rev. B, 28(1983)5019
- [295] H. Schaefer - Stahl, Synth. Met., 8(1983)61
- [296] C. Simon, F. Batallan, I. Rosenman, J. Schweitzer, H. Lauter and R. Vangelisti, Synth. Met., 8(1983)53
- [297] C. Simon, F. Batallan, I. Rosenman, J. Schweitzer, H. Lauter



- and R. Vangelisti, J. Appl. Phys., 55(1984)2441
- [298] R. Yazami, P. Touzain and L. Bonnetain, Synth. Met., 7(1983)169
- [299] P. Touzain, J. Michel and P. Blum, Synth. Met., 8(1983)313
- [300] R. Schloegl, W. Jones, H.P. Boehm and W. Breimeier, Synth. Met., 8(1983)323
- [301] V.L. Solozhenko, I.V. Arkhangel'skii, Ya.A. Kalashnikov and N.A. Chernova, J. Thermal Anal., 28(1983)147
- [302] M. Suzuki, I. Oguro and Y. Jinzaki, J. Phys. C, 17(1984)L575
- [303] J.H. Choy and A. Weiss, Bull. Korean Chem. Soc., 4(1983)262
- [304] G.A. Fatseas, P. Paladeau and J.P. Venien, J. Solid State Chem., 51(1984)17
- [305] A. Weiss and J.H. Choy, Z. Naturforsch. B, 39(1984)1193
- [306] H. Eckert and R. H. Herber, J. Chem. Phys., 80(1984)4526
- [307] S. Kikkawa, F. Kanamaru and M. Koizumi, Inorg. Synth., 22(1983)86
- [308] Z. Zhang, F. Wang and S. Li, Huaxue Xuebao, 42(1984)650; CA, 101(1984)142727v
- [309] M.G. Meirim and E.W. Neuse, Transition Met. Chem., 9(1984)337
- [310] G.A. Bottomley, A.M. Carter, L.M. Engelhardt, F.J. Lincoln, J.M. Patrick and A.H. White, Aust. J. Chem., 37(1984)871
- [311] P. Sobota, T. Pluzinski and T. Lis, Polyhedron, 3(1984)45
- [312] M. Speisser, P. Palvadeau, C. Guillot and J. Cerisier, Solid State Ionics, 9-10(1983)103
- [313] P.J. McCarthy and I.M. Walker, Spectrochim. Acta A, 39(1983)827
- [314] M.M. Yunusov, G.M. Larin and P.M. Solozhenkin, Dokl. Akad. Nauk Tadzh. SSR, 26(1983)436; CA, 100(1984)109889f
- [315] P. Sartori and F.M. Helmy, Chem. - Ztg., 108(1984)149
- [316] A. Radde and L. Kolditz, Z. Chem., 23(1983)436

- [317] S. Pohl and W. Saak, Z. Naturforsch. B, 39(1984) 1236
- [318] G. Algra and S. Balt, Inorg. Chim. Acta, 75(1983)179
- [319] G.P. Algra and S. Balt, Inorg. Chim. Acta, 83(1984)191
- [320] I.S. El - Yamani and El - Said I. Shabana, Transition Met. Chem., 9(1984)199
- [321] A. Ohki, H. Rinoshita, M. Tagaki and K. Ueno, Sep. Sci. Technol., 18(1983)969
- [322] J.C. Jochims, R. Abu - El - Halawa, L. Zsolnai and G. Huttner, Chem. Ber., 117(1984)1161
- [323] S.P. Srinavasan and N.S. Gnanapragasam, J. Indian Chem. Soc., 60(1983)1104
- [324] A.D. Jenkins and B.H. Mustafa, Eur. Polym. J., 19(1983)1009
- [325] K. Oka, H. Maeda and S. Ikeda, Bull. Chem. Soc. Jpn., 57(1984)696
- [326] S.M. Brailovskii, O.M. Temkin and E.S. Klimova, Kinet. Katal., 24(1983)1091
- [327] Yu.V. Pozdnyakovich, R.P. Savyak and S. M. Shein, Zh. Org. Khim., 19(1983)1674; CA, 100(1984)5949j
- [328] J. Yamamoto, T. Inohara, T. Tamura, Y. Michikawa and M. Umezo, Nippon Kagaku Kaishi, (1983)1629; CA, 100(1984)191464v
- [329] Y.H. Kuo and S.T. Lin, Experientia, 39(1983)991
- [330] A.N. Panasencko and N.N. Dykhanov, Zh. Org. Khim., 20(1984)1115; CA, 101(1984)90540g
- [331] J.P. Albertini and A. Garnier - Suillerot, Biochemistry, 23(1984)47
- [332] E. Anachkova and I. Savatinova, Spectrosc. Lett., 16(1983)879
- [333] I. Savatinova and E. Anachkova, J. Mol. Struct., 114(1984)89
- [334] E. Anachkova, I. Savatinova and M. Szostak, J. Mol. Struct., 115(1984)83
- [335] A.S.N. Murthy and A.P. Bhardwaj, Spectrochim. Acta A,

- 40(1984)113
- [336] H.D. Rubin, B.D. Humphrey and A.B. Bocarsly, Nature, 308(1984)339
- [337] T.M. Finogenko, I.D. Isaev, V.A. Fedorov and V.E. Mironov, Russ. J. Inorg. Chem., 29(1984)429
- [338] O. Zakharieva - Pencheva, V.A. Dement'ev and H. Foerster, J. Mol. Struct., 112(1984)273
- [339] O. Zakharieva - Pencheva and V.A. Dement'ev, J. Mol. Struct., 115(1984)75
- [340] V.S. Badik and D.I. Semenishin, Koord. Khim., 9(1983)1241
- [341] K. Gyoryova and B. Mohai, Thermochim. Acta, 71(1983)167
- [342] A.N. Sergeeva, V.V. Dovgei, L.I. Pavlenko, D.I. Zubritskaya, Zh.I. Tkachenko, A.P. Okorskaya and Yu.A. Lyubchenko, Kinet. Katal., 24(1983)1473; CA, 100(1984)109677k
- [343] V.K. Tiwari, Natl. Acad. Sci. Lett. (India), 6(1983)159
- [344] S.A.D. Naman, A.H. Ibrahim and A.A. Matti, Int. J. Hydrogen Energy, 9(1984)405
- [345] W. Beck, W. Weigand, U. Nagel and M. Schaal, Angew. Chem. Int. Ed. Engl., 23(1984)377
- [346] H.E. Toma and M.S. Takasugi, J. Solution Chem., 12(1983)547
- [347] M.J. Blandamer, J. Burgess, K.W. Morcom and R. Sherry, Transition Met. Chem., 8(1983)354
- [348] S. Sinha, B.D. Humphrey, E. Fu and A.B. Bocarsly, J. Electroanal. Chem. Interfacial Electrochem., 162(1984)351
- [349] A.B. Altabef, D.B. Soria and N.E. Katz, An. Asoc. Quim. Argent., 71(1983)547; CA, 101(1984)44048e; S. Bagger and P. Stoltze, Acta Chem. Scand. A, 37(1983)247
- [350] A. Neves, W. Herrmann and K. Wieghardt, Inorg. Chem., 23(1984)3435
- [351] J. Metz, O. Schneider and M. Hanack, Inorg. Chem., 23(1984)1065
- [352] B. Simic - Glavaski, S. Zecevic and E.B. Yeager, J. Raman

Spectrosc., 14(1983)338

- [353] L. Kreja and F. Rozploch, Angew. Makromol. Chem., 121(1984)151
- [354] P. Coppens and L. Li, J. Chem. Phys., 81(1984)1983
- [355] A. Labarta, E. Molins, X. Vinas, J. Tajeda, A. Caubet and S. Alvarez, J. Chem. Phys., 80(1984)444
- [356] C.A. Melendres and S.Q. Xu, J. Electrochem. Soc., 131(1984)466
- [357] M. Yamana, R. Darby and R.E. White, Electrochim. Acta, 29(1984)329
- [358] M.A. Abd El - Ghaffar, M.A. Ahmed and M.S. Rizk, J. Indian Chem. Soc., 60(1983)550
- [359] L. Kreja, Electrochim. Acta, 28(1983)1807
- [360] E. Molins, A. Labarta, J. Tajeda, A. Caubet and S. Alvarez, Transition Met. Chem., 8(1983)377
- [361] O. Schneider and M. Hanack, Z. Naturforsch. B, 39(1984)265
- [362] A. Datz, J. Metz, O. Schneider and M. Hanack, Synth. Met., 9(1984)31
- [363] N.K. Singh, S.C. Srivastva and R.C. Aggarwal, Indian J. Chem. A, 22(1983)704
- [364] V.M. Pinchuk, V.A. Korobskii and L.B. Shevardina, Russ. J. Inorg. Chem., 29(1984)637
- [365] G. Vos, R.A.G. de Graaff, J.G. Haasnoot, A.M. van der Kraan, P. de Vaal and J. Reedijk, Inorg. Chem., 23(1984)2905
- [366] S. Descurtins, P. Guetlich, C.P. Koehler, H. Spiering and A. Hauser, Chem. Phys. Lett., 105(1984)1
- [367] M.R. Chaurasia and P. Shukla, J. Indian Chem. Soc., 60(1983)1011
- [368] K.S. Arulsamy, R.F.N. Ashok and U.C. Agarwala, Indian J. Chem. A, 23(1984)21
- [369] H.E. Toma, F. Gebara Filho, A.A. Batista and E. Giesbrecht, An. Acad. Bras. Cienc., 56(1984)57
- [370] A.T. Baker and H.A. Goodwin, Aust. J. Chem., 37(1984)1157

- [371] H.H. Wei and Y.C. Jean, Chem. Phys. Lett., 106(1984)523
- [372] A. El - Dissouky and L.S. Refaat, Inorg. Chim. Acta, 87(1984)213
- [373] J.A. Real and J. Borrás, Synth. React. Inorg. Met. - Org. Chem., 14(1984)857
- [374] G.D. Tiwari and M.N. Mishra, J. Indian Chem. Soc., 60(1983)698
- [375] I. Havlik, T. Simionescu, I. Havlik, A. Maurer and S. Policec, Bull. Stiint. Teh. Inst. Politeh. "Trian Vuia" Timisoara, Ser. Chim., 28(1983)93; CA, 101(1984)102930z
- [376] T.C. Streckas and S.K. Mandal, J. Raman Spectrosc., 15(1984)109
- [377] R.P. Van Duyne and M. Janik - Czachor, J. Electrochem. Soc., 130(1983)2320
- [378] A.T. Fal'kengof, V.A. Bogdanovskaya, M.R. Tarasevich, V.S. Pervov and Yu.A. Buslaev, Dokl. Akad. Nauk SSSR, 275(1984)388
- [379] S. Tachiyashiki and H. Yamatera, Bull. Chem. Soc. Jpn., 57(1984)1061
- [380] M.J. Blandamer, J. Burgess, S.D. Cope and T. Digman, Transition Met. Chem., 9(1984)347
- [381] F. Ortega and E. Rodenas, Transition Met. Chem., 9(1984)331
- [382] M.J. Blandamer, J. Burgess, P.P. Duce, K.S. Payne, R. Sherry, P. Wellings and M.V. Twigg, Transition Met. Chem., 9(1984)163
- [383] M.J. Blandamer, J. Burgess, B. Clark, P.P. Duce and J.M.W. Scott, J. Chem. Soc. Faraday Trans. 1, 80(1984)739
- [384] L.P. Tikhonova, K.B. Yatsimirskii and V.Ya. Zayats, Teor. Eksp. Khim., 20(1984)317; CA, 101(1984)82996v
- [385] A.K. Pyartman, N.N. Snegirev and V.E. Mironov, Russ. J. Inorg. Chem., 29(1984)431
- [386] S.K. Kulshreshtha and R.M. Iyer, Chem. Phys. Lett., 108(1984)501
- [387] B. Maiti and B.R. McGarvey, J. Magn. Reson., 58(1984)37
- [388] E. Koenig, G. Ritter, S.K. Kulshreshtha and N. Csatory, Inorg.

Chem., 23(1984)1903

- [389] E. Koenig, G. Ritter, S.K. Kulshreshtha, J. Waigel and H.A. Goodwin, Inorg. Chem., 23(1984)1896
- [390] S.K. Kulshreshtha, R.M. Iyer, E. Koenig and G. Ritter, Chem. Phys. Lett., 110(1984)201
- [391] A.T. Kowal and J. Skarzewski, Monatsh. Chem., 115(1984)953
- [392] W. Spahni and G. Calzaferri, Helv. Chim. Acta, 67(1984)450
- [393] A.S. Abushamleh, H.A. Goodwin, C.G. Benson and G.J. Long, Aust. J. Chem., 37(1984)281
- [394] A.T. Baker, H.A. Goodwin and A.D. Rae, Aust. J. Chem., 37(1984)443
- [395] A.A. Schilt and S.W. Wong, J. Coord. Chem., 13(1984)331
- [396] J.C. Thompson and H.A. Mottola, Anal. Chem., 56(1984)755
- [397] K.S. Menon and Y.K. Agrawal, Transition Met. Chem., 8(1983)292
- [398] R. Cini, Inorg. Chim. Acta, 73(1983)147
- [399] L. Casella, M.E. Silver and J.A. Ibers, Inorg. Chem., 23(1984)1409
- [400] C. Ruiz - Valero, A. Monge, E. Gutierrez - Puebla and E. Gutierrez - Rios, Acta Crystallogr. C, 40(1984)811
- [401] L.A. Morino and H.E. Toma, Talanta, 31(1984)224
- [402] A.Yu. Nazarenko and Ya.Z. Voloshin, Russ. J. Inorg. Chem., 29(1984)1019
- [403] S. Chandra and K.K. Sharma, Acta Chim. Hung., 115(1984)9
- [404] D.P. Singh, V.K. Chauhan, V.B. Rana and M.P. Teota, J. Coord. Chem., 13(1984)159
- [405] K.J. Charyulu, P. Ettiyah, K.L. Omprakash, A.V.C. Pal and M.L.N. Reddy, Indian J. Chem. A, 23(1984)668
- [406] S.B. Adekeye, E.O. Erinoso and B.N. Ghose, An. Chim. Ser. B, 79(1983)353
- [407] R.C Aggarwal, N.K. Singh and R.P. Singh, Synth. React. Inorg.

- Met. - Org. Chem., 14(1984)637
- [408] K.S. Arulsamy, R.F.N.Ashok and U.C.Agarwala, Indian J. Chem. A, 23(1984)122
- [409] I.I. Kalinchenko, N.M. Titov and M.G. Ivanov, Russ. J. Inorg. Chem., 29(1984)1149
- [410] Y. Kumar, S. Chandra, R.P. Singh and A.P. Singh, Synth. React. Inorg. Met. - Org. Chem., 14(1984)185
- [411] C. Natarajan and A.N. Hussain, Transition Met. Chem., 9(1984)18
- [412] M.A. Romero Molina, J.M. Salas Peregrin, J.D. Lopez Gonzalez and C. Valenzuela Calahorro, An. Quim. Ser. B, 79(1983)383; CA, 101(1984)203216b
- [413] B. Singh, R.N. Singh and R.C. Aggarwal, Indian J. Chem. A, 23(1984)480
- [414] N.K. Singh, S.C. Srivastava and R.C. Aggarwal, J. Indian Chem. Soc., 60(1983)622
- [415] N.K. Singh, S.C. Srivastava and R.C. Aggarwal, Proc. Indian Acad. Sci. Chem. Sci. A, 92(1983)173
- [416] P. Zanello, R. Cini and A. Cingantini, Inorg. Chim. Acta, 74(1983)89
- [417] S. Komiya, M. Katoh, T. Ikariya, R.H. Grubbs, T. Yamamoto and A. Yamamoto, J. Organomet. Chem., 260(1984)115
- [418] H.H. Karsch, Chem. Ber., 117(1984)783
- [419] Y. Endo, S. Saito and E. Hirota, Astrophys. J., 278(1984)L131
- [420] A.S.C. Cheung, A.M. Lyyra, A.J. Merer and A.W. Taylor, J. Mol. Spectrosc., 102(1983)224
- [421] M.S. Seehra and G. Srinivasan, J. Phys. C, 17(1984)883
- [422] Z. Panek and K. Fitzner, Thermochim. Acta, 78(1984)261
- [423] A.A. Fotiev, V.G. Dobosh and V.L. Kozhevnikov, Russ. J. Inorg. Chem., 29(1984)456
- [424] E. Riedel and H. Anik, Z. Naturforsch. B, 38(1983)1630

- [425] M. Greenblatt, K.R. Nair, W.H. McCarroll and J.V. Wasczak, Mater. Res. Bull., 19(1984)777
- [426] J. Zhang, Y. Ni, Y. Xia and S. Qi, Kexue Tongbao (Foreign Lang. Ed.) 29(1984)53
- [427] N.B. Milic and P. Djurdjevic, Glas. Hem. Drus. Beograd., 49(1984)39; CA, 100(1984)216429c
- [428] A.V. El'tsov, N.V. Lebedeva and V.Yu. Kukushkin, Zh. Obshch. Khim., 53(1983)2752; CA, 100(1984)128650n
- [429] S.S. Sawhney, S.D. Matta, R. Jain, R.K. Kashyap and V.K. Painuli, Thermochim. Acta, 70(1983)367
- [430] T. Yamamoto, K. Imaizumi and Y. Kurata, Inorg. Chim. Acta, 85(1984)175
- [431] R.N.K. Vishwanatham, K. Ram and M.G.R. Reddy, J. Indian Chem. Soc., 61(1984)115
- [432] Ramani, M.P. Sathyavathiamma, N.G. Puttaswamy and R.M. Mallia, Mater. Chem. Phys., 9(1983)539
- [433] X. Wang, R. Wu, X. Xin, A. Dai and Y. Zhang, Fenxi Kexue Yu Huaxue Yanjiu, 3(1983)39
- [434] V.A. Sharov and V.A. Larikova, Russ. J. Inorg. Chem., 29(1984)140
- [435] T. M'hiri, A. Demaret and G. Lapluye, J. Chem. Soc. Tunis, 9(1983)19
- [436] A.K. Jain, K.D. Jain and A.K. Ojha, Indian J. Phys. Nat. Sci., 4(1984)25
- [437] M. Asso, R. Panossian and M. Guiliano, Spectrosc. Lett., 17(1984)271
- [438] I. Ahmed, Arab Gulf J. Sci. Res., 1(1983)121
- [439] J.M. Lopez - Alcala, M.C. Puerta and F. Gonzalez - Vilchez, Polyhedron, 3(1984)623
- [440] J.M. Lopez - Alcala, M.C. Puerta and F. Gonzalez - Vilchez, Thermochim. Acta, 78(1984)135
- [441] E. Narita, T. Sato, T. Shioya, M. Ikari and T. Okabe, Ind. Eng. Chem. Prod. Res. Dev., 23(1984)262



- [442] E. Sada, H. Kumazawa and Y. Takada, Ind. Eng. Chem. Fundam., 23(1984)60
- [443] R.P. Hertzberg and P.B. Dervan, Biochemistry, 23(1984)3934
- [444] Zh.N. Bublik, S.V. Volkov and E.A. Mazurenko, Russ. J. Inorg. Chem., 29(1984)73
- [445] N.A. Bailey, D.E. Fenton and M.S. Leal Gonzalez, Inorg. Chim. Acta, 88(1984)125
- [446] S.M. Peng, J.F. Liu, Y. Wang and C.P. Tang, Bull. Inst. Chem. Acad. Sin., 30(1983)35
- [447] M.S.R. Swamy and T.P. Prasad, J. Therm. Anal., 25(1982)347
- [448] M.S.R. Swamy and T.P. Prasad, J. Therm. Anal., 25(1982)355
- [449] I. Ahmadm A.M.A. Rauf and K.M. Saeed, J. Prakt. Chem., 326(1984)23
- [450] D.St.C. Black and L.M. Johnstone, Aust. J. Chem., 37(1984)109
- [451] W.E. Fristad and J.R. Peterson, Tetrahedron Lett., 24(1983)4547
- [452] E.T. Gorodetskii and P.M. Gorbovoi, Kinet. Katal., 25(1984)764; CA, 101(1984)110449q
- [453] J.G. Santiestlban, P.B. Himelfarb, K. Klier, G.W. Simmons and R.G. Herman, Mater. Lett., 2(1984)344
- [454] S.A.A. Zaidi, M. Shakir, M. Aslam, S.R.A. Zaidi and Z.A. Siddiqui, Indian J. Chem.A, 22(1983)1078
- [455] S.K. Sharma, R.K. Mahajan, B. Kapila and V.P. Kapila, Polyhedron, 2(1983)973
- [456] G. Candrini, W. malavasi, C. Preti, G. Tosi and P. Zennini, Spectrochim. Acta A, 39(1983)635
- [457] G. Venturini, A. Courtois, J. Steinmetz, R. Gerardin and C. Gleitzer, J. Solid State Chem., 53(1984)1
- [458] J.C. Kaell, F. Jeannot and C. Gleitzer, Ann. Chim.(Paris), 9(1984)169
- [459] P. Ghosh, T.K. Mukhopadhyay and A.R. Sarkar, Transition Met.

Chem., 9(1984)46

- [460] M.R. Gandhi and K.N. Munshi, J. Indian Chem. Soc., 59(1982)1290
- [461] L.A. Dominey and K. Kustin, Inorg. Chem., 23(1984)103
- [462] V.V. Skopenko, N.N. Gerasimchuk and E.V. Pol'shin, Dopov. Akad. Nauk Ukr. RSR, B, (1983)39
- [463] R. Doerfler, O. Leupold, D.L. Nagy, G. Ritter, H. Spiering and R. Zimmermann, Hyperfine Interact., 16(1983)831
- [464] P. Orlandi and A. Rigamonti, Phys. Status Solidi, 82(1984)K129
- [465] C. Lupiani and M. Castineira, An. Quim. Ser. A, 80(1984)72
- [466] L. Vincze, B. Kraut and S. Papp, Inorg. Chim. Acta, 85(1984)89
- [467] H. Tamura, K. Sato and M. Nagayama, Nippon Kagaku Kaishi, (1983)1405
- [468] W. Linert, V. Gutmann, O. Baumgartner, G. Wiesinger and H. Kirchmayr, Inorg. Chim. Acta, 74(1983)123
- [469] G. Mennenga, L.J. de Jongh, W.J. Huiskamp and J. Reedijk, J. Magn. Magn. Mater., 43(1984)13
- [470] A.A. Salman, I.M. Abd - Ellah, M.A. Abd El - Ghafar, A.N. Khouzondar and A.M. Naser, J. Chem. Soc. Pak., 5(1983)29
- [471] C.A.L. Filgueiras, Transition Met. Chem., 9(1984)158
- [472] V.A. Maroni, C.E. Philbin and R.M. Yonco, Sep. Sci. Technol., 18(1983)1699
- [473] S.I. Shustova, K.I. Tikhonov and I.A. Bud'ko, Zh. Prikl. Khim. (Leningrad), 57(1984)1617; CA, 101(1984)139645m
- [474] W.E. Aker, N. Margalit, D.P. Johnson, R.J. Ekern, N.A. Fleischer and R.J. Brodd, J. Electrochem. Soc., 131(1984)1839
- [475] Q. Huang and X. Fu, Huaxue Xuebao, 42(1984)130; CA, 100(1984)150099m
- [476] C. Sugiura, J. Chem. Phys., 80(1984)1047
- [477] R.A. Sharma and R.N. Seefurth, [a] J. Electrochem. Soc., 131(1984)1084; [b] Proc. Electrochem. Soc., 84-2(1984)498

- [478] K. Sato, J. Phys. Soc. Japan, 53(1984)1617
- [479] S. Lauer, A.X. Trautwein and F.E. Harris, Phys. Rev. B, 29(1984)6774
- [480] H. Vogt, T. Chattopadhyay and H.J. Stolz, J. Phys. Chem. Solids, 44(1983)869
- [481] W. Jaegermann and H. Tributsch, J. Appl. Electrochem., 13(1983)743
- [482] D.M. Pasquariello, R. Kershaw, K. Dwight and A. Wold, Prepr. Am. Chem. Soc. Div. Pet. Chem., 28(1983)1255
- [483] T. Kanomata, S. Haga, K. Shirakawa and T. Kaneko, Tohoku Gakuin Daigaku Kogakubu Kenkyu Hokoku, 18(1983)65; CA, 100(1984)131072f
- [484] N.G. Guseinov and A.I. Dzhabbarov, Fiz. Met. Metalloved., 56(1983)814
- [485] R.Z. Sadykhov, L.M. Valiev, D.A. Guseinov and A.O. Ismailov, Phys. Status Solidi A, 79(1983)K147
- [486] S.K. Lie and C.A. Taft, Phys. Rev. B, 28(1983)7308
- [487] S. Pohl and W. Saak, Angew. Chem., 96(1984)886
- [488] J.W. McDonald, G.D. Friesen, W.E. Newton, A. Mueller, W. Hellmann, U. Schimanski, A. Trautwein and U. Bender, Inorg. Chim. Acta, 76(1983)L297
- [489] W. Clegg, C.D. Garner and P.A. Fletcher, Acta Crystallogr. C, 40(1984)754
- [490] L.D. Rosenhein, J.W. McDonald and W.E. Newton, Inorg. Chim. Acta, 87(1984)L33
- [491] K.S. Arulsamy, R.F.N. Ashok and U.C. Agarawala, Indian J. Chem. A, 23(1984)127
- [492] X. Lin and J. Huang, Youji Huaxue, (1983)426; CA, 100(1984)202357r
- [493] M.M. Olmstead, W.K. Musker and R.M. Kessler, Acta Crystallogr. C, 40(1984)1172
- [494] N.K. Singh, S.C. Srivastava and R.C. Aggarwal, Indian J.

Chem., 23(1984)664

- [495] Z. Zhang, F. Wang, S. Li, C. Zhao and Y. Li, Gaodeng Xueiao Huaxue Xuebao, 5(1984)276; CA, 100(1984)220545e
- [496] R.M. Toussaint and V. Jaccarino, J. Appl. Phys., 55(1984)2458
- [497] E.F. da Silva, F.L.A. Machado and S.M. Rezende, Solid State Commun., 48(1983)1077
- [498] A.S. Gouveia Neto and C.B. de Araujo, Phys. Rev. B, 28(1983)6532
- [499] W.A.H.M. Vlak, E. Frikkie, A.F.M. Arts and H.W. de Wijn, J. Phys. C, 16(1983)L1015
- [500] S. Galindo, Phys. Rev. B, 29(1984)6369
- [501] H.W. Mayer, D. Reinen and G. Heger, J. Solid State Chem., 50(1983)213
- [502] M.C. Lin and R.G. Barnes, J. Appl. Phys., 55(1984)2294
- [503] M. Grade and W. Rosinger, Ber. Bunsen - Ges. Phys. Chem., 88(1984)767
- [504] D.E. Johnson and J.G. Eden, IEEE J. Quantum Electron., QE-18(1982)1836
- [505] M. Grade, W. Rosinger and P.A. Dowben, Ber. Bunsen - Ges. Phys. Chem., 88(1984)65
- [506] P.Z. Wong and J.W. Cable, Phys. Rev. B, 28(1983)5361
- [507] J. Gelard, F. Bensamka, D. Bertrand, A.R. Fert, J.P. Redoules and S. Legrand, J. Phys. C, 16(1983)L939
- [508] G.C. De Fotis, C. Pohl, S.A. Pugh and E. Sinn, J. Chem. Phys., 80(1984)2079
- [509] B.Y. Enwiya, J. Silver and I.E.G. Morrison, J. Chem. Soc. Dalton Trans., (1983)2581
- [510] B. Yenwiya, J. Silver and I.E.G. Morrison, Inorg. Chim. Acta, 86(1984)113
- [511] E.J. Nijhof, G.J. Gerritsma and J. Flokstra, Physica B+C,

- 122(1983)333
- [512] M.V. Diudea, Rev. Roum. Chim., 28(1983)249
- [513] A. Yoshihara, J.C. Burr and S.M. Mudare, J. Chem. Phys., 80(1984)3816
- [514] M.M. Abdel - Kader, Chem. Phys. Lett., 102(1983)69
- [515] M.A. Ahmed, Helv. Phys. Acta, 56(1983)1088
- [516] M. Yoshizawa, T. Suzuki, T. Goto, T. Yamakami, T. Fujimura, T. Nakajima and H. Yamauchi, J. Phys. Soc. Jpn. 53(1984)261
- [517] E. McCafferty, J. Electrochem. Soc., 131(1984)1842
- [518] M.J. Blandamer, J. Burgess, P.P. Duce, N. Gosal, R. Sherry, P. Guardado and F. Sanchez, Transition Met. Chem., 9(1984)3
- [519] M. Ichihashi, H. Wakka and I. Masuda, J. Solution Chem., 13(1984)505
- [520] W. Linert and V. Gutmann, Rev. Chim. Miner., 20(1983)516
- [521] V.K. Bel'skii, V.M. Ishchenko, B.M. Bulychev, A.N. Protskii, G.L. Soloveichik, O.G. Ellert, Z.M. Seifulina, Yu. V. Rakitin and V.M. Novotortsev, Izv. Akad. Nauk SSSR, Ser. Khim., (1984)954
- [522] V.M. Ishchenko, B.M. Bulychev and G.L. Soloveichik, Koord. Khim., 10(1984)998; CA, 101(1984)192134v
- [523] M. Pasternak, M. van der Heyden and G. Langouche, Chem. Phys. Lett., 104(1984)398
- [524] G.A. Ozin and J.G. McCaffrey, J. Phys. Chem., 88(1984)645
- [525] L.F. Halle, F.S. Klein and J.L. Beauchamp, J. Am. Chem. Soc., 106(1984)2543
- [526] J.J. Didisheim, P. Zolliker, K. Yvon, P. Fischer, J. Schefer, M. Gubelmann and A.F. Williams, Inorg. Chem., 23(1984)1953
- [527] P.B. Klein, J.E. Furneaux and R.L. Henry, Phys. Rev. B, 29(1984)1947
- [528] H.H. Wei and H.L. Shyu, Transition Met. Chem., 9(1984)211
- [529] K. Rachlewicz, J. Pietrzyk, K. Drabent and S. Wajda, Inorg.

Chim. Acta, 77(1983)L189

- [530] C. Dalvit, S. Miura, A. de Young, R.W. Noble, M. Cerdino and C. Ho, Eur. J. Biochem., 141(1984)255
- [531] A. Mayer and H. Eicher, J. Mol. Catal., 23(1984)151
- [532] L.C. Dickinson and M.C.R. Symons, Chem. Soc. Rev., 12(1983)387
- [533] P.V. Argade, M. Sassardi, D.L. Rousseau, T. Inubishi, M. Ikeda - Saito and A. Lapidot, J. Am. Chem. Soc., 106(1984)6593
- [534] O. Bangcharoenpaupong, K.T. Schomacker and P.M. Champion, J. Am. Chem. Soc., 106(1984)5688
- [535] D.L. Rousseau, S.L. Tan, M.R. Ondrias, S. Ogawa and R.W. Noble, Biochemistry, 23(1984)2857
- [536] R. Schweitzer - Stenner, W. Dreybrodt and S. El Naggar, Biophys. Struct. Mech., 10(1984)241
- [537] S. El Naggar, R. Schweitzer - Stenner, W. Dreybrodt and A. Mayer, Biophys. Struct. Mech., 10(1984)257
- [538] A. Desbois, G. Mazza, F. Stetzkowski and M. Lutz, Biochim. Biophys. Acta, 785(1984)161
- [539] M.R. Ondrias, E.W. Findsen, D.F. Gaul and D.B. Knaff, Biochim. Biophys. Acta, 788(1984)87
- [540] A. Levy, K. Alston and J.M. Rifkind, J. Biomol. Struct. Dyn., 1(1984)1299
- [541] H. Roder, J. Berendzen, S.F. Bowne, H. Frauenfelder, T.B. Sauke, E. Shyamsunder and M.B. Weissman, Proc. Natl. Acad. Sci. USA, 81(1984)2359
- [542] J.K. Schreiber and L.J. Parkhurst, Comp. Biochem. Physiol., 78(1984)129
- [543] L.J. Groome and S.D. Groome, J. Theor. Biol., 109(1984)157
- [544] C.A. Sawacki and M.A. Khaleque, Biophys. J., 44(1983)191
- [545] L. Cordone, A. Cupane, F. D'Alia and M.G. de Stefano, Faraday Symp. Chem. Soc., (1982)161
- [546] A. Brzowzki, Z. Derewenda, E. Dodson, G. Dodson, M. Grabowski, R. Liddington, T. Skarzynski and D. Vallely, Nature,

307(1984)74

- [547] S.J. Gill and B. Richey, Nature, 310(1984)160
- [548] [a] B.R. Gelin, A.W.M. Lee and M. Karplus, J. Mol. Biol., 171(1983)489; [b] M.L. Johnson, B.W. Turner and G.K. Ackers, Proc. Natl. Acad. Sci. USA, 81(1984)1093
- [549] M.S. Roberts, R.C. Terwilliger and N.B. Terwilliger, Comp. Biochem. Physiol. B: Comp. Biochem., 77(1984)237
- [550] R. Schweitzer - Stenner, W. Dreybrodt, D. Wedekind and S. El Nagggar, Eur. Biophys. J., 11(1984)61
- [551] J.P. Savicki, G. Lang and M. Ikeda - Saito, Proc. Natl. Acad. Sci. USA, 81(1984)5417
- [552] E. Antonini, F. Ascoli, M. Brunori, E. Chiancone, D. Verzili, R.J. Morris and Q.H. Gibson, J. Biol. Chem., 259(1984)6730
- [553] R.J. Morris and Q.H. Gibson, J. Biol. Chem., 259(1984)365
- [554] R.J. Morris, Q.H. Gibson, M. Ikeda - Saito and T. Yonetani, J. Biol. Chem., 259(1984)6701
- [555] P.A. Adams, C. Adams, M.C. Berman and M.C. Lawrence, J. Inorg. Biochem., 20(1984)291
- [556] E. Tsuchida, H. Nishide, H. Yokoyama, H. Inoue and T. Shirai, Polym. J. (Tokyo), 16(1984)325
- [557] C.K. Chang, B. Ward and S. Ebina, Arch. Biochem. Biophys., 231(1984)366
- [558] N.T. Yu, B. Benko, E.A. Kerr and K. Gersonde, Proc. Natl. Acad. Sci. USA, 81(1984)5106
- [559] D.D. Dlott, H. Fraunfelder, P. Langer, H. Roder and E.E. Di Iorio, Proc. Natl. Acad. Sci. USA, 80(1983)6239
- [560] R.D. Young, J. Chem. Phys., 80(1984)554
- [561] T.G. Traylor, L.A. Dearduff, M. Coletta, P. Ascenzi, E. Antonini and M. Brunori, J. Biol. Chem., 258(1983)12147
- [562] D.M. Mills, A. Lewis, A. Harootunian, J. Huang and B. Smith, Science, 223(1984)811
- [563] H. Tomiyasu, S. Sekii, K. Nishi, H. Ishii and H. Fukutomi,

- Bull. Res. Lab. Nucl. React. (Tokyo Inst. Technol.), 9(1984)29
- [564] N.V. Blough and B.M. Hoffman, Biochemistry, 23(1984)2875; N.V. Blough, H. Zemel and B.M. Hoffman, Biochemistry, 23(1984)2883
- [565] D.C. Doetschman, A.K. Rizos and J. Szumowski, J. Chem. Phys., 81(1984)1185
- [566] B. Benko and N.T. Yu, Proc. Natl. Acad. Sci. USA, 80(1983)7042
- [567] I. Morishima and M. Hara, Biochem. Biophys. Res. Commun., 121(1984)229
- [568] K.D. Whitburn, M.Z. Hoffman and I.A. Taub, Radiat. Phys. Chem., 23(1984)271
- [569] W.N. Aldridge and B.W. Street, Toxicol. Appl. Pharmacol., 73(1984)350
- [570] J. Steczko, B. Axelrod and M. Hermodson, Arch. Biochem. Biophys., 232(1984)597
- [571] M.P. Doyle, S.N. Mahapatro and S.V. Tran, Inorg. Chim. Acta, 92(1984)123
- [572] T. Yamaguchi, T. Nakano and E. Kimoto, Biochem. Biophys. Res. Commun., 120(1984)534
- [573] F. Jung and M. Spolaczyk, Biomed. Biochim. Acta, 42(1983)149
- [574] M. Takeda, K. Aoyagi, Y. Aoyama and H. Ogoshi, Chem. Lett., (1983)1393
- [575] R. Timkovich, M.S. Cork and P.V. Taylor, J. Biol. Chem., 259(1984)1577
- [576] V.G. Artyukhov, Zh. Prikl. Spectrosk., 41(1984)87; CA, 101(1984)106021b
- [577] L.A. Eguchi and P. Saltman, J. Biol. Chem., 259(1984)14337
- [578] M.R. Mauk, L.S. Reid and A.G. Mauk, Biochem. J., 221(1984)297
- [579] R.F. Tilton, I.D. Kuntz and G.A. Petsko, Biochemistry, 23(1984)2849
- [580] G.P. Singh, H.J. Schink, H. von Loehneysen, F. Parak and S. Hunklinger, Z. Phys. B, 55(1984)23



- [581] G. Bemski, M. Novak and O.G. Symko, Phys. Lett. A, 99(1983)62
- [582] J.H. Bradbury and J.A. Carver, Biochemistry, 23(1984)4905
- [583] J.A. Carver and J.H. Bradbury, Biochemistry, 23(1984)4890
- [584] G.N. La Mar and R. Krishnamoorthi, Biophys. J., 44(1983)177
- [585] R. Krishnamoorthi, G.N. La Mar, H. Mizukami and A. Romero, J. Biol. Chem., 259(1984)265
- [586] R. Krishnamoorthi, G.N. La Mar, H. Mizukami and A. Romero, J. Biol. Chem., 259(1984)8826
- [587] R. Krishnamoorthi and G.N. La Mar, Eur. J. Biochem., 138(1984)135
- [588] S. Ramaprasad, R.D. Johnson and G.N. La Mar, J. Am. Chem. Soc., 106(1984)5330
- [589] G.N. La Mar, R. Krishnamoorthi, K.M. Smith, K. Gersonde and H. Sick, Biochemistry, 22(1983)6239
- [590] A. Maldotti, C. Bartocci, R. Amadelli and V. Carassiti, Inorg. Chim. Acta, 74(1983)275
- [591] A.C.I. Anusiem and M. Kelleher, Biopolymers, 23(1984)1147
- [592] J. Peisach, W.B. Mims and J.L. Davis, J. Biol. Chem., 259(1984)2704
- [593] G.M. Alter, J. Biol. Chem., 258(1983)14966
- [594] K. Mawatari, S. Matsukawa and Y. Yoneyama, Biochem. Biophys. Acta, 748(1983)381
- [595] D.A. Juckett and D.E. Hultquist, Biophys. Chem., 19(1984)321
- [596] T.G. Wensel and C.F. Meares, Biochem., 22(1983)6247
- [597] J.F. Rusling and M.Y. Brooks, J. Electroanal. Chem. Interfacial Electrochem., 163(1984)277

- [600] J.S. De Ropp, G.N. La Mar, K.M. Smith and K.C. Langry, J. Am. Chem. Soc., 106(1984)4438
- [601] J. Turner and D.E. Reed, Biochim. Biophys. Acta, 789(1984)80
- [602] J.E. Penner - Hahn, T.J. McMurtry, M. Renner, L. Latos - Grazynsky, K. Smith Eble, I.M. Davis, A.L. Balch, J.T. Groves, J.H. Dawson and K.O. Hodgson, J. Biol. Chem., 258(1983)12761
- [603] M. Sono and J.H. Dawson, Biochim. Biophys. Acta, 789(1984)170
- [604] J. Paul and A. Rosen, Chem. Phys. Lett., 105(1984)197
- [605] A.M. English, D. Sharma and M. Walsh, Inorg. Chim. Acta, 91(1984)L27
- [606] A.M. Lambeir, H.B. Dunford and M.A. Pickard, J. Inorg. Biochem., 19(1983)291
- [607] A. Garnier - Suillerot, L. Tosi and E. Paniago, Biochim. Biophys. Acta, 794(1984)307
- [608] T.G. Traylor, W.A. Lee and D.V. Stynes, Tetrahedron, 40(1984)553
- [609] D.A. Baldwin, V.M. Campbell, H.M. Marques and J.M. Pratt, FEBS Lett., 167(1984)339
- [610] G.R. Moore and G. Williams, Biochim. Biophys. Acta, 788(1984)147
- [611] H. Senn, M. Billeter and K. Wuetrich, Eur. Biophys. J., 11(1984)3
- [612] R. Timkovich, M.S. Cork and P.V. Taylor, Biochemistry, 23(1984)3526
- [613] I. Moura, M.C. Liu, J. Le Gall, H.D. Peck, W. J. Payne, A. V. Xavier and J.J.G. Moura, Eur. J. Biochem., 141(1984)297
- [614] H. Senn, F. Guerlesquin, M. Bruschi and K. Wuetrich, Biochim. Biophys. Acta, 748(1983)194
- [615] R. Timkovich and M.S. Cork, Biochemistry, 23(1984)851
- [616] H. Senn, M.A. Cusanovich and K. Wuetrich, Biochim. Biophys. Acta, 785(1984)46
- [617] J.C. Salerno, J. Biol. Chem., 259(1984)2331

- [618] K.T. Schomacker, O. Bangcharoenpaurpong and P.M. Champion, J. Chem. Phys., 80(1984)4701
- [619] B.R. Stallard, P.R. Callis, P.M. Champion and A.C. Albrecht, J. Chem. Phys., 80(1984)70
- [620] K.K. Andersson, G.T. Babcock and A.B. Hooper, FEBS Lett., 170(1984)331
- [621] A. Schester and W.A. Eaton, Biochemistry, 23(1984)1081
- [622] J. Haladjian and P. Bianco, Bioelectrochem. Bioenerg., 11(1983)319
- [623] E. Margoliash and H.R. Bosshard, Trends Biochem. Sci. (Pers. Ed.), 8(1983)316
- [624] B.C. Hill and C. Greenwood, FEBS Lett., 166(1984)362
- [625] H.E. Toma and A.A. Batista, J. Inorg. Biochem., 20(1984)53
- [626] C.G.S. Eley, E. Raggo and G.R. Moore, J. Inorg. Biochem., 21(1984)295
- [627] E. Ragg and G.R. Moore, J. Inorg. Biochem., 21(1984)253
- [628] K.C. Cho, C.M. Che, F.C. Cheng and C.L. Choy, J. Am. Chem. Soc., 106(1984)6843
- [629] D.G. Nocera, J.R. Winkler, K.M. Yocom, E. Bordinon and H.B. Gray, J. Am. Chem. Soc., 106(1984)5145
- [630] M.A. Harmer and H.A.O. Hill, J. Electroanal. Chem. Interfacial Electrochem., 170(1984)369
- [631] E.F. Bowden, F.M. Hawkridge and H.N. Blount, J. Electroanal. Chem. Interfacial Electrochem., 161(1984)355
- [632] D.M. Kokkinakis and J. Everse, Bioorg. Chem., 12(1983)71
- [633] M.A. Augustin, S.K. Chapman, D.M. Davies, A.D. Watson and A.G. Sykes, J. Inorg. Biochem., 20(1984)281
- [634] Y.P. Myer and S. Kumar, J. Biol. Chem., 259(1984)8144
- [635] A.J. Davison and P. Gee, Arch. Biochem. Biophys., 233(1984)761
- [636] C.H. Kim, C. Balny and T.S. King, Proc. Natl. Acad. Sci. USA,

81(1984)2026

- [637] S.H. Speck, C.A. Neu, M.S. Swanson and E. Margoliash, FEBS Lett., 164(1983)379
- [638] Q.S. Zhu, H. van der Wal, R. van Grondelle and J.H. Berden, Biochim. Biophys. Acta, 725(1983)121
- [639] F. Guerlesquin, M. Bruschi and G. Bovier - Lapierre, Biochimie, 66(1984)93
- [640] Y. Higuchi, M. Kusunoki, Y. Matsura, N. Yasuoka and M. Kakudo, J. Mol. Biol., 172(1984)109
- [641] H. Santos, D.L. Turner, A.V. Xavier and J. Le Gall, J. Magn. Reson., 59(1984)177
- [642] H. Santos, J.J.G. Moura, I. Moura, J. Le Gall and A.V. Xavier, Eur. J. Biochem., 141(1984)283
- [643] K. Zhao and C. Zhu, Kexue Tongbao (Foreign Lang. Ed.), 29(1984)675
- [644] R. Cammack, G. Fauque, J.J.G. Moura and J. Le Gall, Biochim. Biophys. Acta, 784(1984)68
- [645] I. Tabushi, T. Nishiyama, T. Yagi and H. Inokuchi, J. Biochem. (Tokyo), 94(1983)1375
- [646] T.E. Meyer, C.T. Przysiecki, J.A. Watkins, A. Bhattacharyya, R.P. Simonsen, M.A. Cusanovich and G. Tollin, Proc. Natl. Acad. Sci. USA, 80(1983)6740
- [647] M. Bruschi, M. Loutfi, P. Bianco and J. Haladjian, Biochem. Biophys. Res. Commun., 120(1984)384
- [648] L. Sawyer and R.E. Hester, FEBS Lett., 174(1984)219
- [649] Y. Sugimura, Y. Okada, F. Yoshizaki and M. Shimokoriyama, Proc. Jpn. Acad., Ser. B, 59(1983)280
- [650] O. Ristau and H. Rein, Biomed. Biochim. Acta, 42(1983)164
- [651] S.C. Rubinow and R.J. Kassner, Biochemistry, 23(1984)2590
- [652] J. Villalain, I. Moura, M.C. Liu, W.J. Payne, J. Le Gall, A.V. Xavier and J.J.G. Moura, Eur. J. Biochem., 141(1984)305
- [653] W.R. Dunham, R.H. Sands, R.W. Shaw and H. Beinert, Biochim.

- Biophys. Acta, 748(1983)73
- [654] C.P. Scholes, R. Janakiraman, H. Taylor and T.E. King, Biophys. J., 45(1984)1027
- [655] T. Ogura, K. Honnami, T. Oshima, S. Yoshikawa and T. Kitagawa, J. Am. Chem. Soc., 105(1983)7781
- [656] D.G. Eglinton, B.C. Hill, C. Greenwood and A.J. Thomson, J. Inorg. Biochem., 21(1984)1
- [657] B. Chance, C. Kumar, L. Powers and Y.C. Ching, Biophys. J., 44(1983)353
- [658] B.C. Hill and C.T. Greenwood, Biochem. J., 215(1983)659
- [659] M. Brunori and Q.H. Gibson, EMBO J., 2(1983)2025
- [660] P.V. Argade, Y.C. Ching and D.L. Rousseau, Science, 225(1984)329
- [661] R. Boelens, H. Rademaker, R. Wever and B.F. van Gelder, Biochim. Biophys. Acta, 765(1984)196
- [662] N. Ellfolk, M. Roennberg, R. Aasa, L.E. Andreasson and T. Vaenngard, Biochim. Biophys. Acta, 784(1984)62
- [663] H. Baum, J. Elsdon and J.M. Wrigglesworth, Protides Biol. Fluids, 31(1983)35
- [664] C. Kumar, A. Naqui and B. Chance, J. Biol. Chem., 259(1984)2073
- [665] J. M. Wrigglesworth, Biochem. J., 217(1984)715
- [666] J.K. Vandell and T. Yonetani, Biochim. Biophys. Acta, 748(1983)263
- [667] G.D. Jones and M.T. Wilson, J. Inorg. Biochem., 21(1984)159
- [668] P.R. Ortiz de Montellano, Rev. Biochem. Toxicol., 6(1984)1
- [669] F.P. Guengerich and T.L. Macdonald, Acc. Chem. Res., 17(1984)9
- [670] S.G. Sligar, M.H. Gelb and D.C. Heimbrook, Xenobiotica, 14(1984)63
- [671] G.G. Gibson and P.P. Tamburini, Xenobiotica, 14(1984)27

- [672] O. Ristau and H. Rein, Biomed. Biochim. Acta, 42(1983)673
- [673] L.A. Andersson, M. Sono and J.H. Dawson, Biochim. Biophys. Acta, 748(1983)341
- [674] S.A. Usanov, V.A. Figlovskii and A.A. Akhrem, Dokl. Akad. Nauk SSSR, 274(1984)455
- [675] L.A. Andersson and J.H. Dawson, Xenobiotica, 14(1984)49
- [676] S. Narasimhulu and C.R. Eddy, Biochemistry, 28(1984)1109
- [677] L.R. Pohl, R.D. Schulick, R.J. Highet and J.W. George, Mol. Pharmacol., 25(1984)318
- [678] P.G. Gervasi, L. Citti, E. Testai and G. Turchi, Toxicol. Lett., 20(1984)243
- [679] S. Hamill and D.Y. Cooper, Xenobiotica, 14(1984)139
- [680] K.K. Andersson, T.A. Kent, J.D. Lipscomb, A.B. Hooper and E. Munck, J. Biol. Chem., 259(1984)6833
- [681] D.E. Wallick, L.M. Bloom, B.J. Gaffney and S.J. Benkovic, Biochemistry, 23(1984)1295
- [682] J.W. Whittaker and J.D. Lipscomb, J. Biol. Chem., 259(1984)4487
- [683] P.R. Rich, Biochim. Biophys. Acta, 768(1984)53
- [684] T.C. Strekas, Biochim. Biophys. Acta, 765(1984)133
- [685] K.H.D. Le and F. Lederer, EMBO J., 2(1983)1909
- [686] R.C. Prince, C. Larroque and A.B. Hooper, FEBS Lett., 163(1983)25
- [687] T. Yubisui and M. Takeshita, Biochem. Int., 8(1984)319
- [688] J. Bergstroem, L.E. Andreasson and T. Vaenngaard, FEBS Lett., 164(1983)71
- [689] V.S. Kunts, A.A. Konstantinov, L.M. Tsofina and E.A. Liberman, FEBS Lett., 172(1984)261
- [690] M.W. Makinen, S.A. Schichman, S.C. Hill and H.B. Gray, Science, 222(1983)929

- [691] D.J. Davis and K. Hough, Biochem. Biophys. Res. Commun., 116(1983)1000
- [692] S.F. Sontum, D.A. Case and M. Karplus, J. Chem. Phys., 79(1983)2881
- [693] J. Silver, B. Lukas and G. Al - Jaff, Inorg. Chim. Acta, 91(1984)125
- [694] K.M. Kadish, J. Electroanal. Chem. Interfacial Electrochem., 168(1984)261
- [695] L.A. Bottomley, M.R. Deakin and J.N. Gorce, Inorg. Chem., 23(1984)3563
- [696] Y. Matsushita, E. Hasegawa, K. Eshima and E. Tsuchida, Chem. Lett., (1983)1387
- [697] D.K. Geigery, Y.J. Lee and W.R. Scheidt, J. Am. Chem. Soc., 106(1984)6339
- [698] A. Desbois, Y. Henry and M. Lutz, Biochim. Biophys. Acta, 785(1984)161
- [699] T. Watanabe, T. Ama and K. Nakamoto, J. Phys. Chem., 88(1984)440
- [700] K.S. Suslick, M.M. Fox and T.R. Reinert, J. Am. Chem. Soc., 106(1984)4522
- [701] T. Mashiko, C.A. Reed, K.J. Haller and W.R. Scheidt, Inorg. Chem., 23(1984)3192
- [702] M. Fontecave, J.P. Battioni and D. Mansuy, J. Am. Chem. Soc., 106(1984)5217
- [703] P. Doppelt, Inorg. Chem., 23(1984)4009
- [704] K.M. Miller and C.E. Strouse, Inorg. Chem., 23(1984)2395
- [705] W.R. Scheidt, Y.J. Lee, T. Bartczak and K. Hatano, Inorg. Chem., 23(1984)2552
- [706] K.M. Smith and R.K. Pandey, J. Heterocycl. Chem., 20(1983)1383
- [707] G.E. Toney, L.W. ter Haar, J.E. Savrin, A. Gold, W.E. Hatfield and R. Sangaiah, Inorg. Chem., 23(1984)2561

- [708] A.A. Shteinman, Izv. Akad. Nauk SSSR, Ser. Khim., (1983)2415
- [709] D.K. Geiger and W.R. Scheidt, Inorg. Chem., 23(1984)1970
- [710] R. Quinn, J. Mercer - Smith, J.N. Burstyn and J.S. Valentine, J. Am. Chem. Soc., 106(1984)4136
- [711] F.A. Walker, D. Reis and V.L. Balke, J. Am. Chem. Soc., 106(1984)6888
- [712] A.D. Boersma, M.A. Phillippi and H.M. Goff, J. Magn. Reson., 57(1984)197
- [713] T.A. Babushkina, R.E. Garibov, L.B. Lazukova and V.V. Khrapov, Koord. Khim., 10(1984)77
- [714] J. Silver and B. Lukas, Inorg. Chim. Acta, 92(1984)259
- [715] J.W. Buchler and C. Dreher, Z. Naturforsch. B, 39(1984)222
- [716] D. Lancoc and K.M. Kadish, Inorg. Chem., 23(1984)3942
- [717] T. Yoshimura and T. Ozaki, Arch. Biochem. Biophys., 230(1984)466
- [718] B. Lukas, J. Peterson and J. Silver, Inorg. Chim. Acta, 80(1983)245
- [719] T. Otsuka, T. Ohya and M. Sato, Inorg. Chem., 23(1984)1777
- [720] D. Lancon, P. Cocolios, R. Guillard and K.M. Kadish, Organometallics, 3(1984)1164
- [721] D. Lancon, P. Cocolios, R. Guillard and K.M. Kadish, J. Am. Chem. Soc., 106(1984)4472
- [722] A. Jain and R.J. Kassner, J. Biol. Chem., 259(1984)10309
- [723] N. Kobayashi and T. Osa, J. Electroanal. Chem. Interfacial Electrochem., 157(1983)269
- [724] I. Wilson, K.R. Bretscher, C.K. Chea and H.C. Kelly, J. Inorg. Biochem., 19(1983)345
- [725] A.H. Chu and T.J. Williams, Arch. Biochem. Biophys., 234(1984)129
- [726] T. Yoshimura and T. Ozaki, Arch. Biochem. Biophys.,



229(1984)126

- [727] T. Yoshimura, Inorg. Chim. Acta, 83(1984)17
- [728] A. Strich and A. Veillard, Nouv. J. Chim., 7(1983)347
- [729] A.G. Volkov, M.A. Bibikova, A.F. Mironov and L.I. Boguslavskii, Bioelectrochem. Bioenerg., 10(1983)477
- [730] V.A. Zhorin, G.A. Nikiforov and N.S. Enkilopyan, Izv. Akad. Nauk SSSR, Ser. Khim., (1984)1438; CA, 101(1984)102991v
- [731] G.M. Klecka, Chemosphere, 13(1984)391
- [732] D.W. Dixon, M. Barbusch and A. Shirazi, J. Am. Chem. Soc., 106(1984)4638
- [733] D. Brault, R. Santus, E.J. Land and A.J. Swallow, J. Phys. Chem., 88(1984)5836
- [734] D.R. English, D.N. Hendrickson, K.S. Suslick, C.W. Eigenbrot and W.R. Scheidt, J. Am. Chem. Soc., 106(1984)7258
- [735] K.M. Miller and C.E. Strouse, Acta Crystallogr. C, 40(1984)1324
- [736] H.H. Ruf, H. Ahr, W. Nastainczyk, V. Ullrich, D. Mansuy, J.B. Battioni, R. Montiel - Montoya and A. Trautwein, Biochemistry, 23(1984)5300
- [737] J. Silver and B. Lukas, Inorg. Chim. Acta, 91(1984)279
- [738] N. Miyata, T. Santa and M. Hirobe, Chem. Pharm. Bull., 32(1984)377
- [739] J.P. Collman, T. Kodadek, S.A. Raybuck and B. Meunier, Proc. Natl. Acad. Sci. USA, 80(1983)7039
- [740] H. Sakurai, K. Ishizu and K. Okada, Inorg. Chim. Acta, 91(1984)L9
- [741] M. Shimada, T. Habe, T. Umezawa, T. Higuchi and T. Okamoto, Biochem. Biophys. Res. Commun., 122(1984)1247
- [742] G.J. Hamilton, J.W. Owens, C.J. O'Connor and R.J. Kassner, Inorg. Chim. Acta, 93(1984)55
- [743] R.E. Garibov, V.I. Khleskov and A.I. Lumpov, Teor. Eksp. Khim., 20(1984)157; CA, 101(1984)34748u

- [744] A. Gold, W. Ivey, G.E. Toney and R. Sangaiiah, Inorg. Chem., 23(1984)2932
- [745] J.E. Baldwin and P. Perlmutter, Top. Curr. Chem., 121(1984)181
- [746] P. Bertrand, F.X. Theodule, J.P. Gayda, J. Mispelter and M. Mentaou, Chem. Phys. Lett., 102(1983)442
- [747] H. Ogoshi, H. Sugimoto, M. Miyake and Z. Yoshida, Tetrahedron, 40(1984)579
- [748] Y. Matsushita, E. Hasegawa, K. Eshima and E. Tsuchida, Heterocycles, 22(1984)1403
- [749] M. Shimizu, F. Basolo, M.N. Vallejo and J.E. Baldwin, Inorg. Chim. Acta, 91(1984)247
- [750] T.G. Traylor, K. Nolan, R. Hildreth and T.A. Evans, Heterocycles, 21(1984)249
- [751] E. Tsuchida, H. Nishide, M. Sekine and A. Yamagishi, Biochim. Biophys. Acta, 734(1983)274
- [752] L. Ricard, M. Schappacher, R. Weiss, R. Montiel - Montoya, E. Bill, U. Gonser and A. Trautwein, Nouveau J. Chim., 7(1983)405
- [753] T.K. Miyamoto, S. Tsuzuki, T. Hasegawa and Y. Sasaki, Chem. Lett., (1983)1587
- [754] E. Tsuchida, H. Nishide, M. Yuasa and M. Sekine, Bull. Chem. Soc. Jpn, 57(1984)776
- [755] M. Shimizu, F. Basolo, M.N. Vallejo and J.E. Baldwin, Inorg. Chim. Acta, 91(1984)251
- [756] M.J. Gunter, K.J. Berry and K.S. Murray, J. Am. Chem. Soc., 106(1984)4227
- [757] W.E. Newton, Stud. Inorg. Chem., 5(1984)409
- [758] F. Bonomi and D.M. Kurtz, Anal. Biochem., 142(1984)226
- [759] D.M. Kurtz, Inorg. Chim. Acta, 80(1983)L75
- [760] H. Blum, J.R. Bowyer, M.A. Cusanovich, A.J. Waring and T. Ohnishi, Biochim. Biophys. Acta, 748(1983)418
- [761] J.M. Moulis, J. Meyer and M. Lutz, Biochem. J., 219(1984)829

- [762] Y. Ninomiya and S. Sato, Plant Cell Physiol., 25(1984)453
- [763] C.D. Crawley and F.M. Hawkridge, J. Electroanal. Chem. Interfacial Electrochem., 159(1983)313
- [764] M. Hirasawa and G. Tamura, J. Biochem. (Tokyo), 95(1984)983
- [765] M. Nishikimi, Y. Shimomura, H. Yamada and T. Ozawa, Biochem. Biophys. Res. Commun., 120(1984)237
- [766] L.C. Sieker, L.H. Jensen, B.C. Prickril and J. LeGall, J. Mol. Biol., 171(1983)101
- [767] S.W. Ragsdale and L.G. Ljungdahl, J. Bacteriol., 157(1984)1
- [768] T.E. Gorrell, N. Yarlett and M. Mueller, Carlsberg Res. Commun., 49(1984)259
- [769] P.J. Geary, F. Saboowalla, D. Patil and R. Cammack, Biochem. J., 217(1984)667
- [770] T.M. Chan and J.L. Markley, Biochemistry, 22(1983)5996
- [771] T.M. Chan and J.L. Markley, Biochemistry, 22(1983)6008
- [772] T.M. Chan and J.L. Markley, Biochemistry, 22(1983)5982
- [773] T.M. Chan, E.L. Ulrich and J.L. Markley, Biochemistry, 22(1983)6002
- [774] T.M. Chan, M.A. Hermodson, E.L. Ulrich and J.L. Markley, Biochemistry, 22(1983)5988
- [775] I. Bertini, G. Lanini and C. Luchinat, Inorg. Chem., 23(1984)2729
- [776] J. Meyer, J.M. Moulis and M. Lutz, Biochem. Biophys. Res. Commun., 119(1984)828
- [777] I.K. Adzhamli, R.A. Henderson, J.D. Sinclair - Day and A.G. Sykes, Inorg. Chem., 23(1984)3069
- [778] J.A. Fee, K.L. Findling, T. Yoshida, R. Hille, G.E. Tarr, D.O. Hearshen, W.R. Dunham, E.P. Day, T.A. Kent and E. Munck, J. Biol. Chem., 259(1984)124
- [779] L.C. Sieker, E.T. Adman, L.H. Jensen and J. LeGall, J. Mol. Biol., 179(1984)151

- [780] J.P. Gayda, P. Bertrand, B. Gugliarelli and J. Meyer, J. Chem. Phys., 79(1983)5732
- [781] P. Bertrand, B. Guigliarelli, J. Meyer and J.P. Gayda, Biochimie, 66(1984)77
- [782] J.J.G. Moura, J. LeGall and A.V. Xavier, Eur. J. Biochem., 141(1984)319
- [783] M.Lutz, J.M. Moulis and J. Meyer, FEBS Lett., 163(1983)212
- [784] T. Imai, H. Saito, J. Tobari, D. Ohmori and K. Suzuki, FEBS Lett., 165(1984)227
- [785] T.V. Morgan, P.J. Stephens, B.K. Burgess and C.D. Stout, FEBS Lett., 167(1984)137
- [786] M.C. Kennedy, M.H. Emptage and H. Beinert, J. Biol. Chem., 259(1984)3145
- [787] T.V. Morgan, P.J. Stephens, F. Devlin, C.D. Stout, K.A. Melis and B.K. Burgess, Proc. Natl. Acad. Sci. USA, 81(1984)1931
- [788] K. Nagayama, T. Imai, D. Ohmori and T. Oshima, FEBS Lett., 169(1984)79
- [789] E.C. Hatchikian, R. Cammack, D.S. Patil, A.E. Robinson, A.J.M. Richards, S. George and A.J. Thomson, Biochim. Biophys. Acta, 784(1984)40
- [790] B.H. Huynh, M.H. Czechowski, H.J. Kruger, D.V. Der Vartanian, H.D. Peck and J. LeGall, Proc. Natl. Acad. Sci. USA, 81(1984)3728
- [791] P.A. Lindahl, N. Kojima, R.P. Hausinger, J.A. Fox, B.K. Teo, C.T. Walsh and W.H. Orme - Johnson, J. Am. Chem. Soc., 106(1984)3062
- [792] J.M. Moulis, P. Auric, J. Gaillard and J. Meyer, J. Biol. Chem., 259(1984)11396
- [793] N.R. Orme - Johnson, W.B. Mims, W.H. Orme - Johnson, R.G. Bartsch, M.A. Cusanovich and J. Peisach, Biochem. Biophys. Acta, 748(1983)68
- [794] K. Nagayama and D. Ohmori, FEBS Lett., 173(1984)15
- [795] C. Vicente and V. Requena, Photosynthetica, 18(1984)57

- [796] K. Yanada, T. Nagano and M. Hirobe, Chem. Pharm. Bull., 31(1983)4589
- [797] D. Sellmann, U. Kleine - Kleffmann, L. Zapf, G. Huttner and L. Zsolnai, J. Organomet. Chem., 263(1984)321
- [798] M. Nakata, N. Ueyama, T. Terekawa and A. Nakamura, Bull. Chem. Soc. Jpn., 56(1983)3647
- [799] M. Nakata, N. Ueyama, M. Fuji, A. Nakamura, K. Wada and H. Matsubara, Biochim. Biophys. Acta, 788(1984)306
- [800] H. Strasdeit, B. Krebs and G. Henkel, Inorg. Chim. Acta, 89(1984)L11
- [801] N. Ueyama, S. Ueno, M. Nakata and A. Nakamura, Bull. Chem. Soc. Jpn., 57(1984)984
- [802] D. Coucouvanis, A. Salifoglou, M.G. Kanatzidis, A. Simopoulos and V. Papaefthymiou, J. Am. Chem. Soc., 106(1984)6081
- [803] A. Mueller, W. Hellman, C. Roemer, M. Roemer, H. Bogge, R. Jostes and U. Schimanski, Inorg. Chim. Acta, 83(1984)L75
- [804] J.J. Gixerd, G.C. Papaefthymiou, A.D. Watson, E. Gamp, K.S. Hagen, N. Edelstein, R.B. Frankel and R.H. Holm, J. Am. Chem. Soc., 106(1984)5941
- [805] K.S. Hagen, A.D. Watson and R.H. Holm, Inorg. Chem., 23(1984)2984
- [806] M.G. Kanatzidis, N.C. Baenziger, D. Coucouvanis, A. Simopoulos and A. Kostikas, J. Am. Chem. Soc., 106(1984)4500
- [807] A. Simopoulos and A. Kostikas, Sol. Energy R. & D. Eur. Community, 2(1983)239
- [808] J. Nieman, A.J. Naaktgeboren and J. Reedijk, Inorg. Chim. Acta, 93(1984)L9
- [809] N. Dupre, H.M.J. Hendriks, J. Jordanov, J. Gaillard and P. Auric, Organometallics, 3(1984)800
- [810] W. Saak, G. Henkel and S. Pohl, Angew. Chem. Internat. Ed. Engl., 23(1984)150
- [811] M.G. Kanatzidis, W.R. Dunham, W.R. Hagen and D. Coucouvanis, J. Chem. Soc. Chem. Commun., (1984)356

- [812] H. Strasdeit, B. Krebs and G. Henkel, Inorg. Chem., 23(1984)1816
- [813] G.L. Anderson and J.B. Howard, Biochemistry, 23(1984)2118
- [814] R.E. Palermo, R. Singh, J.K. Bashkin and R.H. Holm, J. Am. Chem. Soc., 106(1984)2600
- [815] Y. Fan, P. Lu, F. Wang, X. Liu and J. Xu, Huaxue Xuebao, 41(1983)776; CA, 100(1984)59867z; J. Xu, X. Liu, S. Li, Y. Fan, F. Wang and P. Lu, Kexue Tongbao (Foreign Lang. Ed.), 29(1984)344; CA, 101(1984)142757e
- [816] F. Wang, Z. Zhang, Y. Fan and C. Shen, Jilin Daxue Ziran Kexue Xuebao, (1984)94; CA, 101(1984)120806t
- [817] J. Xu, X. Liu, Z. Zhang, S. Niu, S. Li, Y. Fan, F. Wang, P. Lu and A. Tang, Fenzi Kexue Yu Huaxue Yanjiu, 3(1983)1; CA 100(1984)131398s
- [818] N. Ueyama, M. Nakata, A. Nakamura, M. Kamata and S. Otsuka, Inorg. Chim. Acta, 80(1983)207
- [819] B. Zhuang, J.W. McDonald and W.E. Newton, Inorg. Chim. Acta, 77(1983)L221
- [820] I.M. Klotz and D.M. Kurtz, Acc. Chem. Res., 17(1984)16
- [821] R.G. Wilkins and P.C. Harrington, Adv. Inorg. Biochem., 5(1983)51
- [822] R.E. Stenkamp, L.C. Sieker and L.H. Jensen, J. Inorg. Biochem., 19(1983)247
- [823] R.E. Stenkamp, L.C. Sieker and L.H. Jensen, Acta Cryst. B, 39(1983)697
- [824] M.J. Maroney, R.B. Lauffer, L. Que and D.M. Kurtz, J. Am. Chem. Soc., 106(1984)6445
- [825] A.K. Shienke, T.M. Loehr and J. Sanders - Loehr, J. Am. Chem. Soc., 106(1984)4951
- [826] J.M. Nocek, D.M. Kurtz, R.A. Pickering and M.P. Doyle, J. Biol. Chem., 259(1984)12334
- [827] P.C. Harrington and R.G. Wilkins, J. Inorg. Biochem., 19(1983)339

- [828] C.A. Lewis, A.G. Lappin and W.J. Ingledew, Biochem. Soc. Trans., 12(1984)503
- [829] K.N. Raymond, G. Muller and B.F. Matzanke, Top. Curr. Chem., 123(1984)49
- [830] B.F. Matzanke, G.I. Muller and K.N. Raymond, Biochem. Biophys. Res. Commun., 121(1984)922
- [831] S.J. Barclay, P.E. Riley and K.N. Raymond, Inorg. Chem., 23(1984)2005
- [832] S.J. Barclay, B.H. Huynh and K.N. Raymond, Inorg. Chem., 23(1984)2011
- [833] T. Emery, Anal. Biochem., 139(1984)301
- [834] B. Monzyk and A.L. Crumbliss, J. Inorg. Biochem., 19(1983)19
- [835] T. Emery, L. Emery and R.K. Olsen, Biochem. Biophys. Res. Commun., 119(1984)1191
- [836] S. Wendenbaum, P. Demange, A. Dell, J.M. Meyer and M.A. Abdallah, Tetrahedron Lett., 24(1983)4877
- [837] G.L. Griffiths, S.P. Sigel, S.M. Payne and J.B. Neilands, J. Biol. Chem., 259(1984)383
- [838] M.J. Smith and J.B. Neilands, J. Plant Nutr., 7(1984)449
- [839] M.A.F. Jalal, R. Mocharla, C.L. Barnes, M.B. Hossain, D.R. Powell, D.L. Eng - Wilmot, S.L. Grayson, B.A. Benson and D. Van der Helm, J. Bacteriol., 158(1984)683
- [840] L.P. Evanlyo, S. Kadis and J.R. Maudsley, Can. J. Microbiol., 30(1984)1046
- [841] G. Deml, K. Voges, G. Jung and G. Winkelmann, FEBS Lett., 173(1984)53
- [842] D.J. Schneider, A.L. Roe, R.J. Mayer and L. Que, J. Biol. Chem., 259(1984)9699
- [843] T. Nilsen and I. Romslo, Biochim. Biophys. Acta, 766(1984)233
- [844] W. Lohmann, D. Holz, B. Kiefer and D. Schmidt, Z. Naturforsch. C, 38(1983)926

- [845] G.C. Ford, P.M. Harrison, D.W. Rice, J.M.A. Smith, A. Treffry, J.L. White and J. Yariv, Phil. Trans. R. Soc. London B, 304(1984)551
- [846] E.C. Theil, Adv. Inorg. Biochem., 5(1983)1
- [847] D.W. Rice, G.C. Ford, J.L. White, J.M.A. Smith and P.M. Harrison, Protides Biol. Fluids, 31(1983)177
- [848] M.P. Weir, J.F. Gibson and T.J. Peters, Biochem. Soc. Trans., 12(1984)316
- [849] S.H. Bell, M.P. Weir, D.P.E. Dickson, J.F. Gibson, G.A. Sharp and T.J. Peters, Biochim. Biophys. Acta, 787(1984)227
- [850] P. Gonzalez del Barrio and M.C. Martin Mateo, Comp. Biochem. Physiol. B, 76(1983)567
- [851] A. Treffry and P.M. Harrison, J. Inorg. Biochem., 21(1984)9
- [852] P. Biemond, H.G. van Eijk, A.J.G. Swaak and J.F. Koster, J. Clin. Invest., 73(1984)1576
- [853] K.A. Berg, L.H. Bowen, S.W. Hedges, R.D. Bereman and C.T. Vance, J. Inorg. Biochem., 22(1984)125
- [854] C. Daniel, M. Benard, A. Dedieu, R. Wiest and A. Veillard, J. Phys. Chem., 88(1984)4805
- [855] Y. Langsam and A.M. Ronn, J. Chem. Phys., 80(1984)749
- [856] P.A. Teng, F.D. Lewis and E. Weitz, J. Phys. Chem., 88(1984)4895
- [857] M.S. Paquette, Proc. SPIE - Int. Soc. Opt. Eng., 458(1984)140
- [858] M.A. Moskalenko and A.B. Terent'ev, Izv. Akad. Nauk SSSR, Ser. Khim., (1983), 2322; CA, 100(1984)102690a
- [859] R.J.P. Corriou, V. Huynh, J.J.E. Moreau and M. Pataud - Sat, J. Organomet. Chem., 255(1983)359
- [860] K.J. Fu, R.L. Whetten and E.R. Grant, Ind. Eng. Chem. Prod. Res. Div., 23(1984)33
- [861] M.E. Miller and E.R. Grant, Proc. SPTE - Int. Soc. Opt. Eng., 458(1984)154
- [862] M.A. Rahdi and L. Marko, J. Organomet. Chem., 262(1984)359



- [863] M.A. Radhi, G. Palyi and L. Marko, J. Mol. Catal., 22(1983)195
- [864] T.J. Lynch, M. Banah, H.D. Kaesz and C.R. Porter, J. Org. Chem., 49(1984)1266
- [865] G. Tanguy, B. Weinberger and H. des Abbayes, Tetrahedron Lett., 24(1983)4005
- [866] G.C. Tustin and R.T. Hembre, J. Org. Chem., 49(1984)1761
- [867] L.N. Kiseleva, N.A. Rybakova and R.Kh. Freidlina, Izv. Akad. Nauk SSSR, Ser. Khim., (1983)2408
- [868] T. Bein and P.A. Jacobs, J. Chem. Soc. Faraday Trans. 1, 80(1984)1391
- [869] M.A. de Paoli, S.M. de Oliviera, E. Baggio - Saitovitch and D. Guenzburger, J. Chem. Phys., 80(1984)730
- [870] A.S. Goldman and D.R. Tyler, Organometallics, 3(1984)449
- [871] S.A. Fairhurst, J.R. Morton, R.N. Perutz and K.F. Preston, Organometallics, 3(1984)1389
- [872] E. Bill, N. Blaes, K.F. Fischer, U. Gonser, K.H. Pauly, R. Preston, F. Seel, R. Staab and A.X. Trautwein, Z. Naturforsch., B, 39(1984)333
- [873] R.N. McDonald, P.L. Schell and W.D. McGhee, Organometallics, 3(1984)182
- [874] A. Lapidus, M.M. Savel'ev, S.D. Sominskii and M.V. Tsapkina, Kinet. Katal., 25(1984)636; CA, 101(1984)130004u
- [875] D.B. Beach, S.P. Smit and W.L. Jolly, Organometallics, 3(1984)556
- [876] A.H. Cowley, J.E. Kilduff, J.G. Lasch, S.K. Mehrotra, N.C. Norman, M. Pakulski and C.A. Stewart, Phosphorus Sulfur, 18(1983)3
- [877] J.T. Mague and S.E. Dessens, J. Organomet. Chem., 262(1984)347
- [878] E. Gross, K. Joerg, K. Fiederling, A. Goettlein, W. Malisch and R. Boese, Angew. Chem. Int. Ed. Engl., 23(1984)738
- [879] R.B. King and M. Shimura, J. Organomet. Chem., 256(1983)71

- [880] E.H. Wong, F.C. Bradley, L. Prasad and E.J. Gabe, J. Organomet. Chem., 263(1984)167
- [881] R.H. Neilson, R.J. Thoma, I. Vickovic and W.H. Watson, Organometallics, 3(1984)1132
- [882] A.N. Chenega, M.Yu. Antipin, Yu. T. Struchkov, Yu.V. Balitskii, Yu.G. Gololobov and I.E. Boldeskul, Zh. Obshch. Khim., 54(1984)271
- [883] H. Nakazawa, D.L. Johnson and J.A. Gladysz, Organometallics, 2(1983)1846
- [884] F. Kober and M. Kerber, Z. Anorg. Allg. Chem., 507(1983)119
- [885] A.H. Cowley, N.C. Norman and M. Pakulski, J. Am. Chem. Soc., 106(1984)6844
- [886] H.J. Breunig, Polyhedron, 3(1984)757
- [887] H.J. Breunig and U. Graefe, Z. Anorg. Allg. Chem., 510(1984)104
- [888] E. Guglielminotti, J. Mol. Catal., 24(1984)331
- [889] J.W. Connolly, Organometallics, 3(1984)1333
- [890] D. Melzer and E. Weiss, J. Organomet. Chem., 255(1983)335
- [891] J.M. Barbe, R. Guillard, C. Lecomte and R. Gerardin, Polyhedron, 3(1984)889
- [892] I.L. Eremenko, A.A. Pasynskii, Yu.V. Rakitin, O.G. Ellert, V.M. Novotortsev, V.T. Kalinnikov, V.E. Shklover and Yu.T. Struchkov, J. Organomet. Chem., 256(1983)291
- [893] C. Glidewell and A.R. Hyde, Inorg. Chim. Acta, 87(1984)L15
- [894] J. Takacs and L. Marko, Transition Met. Chem., 9(1984)10
- [895] D.K. Liu, C.G. Brinkley and M.S. Wrighton, Organometallics, 3(1984)1449
- [896] H.W. Fruehauf, J. Chem. Res. Synop., (1983)218
- [897] A. Laporterie, H. Iloughmane and J. Dubac, Tetrahedron Lett., 24(1983)3521
- [898] R.N. McDonald, A.K. Chowdhury and P.L. Schell,

- Organometallics, 3(1984)644
- [899] M.R. Houchin, Inorg. Chim. Acta, 83(1984)103
  - [900] H.C. Ashton and A.R. Manning, Inorg. Chim. Acta, 71(1983)163
  - [901] P. Robert, B. Demerseman and P.H. Dixneuf, Organometallics, 3(1984)1771
  - [902] O. Gambino, G. Gervasio, R. Rossetti and P.L. Stanghellini, Inorg. Chim. Acta, 84(1984)51
  - [903] R. Lissilour, J. Bennaceur and H. Patin, J. Chim. Phys. Phys.-Chim. Biol., 81(1984)225
  - [904] G.J. Kruger, L. Linford, H.G. Raubenheimer and A.A. Chalmers, J. Organomet. Chem., 262(1984)69
  - [905] E.A. Chernyshev, O.V. Kuz'min, A.V. Lebedev, A.I. Gusev, N.I. Kirillova, N.G. Komalenkova, N.S. Nametkin and V.D. Tyurin, Izv. Akad. Nauk SSSR, Ser. Khim., (1984)633; CA, 101(1984)91158a
  - [906] M. Heberhold and W. Buehlmeier, Angew. Chem. Internat. Ed. Engl., 23(1984)80
  - [907] D. Seyferth, G.B. Womack, M. Cowie and B.W. Hames, Organometallics, 3(1984)1891
  - [908] N.S. Nametkin, V.D. Tyurin, V.V. Trusov and A.M. Krapivin, J. Organomet. Chem., 264(1983)243
  - [909] A.L. Rheingold and M.A. Fountain, Organometallics, 3(1984)1417
  - [910] E.P. Kyba, R.E. Davis, K.L. Hassett, J.S. McKennis and B. Sheikh, Phosphorus Sulfur, 18(1983)275
  - [911] H.R. Allcock, P.R. Suszko, L.J. Wagner, R.R. Whittle and B. Boso, J. Am. Chem. Soc., 106(1984)4966
  - [912] W. Petz, J. Organomet. Chem., 270(1984)81
  - [913] A.A. Bahsoun, J.A. Osborn, P.H. Bird, D. Nucciarone and A.V. Peters, J. Chem. Soc. Chem. Commun., (1984)72
  - [914] E.R.F. Gesing, V. Groth and K.P.C. Vollhardt, Synthesis, 4(1984)351
  - [915] T. Fuchikami and I. Ojima, Tetrahedron Lett., 25(1984)307

- [916] J.C. Bricker and S.G. Shore, Organometallics, 3(1984)201
- [917] M. Iwamoto, H. Kusano and S. Kagawa, Chem. Lett., (1983)1483
- [918] W.K. Wong, K.W. Chiu, G. Wilkinson, A.M.R. Galas, M. Thornton - Pett and M.B. Hursthouse, J. Chem. Soc. Dalton Trans., (1983)1557
- [919] L. Marko, J. Takacs, S. Papp and B. Marko - Monostory, Inorg. Chim. Acta, 45(1980)L189
- [920] J.P. Fackler, A.M. Mazany, D. Seyferth, H.P. Withers, T.G. Wood and C.F. Campana, Inorg. Chim. Acta, 82(1984)31
- [921] E. Lindner, G.A. Weiss, W. Hiller and R. Fawzi, J. Organomet. Chem., 255(1983)245
- [922] I.P. Lorenz and J. Messelhaeuser, Z. Naturforsch. B, 39(1984)403
- [923] V.D. Patel, N.J. Taylor and A.J. Carty, J. Chem. Soc. Chem. Commun., (1984)99
- [924] O. Stelzer, S. Hietkamp and H. Sommer, Phosphorus Sulfur, 18(1983)279
- [925] I. Bruedgam, H. Hartl and D. Lentz, Z. Naturforsch. B, 39(1984)721
- [926] M.I. Bruce, T.W. Hambley and B.K. Nicholson, J. Chem. Soc. Dalton Trans., (1983)2285
- [927] J. Vites and T.P. Fehlner, Organometallics, 3(1984)491
- [928] J. Vites, C. Eigenbrot and T.P. Fehlner, J. Am. Chem. Soc., 106(1984)4633
- [929] J.C. Vites, C.E. Housecroft, G.B. Jacobsen and T.P. Fehlner, Organometallics, 3(1984)1591
- [930] A. Winter, L. Zsolnai and G. Huttner, J. Organomet. Chem., 269(1984)C29
- [931] T. Venalainen and T. Pakkanen, J. Organomet. Chem., 266(1984)269
- [932] U. Honrath and H. Vahrenkamp, Z. Naturforsch. B, 39(1984)545

- [933] M. Mueller and H. Vahrenkamp, Chem. Ber., 116(1983)2311
- [934] F. Richter, E. Roland and H. Vahrenkamp, Chem. Ber., 117(1984)2429
- [935] C.P. Horwitz and D.F. Shriver, Organometallics, 3(1984)756
- [936] T.P. Fehlner and C.E. Housecroft, Organometallics, 3(1984)764
- [937] S. Harris and J.S. Bradley, Organometallics, 3(1984)1086
- [938] S.D. Wijeyesekera, R. Hoffmann and C.N. Wilker, Organometallics, 3(1984)962
- [939] K.S. Bose, E. Sinn and B.A. Averill, Organometallics, 3(1984)1126
- [940] C.P. Horwitz, E.M. Holt and D.F. Shriver, Inorg. Chem., 23(1984)2491
- [941] L.Y. Hsu, A.A. Bhattacharyya and S.G. Shore, Acta Crystallogr. C, 40(1984)722
- [942] G.E. Hawkes, E.W. Randall, S. Aime, R. Gobetto and D. Osella, Polyhedron, 2(1983)1235
- [943] A. Ceriotti, P. Chini, A. Fumagalli, T.F. Koetzle, G. Longoni and F. Takusagawa, Inorg. Chem., 23(1984)1363
- [944] H. Lang, L. Zsolnai and G. Huttner, Angew. Chem., 95(1983)1016
- [945] F. Richter, M. Mueller, N. Gaertner and H. Vahrenkamp, Chem. Ber., 117(1984)2438
- [946] P. Braunstein, C. de Meric de Bellefon and M. Ries, J. Organomet. Chem., 262(1984)C14
- [947] P. Braunstein and J. Rose, J. Organomet. Chem., 262(1984)223
- [948] V.E. Lopatin, N.M. Mikova and S.P. Gubin, Koord. Khim., 10(1984)715
- [949] S.P. Gubin, N.M. Mikova, M.Ts. Tsybenov and V.E. Lopatin, Koord. Khim., 10(1984)625
- [950] R.P. Brint, M.P. Collins, T.R. Spalding and F.A. Deeney, J. Organomet. Chem., 258(1983)C57

- [951] G. A. Martin and E. Bordignon, Inorg. Chem., 23(1984)3822
- [952] F.J. Lalor and L.H. Brookes, J. Organomet. Chem., 251(1983)327
- [953] Y.F. Yu, C.N. Chau, A. Wojcicki, M. Calligaris, G. Nardin and G. Balducci, J. Am. Chem. Soc., 106(1984)3704
- [954] M.D. Seidler and R.G. Bergman, Organometallics, 2(1983)1897
- [955] A. Dedieu and S. Nakamura, J. Organomet. Chem., 260(1984)C63
- [956] D.B. Jacobson and B.S. Freiser, J. Am. Chem. Soc., 106(1984)3900
- [957] D.B. Jacobson and B.S. Freiser, Organometallics, 3(1984)513
- [958] B.S. Larsen and D.P. Ridge, J. Am. Chem. Soc., 106(1984)1912
- [959] D.A. Peake, M.L. Gross and D.P. Ridge, J. Am. Chem. Soc., 106(1984)4307
- [960] W.L. Grady and M.M. Bursey, Int. J. Mass Spectrom. Ion Processes, 56(1984)161
- [961] K.J. Lattermann and W. Seidel, Z. Chem., 23(1983)445
- [962] P.J. Krusic, W.J. Cote and A. Grand, J. Am. Chem. Soc., 106(1984)4642
- [963] W. Keim, M. Roeper, H. Strutz and C. Krueger, Angew. Chem. Int. Ed. Engl., 23(1984)432
- [964] G. Bellachioma, G. Cardaci and G. Reichenbach, J. Chem. Soc. Dalton Trans., (1983)2593
- [965] E. Lindner, C.P. Krieg, W. Hiller and R. Fawzi, Angew. Chem., 96(1984)508
- [966] M. Pankowski and M. Bigorgne, J. Organomet. Chem., 251(1983)333
- [967] G. Cardaci, G. Bellachioma and P.F. Zanazzi, J. Chem. Soc. Chem. Commun., (1984)650
- [968] B. Gregory, C.R. Jablonski and Y.P. Wang, J. Organomet. Chem., 269(1984)75
- [969] R.H. Hooker, A.J. Rest and I. Whitwell, J. Organomet. Chem., 266(1984)C27

- [1970] K. Suenkel, U. Nagel and W. Beck, J. Organomet. Chem., 251(1983)227
- [1971] R.S. Bly, G.S. Silverman, M.M. Hossain and R.K. Bly, Organometallics, 3(1984)642
- [1972] T. Forschner, K. Menard and A. Cutler, J. Chem. Soc. Chem. Commun., (1984)121
- [1973] L. Busetto, M. Monari, A. Palazzi, V. Albano and F. Demartin, J. Chem. Soc. Dalton Trans., (1983)1849
- [1974] R.P. Rosen, J.B. Hoke, R.R. Whittle, G.L. Geoffroy, J.P. Hutchinson and J.A. Zubieta, Organometallics, 3(1984)846
- [1975] L.S. Liebskind, M.E. Welker and V. Goedken, J. Am. Chem. Soc., 106(1984)441
- [1976] S.S. Ullah, M.E. Molla and M. Karim, J. Bangladesh Acad. Sci., 8(1984)75
- [1977] D. Seyferth, T.G. Wood, J.P. Fackler and A.M. Mazany, Organometallics, 3(1984)1121
- [1978] G. Lagrange, P. Cocolios and R. Guillard, J. Organomet. Chem., 260(1984)C16
- [1979] I.R. Butler, W.E. Lindsell and M.J.K. Thomas, J. Organomet. Chem., 262(1984)59
- [1980] A.I. Mikaya, V.G. Zaikin, V.V. Trusov, A.I. Nekhaev and V.D. Tyurin, J. Organomet. Chem., 254(1983)339
- [1981] R. Aumann, H. Heinen, G. Henkel, M. Dartmann and B. Krebs, Z. Naturforsch. B, 38(1983)1325
- [1982] M.E. Wright, T.M. Mezza, G.O. Nelson, N.R. Armstrong, V.W. Day and M.R. Thompson, Organometallics, 2(1983)1711
- [1983] E. Lindner, E. Schauss, W. Hiller and R. Fawzi, Angew. Chem., 96(1984)727
- [1984] M.M. Singh and R.J. Angelici, Inorg. Chem., 23(1984)2691
- [1985] R.P. Micciche, J.J. Briguglio and L.G. Sneddon, Organometallics, 3(1984)1396
- [1986] El Khbir Lhadi, H. Patin and A. Darchen, Organometallics,

- 3(1984)1128
- [987] C.P. Casey and W.H. Miles, J. Organomet. Chem., 254(1983)333
- [988] P. Vierling and J.G. Riess, J. Am. Chem. Soc., 106(1984)2432
- [989] D.L. Reger and E. Mintz, Organometallics, 3(1984)1759
- [990] D.L. Reger and C.A. Swift, Organometallics, 3(1984)876
- [991] J. Ros, X. Solans, M. Font - Alba and R. Mathieu, Organometallics, 3(1984)1014
- [992] Y.W. Chen, M.W. Renner and A.L. Balch, Organometallics, 2(1983)1888
- [993] M.F. Semmelhack, R. Tamura, W. Schnatter and J. Springer, J. Am. Chem. Soc., 106(1984)5363
- [994] E.O. Fischer, J. Schneider and K. Ackermann, Z. Naturforsch. B, 39(1984)468
- [995] C.P. Casey and W.H. Miles, Organometallics, 3(1984)808
- [996] G.H. Kuo, P. Helquist and R.C. Kerber, Organometallics, 3(1984)806
- [997] G. Groetsch and W. Malisch, J. Organomet. Chem., 262(1984)C38
- [998] A. Darchen, El Kbir Lhadi and H. Patin, J. Organomet. Chem., 259(1983)189
- [999] El Kbir Lhadi, H. Patin, A. Benoit, J.Y. le Marouille and A. Darchen, J. Organomet. Chem., 259(1983)321
- [1000] E.L. Hoel, G.B. Ansell and S. Leta, Organometallics, 3(1984)1633
- [1001] J.R. Matachek, R.J. Angelici, K.A. Schgart, K.J. Haller and R.F. Fenske, Organometallics, 3(1984)1038
- [1002] M.E. Wright and G.O. Nelson, J. Organomet. Chem., 263(1984)371
- [1003] G. Consiglio, F. Bangerter, C. Darpin, F. Morandini and V. Lucchini, Organometallics, 3(1984)1446
- [1004] J.P. Battioni, D. Dupre, P. Guerin and D. Mansuy, J. Organomet. Chem., 265(1984)53



- [1005] J.W. Kolis, E.M. Holt, J.A. Hriliac and D.F. Shriver, Organometallics, 3(1984)496
- [1006] E.O. Fischer, J. Schneider and D. Neugebauer, Angew. Chem. Int. Ed. Engl., 23(1984)820
- [1007] [a] M.A. Drezdson and D.F. Shriver, J. Mol. Catal., 21(1983)81; [b] C.P. Casey, M.W. Meszaros, S.R. Marder and P.J. Fagan, J. Am. Chem. Soc., 106(1984)3680
- [1008] F.U. Axe and D.S. Marynick, J. Am. Chem. Soc., 106(1984)6230
- [1009] V. Schurig, Tetrahedron Lett., 25(1984)2739
- [1010] S.S. Ullah, S.E. Kabir, M.E. Molla, A.K.F. Rahman and M. Karim, Indian J. Chem. A, 23(1984)637
- [1011] K.C. Brikman, A.J. Blakeney, W. Krone - Schmidt and J.A. Gladysz, Organometallics, 3(1984)1325
- [1012] U. Haenisch, E. Tagliaferri, R. Roulet and P. Vogel, Helv. Chim. Acta, 66(1983)2182
- [1013] M.I. Rybinskaya, L.V. Rybin, N.A. Stelzer, I.A. Garbusova and B.V. Lokshin, J. Organomet. Chem., 265(1984)291
- [1014] H. Fleckner, F.W. Grevels and D. Hess, J. Am. Chem. Soc., 106(1984)2027
- [1015] C.A. Barras, R. Roulet, P.A. Carrupt, F. Berchier and P. Vogel, Helv. Chim. Acta, 67(1984)986
- [1016] G.D. Annis, E.M. Hebblethwaite, S.T. Hodgson, D.M. Hollinshead and S.V. Ley, J. Chem. Soc. Perkin Trans. 1, (1983)2851
- [1017] J. Keijsper, J. Mul, G. van Koten and K. Vrieze, Organometallics, 3(1984)1732
- [1018] H. Brunner, H. Weber, I. Bernal and G.M. Reisner, Organometallics, 3(1984)163
- [1019] G. Cardaci, J. Chem. Soc. Dalton Trans., (1984)815
- [1020] R.S. Bly and G.S. Silverman, Organometallics, 3(1984)1765
- [1021] M. Marsi and M. Rosenblum, J. Am. Chem. Soc., 106(1984)7264
- [1022] G. Cardaci, G. Bellachioma and P. Zanazzi, Polyhedron,

2(1983)967

- [1023] A.M. Horton, D.M. Hollinshead and S.V. Ley, Tetrahedron, 40(1984)1737
- [1024] [a] C.P. Casey, P.J. Fagan, W.H. Miles and S.R. Marder, J. Mol. Catal., 21(1983)173; [b] E.J.S. Vichi, P.R. Raithby and M. McPartlin, J. Organomet. Chem., 256(1983)111
- [1025] J.R. Bleeke and M.K. Hays, Organometallics, 3(1984)506
- [1026] D. Touchard, P.H. Dixneuf, R.D. Adams, B.E. Segmueller, Organometallics, 3(1984)640
- [1027] D.L. Reger, K.A. Belmore, E. Mintz and P.J. McElligott, Organometallics, 3(1984)134
- [1028] J.C. Jeffery, K.A. Mead, H. Razay, F.G.A. Stone, M.J. Went and P. Woodward, J. Chem. Soc., Dalton Trans., (1984)1383
- [1029] D. de Montauzon and R. Mathieu, J. Organomet. Chem., 252(1983)C83
- [1030] V. Busetti, G. Granozzi, S. Aime, R. Gobetto and D. Osella, Organometallics, 3(1984)1510
- [1031] E. Keller and D. Wolters, Chem. Ber., 117(1984)1572
- [1032] F. Dahan and R. Mathieu, J. Chem. Soc., Chem. Commun., (1984)432
- [1033] A.J. Pearson, Pure Appl. Chem., 55(1983)1767
- [1034] A. Salzer and A. Rafner, Helv. Chim. Acta, 66(1983)1774
- [1035] C. Barras, R. Roulet and P. Vogel, Inorg. Chim. Acta, 82(1984)L1
- [1036] B.M.R. Bandara, A.J. Birch and L.F. Kelly, J. Org. Chem., 49(1984)2496
- [1037] A.J. Birch, W.D. Raverty and G.R. Stephenson, Organometallics, 3(1984)1075
- [1038] M.F. Semmelhack, J.W. Herndon and J.K. Liu, Organometallics, 2(1983)1885
- [1039] J.M. McCall, J.R. Morton, Y. LePage and K.F. Preston, Organometallics, 3(1984)1299

- [1040] S. Komiya, H. Minato, T. Ikariya, T. Yamamoto and A. Yamamoto, J. Organomet. Chem., 254(1983)83
- [1041] R. Corriu, N. Escudie and C. Guerin, J. Organomet. Chem., 264(1984)207
- [1042] T. Rukachaisirikul, S. Arabi, F. Hartstock, N.J. Taylor and A.J. Carty, Organometallics, 3(1984)1587
- [1043] F.J. Lalor, L.H. Brookes, G. Ferguson and M. Parvez, J. Chem. Soc. Dalton Trans., (1984)245
- [1044] H.W. Fruehauf, F. Seils, M.J. Romao and R.J. Goddard, Angew. Chem., 95(1983)1014
- [1045] R.D. Ernst, D.R. Wilson and R.H. Herber, J. Am. Chem. Soc., 106(1984)1646
- [1046] M.F. Semmelhack and H.T.M. Le, J. Am. Chem. Soc., 106(1984)2715
- [1047] G.T. Burns, E. Colomer and R.J.P. Corriu, Organometallics, 2(1983)1901
- [1048] J.W. Chinn and M.B. Hall, Organometallics, 3(1984)284
- [1049] B.M.R. Bandara and A.J. Birch, J. Organomet. Chem., 265(1984)C6
- [1050] D.A. Brown, N.J. Fitzpatrick, W.K. Glass and P.K. Sayal, Organometallics, 3(1984)1137
- [1051] M.F. Semmelhack and J.W. Herndon, J. Organomet. Chem., 265(1984)C15
- [1052] L. Harland, G.R. Stephenson and M.J. Whittaker, J. Organomet. Chem., 263(1984)C30
- [1053] J.A.S. Howell, D.T. Dixon and J.C. Kola, J. Organomet. Chem., 266(1984)69
- [1054] J.A.S. Howell, J.C. Kola, D.T. Dixon, P.M. Burkinshaw and M.J. Thomas, J. Organomet. Chem., 266(1984)83
- [1055] P.J. Parsons and E. Mincione, Tetrahedron Lett., 24(1983)5781
- [1056] A.J. Birch, W.D. Raverty, S.Y. Hsu and A.J. Pearson, J. Organomet. Chem., 260(1984)C59

- [1057] L.A. Paquette, R.G. Daniels and R. Gleiter, Organometallics, 3(1984)560
- [1058] R. Yamaguchi, Y. Nakazono and M. Kawanisi, Bull. Chem. Soc. Jpn., 57(1984)2331
- [1059] O. Eisenstein, W.M. Butler and A.J. Pearson, Organometallics, 3(1984)1150
- [1060] M.J. Buckingham, B.W. Fitzsimmons and I. Sayer, J. Chem. Soc. Chem. Commun., (1984)339
- [1061] L.A.P. Kane - Maguire, E.D. Honig and D.A. Sweigart, J. Chem. Soc. Chem. Commun., (1984)345
- [1062] A.J. Pearson, S.L. Kole and T. Ray, J. Am. Chem. Soc., 106(1984)6060
- [1063] Z. Goldschmidt and Y. Bakal, J. Organomet. Chem., 269(1984)191
- [1064] Z. Goldschmidt, S. Antebi and I. Goldberg, J. Organomet. Chem., 260(1984)105
- [1065] K. Broadley, N.G. Connelly, R.M. Mills, M.W. Whiteley and P. Woodward, J. Chem. Soc. Dalton Trans., (1984)683
- [1066] R.C. Kerber, D. Glasser and B. Luck, Organometallics, 3(1984)840
- [1067] S.S. Ullah, S.E. Kabir, A.K.F. Rahman and M. Karim, Indian J. Chem. B, 23(1984)103
- [1068] A. Madonik and D. Astruc, J. Am. Chem. Soc., 106(1984)2437
- [1069] N.G. Connelly, A.R. Lucy, R.M. Mills, M.W. Whiteley and P. Woodward, J. Chem. Soc. Dalton Trans., (1984)161
- [1070] E. Meier, R. Roulet and A. Scrivanti, Inorg. Chim. Acta, 45(1980)L267
- [1071] D. Kawka, P. Mues and E. Vogel, Angew. Chem., 95(1983)1006
- [1072] Z. Goldschmidt, Y. Bakal, D. Hoffer, A. Eisenstadt and P.F. Lindley, J. Organomet. Chem., 258(1983)53
- [1073] S.S. Lullah, S.E. Kabir, M.E. Molla and M. Karim, Indian J. Chem. A, 23(1984)404

- [1074] S.S. Ullah, S.E. Kabir, M.E. Molla and S.M. Wahiduzzaman, Indian J. Chem. A, 23(1984)180
- [1075] H. Butenschoen and A. de Meijere, Angew. Chem., 96(1984)722
- [1076] L.V. Rybin, N.A. Stelzer, I.A. Garbusova, B.V. Lokshin and M.I. Rybinskaya, J. Organomet. Chem., 265(1984)295
- [1077] C.C. Santini and F. Mathey, J. Organomet. Chem., 266(1984)285
- [1078] Y.F. Yu, J. Gallucci and A. Wojcicki, J. Chem. Soc. Chem. Commun., (1984)653
- [1079] J.C. Daran and Y. Jeannin, Organometallics, 3(1984)1158; J. Ros, R. Mathieu, X. Solans and M. Font - Alba, J. Organomet. Chem., 260(1984)C40
- [1080] A. Zimniak, W. Jasobedzki and J. Kiljanska, Bull. Pol. Acad. Sci. Chem., 31(1983)9
- [1081] W. Kuei, Huaxue Shijie, 25(1984)164; CA, 101(1984)171444f
- [1082] D. Catheline and D. Astruc, Nouv. J. Chim., 8(1984)381
- [1083] L.Y. Hsu, S.J. Hathaway and L.A. Paquette, Tetrahedron Lett., 25(1984)259
- [1084] M. Heberhold and L. Haumaier, Angew. Chem. Int. Ed. Engl., 24(1984)521
- [1085] R.G. Sutherland, A. Piorko and C.C. Lee, Organometallics, 3(1984)331
- [1086] V. Guerschais and D. Astruc, J. Chem. Soc. Chem. Commun., (1983)1115
- [1087] C.C. Lee, U.S. Gill and R.G. Sutherland, J. Organomet. Chem., 267(1984)157
- [1088] A.A. Erdman, V.L. Shirokii, Z.P. Zubroichuk, N.A. Maier and Yu.A. Ol'dekop, Vestsi Akad. Navuk BSSR. Ser. Khim. Navuk, (1984)74; CA, 101(1984)171445g
- [1089] R.P. Micciche, J.J. Briguglio and L.G. Sneddon, Inorg. Chem., 23(1984)3992
- [1090] F. Mercier and F. Mathey, J. Organomet. Chem., 263(1984)55

- [1091] M.F. Daniel, A.J. Leadbetter, M.A. Mazid and J. Piper, J. Chem. Soc. Faraday Trans. 2, 79(1983)1663
- [1092] S. Ikuta, I. Motoyama and H. Sano, Radiochem. Radioanal. Lett., 58(1983)329
- [1093] R. Gleiter, M.C. Boehm and R.D. Ernst, J. Electron Spectrosc. Relat. Phenom., 33(1984)269
- [1094] C. Guimon, D. Gonbeau, G. Pfister - Guillouzo, G. de Lauzon and F. Mathey, Chem. Phys. Lett., 104(1984)560
- [1095] F. Delgado - Pena, D.R. Talham and D.O. Cowan, J. Organomet. Chem., 253(1983)C43
- [1096] B. Lukas, R.M.G. Roberts, J. Silver and A.S. Wells, J. Organomet. Chem., 256(1983)103
- [1097] A. Stebler, A. Furrer and J.H. Ammeter, Inorg. Chem., 23(1984)3493
- [1098] J.P. Mariot, P. Michaud, S. Lauer, D. Astruc, A.X. Trautwein and F. Varret, J. Phys. (Les Ulis, Fr.), 44(1983)1377
- [1099] G. Neshvad, R.M.G. Roberts and J. Silver, J. Organomet. Chem., 260(1984)319
- [1100] I. Motoyama, K. Suto, M. Katada and H. Sano, Chem. Lett., (1983)1215
- [1101] M.J. Cohn, M.D. Timken and D.N. Hendrickson, J. Am. Chem. Soc., 106(1984)6683
- [1102] R.M.G. Roberts and J. Silver, J. Organomet. Chem., 263(1984)235
- [1103] K.E. Lewis and G.P. Smith, J. Am. Chem. Soc., 106(1984)4650
- [1104] C. Lo Sterzo and G. Ortaggi, Gazz. Chim. Ital., 113(1983)721
- [1105] A.M. McNair, J.L. Schrenk and K.R. Mann, Inorg. Chem., 23(1984)2633
- [1106] P. Michaud, C. Lapinte and D. Astruc, Ann. N.Y. Acad. Sci., 415(1983)97
- [1107] M.A. El - Hashash, S. El - Nagdy and R. Saleh, Indian J. Chem. A, 22(1983)605

- [1108] A.G. Nagy and S. Toma, J. Organomet. Chem., 266(1984)257
- [1109] P. Lemoine, M. Gross, P. Braunstein, F. Mathey, B. Deschamps and J.H. Nelson, Organometallics, 3(1984)1303
- [1110] W.J. Bowyer, W.E. Geiger and V. Boekelheide, Organometallics, 3(1984)1079
- [1111] A. Grimes and S.R. Logan, Inorg. Chim. Acta, 45(1980)L223
- [1112] E.W. Neuse, R. Loonat and J.C.A. Boeyens, Transition Met. Chem., 9(1984)12
- [1113] B. Czech, A. Czech, S.I. Kang and R.A. Bartsch, Chem. Lett., (1984)37
- [1114] L. Zhang, P. Hu and G. Zhang, Youji Huaxue, (1983)277
- [1115] J.C. Boeyens, P.C. Lalloo, E.W. Neuse and H.H. Wei, S. Afr. J. Chem., 37(1984)32
- [1116] A.A. Koridze, N.M. Astakhova and P.V. Petrovskii, J. Organomet. Chem., 254(1983)345
- [1117] G.Z. Suleimanov, P.V. Petrovskii, Yu.S. Bogachev, I.L. Zhuravleva, E.I. Fedin and I.P. Beletskaya, J. Organomet. Chem., 262(1984)C35
- [1118] B. Gautheron and G. Tainturier, J. Organomet. Chem., 262(1984)C30
- [1119] P.T. Barger and J.E. Bercaw, Organometallics, 3(1984)278
- [1120] F.H. Koehler, W.A. Geike, P. Hofmann, U. Schubert and P. Stauffert, Chem. Ber., 117(1984)904
- [1121] D. Greissinger, W. Malisch and H.A. Kaul, J. Organomet. Chem., 252(1983)C23
- [1122] G.V. Gridunova, V.E. Shklover, Yu.T. Struchkov and L.P. Asatiani, Zh. Strukt. Khim., 24(1983)93
- [1123] B. Deschamps, F. Mathey, J. Fischer and J.H. Nelson, Inorg. Chem., 23(1984)3455
- [1124] G.Y. Lin and J. Takats, J. Organomet. Chem., 269(1984)C4
- [1125] T. Hayashi, K. Kabeta and M. Kumada, Tetrahedron Lett.,

25(1984)1499

- [1126] M. Onishi, K. Hiraki and A. Iwamoto, J. Organomet. Chem., 262(1984)C11
- [1127] E.G. Perevalova, E.I. Smyslova, V.P. Dyadchenko and K.I. Granberg, Izv. Akad. Nauk SSSR, Ser. Khim., (1984)956; CA, 101(1984)91116k
- [1128] M.M. Kubicki, R. Kergoat, J.Y. de Vanssay, J.Y. Le Gall and J.E. Guerschais, Polyhedron, 3(1984)365
- [1129] M. Watanabe, H. Ichikawa, I. Motoyama and H. Sano, Bull. Chem. Soc. Jpn, 56(1983)3291
- [1130] D.A. Lemonovskii, I.F. Urazovskii, T.V. Baukova, I.L. Arkhipov, R.A. Stukan and E.G. Perevalova, J. Organomet. Chem., 264(1984)283
- [1131] R. Yin and X. Zhen, Lanzhou Daxue Xuebao, Ziran Kexueban, 19(1983)41; CA, 100(1984)103578g
- [1132] A.J. Lindsay, S. Kim, R.A. Jacobson and R.J. Angelici, Organometallics, 3(1984)1523
- [1133] S.G. Davies, S.J. Simpson and S.E. Thomas, J. Organomet. Chem., 254(1983)C29
- [1134] K.H. Pannell and R.N. Kapoor, J. Organomet. Chem., 269(1984)59
- [1135] J.C. Watkins and M. Rosenblum, Tetrahedron Lett., 25(1984)2097
- [1136] S.B. Fergusson, L.J. Sanderson, T.A. Shackleton and M.C. Baird, Inorg. Chim. Acta, 83(1984)L45
- [1137] D. Miholova and A.A. Vlcek, Inorg. Chim. Acta, 73(1983)249
- [1138] L. Parkanyi, K.H. Pannell and C. Hernandez, J. Organomet. Chem., 252(1983)127
- [1139] W. Angerer, K. Fiederling, G. Groetsch and W. Malisch, Chem. Ber., 116(1983)3947
- [1140] A.R. Manning, G. McNally, R. Davis and C.C. Rowland, J. Organomet. Chem., 259(1983)C15
- [1141] L.I. Zakharkin, A.I. Kovredov, V.A. Ol'Shevskaya and V.A. Antonovich, J. Organomet. Chem., 267(1984)81



- [1142] T.C.T. Chang and M. Rosenblum, Isr. J. Chem., 24(1984)99
- [1143] C. Lapinte and D. Astruc, J. Organomet. Chem., 260(1984)C13
- [1144] D. Catheline and D. Astruc, Organometallics, 3(1984)1094
- [1145] D. Catheline and D. Astruc, J. Organomet. Chem., 266(1984)C11
- [1146] T.Yu. Orlova, V.N. Setkina and D.N. Kursanov, J. Organomet. Chem., 267(1984)309
- [1147] W. Angerer, M. Luksa and W. Malisch, J. Organomet. Chem., 253(1983)C36
- [1148] H. Nakazawa, W.E. Buhro, G. Bertrand and J.A. Gladysz, Inorg. Chem., 23(1984)3431
- [1149] E. Lindner and C.P. Krieg, J. Organomet. Chem., 269(1984)65
- [1150] C. Combes, R.J.P. Corriu, G. Dabosi, B.J.L. Henner and M. Martineau, J. Organomet. Chem., 270(1984)141
- [1151] R.S. Glass and W.M. McConnell, Organometallics, 3(1984)1630
- [1152] H. Stolzenberg, W.P. Fehlhammer, M. Monari, V. Zanotti and L. Busetto, J. Organomet. Chem., 272(1984)73
- [1153] T.C. Flood and K.D. Campbell, J. Am. Chem. Soc., 106(1984)2853
- [1154] N. Aktogu, H. Felkin, G.J. Baird, S.G. Davies and O. Watts, J. Organomet. Chem., 262(1984)49
- [1155] S.G. Davies, J. Hibberd, S.J. Simpson, S.E. Thomas and O. Watts, J. Chem. Soc. Dalton Trans., (1984)701
- [1156] S.G. Davies and S.J. Simpson, J. Organomet. Chem., 268(1984)C53
- [1157] C. Lapinte, D. Catheline and D. Astruc, Organometallics, 3(1984)817
- [1158] I.R. Butler and W.R. Cullen, Organometallics, 3(1984)1846
- [1159] A. Stasunik and W. Malisch, Z. Naturforsch. B, 39(1984)56
- [1160] P.M. Treichel and L.D. Rosenhein, Inorg. Chem., 23(1984)4018
- [1161] A.S. Batsanov and Yu.T. Struchkov, J. Organomet. Chem., 266(1984)295

- [1162] K.E. Schwarzhaus and H. Schottenberger, Z. Naturforsch. B, 38(1983)1493
- [1163] M.L. Ziegler and R. Korswagen, Bol. Soc. Quim. Peru, 48(1982)135
- [1164] G.J. Baird, J.A. Bandy, S.G. Davies and K. Prout, J. Chem. Soc. Chem. Commun., (1983)1202
- [1165] P.M. Treichel, M.S. Schmidt and S.D. Koehler, J. Organomet. Chem., 258(1983)209
- [1166] D. Catheline and D. Astruc, J. Organomet. Chem., 269(1984)C33
- [1167] A. Castel, P. Riviere, J. Satge, J.J.E. Moreau and R.P. Corriu, Organometallics, 2(1983)1498
- [1168] A.F. Hepp, J.P. Blaha, C. Lewis and M.S. Wrighton, Organometallics, 3(1984)174
- [1169] R.H. Hooker, K.A. Mahmoud and A.J. Rest, J. Chem. Soc. Chem. Commun., (1983)1022
- [1170] K.D. Janda, W.W. McConnell, G.O. Nelson and M.E. Wright, J. Organomet. Chem., 259(1983)139
- [1171] G. Thum and W. Malisch, J. Organomet. Chem., 264(1984)C5
- [1172] H.U. Wekel and W. Malisch, J. Organomet. Chem., 264(1984)C10
- [1173] J.F. Harrod and E. Pelletier, Organometallics, 3(1984)1064
- [1174] S.G. Davies, S.J. Simpson and V.D. Parker, J. Chem. Soc. Chem. Commun., (1984)352
- [1175] H.H. Wei and K.T. Dai, J. Chin. Chem. Soc. (Taipei), 30(1983)131
- [1176] M. Cooke, N.J. Forrow and S.A.R. Knox, J. Chem. Soc. Dalton Trans., (1983)2435
- [1177] B.H. Chang, P.C. Coil, M.J. Brown and K.W. Barnett, J. Organomet. Chem., 270(1984)C23
- [1178] J.A.S. Howell, P. Mathur, R. Kumar, A.R. Manning and F.S. Stephens, J. Organomet. Chem., 262(1984)227
- [1179] R.B. English, Acta Crystallogr. C, 40(1984)1567

- [1180] D. Bickar, B. Lukas, G. Neshvad, R.M.G. Roberts and J. Silver, J. Organomet. Chem., 263(1984)225
- [1181] T. Yamamoto, K. Sanechika, A. Yamamoto, M. Takada, I. Motoyama and H. Sano, IOorg. Chim. Acta, 73(1983)75
- [1182] A. Darchen, J. Chem. Soc. Chem. Commun., (1983)768
- [1183] E. Deschamps, F. Mathey, C. Knobler and Y. Jeannin, Organometallics, 3(1984)1144
- [1184] M. Sato, M. Kubo, S. Ebine and S. Akabori, Bull. Chem. Soc. Jpn., 57(1984)421
- [1185] G.J. Baird and S.G. Davies, J. Organomet. Chem., 262(1984)215
- [1186] V.I. Boev and A.V. Dombrovskii, Zh. Obshch. Khim., 54(1984)1617; CA, 101(1984)211396a
- [1187] G.J. Baird, S.G. Davies and T.R. Maberly, Organometallics, 3(1984)1764
- [1188] W. Kalcher, W.A. Herrmann, C. Pahl and M.L. Ziegler, Chem. Ber., 117(1984)69
- [1189] V.G. Adrianov, Yu. T. Struchkov, V.N. Postnov, E.I. Klimova and V.A. Sazonova, J. Organomet. Chem., 272(1984)81
- [1190] K. Sato, M. Konno and H. Sano, Chem. Lett., (1984)17
- [1191] R.J. Ranson and R.M.G. Roberts, J. Organomet. Chem., 260(1984)307
- [1192] A. Perjessy and S. Toma, Collect. Czech. Chem. Commun., 48(1983)1635
- [1193] D.R. Talham and D.O. Cowan, Organometallics, 3(1984)1712
- [1194] A.G. Nagy, J. Organomet. Chem., 270(1984)327
- [1195] D. Astruc, Tetrahedron, 39(1983)4027
- [1196] H.R. Allcock, K.D. Lavin, G.H. Riding and R.R. Whittle, Organometallics, 3(1984)663
- [1197] J.C. Barborak and J.M. Chance, J. Org. Chem., 49(1984)703
- [1198] Z.D. Belykh, E.A. Mal'tseva and M.D. Reshetova, Sint. Metody

Osn. Elementoorg. Soedin., (1982)66

- [1199] P. Bickert, B. Hildebrandt and K. Hafner, Organometallics, 3(1984)653
- [1200] T.E. Bitterwolf and M.J. Golightly, Inorg. Chim. Acta, 84(1984)153
- [1201] V.I. Boev and A.V. Dombrovskii, Zh. Obshch. Khim., 54(1984)970; CA, 101(1984)55289k
- [1202] V.I. Boev and A.V. Dombrovskii, Zh. Obshch. Khim., 54(1984)1192; CA, 101(1984)21392w
- [1203] V.I. Boev and A.V. Dombrovskii, Zh. Obshch. Khim., 54(1984)1863; CA, 101(1984)211402z
- [1204] I.R. Butler and W.H. Cullen, Can. J. Chem., 61(1983)2354
- [1205] Y.K. Chung, E.D. Honig and D.A. Sweigart, J. Organomet. Chem., 256(1983)277
- [1206] A. Clearfield, C.J. Simmons, H.P. Withers and D. Seyferth, Inorg. Chim. Acta, 75(1983)139
- [1207] P.J. Hammond, P.D. Beer and C.D. Hall, J. Chem. Soc. Chem. Commun., (1983)1161
- [1208] J.F. Helling and U.S. Gill, J. Organomet. Chem., 264(1984)353
- [1209] E.A. Kalennikov, Khim. Khim. Tekhnol. (Minsk), 16(1983)77; CA, 100(1984)139293z
- [1210] A. Kasahara, T. Izumi and K. Saito, Bull. Chem. Soc. Jpn., 56(1983)2865
- [1211] E.I. Klimova, A.N. Pushin and V.A. Sazonova, J. Organomet. Chem., 270(1984)319
- [1212] I.Yu. Kozak, G.N. Yashchenko and A.K. Sheinkman, Khim. Geterotsikl. Soedin., (1984)996; CA, 101(1984)171451f
- [1213] C.C. Lee, A. Piorkjo, B.R. Steele, U.S. Gill and R.G. Sutherland, J. Organomet. Chem., 256(1983)303
- [1214] S.R. Logan and G.A. Salmon, J. Chem. Soc. Perkin Trans. 2, (1983)1781
- [1215] A.M. Madonik, D. Mandon, P. Michaud, C. Lapinte and D. Astruc,

- J. Am. Chem. Soc., 106(1984)3381
- [1216] H. Martin, R. Herrmann and I. Ugi, J. Organomet. Chem., 269(1984)87
- [1217] B. McCulloch and C.H. Brubaker, Organometallics, 3(1984)1707
- [1218] J. Mirek and B. Kawalek, Pol. J. Chem., 56(1982)1121
- [1219] T.Yu. Orlova, V.N. Setkina, V.F. Sizoi and D.N. Kursanov, J. Organomet. Chem., 252(1983)201
- [1220] S.R. Patil, U.N. Kantak and D.N. Sen, J. Indian Chem. Soc., 59(1982)1300
- [1221] B.V. Polyakov, V.P. Tverdokhlebov, T.V. Subbotina and I.V. Tselinskii, Zh. Obshch. Khim., 53(1983)2392
- [1222] B.V. Polyakov, V.P. Tverdokhlebov, I.V. Tselinskii, N.M. Bakstova and G.M. Frolova, Zh. Obshch. Khim., 53(1983)2046
- [1223] V.N. Postnov, Yu.N. Polivin and V.A. Sazonova, Dokl. Akad. Nauk SSSR, 271(1983)113
- [1224] V.N. Postnov, Yu.N. Polivin and V.A. Sazonova, Dokl. Akad. Nauk SSSR, 271(1983)1399
- [1225] V.N. Postnov, Yu.N. Polivin and V.A. Sazonova, Dokl. Akad. Nauk SSSR, 276(1984)617; CA, 101(1984)211394y
- [1226] V. Rapic and I. Habus, Croat. Chem. Acta, 57(1984)265
- [1227] V. Rapic and L. Korontos, J. Organomet. Chem., 260(1984)219
- [1228] N.J. Singletary, M. Hillman, H. Dauplaise, A. Kvik and R.C. Kerber, Organometallics, 3(1984)1427
- [1229] M. Vlcek, K. Handlir, J. Holecek and I. Kolb, Z. Chem., 23(1983)294
- [1230] M. Hillman and A. Kvik, Organometallics, 2(1983)1780
- [1231] L.G. Kuz'mina, Yu.T. Struchkov, D.A. Lemenovskii, I.F. Urazovskii, I.E. Nifant'ev and E.G. Perevalova, Koord. Khim., 9(1983)1212
- [1232] M. Hillman, E. Fujita, H. Dauplaise, A. Kvik and R.C. Kerber, Organometallics, 3(1984)1170

- [1233] E. Solcaniova, S. Toma, M. Salisova and T. Liptaj, Org. Magn. Reson., 21(1983)429
- [1234] H.H. Wei and S.J. Chang, J. Chin. Chem. Soc. (Taipei), 31(1984)9
- [1235] M. Watanabe, K. Sato, I. Motoyama and H. Sano, Chem. Lett., (1983)1775
- [1236] M.G. Peterleitner, L.I. Denisovich, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya and D.V. Zagorevskii, Izv. Akad. Nauk SSSR, Ser. Khim., (1983)2814; CA, 100(1984)175020s
- [1237] S. Akabori, Y. Habata and M. Sato, Bull. Chem. Soc. Jpn, 57(1984)68
- [1238] S. Akabori, S. Shibahara, Y. Habata and M. Sato, Bull. Chem. Soc. Jpn, 57(1984)63
- [1239] B. Czech, A. Piorko and R. Annunziata, J. Organomet. Chem., 255(1983)365
- [1240] M. Hisatome, Y. Kawajiri, J. Watanabe, M. Yoshioka and K. Yamakawa, J. Organomet. Chem., 266(1984)147
- [1241] M. Hisatome, J. Watanabe and K. Yamakawa, J. Organomet. Chem., 266(1984)159
- [1242] M. Hisatome, J. Watanabe, K. Yamakawa, K. Kozawa and T. Uchida, J. Organomet. Chem., 262(1984)365
- [1243] U.T. Mueller - Westerhoff and A. Nazzari, J. Am. Chem. Soc., 106(1984)5381
- [1244] T. Nomura, Bull. Chem. Soc. Jpn, 56(1983)2937
- [1245] H. Paulus, K. Schloegl and W. Weissensteiner, Monatsh. Chem., 114(1983)799
- [1246] M. Sato, S. Tanaka, S. Ebine and S. Akabori, Bull. Chem. Soc. Jpn, 57(1984)1929
- [1247] D. Tanner and O. Wennerstroem, Acta Chem. Scand. B, 37(1983)693
- [1248] C.A. Bunton and W.E. Watts, J. Chem. Soc. Perkin Trans. 2, (1983)1591
- [1249] A. Waleh, G.H. Loew and U.T. Mueller - Westerhoff, Inorg. Chem., 23(1984)2859