

12. Platinum 1993

Peter K. Byers

CONTENTS


INTRODUCTION	431
12.1 PLATINUM(IV)	431
12.2 PLATINUM(II)	432
12.2.1 Complexes with phosphine donor ligands	432
12.2.2 Complexes with nitrogen donor ligands	441
12.2.3 Complexes with sulfur group donor ligands	443
12.2.4 Complexes with mixed donor ligands	444
12.3 PLATINUM(I)	447
12.4 PLATINUM(0)	447
REFERENCES	448

INTRODUCTION


The coordination chemistry of platinum reported during the year 1993 has been reviewed. Although this account does not intend or claim to be comprehensive, examples have been selected to represent important features. Only the primary journals are covered and were searched using both platinum (and related) as a keyword and being present in the title. The review covers the chemistry of platinum in its 0 → +4 oxidation states, but does not include the catalytic activity of platinum complexes, complexes containing only Pt-C bonds or complexes containing nucleotide or nucleoside bases. Classification of the complexes is according to the oxidation state and donor atom. For complexes containing mixed donor sets the priority ordering P>N>S applies, and neutral donors have priority over charged donor atoms; mixed donor didentate ligands are treated separately.

12.1 PLATINUM(IV)

Reaction of [PtMe₂(phen)] with RE-ER (ER = OH, OC(O)Ph, SMe, SPh, SePh) gives [Pt(Me)₂(ER)₂(phen)] (1) as a mixture of *trans*- (major) and *cis*- (minor) isomers. For (1; ER = SePh) the solid state structure was determined by X-ray crystallography [1]. Treatment of the platinum(IV) iodide [PtMe₃(μ-I)]₄ with the aminothiols HS(CH₂)₂NEt₂ in methanol afforded the trinuclear species [(PtMe₃)₃{(μ-I), μ-S(CH₂)₂NEt₂}₂] (2). The complex displays a cubane type structure but with one less PtMe₃ vertex, as revealed by an X-ray analysis [2].



A novel *tetrakis*(arylalkyltriazenido) derivative of platinum(IV) has been reported. The complex [C₆H₅NNN(CH₂)₅CH₃]₄Pt (3) was characterised by ³¹Pt NMR spectroscopy and single crystal X-ray crystallography [3]. The first crystal structure of a platinum(IV)-cyclobutane bearing an electron withdrawing aldehyde functionality (4) has been described [4].





Electrospray mass spectra have been obtained for a variety of organo-platinum(IV) compounds containing polydentate *N*-donor ligands. In every case the intact ion was observed as the base peak at low ion source energies [5]. The compounds Cs₂[MCl₆] (M = Rh, Pt, Ir) have been shown to be isomorphous by powder diffraction [6].

12.2 PLATINUM(II)



12.2.1 Complexes with phosphine donor ligands

The dinuclear, double-bridged acetylide complexes [Pt(μ-C≡CR)(C₆F₅)(PPh₃)₂]₂ (R = Ph, Bu, SiMe₃) (5) have been synthesised from *trans*-[Pt(C≡CR)₂(PPh₃)₂] and *cis*-[Pt(C₆F₅)₂(thf)₂]. Treatment of the dimeric complexes with neutral ligands (L = py, PPh₃) produced the mononuclear


species *trans*-[Pt(C₆F₅)(C≡CR)(PPh₃)(L)]. The solid state structure of (5; R = Ph) was established by X-ray diffraction methods [7]. The dinuclear platinum(II) complexes [Pt₂(μ-dppm)₂(μ-C≡CPh)(C≡CPh)₂]⁺ and [Pt₂(μ-dppm)₂(μ-C≡C^tBu)(C≡C^tBu)Cl]⁺ (6) containing bridging and terminal acetylide ligands have been prepared by reaction of [Pt(dppm-*P,P'*)₂]Cl₂ with [Hg(PhC≡C)₂] or [Hg(^tBuC≡C)₂] respectively. The crystal structures of both complexes were reported [8].



Reaction of [Pt₂(C₆F₅)₄(μ-X)₂]²⁻ (X = Cl, Br, I) with dppm at room temperature affords [(C₆F₅)₂XPt(μ-dppm)PtX(C₆F₅)₂]²⁻ as the kinetic products. Subsequent reaction of these di-anions with AgClO₄ gave the mono-anions (7). The bridging system in these complexes is exceptionally stable, and was not cleaved by anionic (X⁻) or neutral ligands (*e.g.* py, PPh₃) [9]. Symmetrical and unsymmetrical dppm-bridged platinum(II) complexes (8) have also been prepared, in this instance by reaction of compounds of the form [PtR(dppm-*P,P'*)(dppm-*P*)]⁺ with [PtRCl(cod)] and [PtR'Cl(cod)], respectively. In a similar series of reactions, the mixed metal complexes [PtPdRMe(μ-Cl)(μ-dppm)₂]PF₆ were obtained upon addition of [PdMeCl(cod)] to [PtR(dppm-*P,P'*)(dppm-*P*)]⁺ [10].




The formylketenyl platinum complex *trans*-[PtH{η¹-C(CHO)CO}(PCy₃)₂] reacts with acids to cleave the Pt-C bond and generate the species C₃H₂O₂, which is subsequently trapped by an excess of the starting material. The reaction gave rise to *trans*-[PtH(C₆H₃O₄)(PCy₃)₂] (9), where C₆H₃O₄ is an α-pyrone ligand σ-bonded to platinum; the solid state structure of (9) was established [11].




(9)


The reaction of (2-propenyl)*bis*(triphenylphosphine)platinum(II) triflate with unsaturated organic substrates has been investigated. Alkynols {3-buten-1-ol and (\pm)-4-pentyn-2-ol} produce the vinylcarbene species (10), whilst reaction with nitriles (acrylonitrile and ethyl cyanoacetate) affords the *N*-bound nitrile complexes (11). Reaction with methyl- and ethyl acrylates, on the other hand, results in the formation of the chelated compounds (12); the structure of (12; R = Me) was determined using X-ray diffraction [12].



(10*; R=H, Me)




(11*)




(12*; R=Me, Et)


Insertion reactions of isocyanides into the M-C bond of (σ -allenyl)platinum(II) complexes have been reported. When the R groups of the allenyl ligand are small, new (metallovinyl)ketenimine compounds *trans*-[PtX{C(CRR')(CH=C=NR'')}](PPh₃)₂ (13) are formed, whilst with large R groups the insertion of the isocyanide is sterically hindered and ionic *trans*-[Pt{C(H)=C=CRR'}](CN'Bu)(PPh₃)₂]Br results [13]. A new type of ligating behaviour of the phosphirene ring system (14) has been described, in which novel dimeric platinum(II) chlorophosphane complexes (15) result from Pt-Cl bond insertion and phosphirene ring opening [14].



(13)




(14)




(15)

Dichloroplatinum(II) complexes containing chelating *bis*-phosphine ligands react with diazo-compounds to give Pt-Cl insertion products. Thus, $[\text{PtCl}_2(\text{S,S-skewphos})]$ reacted with $\text{Me}_3\text{SiCHN}_2$ to give $[\text{PtCl}(\text{S-CHClSiMe}_3)(\text{S,S-skewphos})]$ with high diastereoselectivity [15]. In a similar series of reactions, $[\text{PtCl}_2(\text{R,R-skewphos})]$ reacted with RCHN_2 ($\text{R} = \text{SiMe}_3$, $\text{P}(\text{O})(\text{OMe})_2$, CO_2Me) to give mono-insertion products (16) in which (*R*) stereochemistry is preferred for the newly created chiral centre [16].




Acetylenes, $\text{RC}\equiv\text{CR}$ ($\text{R} = \text{Ph}$, Pr), insert into the Pt-Si bond of *trans*- $[\text{PtX}(\text{SiMe}_3)(\text{PEt}_3)_2]$ ($\text{X} = \text{Br}$, I) to give *trans*- $[\text{PtX}\{\text{C}(\text{R})=\text{C}(\text{R})\text{SiMe}_3\}(\text{PEt}_3)_2]$; the structure of *trans*- $[\text{PtX}\{\text{C}(\text{Ph})=\text{C}(\text{Ph})\text{SiMe}_3\}(\text{PEt}_3)_2]$ was determined by X-ray crystallography [17]. A Fischer-type silylene complex of platinum, *trans*- $[\text{Pt}(\text{H})\{\text{Si}(\text{SEt})_2\}(\text{PCy}_3)_2]\text{BPh}_4$ (17), has been prepared from *cis*- $[\text{Pt}(\text{H})\{\text{Si}(\text{SEt})_3(\text{PCy}_3)_2\}]$ upon reaction with Me_3SiOTf and NaBPh_4 . The solid state structure of the silylene complex was reported [18].







The syntheses of two series of cyclopentadienyl complexes have appeared. In the first account, reaction of $[\text{Pt}(\text{OTf})_2(\text{P-P})]$ ($\text{P-P} = \text{dppm}$, dppe , dppp) with $\text{Ti}(\text{C}_5\text{H}_5)$ to produce the ionic compounds $[\text{Pt}(\eta^5\text{-C}_5\text{H}_5)(\text{P-P})\text{OTf}]$ was described [19]. In the second report, the neutral compounds $[\text{Pt}(\eta^5\text{-C}_5\text{H}_4\text{R})(\text{Ph})(\text{L})]$ ($\text{Y} = \text{H}$, Me ; $\text{L} = \text{PCy}_3$, PPh_3) were obtained directly on treatment of $[\text{Pt}(\text{Ph})(\mu\text{-OH})(\text{L})_2]$ with cyclopentadiene or methylcyclopentadiene [20]. A range of complexes of the type *cis*- $[\text{Pt}(\text{Ar})_2(\text{PPh}_3)_2]$ ($\text{Ar} = \text{aryl group}$) with substituents of different van der Waals volumina and electronic character in both *ortho*-positions of each platinum bonded phenyl ring have been synthesised, and studied with respect to substituent influences on their photochemical reactivity [21].

The *trans*-spanning ligand $1,3\text{-C}_6\text{H}_4(\text{CH}_2\text{PPh}_2)_2$ undergoes cyclometallation with either $[\text{PtMeCl}(\text{cod})]$ or $[\text{PtCl}_2(\text{cod})]$ to give (18; $\text{X} = \text{Cl}$), containing the ligand bound to platinum in a


P,C,P'-tridentate fashion. Successive treatment of (18; X = Cl) with AgBF_4 and KOH gives (18; X = OH) which reacts with CO to produce the hydroxycarbonyl (18; X = CO_2H). The solid state structure of the hydroxycarbonyl species was determined using X-ray diffraction methods [22].




Treatment of π -allyl platinum(II) complexes bearing a methoxymethoxy group at the 2-position with base gives oxodimethylmethane complexes (19) in contrast to 2-hydroxy substituted π -allyl complexes *via* acidic hydrolysis (20) [23]. A versatile route to 2-substituted π -allyl complexes, *e.g.* (21), from addition of hard and soft nucleophiles to the cationic η^3 -propargyl complex (22) has been described [24].





A series of dinuclear pyrazolato-bridged platinum(II) complexes of the form $[\text{Pt}_2\text{Cl}_2(\mu\text{-L})_2(\text{PR}_3)_2]$ (HL = pz, 3,5-Me₂pz; PR_3 = PEt_3 , PMe_2Ph , PMePh_2) have been prepared by reaction of the corresponding acetate-bridged analogue with HL. A single crystal structure analysis of a representative complex, $[\text{Pt}_2\text{Cl}_2(\mu\text{-pz})_2(\text{PMe}_2\text{Ph})_2]$, was also reported [25]. In a similar account, the synthesis of a second series of pyrazolato-bridged platinum(II) dimers was described. Thus, treatment of $[\text{Pt}_2(\text{Ar})_2(\mu\text{-Cl})_2(\text{PR}_3)_2]$ (Ar = Ph, $\text{C}_6\text{H}_4\text{Me-4}$, $\text{C}_4\text{H}_3\text{S-2}$; PR_3 = PBu_3 , PMe_2Ph , PMePh_2) with HL (L = pz, 3,5-Me₂pz, 3,4,5-Me₃pz) and NaOH gave $[\text{Pt}_2(\text{Ar})_2(\mu\text{-L})_2(\text{PR}_3)_2]$ (23) [26].



Reaction of *cis*-[PtCl₂(dmsO)₂] with Ph₂Ppy gave as the main product *cis*-[PtCl₂(Ph₂Ppy)(dmsO)], in which Ph₂Ppy is η¹-bonded through phosphorus to platinum. However, treatment of *cis*-[PtCl₂(Ph₂Ppy)(dmsO)] with [PtMe₂(dmsO)₂] gave the dinuclear platinum(II) compound [MeClPt(μ-Cl)(μ-Ph₂Ppy)PtMe(dmsO)] (24) and the Pt(I)-Pt(I) dimer [Pt₂Cl₂(μ-Ph₂Ppy)₂]; both were structurally characterised by an X-ray diffraction study [27].




The preparation and characterisation (by X-ray diffraction and multinuclear NMR spectroscopy) of mono- and bis-platinated derivatives of two diaziridines (25) has been reported [28]. Reaction of 1,5-Ph₄P₂N₄Se₂ with [PtCl₂(PEt₃)₂] produced the η¹-N-bonded adducts [PtCl₂(PEt₃)]_n(1,5-Ph₄P₂N₄Se₂) [n = 1; 2 (26)] [29].




Treatment of $[\text{PtX}_2\{\text{P}(\text{CH}_2\text{OH})_3\}_2]$ ($\text{X} = \text{Cl}, \text{Br}$) with AgY ($\text{Y} = \text{NO}_3, \text{BF}_4, \text{PF}_6, \text{ClO}_4$) in H_2O gave three species, which were identified as $[\text{Pt}(\text{OH}_2)_2\{\text{P}(\text{CH}_2\text{OH})_3\}_2]^{2+}$, $[\text{Pt}_2(\mu\text{-OH})_2\{\text{P}(\text{CH}_2\text{OH})_3\}_4]^{2+}$ and $[\text{Pt}_3(\mu\text{-OH})_3\{\text{P}(\text{CH}_2\text{OH})_3\}_6]^{3+}$. The mononuclear product $[\text{Pt}(\text{OH}_2)_2\{\text{P}(\text{CH}_2\text{OH})_3\}_2]^{2+}$ reacted further with *N*-donor ligands to afford complexes of the form $[\text{PtL}_2\{\text{P}(\text{CH}_2\text{OH})_3\}_2]^{2+}$ ($\text{L}_2 = 2\text{py}, \text{bpy}, \text{phen}$) [30].

A series of cationic platinum(II) complexes $[\text{PtCl}(\text{dppe})\{\text{OC}(\text{R})\text{CH}(\text{AsPh}_3)\}]\text{BF}_4$ ($\text{R} = \text{Me}, \text{Ph}, \text{OMe}$) (27) in which the carbonyl stabilised arsonium ylides are coordinated to the metal centre have been described [31]. Reaction of *cis*- $[\text{PtCl}_2(\text{PPh}_3)_2]$ with dialkyl malonate in the presence of Ag_2O gave new platinalactone complexes (28), *via* a cyclisation reaction [32].




(27)




(28; R=Me, Et)

The crystal structures of the complexes $[\text{Pt}(\text{Et})(\beta\text{-dik})(\text{PPh}_3)]$ ($\beta\text{-dik} = \text{acac}$, monothioacetylacetonate) have been determined, and compared to those for $[\text{Pd}(\text{Me})(\beta\text{-dik})(\text{PPh}_3)_2]$ [33]. In a series of related papers, the preparation and characterisation of *bis*(phosphine)platinum(II) complexes containing group 16 ligands has appeared [34-36]. Thus, for example, $[\text{Pt}(\text{EPh})_2(\text{dppe})]$ ($\text{E} = \text{O}, \text{S}$) have been prepared by reaction of the parent chloro-complex with NaOPh or $\text{PhSH}/\text{Bu}_2\text{NH}$, respectively, while $[\text{Pt}(\text{SePh})_2(\text{dppe})]$ was formed by oxidative addition of Ph_2Se_2 to $[\text{Pt}(\text{C}_2\text{H}_4)(\text{dppe})]$ [34]. The ability of complexes of this type to act as didentate 'ligands' towards other metal species was demonstrated [35, 36]. For example, *cis*- $[\text{Pt}(\text{SC}_6\text{F}_5)_2\text{L}_2]$ ($\text{L}_2 = \text{dppm}, \text{dppe}, 2\text{PPh}_3$) when treated with *cis*- $[\text{M}(\text{C}_6\text{F}_5)_2(\text{thf})_2]$ ($\text{M} = \text{Pd}, \text{Pt}$) yielded the geminal homo- or heterodinuclear compounds (29) [36].




(29; M=Pd, Pt)

The complex $[\text{Pt}(\text{H}_4\text{dmsucc-}S,S')(\text{L-L})]$ ($\text{H}_4\text{dmsucc} = \text{dimercaptosuccinic acid}$; $\text{L-L} = \text{dppe}$, 1-diphenylarsino-2-diphenylphosphino ethane) (30) have been prepared and characterised [37]. The structure of $[\text{Pt}(\text{Et}_2\text{dtc})(\text{P}_2\text{P}')]\text{BF}_4$ ($\text{Et}_2\text{dtc} = \text{S}_2\text{CNEt}_2$; $\text{P}_2\text{P}' = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$) (31) has been established using X-ray diffraction techniques. The short non-bonding distance between platinum and sulfur (2.753(3) Å) indicated the possibility of some interaction between the two atoms in the solid state [38].




The platinum(0) complex $[\text{Pt}(\text{PEt}_3)_3]$ inserts (reversibly) into the C-S bond of dibenzothiophene to give the compound (32) containing a Pt-S-C linkage. The new complex was characterised by NMR spectroscopy and an X-ray crystal structure determination [39]. The solid state structure of *cis*- $[\text{PtCl}_2(\text{SEt}_2)\{\text{P}(\text{CH}_2\text{CH}_2\text{CN})_3\}]$ and *trans*- $[\text{PtCl}_2\{\text{P}(\text{CH}_2\text{CH}_2\text{CN})_3\}_2]$ have been determined crystallographically. In the latter, the $\text{P}(\text{CH}_2\text{CH}_2\text{CN})_3$ ligands were found to form an 'inverted' umbrella relative to platinum [40]. Reaction of *cis*- $[\text{PtCl}_2(\text{PPh}_3)_2]$ with (2,4,6- $\text{tBu}_3\text{C}_6\text{H}_2$) $\text{TeLi}\cdot 3\text{thf}$ or (2,4,6- $\text{iPr}_3\text{C}_6\text{H}_2$) $\text{TeLi}\cdot 3\text{thf}$ gave the corresponding, air stable, *cis*-platinum(II) tellurolates; for example (33) whose structure was established using X-ray diffraction [41].



A convenient one step method for the synthesis of dinuclear platinum(II) phosphine complexes has been reported. The procedure involves direct reaction of PtCl_2 and the appropriate phosphine in *p*-chlorotoluene [42]. Addition of 3 equivalents of $\text{P}(\text{CH}_2\text{OH})_3$ to $\text{Na}[\text{PtCl}_6]$ or 2 equivalents to $\text{K}_2[\text{PtCl}_4]$ results in the formation of the platinum(II) complex $[\text{PtCl}_2\{\text{P}(\text{CH}_2\text{OH})_3\}_2]$. Addition of a further equivalent of $\text{P}(\text{CH}_2\text{OH})_3$ to this complex gives $[\text{PtCl}\{\text{P}(\text{CH}_2\text{OH})_3\}_3]\text{Cl}$, while reaction with 2 equivalents gives an unstable species tentatively assigned as $[\text{PtCl}\{\text{P}(\text{CH}_2\text{OH})_3\}_4]\text{Cl}$. In the presence of further $\text{P}(\text{CH}_2\text{OH})_3$, $[\text{PtCl}\{\text{P}(\text{CH}_2\text{OH})_3\}_4]\text{Cl}$ gives, amongst other things, the bis(chelate) salts *trans*- and *cis*- $[\text{Pt}\{(\text{HOCH}_2)_2\text{PCH}_2\text{OP}(\text{CH}_2\text{OH})_2\}_2]\text{Cl}_2$ (34) [43]. The related phosphine ligand $\text{Ph}_2\text{PCH}_2\text{OH}$ reacted with $[\text{PtCl}_2(\text{cod})]$ to give *cis*- $[\text{PtCl}_2(\text{Ph}_2\text{PCH}_2\text{OH})_2]$, the structure of which has been determined by X-ray crystallography [44].


The new *meso*-ligand (R^*,S^*)-2,3-bis(diphenylphosphino)butane (achiraphos) has been synthesised and used to prepare the prochiral complex $[\text{PtCl}_2(\text{achiraphos})]$. Displacement of one of the enantiotopic chlorine atoms with $[1R-(1\alpha,2\beta,5\alpha)]$ -diphenylmenthylphosphine (R -PMenPh₂)

and subsequent treatment with NH_4PF_6 gave diastereomeric complexes of the form $[\text{PtCl}(\text{achiraphos})(R\text{-PMenPh}_2)]\text{PF}_6$ (35). The solid state structure of one of the epimers has been determined by X-ray crystallography [45].




(35)

Preparation of the diphenyl(phenylethynyl)phosphine complexes (36) has been reported. For (36) intramolecular coupling of the phosphinoacetylene ligands occurs on heating to form (37), and from (37; $\text{X} = \text{Cl}$) the free unsymmetrical diphosphine $[o\text{-C}_{16}\text{H}_{10}(\text{PPh}_2)_2]$, generated by this coupling step, has been isolated [46].




(36)




(37)

The series of complexes $[\text{PtXX}'\{\eta^2\text{-(PPh}_2)_3\text{CH}\}]$ ($\text{X} = \text{X}' = \text{Cl}$, C_6F_5 ; $\text{X} = \text{Cl}$, $\text{X}' = \text{C}_6\text{F}_5$) have been prepared, and $31\text{P}\{1\text{H}\}$ NMR spectra indicate that one of the phosphorus atoms of the triphosphine ligand remains uncoordinated [47]. The new diphosphine $\text{PPh}_2\text{CH}_2\text{C}(\text{tBu})=\text{N}-\text{N}(\text{Ph})\text{PPh}_2$ has been synthesised, and treatment with $[\text{PtMe}_2(\text{cod})]$ affords the expected dimethylplatinum(II) species (38) [48].



(38)




(39)

Treatment of $\text{cis-}[\text{PtCl}_2(\text{PR}_3)_2]$ ($\text{PR}_3 = \text{PMe}_3, \text{PMe}_2\text{Ph}, \text{PMePh}_2, \text{P}(\text{nBu})_3, \text{PPh}_3, \text{PEt}_3$) with dppa ($\text{Ph}_2\text{PN}(\text{H})\text{PPh}_2$) afforded the cationic complexes $[\text{PtCl}(\text{PR}_3)(\text{dppa})]\text{Cl}$ (39), whilst


reaction of $[\text{PtCl}_2(\text{dppe})]$ with dppa gave $[\text{Pt}(\text{dppa})(\text{dppe})]\text{Cl}_2$. The molecular structures of $[\text{PtCl}(\text{PMe}_2\text{Ph})(\text{dppa})]\text{Cl}$, $[\text{PtCl}\{\text{P}(\text{nBu})_3\}(\text{dppa})]\text{Cl}$ and $[\text{Pt}(\text{dppa})(\text{dppe})]\text{Cl}_2$ have been determined by X-ray crystallography and in each instance, reveal dppa acting as a chelate through both phosphorus atoms [49]. The reaction of $[\text{PtCl}_2(\text{cod})]$ with $\text{RN}[\text{P}(\text{OPh})_2]_2$ ($\text{R} = \text{Me}, \text{Ph}$) affords similar P,P' -chelate complexes $[\text{PtCl}_2\{\text{RN}[\text{P}(\text{OPh})_2]_2\}]$ [50].

12.2.2 Complexes with nitrogen donor ligands

Reaction of organomercury halides $\text{HgR}'\text{Cl}$ ($\text{R}' = \text{Me}, \text{Et}, \text{iBu}, \text{Ph}$) with the platinum(0) complexes $[\text{Pt}(\text{dmphen})\{(Z)\text{-RO}_2\text{CCH=CHCO}_2\text{R}\}]$ ($\text{dmphen} = 2,9\text{-Me}_2\text{-1,10-phenanthroline}$; $\text{R} = \text{Me}, \text{Et}, \text{iBu}$) produced 5 coordinate platinum(II) compounds of the form $[\text{PtCl}(\text{HgR}')(\text{dmphen})\{(Z)\text{-RO}_2\text{CCH=CHCO}_2\text{R}\}]$ (40); the solid state structure of (40; $\text{R} = \text{Me}, \text{R}' = \text{Me}$) was established using X-ray diffraction methods [51]. A second series of 5-coordinate platinum(II) complexes of general formula $[\text{PtR}(\text{Cl})(\text{N-N})(\text{YC=CY})]$ ($\text{N-N} = \text{didentate N-donor ligand}$; $\text{Y} = \text{CF}_3, \text{CO}_2\text{Me}, \text{CH}_2\text{Cl}$; $\text{R} = \text{Me}, 4\text{-MeOC}_6\text{H}_4$), e.g. (41), have been prepared by treatment of $[\text{PtR}(\text{Cl})(\text{N-N})]$ with alkynes. Using a 1:2 alkyne: $[\text{PtMe}(\text{Cl})(\text{dmphen})]$ ratio dinuclear species $[\text{Pt}_2\text{Me}_2(\text{Cl})_2(\text{YC=CY})(\text{dmphen})]$ were isolated, and for the but-2-yne derivative the molecular structure determined crystallographically [52].




(40)




(41)

Cyclometallation reactions of N -benzyl and N -(phenylethyl)- α -benzoylbenzylideneamines with platinum(II) salts have been reported, and the structure of $[\text{Pt}\{\text{C}_6\text{H}_4(\text{C}_6\text{H}_5\text{C=O})\text{C=N-CH}_2\text{C}_6\text{H}_5\}(\mu\text{-OAc})_2]$ (42) established [53]. Reaction of $[\text{Pt}(\text{MeCN})_4]^{2+}$ with dpphen ($\text{dpphen} = 2,9\text{-diphenyl-1,10-phenanthroline}$) gave the cyclometallated complex (43). The acetonitrile ligand in (43) is substitution labile, and can be replaced by nitrogen bases [54]. The synthesis of the cyclometallated derivative $[\text{PtLCl}]$ ($\text{HL} = 6\text{-iBu-2,2'-bipyridine}$) has been reported [55].



(42)




(43)

The mixed alkylplatinum(II) complex $[\text{Pt}(\text{CF}_3)(\text{CH}_3)(\text{nbd})]$ reacts with neutral ligands L_2 ($\text{L}_2 = \text{e.g. tmen, 2py, bpy}$) to afford the corresponding complexes $[\text{Pt}(\text{CF}_3)(\text{CH}_3)(\text{L}_2)]$, whilst with MeCN , $[\text{Pt}(\text{CF}_3)(\text{CH}_3)(\text{MeCN})_2]$ is formed in equilibrium with starting materials [56]. The kinetics of CN^- substitution at platinum(II) in the series *cis*- and *trans*- $[\text{PtCl}_2(\text{am})_2]$ ($\text{am} = \text{e.g. Me}_2\text{NH, py, 4-CNpy, 4-Clpy, 2-Mepy}$) has been investigated for the slow step in equation (i) [57].




Treatment of *cis*- $[\text{Pt}(\text{C}_6\text{F}_5)_2(\text{thf})_2]$ with 2-benzyl pyridine gave *cis*- $[\text{Pt}(\text{C}_6\text{F}_5)_2(\text{PhCH}_2\text{py})]$ (44), which in the solid state, as established by X-ray diffraction, exhibits a weak η^2 -benzyl platinum interaction. Subsequent reaction of (44) with CO yielded $[\text{Pt}(\text{C}_6\text{F}_5)_2(\text{PhCH}_2\text{py-}N)(\text{CO})]$ which did not contain a Pt-benzyl interaction [58].




(44)

Three platinum(II) complexes of 1,1'-biisoquinoline (biq) have been prepared and their molecular structures determined using X-ray crystallography. In one of these, $[\text{PtCl}_2(\text{biq})]$ (45) the ligand was found to be didentate, whilst in the complexes $[\text{Pt}(\text{biq})(\text{L})_3]$ ($\text{L} = \text{py, 4-NMe}_2\text{py}$) 1,1'-biisoquinoline adopted a unidentate bonding mode [59]. Mono- and bis-nitrile complexes $[\text{PtCl}_3(\text{NCR})]^-$ and *cis*- and *trans*- $[\text{PtCl}_2(\text{NCR})_2]$ ($\text{R} = \text{tBu, Me, Ph}$) hydrolyse to amidate-amide and mixed amide-nitrile complexes, with the amide ligands in the enol form and *N*-bound to platinum. The complexes are composed of two platinum(II) monomeric units, with a Pt-Pt intermetallic interaction. For example, *cis*- $[\text{PtCl}_2\{\text{HN}=\text{C}(\text{OH})\text{tBu}\}_2]$ (46) possesses a Pt-Pt distance of 3.165(1) Å [60].




(45)



(46)

Reaction of the complex $[\text{PtCl}(\eta^2\text{-C}_2\text{H}_4)(\text{tmen})]^+$ with acridines {3,6-diaminoacridine and 3,6-bis-(dimethylamino)acridine} afforded complexes in which the metal is linked to the polycyclic molecule by an ethylene chain. For 3,6-bis-(dimethylamino)acridine attack occurred at the endocyclic nitrogen to give (47) whilst for 3,6-diaminoacridine attack occurred at the exocyclic aminic groups to afford (48); the structure of the former was established using X-ray diffraction [61].




Treatment of *cis*- and *trans*- $[\text{PtCl}_2(\text{NCR})_2]$ ($\text{R} = 4\text{-MePh}, 4\text{-CF}_3\text{Ph}, 2\text{-MePh}, \text{Et}, n\text{Pr}, i\text{Pr}, t\text{Bu}$) with 2 equivalents of $[\text{OCH}_2\text{CH}_2\text{Cl}]^-$ affords the *bis*(Δ^2 -1,3-oxazoline) derivatives *cis*- and *trans*- $[\text{PtCl}_2[\text{N}=\text{C}(\text{R})\text{OCH}_2\text{CH}_2]_2]$ in high yield [62]. A series of platinum-terpy complexes $[\text{Pt}(\text{terpy})\text{L}]^{n+}$ ($\text{L} = \text{Cl}, \text{Br}, \text{I}, \text{N}_3, \text{SCN}$ ($n = 1$); $\text{L} = \text{NH}_3$ ($n = 2$)) have been prepared and their spectroscopic and emission properties studied [63]. The ^1H NMR and vibrational spectra for a range of complexes of the general formula $[\text{PtCl}_2(\text{am})]$ ($\text{am} = \text{cycloalkylamine } \text{C}_n\text{H}_{2n-1}\text{NH}_2$; $n = 3 \rightarrow 8$) have been reported [64].

12.2.3 Complexes with sulfur group donor ligands


The series of complexes *trans*- $[\text{PtCl}_2(\text{C}_2\text{H}_4)\text{L}]$ ($\text{L} = \text{quinoline } N\text{-oxide (QNO)}, \text{isoquinoline-}N\text{-oxide}, 4\text{-ClQNO}, 2\text{-MeQNO}, 4\text{-MeQNO}$) have been prepared by addition of L to Zeise's salt. Subsequent reaction of $[\text{PtCl}_2(\text{C}_2\text{H}_4)(4\text{-MeQNO})]$ with L' ($\text{L}' = \text{CO}, \text{PPh}_3, \text{py}$) afforded *trans*- $[\text{PtCl}_2(\text{L}')(4\text{-MeQNO})]$ [65].

An account detailing the synthesis of a range of complexes containing the potentially tridentate ligand 1,4,7-trithiacyclononane, $[\text{9}] \text{aneS}_3$ (49), has appeared. The complexes $[\text{PtRR}'\{[\text{9}] \text{aneS}_3\}]$ ($\text{R} = \text{R}' = \text{Me}, \text{Et}, \text{CH}_2\text{CMe}_3, \text{CH}_2\text{SiMe}_3, \text{Ph}$; $\text{R} = \text{Me}, \text{R}' = \text{CH}_2\text{SiMe}_3$; $\text{R} = \text{Cl}, \text{R}' = \text{CH}_2\text{SiMe}_3$) were prepared by displacement of *cod* or *nbd* from appropriate precursors, and for $[\text{PtPh}_2\{[\text{9}] \text{aneS}_3\}]$ the molecular structure was established by X-ray crystallography [66]. In two

related reports, the complexes $[\text{Pt}\{(\mathbf{50})\}_2]^{2+}$ [67] and $[\text{Pt}\{(\mathbf{51})\}]^{2+}$ [68] were prepared and structurally characterised using X-ray diffraction methods.




The synthesis, characterisation, X-ray structure determination and emission spectroscopy of $(\mathbf{52})$ and $(\mathbf{53})$ has been reported. Both complexes were obtained upon reaction of $[\text{PtCl}_2(\text{cod})]$ and the corresponding dianionic dithiolate salt [69]. The platinum(II) complexes $[\text{Pt}(\text{S}_2\text{C})]_2$, $[\text{Pt}(\text{S}_2\text{C})(\text{py})]$ and $[\text{Pt}(\text{S}_2\text{C})(\text{PMe}_3)]$ ($\mathbf{54}$) ($\text{S}_2\text{C} = 1,3\text{-imidazolidinyl-}N,N'\text{-bis}(2\text{-benzenethiolate})$) have been prepared and fully characterised [70].





12.2.4 Complexes with mixed donor ligands


The new P,N -didentate ligand ($\mathbf{55}$), derived from valine, has been synthesised and used to prepare the complexes $[\text{PtX}_2(\mathbf{55})]$ ($\text{X} = \text{Cl}, \text{I}$) [71]. Reaction of $\text{trans-}[\text{PtCl}_2\{\text{PPh}_2\text{NC}(4\text{-Ar})\text{N}(\text{SiMe}_3)_2\}(\text{PEt}_3)]$ ($\text{Ar} = 4\text{-MeC}_6\text{H}_4$) with PhSeCl was reported to give ($\mathbf{56}$), containing the N,P -chelating ligand $\text{PPh}_2\text{N}(\text{H})\text{C}(\text{Ar})=\text{N}(\text{H})$. The structure of ($\mathbf{56}$) was established using NMR spectroscopy and X-ray crystallography [72]. The dinuclear platinum compounds ($\mathbf{57}$) and ($\mathbf{58}$) have been isolated from the reaction of $\text{cis-}[\text{PtCl}_2(\text{dmso})(\text{PPh}_2\text{py})]$ with an equimolar quantity of $\text{cis-}[\text{PtMe}_2(\text{dmso})_2]$; both complexes were the subject of an X-ray diffraction study [27].



(55)

(56; Ar=4-MeC₆H₄)


(57; S=dmsO)




(58)

The ligands $\text{Ph}_2\text{PN}(\text{Ph})\text{PPh}_2(\text{E})$ ($\text{E} = \text{S}, \text{Se}$) (P-E) have been used to prepare the mixed donor complexes $[\text{PtCl}_2(\text{P-E})]$ (59), and for $\text{E} = \text{S}$ the solid state structure has been established [73]. The neutral, dimeric complexes $[\{\text{Pt}(\text{C}_6\text{F}_5)(\mu\text{-SPPH}_2)(\text{PR}_3)\}_2]$ ($\text{PR}_3 = \text{PPh}_3, \text{PPh}_2\text{Et}$) and cationic complex $[\text{Pt}(\text{SPPH}_2)(\text{PPh}_3)_2]\text{ClO}_4$ have been reported. For the former the $[\text{SPPH}_2]^-$ ligand is present as a P,S -bridge whilst in the latter it adopts a P,S -chelating bonding mode [74].

Treatment of the chloro-bridged complexes $[\text{Pt}_2\text{Cl}_4(\text{PEt}_3)_2]$ and $[\text{Pt}_2\text{Cl}_2(\text{MeOcod})_2]$ with the anion $[\text{C}(\text{PPh}_2)\{\text{P}(\text{S})\text{Ph}_2\}_2]^-$ affords the complexes $[\text{PtCl}(\text{PEt}_3)\{\text{C}(\text{PPh}_2)\{\text{P}(\text{S})\text{Ph}_2\}_2-P,S\}]$ (60) and $[\text{Pt}(\text{MeOcod})\{\text{C}(\text{PPh}_2)\{\text{P}(\text{S})\text{Ph}_2\}_2-P,S\}]$, respectively, as a mixture of *cis*- and *trans*-isomers [75]. The synthesis of $[\text{PtCl}_2(\text{E-S})]$ [$\text{E-S} = (\pm)\text{Ph}_2\text{ECH}_2\text{CH}_2\text{S}(\text{O})\text{Me}$ ($\text{E} = \text{P}, \text{As}$)] complexes has been reported. In both cases, the ligand acts as an E,S -didentate ligand, and M-O bonds are not involved [76].




(59)




(60)

Reaction of $[\text{Pt}(\text{C}_6\text{F}_5)_2(\mu\text{-Cl})_2]^{2-}$ with $[\text{C}_5\text{H}_4\text{NS}]^-$ (pyridine-2-thiolate) affords the monomeric complex $[\text{Pt}(\text{C}_6\text{F}_5)_2(\text{C}_5\text{H}_4\text{NS})]^-$, possessing a $[\text{N-S}]^-$ chelate. Upon reaction of this mono-anionic complex with PPh_3 the species $[\text{Pt}(\text{C}_6\text{F}_5)_2(\text{C}_5\text{H}_4\text{NS})(\text{PPh}_3)]^-$, possessing the ligand


in a monodentate *S*-binding mode is obtained [77]. The crystal structures of four platinum(II) complex cations (61) containing the organic ligand 2,6-bis(methylthiomethyl)pyridine have been determined by X-ray diffraction [78]. Reaction of *trans*-[PhSNC(MeC₆H₄)N=NC(MeC₆H₄)-NSPh] with [Pt(C₂H₄)(PPh₃)₂] gave the platinum(II) complex (62) in which the ligand is bonded in a *N,N',S*-tridentate fashion [79]




(61; X=Cl, Br, SCN)

(62; Ar=4-MeC₆H₄)

The thioether and sulphinyl azo-benzene complexes (63) and (64), respectively, slowly insert oxygen into the Pt-O bond when treated with *m*-chloroperbenzoic acid to produce species containing the *O,N,S*-donor set [80].




(63)



(64)


Treatment of platinum salts with L²⁻ (65) affords the complex [Pt₂(L)]²⁺. For the analogous palladium(II) salt, [Pd₂(L)]²⁺, the solid state structure showed the dinucleating macrocycle to be planar with inter-molecular π -ring stacking [81].




(65)

12.3 PLATINUM(I)

The first example of a μ -parallel ligated phospho-alkyne has been reported for the complex $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\mu\text{-}^t\text{BuCP})]$ (66). The structure of the molecule was elucidated by ^{31}P and ^{195}Pt NMR spectroscopy [82].

(66; $R = ^t\text{Bu}$)

Reaction of $[\text{Pt}_2\text{Cl}_2(\text{dppm})_2]$ with 2 L ($L = \text{quinoline}, N\text{-methylimidazole}, 4\text{-}^t\text{butylpyridine}$) afforded the new compounds $[\text{Pt}_2(\text{dppm})_2\text{L}_2]^{2+}$ (67), whilst reaction with $L' = 2,4,6\text{-Me}_3\text{pyridine}$ gave $[\text{Pt}_2(\text{dppm})_2(L')\text{Cl}]^+$. For (67; $L = \text{quinoline}, N\text{-methylimidazole}$) and $[\text{Pt}_2(\text{dppm})_2(L')\text{Cl}]^+$ the solid state structure was determined crystallographically, and the Pt-Pt distances found to be 2.615(1), 2.580(1) and 2.627(2) Å, respectively, thus establishing the presence of a Pt-Pt bond [83].


(67²⁺)

12.4 PLATINUM(0)


Reaction of 1-bromocycloheptene or a mixture of 1-, 2- and 3-bromocycloheptatrienes with $^t\text{BuOK}$ in the presence of $[\text{Pt}(\text{PPh}_3)_3]$ led to the predominant formation of $\text{Pt}(\text{PPh}_3)_2$ π -complexes of the corresponding cyclic allenes, e.g. (68). In contrast, reaction of the same halides with LDA in thf in the presence of $[\text{Pt}(\text{PPh}_3)_3]$ gave exclusively $\text{Pt}(\text{PPh}_3)_2$ complexes of the corresponding cyclic alkynes, e.g. (69) [84].

The preparation of $[\text{Pt}\{\text{trans-CH}(\text{CO}_2\text{Et})=\text{CH}(\text{CO}_2\text{Et})\}(\text{PPh}_3)_2]$ and $[\text{Pt}\{\text{cis-CH}(\text{CO}_2\text{Et})=\text{CH}(\text{CO}_2\text{Et})\}(\text{PPh}_3)_2]$ has been described. The complexes were prepared by reaction of $[\text{Pt}(\eta^2\text{-CH}_2\text{CH}_2)(\text{PPh}_3)_2]$ with *trans*- and *cis*- $\text{CH}(\text{CO}_2\text{Et})=\text{CH}(\text{CO}_2\text{Et})$, respectively, and for $[\text{Pt}\{\text{trans-CH}(\text{CO}_2\text{Et})=\text{CH}(\text{CO}_2\text{Et})\}(\text{PPh}_3)_2]$ the structure was determined from an X-ray analysis [85]. The platinum(0) dimer $[\text{Pt}_2(\mu\text{-CO})(\text{CO})_2(\mu\text{-dppm})_2]$ (70) has been prepared from $[\text{Pt}_2\text{Cl}_2(\mu\text{-}$


dppm)₂] via NaBH₄ reduction under a CO atmosphere and by substitution of a dppm ligand in [Pt₂(μ-dppm)₃] with CO [86].



(68)



(69)



(70)

REFERENCES

1. K.-T. Aye, J.J. Vittal and R.J. Puddephatt, *J. Chem. Soc., Dalton Trans.*, (1993) 1835.
2. W. Clegg, K.A. Fraser, N. Duran and P. Gonzalez-Duarte, *Polyhedron*, 12 (1993) 2737.
3. J. Dauth, B. Deubzer and J. Weis, *J. Organomet. Chem.*, 459 (1993) 359.
4. F.F. Stewart, W.D. Neilsen, R.E. Ekeland, R.D. Larsen and P.W. Jennings, *Organometallics*, 12 (1993) 4585.
5. A.J. Canty, P.R. Traill, R. Colton and I.M. Thomas, *Inorg. Chim. Acta*, 210 (1993) 91.
6. A.N. Buckley, J.A. Busby, I.J. Ellison and R.D. Gillard, *Polyhedron*, 12 (1993) 247.
7. J.R. Berenguer, J. Fornies, F. Martinez, J.C. Cubero, E. Lalinde, M.T. Moreno and A.J. Welch, *Polyhedron*, 12 (1993) 1797.
8. V. W.-W. Yam, L.-P. Chan and T.-F. Lai, *Organometallics*, 12 (1993) 2197.
9. J.M. Casas, L.R. Falvello, J. Fornies, A. Martin and M. Tomas, *J. Chem. Soc., Dalton Trans.*, (1993) 1107.
10. K.A. Fallis, C. Xu and G.K. Anderson, *Organometallics*, 12 (1993) 2243.
11. L. Pandolfo, G. Paiaro, P. Ganis, G. Valle and P. Traldi, *Inorg. Chim. Acta*, 210 (1993) 39.
12. R.J. Hinkle, P.J. Stang and A.M. Arif, *Organometallics*, 12 (1993) 3510.
13. J.M.A. Wouters, R.A. Klein, C.J. Elsevier, M.C. Zoutberg and C.H. Stam, *Organometallics*, 12 (1993) 3864.
14. F.A. Ajulu, P.B. Hitchcock, F. Mathey, R.A. Michelin, J.F. Nixon and A.J.L. Pombeiro, *J. Chem. Soc., Chem. Commun.*, (1993) 142.
15. P. Bergamini, E. Costa, S. Sostero, C. Ganter, J. Hogg, A.G. Orpen and P.G. Pringle, *J. Organomet. Chem.*, 455 (1993) C13.
16. R. McCrindle and A.J. McAlees, *Organometallics*, 12 (1993) 2445.
17. H. Yamashita, M. Tanaka and M. Goto, *Organometallics*, 12 (1993) 988.
18. S.D. Grumbine and T.D. Tilley, *J. Am. Chem. Soc.*, 115 (1993) 7884.
19. S. Fallis, L. Rodriguez, G.K. Anderson and N.P. Rath, *Organometallics*, 12 (1993) 3851.
20. V.V. Grushin, C. Bensimon and H. Alper, *Organometallics*, 12 (1993) 2737.
21. H.-A. Brune, J. Lerche, G. Schmidtberg and A. Baur, *J. Organomet. Chem.*, 450 (1993) 269.
22. M.A. Bennett, H. Jin and A.C. Willis, *J. Organomet. Chem.*, 451 (1993) 249.
23. A. Ohsuka, T. Fujimori, T. Hirao, H. Kurosawa and I. Ikeda, *J. Chem. Soc., Chem. Commun.*, (1993) 1039.
24. T.-M. Huang, J.-T. Chen, G.-H. Lee and Y. Wang, *J. Am. Chem. Soc.*, 115 (1993) 1170.
25. V.K. Jain, S. Kannan and E.R.T. Tiekink, *J. Chem., Soc., Dalton Trans.*, (1993) 3625.
26. A. Singhal, V.K. Jain and S. Kannan, *J. Organomet. Chem.*, 447 (1993) 317.
27. C.G. Arena, G. Bruno, G. De Munno, E. Rotondo, D. Drommi and F. Faraone, *Inorg. Chem.*, 32 (1993) 1601.
28. A. Adedapo, A.G. Avent, D. Carmichael, P.A. Chaloner, P.B. Hitchcock and A. Wagenaar, *J. Chem. Soc., Chem. Commun.*, (1993) 186.
29. T. Chivers, D.D. Doxsee and R.W. Hilt, *Inorg. Chem.*, 32 (1993) 3244.
30. E. Costa, M. Murray, P.G. Pringle and M.B. Smith, *Inorg. Chim. Acta*, 213 (1993) 25.
31. G. Facchin, L. Zanutto, R. Bertani, L. Canovese and P. Uguagliati, *J. Chem. Soc., Dalton Trans.*, (1993) 2871.
32. W. Henderson, R.D.W. Kemmitt and A.L. Davis, *J. Chem. Soc., Dalton Trans.*, (1993) 2247.

33. K.J. Cavell, H. Jin, B.W. Skelton and A.H. White, *J. Chem. Soc., Dalton Trans.*, (1993) 1973.
34. C. Xu, J.W. Siria and G.K. Anderson, *Inorg. Chim. Acta*, 206 (1993) 123.
35. V.K. Jain, S. Kannan, R.J. Butcher and J.P. Jasinski, *J. Chem. Soc., Dalton Trans.*, (1993) 1509.
36. R. Usón, J. Forniés, M.A. Usón and S. Herrero, *J. Organomet. Chem.*, 447 (1993) 137.
37. P.S. Jarrett, O.M. Ni Dhubhghaill and P.J. Sadler, *J. Chem. Soc., Dalton Trans.*, (1993) 1863.
38. R. Colton, M.F. Mackay and V. Tedesco, *Inorg. Chim. Acta*, 207 (1993) 227.
39. J.J. Garcia and P.M. Maitlis, *J. Am. Chem. Soc.*, 115 (1993) 12200.
40. M.N.I. Khan, C. King, J.P. Fackler and R.E.P. Winpenny, *Inorg. Chem.*, 32 (1993) 2502.
41. P.J. Bonasia and J. Arnold, *J. Organomet. Chem.*, 449 (1993) 147.
42. W. Baratta and P.S. Pregosin, *Inorg. Chim. Acta*, 209 (1993) 85.
43. P.A.T. Hoye, P.G. Pringle, M.B. Smith and K. Worboys, *J. Chem. Soc., Dalton Trans.*, (1993) 269.
44. D.L. Davies, J. Neild, L.J.S. Prouse and D.R. Russell, *Polyhedron*, 12 (1993) 2121.
45. P. Gugger, S.O. Limmer, A.A. Watson, A.C. Willis and S.B. Wild, *Inorg. Chem.*, 32 (1993) 5692.
46. D.K. Johnson, T. Rukachaisirikul, Y. Sun, N.J. Taylor, A.J. Canty and A.J. Carty, *Inorg. Chem.*, 32 (1993) 5544.
47. J. Forniés, R. Navarro and E.P. Urriolabeitia, *J. Organomet. Chem.*, 452 (1993) 241.
48. N. Iranpoor and B.L. Shaw, *Inorg. Chim. Acta*, 204 (1993) 39.
49. P. Bhattacharyya, R.N. Sheppard, A.M.Z. Slawin, D.J. Williams and J.D. Woollins, *J. Chem. Soc., Dalton Trans.*, (1993) 2393.
50. M.S. Balakrishna, S.S. Krishnamurthy, R. Murugavel, M. Nethaji and I.I. Mathews, *J. Chem. Soc., Dalton Trans.*, (1993) 477.
51. M.E. Cucciolito, F. Giordano, A. Panunzi, F. Ruffo and V. De Felice, *J. Chem. Soc., Dalton Trans.*, (1993) 3421.
52. V. De Felice, A. De Renzi, F. Giordano and D. Tesauero, *J. Chem. Soc., Dalton Trans.*, (1993) 1927.
53. C. Navarro-Ranninger, I. López-Solera, A. Alvarez-Valdés, J.H. Rodríguez-Ramos, J.R. Masaguer and J.L. García-Ruano, *Organometallics*, 12 (1993) 4104.
54. C.-W. Chan, T.-F. Lai, C.-M. Che and S.-M. Peng, *J. Am. Chem. Soc.*, 115 (1993) 11245.
55. S. Stoccoro, G. Chelucci, M.A. Cinellu, A. Zucca and G. Minghetti, *J. Organomet. Chem.*, 450 (1993) C15.
56. T.G. Appleton, J.R. Hall, M.T. Mathieson and D.W. Neale, *J. Organomet. Chem.*, 453 (1993) 307.
57. L. Cattalini, F. Guidi and M.L. Tobe, *J. Chem. Soc., Dalton Trans.*, (1993) 233.
58. J.M. Casas, J. Forniés, A. Martín and B. Menjón, *Organometallics*, 12 (1993) 4376.
59. L.-K. Cheng, K.-S. Yeung and C.-M. Che, *Polyhedron*, 12 (1993) 1201.
60. R. Cini, F.P. Fanizzi, F.P. Intini, L. Maresca and G. Natile, *J. Am. Chem. Soc.*, 115 (1993) 5123.
61. E. Ceci, R. Cini, A. Karaulov, M.B. Hursthouse, L. Maresca and G. Natile, *J. Chem. Soc., Dalton Trans.*, (1993) 2491.
62. R.A. Michelin, R. Bertani, M. Mozzon, G. Bombieri, F. Benetollo and R.B. Angelici, *J. Chem. Soc., Dalton Trans.*, (1993) 959.
63. H.-K. Yip, L.-K. Cheng, K.-K. Cheung and C.-M. Che, *J. Chem. Soc., Dalton Trans.*, (1993) 2933.
64. J. Kritzenberger, F. Zimmermann and A. Wokaun, *Inorg. Chim. Acta*, 210 (1993) 47.
65. M.-A.D. Stiakaki and A. Christofides, *Polyhedron*, 12 (1993) 661.
66. M.A. Bennett, A.J. Canty, J.K. Felixberger, L.M. Rendina, C. Sunderland and A.C. Willis, *Inorg. Chem.*, 32 (1993) 1951.
67. A.J. Blake, R.D. Crofts and M. Schröder, *J. Chem. Soc., Dalton Trans.*, (1993) 2259.
68. M.A. Watzky, D. Waknine, M.J. Heeg, J.F. Endicott and L.A. Ochrymowycz, *Inorg. Chem.*, 32 (1993) 4882.
69. J.M. Bevilacqua, J.A. Zuleta and R. Eisenberg, *Inorg. Chem.*, 32 (1993) 3689.
70. D. Sellmann, W. Prechtel, F. Knoch and M. Moll, *Inorg. Chem.*, 32 (1993) 538.
71. A. Albinati, F. Lianza, H. Berger, P.S. Pregosin, H. Rüegger and R.W. Kunz, *Inorg. Chem.*, 32 (1993) 478.
72. T. Chivers, K. McGregor and M. Parvez, *Inorg. Chem.*, 32 (1993) 5119.
73. M.S. Balakrishna, R. Klein, S. Uhlenbrock, A.A. Pinkerton and R.G. Cavell, *Inorg. Chem.*, 32 (1993) 5676.

74. J. Forniés, F. Martínez, R. Navarro, E.P. Urriolabeitia and A.J. Welch, *J. Chem. Soc., Dalton Trans.*, (1993) 2147.
75. J. Browning, K.R. Dixon, N.J. Meanwell and F. Wang, *J. Organomet. Chem.*, 460 (1993) 117.
76. S.Y.M. Chooi, P.-H. Leung and K.F. Mok, *Inorg. Chim. Acta*, 205 (1993) 245.
77. J. Forniés, C. Fortuño, M.A. Gómez, B. Menjón and E. Herdtweck, *Organometallics*, 12 (1993) 4368.
78. G. Marangoni, B. Pitteri, V. Bertolasi, V. Ferretti and P. Gilli, *Polyhedron*, 12 (1993) 1669.
79. T. Chivers, K. McGregor and M. Parvez, *J. Chem. Soc., Chem. Commun.*, (1993) 1021.
80. C. Sinha, *Polyhedron*, 12 (1993) 2363.
81. A.J. Atkins, A.J. Blake and M. Schröder, *J. Chem. Soc., Chem. Commun.*, (1993) 353.
82. S.I. Al-Resayes and J.F. Nixon, *Inorg. Chim. Acta*, 212 (1993) 265.
83. H.-K. Yip, C.-M. Che and S.-M. Peng, *J. Chem. Soc., Dalton Trans.*, (1993) 179.
84. Z. Lu, K.A. Abboud and W.M. Jones, *Organometallics*, 12 (1993) 1471.
85. M.F.C. Guedes da Silva, J.J.R. Fraústo da Silva, A.J.L. Pombeiro, R. Bertani, R.A. Michelin, M. Mozzon, F. Benetollo and G. Bombieri, *Inorg. Chim. Acta*, 214 (1993) 85.
86. S. Schreiner and T.N. Gallaher, *Organometallics*, 12 (1993) 4201.