

# Advances in the chemistry of bis-cyclopentadienyl hydride derivatives of niobium and tantalum

Antonio Antiñolo <sup>a</sup>, Fernando Carrillo-Hermosilla <sup>a</sup>,  
Mariano Fajardo <sup>b</sup>, Juan Fernández-Baeza <sup>a</sup>,  
Santiago García-Yuste <sup>a</sup>, Antonio Otero <sup>a,\*</sup>

<sup>a</sup> *Departamento de Química Inorgánica, Orgánica y Bioquímica, Facultad de Químicas,  
Campus Universitario, Universidad de Castilla-La Mancha, 13071 Ciudad Real, Spain*

<sup>b</sup> *Departamento de Química Inorgánica, Campus Universitario, Universidad de Alcalá,  
28871 Alcalá de Henares, Spain*

Received 28 September 1998; accepted 30 April 1999

## Contents

|                                                                                                                    |    |
|--------------------------------------------------------------------------------------------------------------------|----|
| Abstract . . . . .                                                                                                 | 44 |
| 1. Introduction . . . . .                                                                                          | 44 |
| 2. Hydrogen exchange coupling in metallocene trihydride complexes . . . . .                                        | 45 |
| 3. $\sigma$ -Bond activation by niobocene and tantalocene hydrides . . . . .                                       | 50 |
| 4. The chemistry of $d^2$ hydride niobocene and tantalocene derivatives . . . . .                                  | 55 |
| 4.1 Preparations . . . . .                                                                                         | 55 |
| 4.2 Reactivity of the $Cp_2MHL$ and related species in insertion processes . . . . .                               | 57 |
| 4.2.1 Insertion processes with heterocumulene molecules . . . . .                                                  | 57 |
| 4.2.2 Insertion processes with alkynes . . . . .                                                                   | 58 |
| 4.3 Reactivity of the $Cp_2MHL$ complexes in protonation processes: synthesis of dihydrogen<br>complexes . . . . . | 59 |
| 4.4 The hydride–olefin niobocene and tantalocene complexes . . . . .                                               | 61 |
| 4.4.1 Preparations . . . . .                                                                                       | 61 |
| 4.4.2 Reactivity studies . . . . .                                                                                 | 64 |
| 5. Heterobimetallic complexes containing cyclopentadienyl–hydride niobium and tantalum<br>moieties . . . . .       | 65 |
| Acknowledgements . . . . .                                                                                         | 68 |
| References . . . . .                                                                                               | 68 |

\* Corresponding author. Tel.: +34-926-295-326; fax: +34-926-295-318.

E-mail address: aotero@qino-cr.uclm.es (A. Otero)

## Abstract

The bis-cyclopentadienyl hydride chemistry of the elements niobium and tantalum has been dominated by the family of trihydride complexes, namely  $\text{Cp}_2\text{MH}_3$  ( $\text{M} = \text{Nb, Ta}$ ,  $\text{Cp} = \text{C}_5\text{H}_5$ ,  $\text{C}_5\text{Me}_5$ ,  $\text{C}_5\text{H}_4\text{SiMe}_3$ ,  $\text{C}_5\text{H}_3(\text{SiMe}_3)_2$  and related substituted cyclopentadienyl rings). The chemical and spectroscopic properties of these complexes are strongly influenced by both the nature of the Cp ring and the metal center. In some cases, anomalous NMR spectroscopic behavior has been observed, and values of  $J_{\text{HH}}$  that are largely dependent on the temperature have been found. This behavior can be modulated by means of the interaction of the trihydride complexes with Group 11 metal fragments, and in some cases HH interactions in the molecules can be observed. Several X-ray crystal structure determinations of this kind of complex, as well as different theoretical calculations, have been performed in order to gain a deeper insight into these systems. The reactivity of the trihydride metallocene complexes has been extensively considered. Thus, the easy elimination of  $\text{H}_2$  from the  $\text{Cp}_2\text{MH}_3$  species, giving rise to the unsaturated species  $\text{Cp}_2\text{MH}$ , has been used to study the activation processes of several kinds of H–X bond, in hydrosilanes, hydrogermananes, etc., which allows the synthesis of complexes such as  $\text{Cp}_2\text{MH}_2\text{X}$ . This type of complex has been extensively studied, especially when  $\text{X} = \text{SiR}_3$ , because in several cases these complexes exhibit a special hypervalence behavior. A large family of complexes of stoichiometry  $\text{Cp}_2\text{MHL}$  was prepared from the reaction of the  $\text{Cp}_2\text{MH}_3$  complexes with different classes of  $\pi$ -acid ligands, L. The  $\text{Cp}_2\text{MHL}$  complexes can undergo insertion reactions into the M–H bond with several classes of unsaturated molecules, and such processes are of interest in the field of organic synthesis. Furthermore, the protonation processes of the  $\text{Cp}_2\text{MHL}$  complexes give rise to a new family of  $\eta^2\text{-H}_2$ -containing cationic species,  $\text{Cp}_2\text{M}(\eta^2\text{-H}_2)\text{L}^+$ , which are stable at low temperatures. The NMR data of such compounds indicate that it is possible to block the rotation of the  $\text{H}_2$  molecule at low temperatures. Of particular interest are the  $\text{Cp}_2\text{MH}(\text{olefin})$  complexes, and these have been widely studied because they can be considered as excellent models for the study of the  $\beta$ -elimination reaction and the reverse olefin insertion process. Finally, from the trihydride derivatives  $\text{Cp}_2\text{MH}_3$  and related species, several heterometallic complexes have been prepared and characterized. © 1999 Elsevier Science S.A. All rights reserved.

**Keywords:** Niobocene; Tantalocene; Hydride; Cyclopentadienyl; Synthesis; Reactivity

## 1. Introduction

Hydride complexes of transition metals represent one of the most important classes of compounds because of their chemical reactivity and their important role as intermediates in a number of catalytic hydrogenation processes [1]. This review is concerned with both the spectroscopic and chemical behavior of bis-cyclopentadienyl hydride complexes of niobium and tantalum since they constitute a special family of hydride-containing organometallics of early transition metals [2]. Bis-cyclopentadienyl niobium or tantalum hydride derivatives have long been of interest for both their high reactivity and their spectroscopic properties. In particular, the spectroscopic properties of the niobocene and tantalocene trihydrides, namely

$\text{Cp}_2\text{MH}_3$  ( $\text{M} = \text{Nb}, \text{Ta}$ ,  $\text{Cp} = \text{C}_5\text{H}_5$ ,  $\text{C}_5\text{Me}_5$ ,  $\text{C}_5\text{H}_4(\text{SiMe}_3)$ ,  $\text{C}_5\text{H}_3(\text{SiMe}_3)_2$ , etc.), are of great interest. Thus, bis(cyclopentadienyl)niobium trihydrides exhibit quantum mechanical exchange couplings, and this phenomenon, also observed with other transition metal complexes, is characterized by the presence of large temperature-dependent H–H coupling between the hydrides. Nevertheless, the analogous tantalocene trihydrides exhibit the behavior of classical hydrides. A rich chemistry of hydride complexes has been developed from studies of the reactivity of the trihydride complexes  $\text{Cp}_2\text{MH}_3$  towards several types of reagents. Thus, an extensive family of 18-electron niobium (III) and tantalum (III) hydrides,  $\text{Cp}_2\text{MHL}$ , has been reported from the reactions of the trihydride complexes with  $\pi$ -acid ligands, namely phosphine, phosphite, olefin, isocyanide, etc. Some of these complexes are of special interest; for example, the bis(cyclopentadienyl)–hydride–olefin species are among the more interesting hydride–olefin complexes and they are especially appropriate for the study of the insertion of olefins into metal–hydride bonds and its microscopic reverse, beta-elimination, which are very common steps in industrial catalytic reactions such as olefinic isomerization, hydrogenation, and hydroformylation [3]. Furthermore, the discovery by Kubas [4] of dihydrogen coordination on a transition metal center has led to intense research activity in the chemical, theoretical, and spectroscopic aspects of the synthesis of dihydrogen complexes. As a consequence, it has been possible to easily carry out the synthesis of Group 5 dihydrogen complexes,  $\text{Cp}_2\text{M}(\eta^2\text{-H}_2)\text{L}^+$  [5], by protonation at low temperature of monohydride complexes,  $\text{Cp}_2\text{MHL}$ . These complexes also exhibit interesting spectroscopic properties. The oxidative addition of organo-Group 14 element hydrides to coordinatively unsaturated niobium or tantalum hydride complexes has also been applied to the preparation of interesting Group 14 fragments containing niobium or tantalum organometallics. The reactivity of these compounds has been widely studied. In other fields, insertion processes of unsaturated molecules, such as carbon disulfide, carbon dioxide, alkynes, heterocumulenes etc., into the hydride niobium and tantalum bond, which constitute fundamental chemical steps in several catalytic cycles and frequently lead to reactive intermediates [3b], have been considered. Finally, in another interesting aspect of the reactivity, the selective substitution of protons in the coordination sphere of niobium or tantalum hydride complexes by metallic fragments occurred. This gave rise to the synthesis of a series of heterometallic hydride species associating several transition metals and Group 5 metals.

This review will summarize, in a comprehensive fashion, the synthesis, characterization, structure and reactivity of bis-cyclopentadienyl hydride complexes of niobium and tantalum. Certain areas, however, have received a good deal more attention than others for various reasons, and in these cases we have tended to adhere to the pattern of the original studies. Coverage of the literature is up to the middle of 1998.


## 2. Hydrogen exchange coupling in metallocene trihydride complexes

The trihydride niobocene or tantalocene complexes,  $\text{Cp}_2\text{MH}_3$  ( $\text{M} = \text{Nb}, \text{Ta}$ ,

$\text{Cp} = \text{C}_5\text{H}_5$ ,  $\text{C}_5\text{Me}_5$ ,  $\text{C}_5\text{H}_4(\text{SiMe}_3)$ ,  $\text{C}_5\text{H}_3(\text{SiMe}_3)_2$  and related substituted cyclopentadienyl rings), are probably the most useful starting materials in the bis-cyclopentadienyl hydride chemistry of these elements. References to various preparations of these compounds and selected references for their use in synthesis can be found within Ref. [2] of this review. In particular, the preparation of niobium trihydride containing silylated cyclopentadienyl rings has been described [6] using the reaction of  $\text{Cp}'_2\text{NbCl}_2$  or  $\text{Cp}'_2\text{NbCl}_2$  ( $\text{Cp}' = \text{C}_5\text{H}_4\text{SiMe}_3$ ,  $\text{Cp}' = \text{C}_5\text{H}_3(\text{SiMe}_3)_2$ ) with  $\text{NaAlH}_2(\text{OCH}_2\text{CH}_2\text{OMe})_2$  ('Red-Al'). This method, i.e. the reaction of a dichloride metallocene of niobium or tantalum with an excess of a hydride-containing aluminum reagent, has been widely employed for the synthesis of metallocene trihydride complexes. The properties of the niobocene trihydride,  $\text{Cp}_2\text{NbH}_3$ , have been of great interest ever since Parshall and Tebbe [7] demonstrated its activation of carbon–hydrogen bonds. In particular, the  $^1\text{H}$  NMR properties of this complex have been found to be anomalous in many respects. The first study reported that the hydride gave two separate broad resonances at room temperature (r.t.), which sharpened at low temperature [8]. Subsequently, Curtis et al. observed a small temperature-dependence of the coupling constant  $J(\text{H},\text{H})$  of the hydrido hydrogen that was not explained [9]. Similarly, the adduct of  $\text{Cp}_2\text{NbH}_3$  with  $\text{AlEt}_3$  exhibits a surprisingly large value of  $J(\text{H},\text{H})$  [8]. This anomalous behavior was found to be due to a phenomenon of hydrogen exchange coupling 20 years ago, a situation that is present in several trihydrides of Ru [10], Ir [11], Nb [6,12], Os [13] and Mo [12], all of which adopt the same geometry, implying a planar array for the three hydrides and the metal. Exchange coupling between only two hydrides has been observed in a tantalum dihydride [14]. The quantum mechanical nature of the coupling was recognized by Zilm [15] and Weitekamp [16], who proposed the existence in these compounds of exchange coupling between two hydrogen atoms. Exchange couplings are known in EPR, but their observation in 'heavy' particles such as protons was novel in coordination chemistry. Heinekey [12] has analyzed the effect of the exchange coupling in the metallocene trihydride complexes  $\text{Cp}_2\text{MH}_3$  ( $\text{M} = \text{Nb}, \text{Ta}$ ;  $\text{Cp} = \text{C}_5\text{H}_5$ ) by  $^1\text{H}$  NMR spectroscopy. In this study, moderate values of the  $J_{\text{A-B}}$  coupling were observed at r.t., a fact consistent with an  $\text{AB}_2$  spin system for the hydride ligands. The Ta complex exhibits a proton coupling of 9.5 Hz, which is independent of the temperature in the range 203–303 K. Therefore, no significant exchange coupling was found in this metallocene. Conversely, for the Nb complex, a clear temperature dependence is found for the observed coupling that decreases with decreasing temperature, practically vanishes at 243 K and then reappears at 173 K. This phenomenon was attributed to a modest quantum mechanical exchange coupling, which is a sensitive function of temperature, that has a value of 26.4 Hz at 303 K and a value of zero at 173 K. In these complexes that exhibit hydrogen exchange coupling, the observed constant,  $J_{\text{obs}}$ , is given by the relationship  $J_{\text{obs}} = -2J_{\text{exch}} + J_{\text{magn}}$ , where  $J_{\text{exch}}$  = exchange coupling and  $J_{\text{magn}}$  = magnetic coupling, and in the studied  $\text{Cp}_2\text{NbH}_3$  the two constants are of opposite sign. An exhaustive study [17] has been also carried out for the tantalum and niobium trihydrides containing silylated cyclopentadienyl

rings, namely  $\text{Cp}'_2\text{MH}_3$  and  $\text{Cp}''_2\text{MH}_3$  ( $\text{M} = \text{Nb}, \text{Ta}$ ). As in  $\text{Cp}_2\text{NbH}_3$ , but unlike the corresponding tantalum derivatives, the niobium trihydrides show exchange coupling in their high-field  $^1\text{H}$  NMR spectra due to a hydride tunneling phenomenon. The magnitudes of the exchange coupling are larger in the cases of  $\text{Cp}'_2\text{NbH}_3$  and  $\text{Cp}''_2\text{NbH}_3$  than in the case of  $\text{Cp}_2\text{NbH}_3$ , and this is a result of the decrease in electron density upon increasing the number of  $\text{SiMe}_3$  substituents on the Cp ring. The tantalum trihydride derivatives show an  $\text{AB}_2$  pattern at high field for the hydride ligands with coupling constants of ca. 10 Hz that are not dependent upon the temperature. In contrast, the niobium complexes  $\text{Cp}'_2\text{NbH}_3$  and  $\text{Cp}''_2\text{NbH}_3$  show an  $\text{AB}_2$  pattern at r.t. for the hydrides, but in this case they are characterized by large coupling constants (36.5 and 70 Hz, respectively) that vary with temperature but not with the magnetic field. Further lowering of the temperature did not lead to the reappearance of this coupling. The longitudinal relaxation time,  $T_1$ , for the trihydrides  $\text{Cp}'_2\text{NbH}_3$  and  $\text{Cp}''_2\text{NbH}_3$  were measured. It was found that all hydrides show the same minimum of 40 and 55 ms, respectively, at 205 K (the value should be corrected to 100 ms when the influence of the Nb–H dipole–dipole relaxation is accounted for). In spite of the large number of studies performed on hydrogen exchange coupling in several polyhydride complexes, the exact physical processes accounting for the observation of the phenomenon still remain open to discussion. Zilm and colleagues have proposed a vibrational model to explain this coupling [15] whereas other authors have proposed that rotational tunneling of dihydrogen in a thermally accessible dihydrogen state could account for it [18]. Indeed, ab initio calculations show that niobium trihydrides possess a hydrido dihydrogen state that is almost degenerate in energy, whereas such a state is much higher in energy in the case of tantalum trihydrides [19]. Furthermore, a number of ab initio electronic energy calculations using a one-dimensional tunneling model have recently been carried out on metallocenes  $[\text{Cp}_2\text{MH}_3]^n+$  [ $\text{M} = \text{Mo}, \text{W}$ , ( $n = 1$ ) and  $\text{M} = \text{Nb}, \text{Ta}$  ( $n = 0$ )], establishing that the stability of the  $\eta^2\text{-H}_2$  structure relative to the minimum energy trihydride is the main parameter governing the magnitude of the exchange coupling [20]. In the search for a chemical discrimination between a purely vibrational mechanism and one implying rotational tunneling of dihydrogen, the addition of a Lewis acidic coinage cation to the niobocene trihydrides containing silylated cyclopentadienyl rings was extensively considered. Such additions were expected to inhibit the vibrational mode proposed to be at the origin of exchange coupling in Zilm's model. In particular, the addition of copper, silver or gold cations to the trihydride niobocene complexes  $\text{Cp}_2\text{NbH}_3$ ,  $\text{Cp}'_2\text{NbH}_3$  or  $\text{Cp}''_2\text{NbH}_3$ , was considered [17,21]. In fact, these complexes react with  $[\text{Cu}(\text{MeCN})_4]\text{PF}_6$ ,  $\text{AgBF}_4$ , and  $[\text{Au}(\text{THT})\text{PF}_6]$ , in a 2:1 ratio to yield the adducts  $[\{\text{Nb}(\text{C}_5\text{H}_3\text{RR}')_2\text{H}_3\}\text{M}]^+$  ( $\text{M} = \text{Cu}$ ,  $\text{R} = \text{R}' = \text{H}$ ;  $\text{R} = \text{H}$ ,  $\text{R}' = \text{SiMe}_3$ ;  $\text{R} = \text{R}' = \text{SiMe}_3$ ;  $\text{M} = \text{Ag}$ ,  $\text{R} = \text{H}$ ,  $\text{R}' = \text{SiMe}_3$ ;  $\text{R} = \text{R}' = \text{SiMe}_3$ ;  $\text{M} = \text{Au}$ ,  $\text{R} = \text{R}' = \text{H}$ ;  $\text{R} = \text{H}$ ,  $\text{R}' = \text{SiMe}_3$ ;  $\text{R} = \text{R}' = \text{SiMe}_3$ ) (Scheme 1).


The addition of a Lewis acid cation results in the observation of an  $\text{AB}_2$  pattern for the hydrides at r.t., which is split into an ABC pattern at low temperature. This




Scheme 1. Adapted from Ref. [17].

fact is in agreement with a fluxional behavior of the cation that binds to two hydrides of each niobium center. This structural situation has been confirmed by means of an X-ray crystal structure determination for the gold adduct with the  $\text{C}_5\text{H}_3(\text{SiMe}_3)_2$  ring. The activation energies of these fluxional processes are close to  $42\text{--}45 \text{ kJ mol}^{-1}$  in the case of Cu adducts,  $37 \text{ kJ mol}^{-1}$  in the case of Ag adducts, and  $40 \text{ kJ mol}^{-1}$  in the case of Au adducts. The magnitude of the exchange coupling is reduced upon addition of the copper cation, it is of the same order of magnitude after addition of a silver cation, and is greatly increased by addition of a gold cation (Scheme 2).

A model has been proposed to explain these variations which involves two isomeric states that are close in energy, one involving two bridging and one terminal hydride ions on niobium and one involving one bridging hydride and a dihydrogen molecule. The difference in the magnitude of the exchange coupling between the coinage elements has been explained as follows. The fact that the presence of copper inhibits exchange coupling could arise from the stabilization of the dihydrido-bridged form, which could result from the good Lewis acid character of the copper cation. Furthermore, a coordination number of 2, although not impossible, is unlikely for copper. When the copper is substituted by silver or gold, the presence of a more extended electronic cloud will render the coinage cations less acidic and therefore less strongly bound to the hydrides. This could explain both the higher classical fluxionality and the increase in exchange coupling for silver compared to copper. However, the very significant increase in exchange coupling, but decrease in fluxionality, of the gold adducts compared to the silver ones looks anomalous. It has been proposed that this probably results from a stabilization of the dihydrogen form, perhaps through electron donation from the niobium(III) center to the gold cation. This type of interaction was previously observed in a related niobium/gold raft cluster [22]. Furthermore, an excellent model of the isomeric state with a bridging hydride and a dihydrogen molecule with copper, in which  $\text{H}_2$  was substituted by CO, has been isolated and characterized by an X-ray crystal structure determination [23]. In order to theoretically study the stabilization of the  $\eta^2\text{-H}_2$ -containing state as a thermally accessible state in the trihydride niobocenes, different studies have been carried out more recently. For the first time,



Scheme 2. Adapted from Ref. [17].




Scheme 3. Adapted from Ref. [24].

ab initio DFT calculations have been presented [24] that indicate that the formation of an adduct between a trihydride complex,  $\text{Cp}_2\text{NbH}_3$ , and a Lewis acid,  $\text{BH}_3$ , converts the dihydrogen structure into a stable complex. In fact, very interestingly, the interaction of  $\text{BH}_3$  with a lateral hydride leads to the formation of a dihydrogen complex. This structure is only  $3.3 \text{ kcal mol}^{-1}$  above the trihydride structure in energy (Scheme 3).

The formation of this stable  $\eta^2\text{-H}_2$  complex can be attributed to the diminution of the electronic density at the niobium metal caused by the presence of the Lewis acid attached to one of its hydride ligands. The authors have also considered the elimination of molecular hydrogen from the original trihydride complex. The final product of this process confirms the formation of the  $\text{Cp}_2\text{Nb}(\eta^2\text{-BH}_4)$  (Scheme 4).

Following the study on the effect of a Lewis acid on the quantum exchange coupling in niobocene trihydrides, by means of the density functional methodology (DFT), this effect has also been calculated for the different adducts and transition states that appear in the  $\text{Cp}_2\text{NbH}_3\cdot\text{AlH}_3$  system [25]. A Lewis acid of the  $\text{AlH}_3$ -type may increase the exchange coupling through formation of an adduct where the Lewis acid is only bonded to the outer hydride. In this structure, which is not the absolute minimum but is close enough in energy to be significantly populated at the experimental range of temperatures, the two 'free' hydrides came closer to each other so that an incipient dihydrogen ligand that favors the exchange tends to be formed. Finally, the incoherent contribution to the tunneling exchange of hydrogen pairs in the complex  $\text{Cp}'_2\text{NbH}_3$  has been investigated [26] by  $^1\text{H}$  NMR line-shape analysis and two-dimensional  $^1\text{H}$  NMR exchange spectroscopy. The experimental data were analyzed by fitting them with simulated one- and two-dimensional spectra, taking into account the coherent and incoherent tunneling contributions and nuclear cross relaxation. The dissipative exchange process shows an Arrhenius-like temperature dependence with an activation energy  $E_a = 69(2) \text{ kJ mol}^{-1}$  over the temperature range 246–341 K.




Scheme 4. Adapted from Ref. [24].

### 3. $\sigma$ -Bond activation by niobocene and tantalocene hydrides

Activation of  $\sigma$ -bonds, principally C–H bonds, by transition metal complexes is currently an area of intense research activity, and several important advances in this area have recently been reported [27]. Although late metal complexes are usually applied to these reactions,  $\text{Cp}_2\text{MH}_3$  complexes ( $\text{Cp} = \text{C}_5\text{H}_5$ ;  $\text{M} = \text{Nb, Ta}$ ) catalyze [7a,28] H/D exchange between  $\text{H}_2$  and  $\text{C}_6\text{D}_6$ , as well as the exchange between  $\text{PhH}$  and  $\text{AlEt}_3$ . In a similar way [9], these trihydride derivatives react with hydrosilanes,  $\text{PhMe}_2\text{SiH}$ , to give the appropriate silyl complexes  $\text{Cp}_2\text{M}(\text{H})_2(\text{SiMe}_2\text{Ph})$ . This method was extended [29] to niobocene trihydride complexes bearing electron-withdrawing substituents,  $\text{SiMe}_3$ , in the cyclopentadienyl rings (Scheme 5), and gave rise to the corresponding niobocene silyl dihydride species.


The initial step, presumably, is the formation of the  $16e^-$  coordinatively unsaturated  $\text{Cp}'_2\text{NbH}$  by thermolytic loss of  $\text{H}_2$  (see Scheme 5). This process is then followed by oxidative addition of the organosilicon hydride, so that a closed-shell  $18e^-$  configuration in the silyl dihydride derivative is achieved. A similar intermediate was proposed by Berry et al. [30] for the thermolysis ( $> 100^\circ\text{C}$ ), or photolysis (350 nm) of  $\text{Cp}_2\text{Ta}(\text{L})(\text{H})$  ( $\text{L} = \text{CO, PMe}_3, \text{CH}_2=\text{CH}_2, \text{H}_2$ ) in the presence of hydrosilanes, to afford the corresponding tantalum silyl dihydrides  $\text{Cp}_2\text{Ta}(\text{H})_2(\text{SiR}_3)$  ( $\text{SiR}_3 = \text{SiH}_3, \text{SiMe}_2\text{H, SiMe}_3, \text{SiMe}_2\text{Cl, Si(OMe)}_3$ ).  $\text{Cp}_2\text{NbH}_3$  has been found [9] to be an excellent catalyst for the redistribution of groups on silicon. Heating a solution of  $\text{HSiMe}_2\text{OSiMe}_3$  and  $\text{Cp}_2\text{NbH}_3$  in benzene gave rise to a small amount of conversion of the pentamethylsiloxane into higher polysiloxanes. The reaction presumably involves oxidative addition of Si–H, Si–C, and Si–O bonds onto a low-valent metal species. In a stoichiometric reaction, the compounds isolated were the starting materials and  $\text{Cp}_2\text{Nb}(\text{H})_2(\text{SiMe}_2\text{OSiMe}_3)$ . It was expected that the complex  $\text{Cp}_2\text{NbH}_3$  could catalyze the formation of phenylsilanes in a productive activation process according to Scheme 6 [9], due to the fact that it catalyzes H/D exchange between  $\text{H}_2$  and  $\text{C}_6\text{D}_6$  [7a].

However, heating a 1:10 mixture of  $\text{Cp}_2\text{NbH}_3$  and  $\text{PhMe}_2\text{SiH}$  in  $\text{C}_6\text{D}_6$  to  $60^\circ\text{C}$  for 16 h gave a quantitative conversion of  $\text{Cp}_2\text{NbH}_3$  into  $\text{Cp}_2\text{Nb}(\text{H})_2(\text{SiPhMe}_2)$  and the excess  $\text{PhMe}_2\text{SiH}$  was found to be ca. 2% monodeuterated. When  $\text{Et}_3\text{SiH}$  was used, 20% of  $d^1 \text{Et}_3\text{SiH}$  was recovered. To explain why phenylsilanes are not formed even though both the C–D and Si–H bonds are activated by the niobium complex, the stereochemistry of the oxidative addition on the monohydride niobocene intermediate must be considered. This oxidative addition can give rise to two isomers, i.e. symmetrical and asymmetrical (Scheme 7).



Scheme 5. Adapted from Ref. [29].






Scheme 6. Adapted from Ref. [9].


The silyl derivatives described above were widely characterized by NMR and X-ray diffraction studies. These techniques showed the presence of only the symmetrical isomer with the silyl group in the central position replacing the central hydrogen. As the asymmetrical isomer was not formed, productive activation was not observed. The asymmetrical complexes could eliminate the *cis* hydrides as  $H_2$  and form intermediates that could add new Si-H or C-H bonds to give a productive activation process (Scheme 6). Although stable  $Cp_2Nb(H)_2(Ar)$  complexes have not been isolated, it is reasonable to assume that the Ar group will also occupy the central position in this process. This would give the symmetrical isomer and non-productive activation would also take place.

However, a mixture of symmetrical and asymmetrical isomers was obtained [30] in the photochemical reactions between  $Cp_2Ta(H)(L)$  and hydrosilanes. The minor asymmetrical isomer could not be isolated from the mixture and isomerizes, by means of an intramolecular exchange, to the thermodynamically stable symmetrical isomer. Alternative routes have been described [31] to obtain asymmetrical dihydride silyl complexes. The transfer of a silylene to tantalum hydride complexes proceeds by the insertion of this silylene into the Ta-H bond (Eqs. (1 and 2)).



Scheme 7.

In the reaction shown in Eq. (2), a 9:1 mixture of symmetrical and asymmetrical complexes was obtained. In order to complete the series of dihydride niobocene derivatives containing Group 14 fragments, analogous  $d^0$  niobocene germyl and stannyl complexes,  $\text{Cp}_2\text{Nb}(\text{H})_2(\text{ER}_3)$  ( $\text{ER}_3 = \text{SnPh}_3, \text{GePh}_3, \text{GePh}_2\text{H}, \text{GeEt}_3, \text{Ge}(\text{C}_6\text{H}_{13})_3, \text{Ge}[(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2]_3, \text{Ge}(\text{C}_6\text{H}_{13})_2\text{Cl}, \text{Ge}[(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2]_2\text{Cl}, \text{Ge}(\text{C}_6\text{H}_{13})_2\text{H}, \text{Ge}[(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2]_2\text{H}$ ) have been prepared [32] by the reaction between  $\text{Cp}_2\text{NbH}_3$  and several organogermanium hydrides,  $\text{HGeR}_3$ , or  $\text{HSnPh}_3$ , in toluene at  $65^\circ\text{C}$ . This is the first time that  $d^0$  early metal germyl complexes have been prepared by oxidative addition of organogermanium hydrides to coordinatively unsaturated hydride substrates. In a similar way to the silyl niobocene complexes described above, these germyl or stannyl derivatives show a symmetrical disposition with respect to the niobium center. Although the germyl niobocene complexes are the first described, precedents of the analogous stannyl complex were reported by Lappert [33] and Green [34]. Thus, the compound  $\text{Cp}_2\text{Ta}(\text{H})_2(\text{SnMe}_3)$  was prepared by the reaction of the trihydride with  $\text{SnMe}_3(\text{NMe}_2)$ . In an alternative approach, the addition of  $\text{BuLi}$  to a toluene solution of  $\text{Cp}_2\text{NbH}_3$  gave rise to a solid described as  $[\{\text{Cp}_2\text{NbH}_2\text{Li}\}_x]$ , which can be isolated as a stable  $[\text{Li}(12\text{-crown-4})][\text{NbCp}_2\text{H}_2]$  salt. Addition of one equivalent of  $\text{SnMe}_3\text{Cl}$  to this niobium complex led to the isolation of the complex  $\text{Cp}_2\text{Nb}(\text{H})_2(\text{SnMe}_3)$ . The possibility that the two hydrogens would be present as a dihydrogen ( $\eta^2\text{-H}_2$ ) ligand in this last complex was ruled out by the preparation of the deuterium-substituted complex  $\text{Cp}_2\text{Nb}(\text{H})(\text{D})(\text{SnMe}_3)$  by the reaction of  $\text{Cp}_2\text{NbH}_3$  with  $\text{SnMe}_3\text{D}$ . The  $^1\text{H}$  NMR spectrum of the complex shows an unresolved H–D coupling in the broad hydride signal, with  $J(\text{H–D}) < 10$  Hz. This may be compared with the typical values of molecular hydrogen complexes ( $22 < J(\text{H–D}) < 36$  Hz). The hydride  $T_1$  relaxation time reaches a minimum of 116 ms at 183 K, indicative of a classical dihydride structure. This classical behavior is in contrast with the reactivity of this compound. In fact, the smooth reaction of  $\text{Cp}_2\text{Nb}(\text{H})_2(\text{SnMe}_3)$  with two-electron donor ligands,  $\text{PMe}_3$  and styrene, occurred and resulted in the addition of the ligand and displacement of  $\text{H}_2$ . The elimination of  $\text{H}_2$  would presumably require initial interchange of position of one hydrogen atom with the  $\text{SnMe}_3$  ligand, possibly via an  $\eta^2\text{-H}_2$  or  $\eta^2\text{-HSnMe}_3$  intermediate. In the silyl dihydride derivatives that have been described,  $^{29}\text{Si}$  NMR data can be used to detect the presence of a  $\text{H}\cdots\text{Si}$  interaction. Corriu and co-workers [35] showed that the  $^1J(\text{Si–H})$  value for the agostic-like  $\text{H}\cdots\text{Si}$  interaction is near to 65 Hz, thus appearing between that characteristic of classical silyl hydrides ( $\approx 6$  Hz) and that of free silanes ( $\approx 200$  Hz). The  $^{29}\text{Si}$  NMR spectrum of  $\text{Cp}_2\text{Nb}(\text{H})_2(\text{SiPh}_2\text{H})$  [29] shows a doublet at  $\delta = 23.8$  for the silyl group with  $^1J(\text{Si–H}) = 17$  Hz. This coupling disappeared when the hydrogen of the silyl group was irradiated. In the  $^1\text{H}$  NMR spectrum, the coupling is also observed for the resonance of this hydrogen atom and, in addition, the line width of the signals for the two hydride ligands (6 Hz) is characteristic of a classical silyl hydride. These spectroscopic data establish the classical nature of these silyl hydrides. However, in the solid state the molecular structure cannot exclude any interaction between the hydrides and the silicon atom on the basis of van der Waals radii, in a hypervalence mode for the silicon atom [36].




Scheme 8. Adapted from Ref. [29].

The reactivity of hydride-, silyl-, germyl- or stannyl-containing niobocene complexes towards the hydrides  $\text{HEPh}_3$  ( $\text{E} = \text{Si, Ge, Sn}$ ) in  $\sigma$ -bond metathesis reactions of  $d^0$  Nb–E bonds with H–E'  $\sigma$  bonds have been studied [32] (Scheme 8). The reactions were monitored by  $^1\text{H}$  NMR spectroscopy. From these studies the activation parameters  $\Delta H^\ddagger$  and  $\Delta S^\ddagger$  were obtained. These parameters allow the estimation of the bond dissociation enthalpies, which were found to be comparable to Nb–H, Nb–Si, Nb–Ge and Nb–Sn bonds, with values near to  $50 \text{ kcal mol}^{-1}$ .

However, the experimental results indicate the following reactivity trend:  $\text{Nb–H} \approx \text{Nb–Si} > \text{Nb–Ge} \gg \text{Nb–Sn}$  for the  $\sigma$ -bond metathesis processes. In fact,  $\text{Cp}'_2\text{NbH}_3$  reacts with  $\text{HEPh}_3$  ( $\text{E} = \text{Si, Ge, Sn}$ ),  $\text{Cp}'_2\text{Nb}(\text{H})_2(\text{SiPh}_3)$  reacts with  $\text{H}_2$ ,  $\text{HGePh}_3$  or  $\text{HSnPh}_3$ ,  $\text{Cp}'_2\text{Nb}(\text{H})_2(\text{GePh}_3)$  reacts with  $\text{HSnPh}_3$ , but  $\text{Cp}'_2\text{Nb}(\text{H})_2(\text{SnPh}_3)$  does not react with either  $\text{H}_2$ ,  $\text{HSiPh}_3$  or  $\text{HGePh}_3$ . The results can be rationalized in terms of differences in reactant and product bond strengths. The energy required to break the Nb–H, Nb–Si or Nb–Ge bonds is compensated by formation of relatively strong H–H, H–Si and H–Ge bonds in the  $\text{HX}$  molecule eliminated in the first step (see Scheme 8). In contrast, the formation of the weaker H–Sn bond does not compensate the energy required to break the strong Nb–Sn bond. A similar explanation has been postulated by Tilley et al. [37] to rationalize the reactivity trends of analogous  $d^0$  zirconocene complexes in bond metathesis reactions that parallel those described above in the niobium system.

Recently, a similar dihydride niobocene containing a  $\text{PbMe}_3$  moiety, namely  $\text{Cp}_2\text{NbH}_2(\text{PbMe}_3)$ , was described [38a] to complete the family of niobocene dihydride-containing Group 14 fragments. This compound was prepared by reacting  $\text{Cp}_2\text{NbH}_3$  with  $\text{ClPbMe}_3$  in the presence of amine to eliminate  $\text{HCl}$ .

A number of other main group element–hydrogen bonds can be activated by the trihydride niobocene complex. Thermolysis of  $\text{Cp}'_2\text{NbH}_3$  in the presence of pyrazole gives rise to a pyrazolyl–pyrazole niobocene,  $\text{Cp}'_2\text{Nb}(\text{pzH})(\text{pz})$  [39a] (Scheme 9).




Scheme 9. Adapted from Ref. [39a].

Two mechanisms can be envisaged for the formation of this complex. In the first possibility (Scheme 9, route I), N–H bond activation leads to an asymmetrical dihydride intermediate, which loses  $\text{H}_2$  and coordinates the second molecule of pyrazole. A second possibility (Scheme 9, route II) involves pyrazole coordination by a pyridine nitrogen as the first step. Subsequently, the monohydride–pyrazole complex evolves with protonation, probably in an intramolecular mode with the loss of  $\text{H}_2$ . Finally, coordination of the second pyrazole substituent takes place. Activation of P–H bonds was also observed [39b] in the reactions of  $\text{Cp}_2\text{MH}_3$  ( $\text{M} = \text{Nb}$ ,  $\text{Ta}$ ) with an excess of primary phosphines. These reactions yield the diphosphonato complexes  $\text{Cp}_2\text{Nb}(\text{H})\{(\text{PPh})_2\}$ ,  $\text{Cp}_2\text{Nb}(\text{H})\{(\text{PPh})_2\}$ , and  $\text{Cp}_2\text{Ta}(\text{H})\{(\text{PPh})_2\}$ . The insertion of Group 15 halogenides into the M–H bond is an alternative route to bond activation for the synthesis of main-group element-substituted niobocene and tantalocene hydrido complexes. Phosphorus or arsenic monohalogenides  $\text{ClPPh}_2$  [38b,c], or  $\text{BrAsR}_2$  ( $\text{R} = \text{Me}$ ,  $\text{Et}$ ) [38d], insert easily into the M–H bond of  $\text{Cp}_2\text{MH}_3$  to afford the ionic complexes  $[\text{Cp}_2\text{MH}_2(\text{HER}_2)]\text{X}$  ( $\text{M} = \text{Nb}$ ,  $\text{Ta}$ ,  $\text{E} = \text{P}$ ,  $\text{R} = \text{Ph}$ ,  $\text{X} = \text{Cl}$ ;  $\text{M} = \text{Nb}$ ,  $\text{E} = \text{As}$ ,  $\text{R} = \text{Me}$ ,  $\text{Et}$ ,  $\text{X} = \text{Br}$ ). Deprotonation of these complexes with a base, i.e.  $\text{NEt}_3$  or aqueous  $\text{NaOH}$ , gives rise to the hydride phosphino-, arsenido- or phosphido derivatives  $\text{Cp}_2\text{NbH}(\text{PPh}_3)$ ,  $\text{Cp}_2\text{NbH}_2(\text{AsR}_2)$  and  $\text{Cp}_2\text{TaH}_2(\text{PPh}_2)$ .


In recent years, niobocene complexes containing metal–boron bonds,  $\text{Cp}_2\text{Nb}(\text{H}_2\text{BR}_2)$  ( $\text{R}_2 = \text{O}_2\text{C}_6\text{H}_4$ ,  $\text{C}_8\text{H}_{14}$ ), were prepared [40] by the reaction between  $\text{Cp}_2\text{NbH}_3$  and hydroboranes. Analogous compounds,  $\text{Cp}_2^*\text{M}(\text{H}_2\text{B}(\text{O}_2\text{C}_6\text{H}_4))$ , were obtained from the reaction between  $\text{Cp}_2^*\text{M}(\text{CH}_2=\text{CH}(\text{R}))(\text{H})$  ( $\text{Cp}^* = \text{C}_5\text{Me}_5$ ,  $\text{M} = \text{Nb}$ ,  $\text{Ta}$ ) and two equivalents of catecholborane. In the last case, experimental results are consistent with a mechanism involving an unsaturated alkyl intermediate (Scheme 10) that activates the H–B bond.

As an alternative, analogous tantalocene complexes  $\text{Cp}_2\text{Ta}(\text{H})_2(\text{B}(\text{O}_2\text{C}_6\text{H}_4))$  in the form of their symmetrical and asymmetrical isomers can be obtained [42] by the reaction between  $[\{\text{Cp}_2\text{TaH}_2\text{Li}\}_x]$  and  $\text{ClBO}_2\text{C}_6\text{H}_4$ . For these  $\text{Cp}_2\text{MH}_2\text{BX}_2$  ( $\text{Cp} = \text{C}_5\text{H}_5$  or  $\text{C}_5\text{Me}_5$ ) complexes, three limiting forms can be used to describe the  $\text{H}_2\text{BX}_2$  coordination. These include  $d^0$  boryl (A),  $d^2$  borohydride (B), and  $d^2$  agostic borane (C) structures (Scheme 11).

While tantalum complexes are clearly characterized as boryl complexes of type A, the niobium ones show a pronounced Nb(III) character. For catechol-containing niobocene derivatives, equilibria between situations A and B were proposed on the



Scheme 10. Adapted from Ref. [41].



Scheme 11.

basis of spectroscopic data and isotopic perturbations of their NMR spectra. Spectroscopic data support some B–H interaction for *exo*-Cp<sub>2</sub>TaH<sub>2</sub>(BO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>), although the sum of the angles about boron, from the molecular structure, is consistent with an sp<sup>2</sup> hybridization at the boron atom and suggests that this compound is a tantalum(V) boryl dihydride rather than an agostic borane of tantalum(III).

#### 4. The chemistry of d<sup>2</sup> hydride niobocene and tantalocene derivatives

In this chapter the most extensive family of hydride metallocene complexes of niobium and tantalum will be considered, and this family corresponds to the 18-electron d<sup>2</sup> Cp<sub>2</sub>MHL (M = Nb, Ta, L = π-acid ligand). Complexes have been described with different classes of π-acid ligands obtained by several types of preparation processes. In the following discussion, the hydride–olefin metallocenes have been treated in a different section than the rest of complexes due to their special significance in catalysis.

##### 4.1. Preparations

Table 1 includes references to various preparations of these complexes with different types of cyclopentadienyl rings.

The thermal treatment of Cp<sub>2</sub>MH<sub>3</sub> species in the presence of L constitutes the most important process for the preparation of a huge family of Cp<sub>2</sub>MHL complexes (L = π-acid ligand). In most of the cases, the processes are not facile because harsh conditions (reflux in toluene during several days) for the elimination of H<sub>2</sub> and the subsequent coordination of the π-acid ligand are necessary. Moreover, the substitution is frequently incomplete so that the new Cp<sub>2</sub>MHL derivative contains impurities. Extensive families of complexes with different classes of both cyclopentadienyl rings and π-acid ligands, especially carbon monoxide or phosphine, have been thoroughly described and these are collected in Table 1. However, the Cp<sub>2</sub>'MH<sub>3</sub> (M = Nb, Ta, Cp' = η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>) complexes react easily [48,49,56] with π-acid ligands, using THF as solvent, to quantitatively give several hydride complexes, Cp<sub>2</sub>'MHL. Thus, Cp<sub>2</sub>'NbH<sub>3</sub> reacts under mild conditions in THF with phosphites to give new hydride niobocene complexes Cp<sub>2</sub>'NbH(P(OR)<sub>3</sub>). Similarly, the easy preparation in high yields (95%) [56] of hydride–isocyanide complexes has recently been reported: Cp<sub>2</sub>'NbH(CNR) (R = <sup>t</sup>Bu, Cy, Xyllyl) by thermal treatment in THF of Cp<sub>2</sub>'NbH<sub>3</sub> in the presence of the corresponding isocyanide. No subse-

Table 1

d<sup>2</sup> Hydride complexes of niobium and tantalum, Cp<sub>2</sub>MHL, M = Nb, Ta, L =  $\pi$ -acid ligand

| Cp <sub>2</sub> MH <sub>3</sub> + L → Cp <sub>2</sub> M(H)L |                                                                                                                 |                                                                                                                                                                                                                                                                                                               |
|-------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (a) M = Nb                                                  | Cp = C <sub>5</sub> H <sub>5</sub>                                                                              | L = CO, PMe <sub>3</sub> , PEt <sub>3</sub> [43], Ph <sub>2</sub> H [45], N <sub>2</sub> CPh <sub>2</sub> [46], PMe <sub>2</sub> Ph [47], <sup>n</sup> PrCC <sup>n</sup> Pr, MeCC <sup>n</sup> Pr, MeCCMe, MeCC <sup>n</sup> Pr [53], EtCCEt, PhCCPh, TMSCTMS [50,54], CF <sub>3</sub> CCCF <sub>3</sub> [55] |
|                                                             | Cp = C <sub>5</sub> H <sub>4</sub> SiMe <sub>3</sub>                                                            | L = CO, P(OMe) <sub>3</sub> , PhCCPh [48], P(OEt) <sub>3</sub> , P(OPh) <sub>3</sub> [49], CN <sup>n</sup> Bu, CNCy, CNXylyl [56]                                                                                                                                                                             |
|                                                             | Cp = C <sub>5</sub> H <sub>4</sub> <sup>n</sup> Bu                                                              | L = CO [57]                                                                                                                                                                                                                                                                                                   |
|                                                             | Cp = C <sub>5</sub> Me <sub>5</sub>                                                                             | L = CO [60], MeCCMe [61]                                                                                                                                                                                                                                                                                      |
|                                                             | Cp = C <sub>5</sub> H <sub>3</sub> (SiMe <sub>3</sub> ) <sub>2</sub>                                            | L = CO [6]                                                                                                                                                                                                                                                                                                    |
|                                                             | CpCp' = (C <sub>5</sub> H <sub>5</sub> )<br>(C <sub>5</sub> Me <sub>5</sub> )                                   | L = CO, MeCCMe [61]                                                                                                                                                                                                                                                                                           |
| (b) M = Ta                                                  | Cp = C <sub>5</sub> H <sub>5</sub>                                                                              | L = CO, PEt <sub>3</sub> , MeCCMe, PhCCPh, TMSCTMS [43], PMe <sub>2</sub> Ph, P(OMe) <sub>3</sub> [44], N <sub>2</sub> CPh <sub>2</sub> [46], <sup>n</sup> PrCC <sup>n</sup> Pr [51], <sup>n</sup> BuCC <sup>n</sup> Bu [52]                                                                                  |
|                                                             | Cp = C <sub>5</sub> H <sub>4</sub> SiMe <sub>3</sub>                                                            | L = CO [57b]                                                                                                                                                                                                                                                                                                  |
|                                                             | Cp = C <sub>5</sub> H <sub>4</sub> <sup>n</sup> Bu                                                              | L = CO [57]                                                                                                                                                                                                                                                                                                   |
|                                                             | Cp = C <sub>5</sub> Me <sub>5</sub>                                                                             | L = MeCCMe [61]                                                                                                                                                                                                                                                                                               |
|                                                             | Cp = C <sub>5</sub> H <sub>4</sub> <sup>n</sup> Pr                                                              | L = CO [58]                                                                                                                                                                                                                                                                                                   |
|                                                             | Cp = C <sub>5</sub> H <sub>4</sub> R                                                                            | L = CO [57b,59]                                                                                                                                                                                                                                                                                               |
|                                                             | R = CMe <sub>2</sub> Ph, CHMePh                                                                                 |                                                                                                                                                                                                                                                                                                               |
|                                                             | CpCp' = (C <sub>5</sub> Me <sub>5</sub> )<br>(C <sub>5</sub> H <sub>3</sub> (SiMe <sub>3</sub> ) <sub>2</sub> ) | L = CO, CNXylyl [62]                                                                                                                                                                                                                                                                                          |

quent insertion process was observed even when the process was carried out with an excess of ligand in refluxing THF for a long time. This behavior contrasts with that found for hydride Group 4 metallocenes, where the insertion process to give iminoacyl derivatives is especially favoured [63]. In these processes, the initial step would be the formation of the coordinatively unsaturated 16-electron species 'Cp<sub>2</sub>MH' by thermolytic loss of H<sub>2</sub> followed by the coordination of the  $\pi$ -acid ligand. The mechanism for the formation could be explained in terms of a model, which implies a consecutive reaction with a reversible step and a first-order rate dependence with respect to Cp<sub>2</sub>MH<sub>3</sub> [64]. On the other hand, an alternative preparation of Cp<sub>2</sub>MHL complexes has been described [65] involving the reaction of Cp<sub>2</sub>MCl<sub>2</sub> with NaBH<sub>4</sub> followed by the addition of triethylamine in the presence of the appropriate  $\pi$ -acid ligand. Likewise, the reaction of Cp<sub>2</sub>MCl<sub>2</sub> with LiAlH<sub>4</sub> in the presence of a phosphine (PMe<sub>2</sub>Ph, PMe<sub>3</sub>) allows the synthesis of the same class of complexes Cp<sub>2</sub>MHL [47,66].

Alkyne-hydride derivatives constitute a special family of this type of complex (see Table 1). However, few alkyne-hydride complexes of niobium and tantalum have been isolated. Most of those that have, have been prepared by the reaction of Cp<sub>2</sub>MH<sub>3</sub> complexes with the appropriate alkyne ligand.


#### 4.2. Reactivity of the $\text{Cp}_2\text{MHL}$ and related species in insertion processes

In this section the reactivity of the hydride niobocene and tantalocene species, the  $d^2$   $\text{Cp}_2\text{MHL}$  complexes, will be discussed. In fact, many examples have been described of the insertion of unsaturated molecules into the M–H bond in hydride metallocenes of niobium and tantalum. In order to rationalize the discussion, the reactions with heterocumulene molecules will be considered first, and afterwards a specific group of insertion processes with alkynes will also be described.

##### 4.2.1. Insertion processes with heterocumulene molecules

Insertion reactions of heterocumulenes, like that carbon disulfide, isocyanate or isothiocyanate, are especially interesting because they allow the modelling of the insertion reactions of carbon dioxide, a process that constitutes an attractive goal in the field of organometallic chemistry. The reaction of  $\text{Cp}_2\text{MHL}$  ( $\text{Cp} = \text{C}_5\text{H}_5$ ,  $\text{M} = \text{Nb}$ ,  $\text{Ta}$ ,  $\text{L} = \text{CO}$  [67],  $\text{Cp} = \text{Cp}' = \text{C}_5\text{H}_4\text{SiMe}_3$ ,  $\text{M} = \text{Nb}$ ,  $\text{L} = \text{CO}$ ,  $\text{P}(\text{OR})_3$  [49]) with  $\text{CS}_2$  proceeds with the expected insertion of the heterocumulene into the M–H bond to give a dithioformate ligand,  $\text{S}(\text{S})\text{CH}$ . The nature of the products for the  $\text{Cp}_2\text{NbHL}$  is dependent on the nature of L. When  $\text{L} = \text{CO}$ , the new ligand exhibits an  $\eta^1$ -dithioformate coordination, whereas when  $\text{L} = \text{P}(\text{OR})_3$  an  $\eta^2$ -dithioformate species is present in the reaction product. In the latter case, an initial  $\eta^1$ -dithioformate complex is probably formed, with loss of phosphite, to give the final  $\eta^2$ -dithioformate ligand, which is, however, unusual in this type of insertion process. When the heterocumulene is isocyanate ( $\text{RNCO}$ ) or isothiocyanate ( $\text{RNCS}$ ) the insertion reaction into the M–H bond gives the new formamido- or thioformamido-containing  $\text{Cp}_2\text{M}(\eta^1\text{-XC}(\text{H})\text{NR})(\text{L})$  complexes ( $\text{Cp} = \text{C}_5\text{H}_5$ ,  $\text{M} = \text{Nb}$ ,  $\text{Ta}$ ,  $\text{L} = \text{CO}$ ,  $\text{PMePh}_2$  [67,68],  $\text{Cp} = \text{Cp}' = \text{C}_5\text{H}_4\text{SiMe}_3$ ,  $\text{M} = \text{Nb}$ ,  $\text{L} = \text{CO}$ ,  $\text{CNR}$  [69]). The subsequent protonation of  $\text{Cp}_2\text{Nb}(\eta^1\text{-XC}(\text{H})\text{NR})(\text{L})$  complexes using  $\text{CF}_3\text{COOH}$  yields the trifluoroacetato complexes  $\text{Cp}_2\text{Nb}(\eta^4\text{-OOC}(\text{CF}_3)_3)(\text{L})$  and the imine derivatives  $\text{HOC}(\text{H})=\text{NR}$  and  $\text{HSC}(\text{H})=\text{NR}$ . The interaction of a trihydride niobocene,  $\text{Cp}_2'\text{NbH}_3$ , with carbon dioxide was investigated because metalloformates  $\text{M-OC}(\text{O})\text{H}$  derived from carbon dioxide insertion into M–H bonds are known to be involved in the catalytic hydrogenation of these compounds to formic acid and its derivatives [70]. The complex  $\text{Cp}_2'\text{NbH}_3$  reacted with  $\text{CO}_2$  to give the formate complex  $\text{Cp}_2'\text{Nb}(\text{OC}(\text{O})\text{H}, \text{O}, \text{O}') [71]$ . The isolation of this complex can be understood in terms of Scheme 12.

Step (a) corresponds to the elimination of  $\text{H}_2$  and the formation of a very reactive 16-electron monohydride niobocene species, which has previously been proposed in several processes [64]. Coordination of  $\text{CO}_2$  to this unsaturated species to give an




Scheme 12. Adapted from Ref. [71].

18-electron hydride–carbon dioxide intermediate is suggested in step (b). The final step (c) corresponds to the insertion of the coordinated  $\text{CO}_2$  into the Nb–H bond to give the formate ligand. The formation of hydride–carbon dioxide tantalocene intermediate at  $-40^\circ\text{C}$ , as well as the subsequent insertion to give formate tantalocene derivatives, has also been published [72]. Reactivity studies on the complex  $\text{Cp}_2\text{Nb}(\text{OC}(\text{O})\text{H}, \text{O}, \text{O}')$  have been carried out. Reaction with  $\text{CS}_2$  or the  $\pi$ -acid ligands CO and 2,6- $\text{Me}_2\text{C}_6\text{H}_3\text{NC}$  gave rise to the transformation from a bidentate to a monodentate formate as a consequence of coordination of the incoming ligand, and the  $\text{Cp}_2\text{Nb}(\text{OC}(\text{O})\text{H}, \text{O})\text{L}$  complexes were isolated. However, with other L, such as phosphites  $\text{P}(\text{OR})_3$ , the transformation has never been observed; this behavior suggests that the presence of a heterocumulene like  $\text{CS}_2$ , which is able to coordinate strongly to the niobium atom as a bidentate ligand or strong  $\pi$  acceptors, is necessary to break an Nb–O bond and to transform bidentate into monodentate formate.

#### 4.2.2. Insertion processes with alkynes

The insertion of alkynes into transition metal–hydrogen bonds is a key elementary step in catalytic polymerization and hydrogenation processes [3b]. In the last few years, the stereochemistry and mechanism involved in the insertion process have received considerable attention [73]. Insertion results in the formation of alkenyl products and these show that *cis* and *trans*-insertion yield the (*E*)-alkenyl and (*Z*)-alkenyl products, respectively, and that the importance of the *cis/trans* stereochemistry depends on several factors, such as the nature of the metal center, ancillary ligands, and substituents on the alkyne. Electron-rich, coordinatively saturated, metallocene–hydride complexes of early transition metals react readily with activated alkynes bearing electron-withdrawing groups to give alkenyl complexes. Thus, in earlier work the behavior of niobocene–hydrides, especially  $\text{Cp}_2\text{NbH}(\text{CO})$ , was extensively studied [73i,74] and most of the processes of hydride metallocenes with disubstituted activated alkynes occur by stereospecific *trans*-insertion reactions that give rise to kinetic products of (*Z*)-configuration [75], see Scheme 13. However, the hydride–isocyanide complexes  $\text{Cp}'\text{NbH}(\text{CNR})$  ( $\text{R} = \text{xylyl} = 2,6\text{-dimethylphenyl}$ , Cy,  $t\text{Bu}$ ,  $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$ ) react with several activated alkynes to afford the corresponding alkenyl–isocyanide complexes  $\text{Cp}'\text{Nb}(\text{CNR})(\text{C}(\text{R}')=\text{CH}(\text{R}''))$  though a stereospecific *cis*-insertion (see Scheme 13) resulting in the formation of (*E*)-alkenyl products [76].



Scheme 13. Adapted from Ref. [76].



As a consequence of these results, it has been established that in the presence of bulky ancillary ligands, such as some isocyanides, a primary *cis* stereospecific insertion under kinetic control affords the (*E*)-isomer, probably via the four-centered transition state shown in Scheme 13, which is favoured by steric arguments. In the same reaction for  $\text{Cp}_2\text{NbH}(\text{CO})$  with the activated alkynes, under kinetic control, a mixture of (*E*)- and (*Z*)-alkenyl niobocene complexes,  $\text{Cp}_2'\text{Nb}(\text{CO})\text{-(C(R')=CH(R''))}$ , was obtained. It has been proposed that a primary *trans* stereospecific insertion would occur via a four-centered transition state to give the kinetic (*Z*)-isomer and a subsequent isomerization would give the (*E*)-isomer, resulting in the mixture of (*E*)- and (*Z*)-isomers. The results obtained in the reaction of the trihydride metallocenes, namely  $\text{Cp}_2\text{TaH}_3$  and  $\text{Cp}_2\text{NbH}_3$ , with activated alkynes are particularly interesting [50,76]. In fact, these complexes react with activated alkynes, such as  $\text{RO}_2\text{CC}\equiv\text{CCHR}'$ , to afford new hydride–olefin complexes,  $\text{Cp}_2\text{Ta}(\text{H})(\text{trans-RO}_2\text{CCH=CHCO}_2\text{R}')$  and  $\text{Cp}_2\text{Nb}(\text{H})(\text{trans-RO}_2\text{CCH=CHCO}_2\text{R}')$ . This behavior is surprising because reactions of these trihydride metallocenes with alkynes are well known and generally give rise to alkyne–hydride metallocenes [53,55,77]. In order to explain the formation of these complexes, a stereospecific *trans*-insertion to give a dihydride alkenyl intermediate is suggested in the first step, followed by an intermolecular rearrangement that leads to the final hydride–olefin complexes.

#### 4.3. Reactivity of the $\text{Cp}_2\text{MHL}$ complexes in protonation processes: synthesis of dihydrogen complexes

The  $\text{d}^2$   $\text{Cp}_2\text{MHL}$  complexes behave as Lewis bases and therefore they can be protonated at r.t. by acids to give the corresponding cationic complexes, which have been isolated as the *trans*- $[\text{Cp}_2\text{MH}_2\text{L}]^+$  cationic dihydride species [44,78]. Thus, the complexes  $\text{Cp}_2'\text{NbHL}$  ( $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$ ) were protonated with  $[\text{PhNH}_3]\text{PF}_6$  in THF, allowing the isolation of the corresponding transoid cationic niobocene dihydrides *trans*- $[\text{Cp}_2'\text{Nb}(\text{H})_2\text{L}]\text{PF}_6$ . In the same way, *trans*- $[\text{Cp}_2\text{Ta}(\text{H})_2(\text{CO})]\text{PF}_6$  was obtained by protonation of  $\text{Cp}_2\text{TaH}(\text{CO})$  ( $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ ) with aqueous HCl at r.t. However, different results have been found in recent years when the protonation processes were carried out at low temperatures, since dihydrogen-containing metallocene species were stabilized. Protonation of a neutral hydride complex to give a cationic dihydrogen complex was first reported by Crabtree et al. [79]. Subsequently, numerous examples have been described [5]. A number of different families of dihydrogen-containing niobocene and tantalocene complexes have been described following this method. The tantalocene complex  $[\text{Cp}_2\text{Ta}(\eta^2\text{-H}_2)(\text{CO})]^+$  was prepared [80] by protonation of  $\text{Cp}_2\text{TaH}(\text{CO})$  ( $\text{Cp} = \text{C}_5\text{H}_5$ ,  $\text{C}_5\text{H}_4\text{Bu}$ ) at low temperature. The relaxation time,  $T_1$ , of the hydrides shows a minimum of 9 ms (for  $\text{C}_5\text{H}_5$ ) and 12 ms (for  $\text{C}_5\text{H}_4\text{Bu}$ ) at 178 K, which corresponds to dihydrogen complexes. Halpern's approximate equation [81] leads to a H–H distance of  $0.84 \pm 0.05$  Å (for  $\text{C}_5\text{H}_5$ ) and  $0.89 \pm 0.05$  Å (for  $\text{C}_5\text{H}_4\text{Bu}$ ). It is interesting to note the variation of the relaxation time of the hydride signals of both isomers when  $\text{C}_5\text{H}_5$  is replaced by  $\text{C}_5\text{H}_4\text{Bu}$ . This is due to the difference in the electronic properties of the Cp ligands:

the more electron releasing ligand  $C_5H_4Bu$  induces a lengthening of the H–H bands. In the deuterated complex  $[Cp_2Ta(\eta^2-HD)(CO)]^+$  the  $J_{HD}$  value is 26.5 Hz, which is in agreement with an unstretched dihydrogen structure. When  $CF_3CO_2H$  was added to an acetone- $d^6$  solution of  $Cp'_2NbHL$  ( $Cp' = \eta^5-C_5H_4SiMe_3$ ,  $L = P(OEt)_3$ ,  $PMe_2Ph$ ) at  $-90^\circ C$  in an NMR tube, dihydrogen-containing niobocene complexes  $[Cp_2Nb(\eta^2-H)L]^+$  were stabilized [78c]. When the temperature was increased the complex with  $L = P(OEt)_3$  transformed into its transoid isomer through an irreversible process. Increasing the temperature did not produce the transoid form of the complex with  $L = PMe_2Ph$ , but its qualitative transformation at  $20^\circ C$  into  $[Cp'_2Nb(CF_3CO_2)(PMe_2Ph)]$ . The analogous  $[Cp'_2Nb(CF_3CO_2)(CO)]$  is the only niobium compound observed at all temperatures when the complex  $Cp'_2NbH(CO)$  is reacted under similar conditions with  $CF_3CO_2H$ , probably because the large  $\pi$ -accepting character of the CO ligand prevents stabilization. The  $T_1$  minima were found to be 24 ms at 188 K and 19 ms at 196 K for the dihydrogen complexes with  $L = P(OEt)_3$  and  $L = PMe_2Ph$ , respectively. H–H distances of 0.97 Å, assuming rapid rotation of dihydrogen, and 1.22 Å, assuming a slow rotation, and 1.17 Å for the aforementioned complexes were also calculated. A value of  $J_{HD} = 18.2$  Hz was also found in the deuterated complex  $[Cp'_2Nb(\eta^2-HD)P(OEt)_3]^+$ , and this value is in good agreement with the distance calculated from  $T_1$  data. One of the most interesting points of the hydrogen-containing tantalocene and niobocene complexes described [78c,80] lies in the fact that activation barriers for the rotation of  $H_2$  are surprisingly high (ca. 10 kcal mol $^{-1}$ ), therefore providing an intermediate case between the normal dihydrogen complex and the polyhydride complexes. As a consequence of the high barrier, rotation of the dihydrogen molecule seems to be blocked from the NMR line scale [78c,82]. For these complexes a single resonance is obtained for the  $^1H$  NMR spectra at high temperatures. When the temperature is lowered, decoalescence is not observed for the nonisotopically substituted  $H_2$  isomer, but it is observed for the partially deuterated HD species. This has allowed estimation of the activation barriers for these complexes. This surprising result has been explained by the postulation of a large exchange coupling in the  $H_2$  case. It is known [82] that for  $J/\Delta\delta$  ratios higher than 10, the expected AB-type spectrum becomes a single line. Although it is expected that H–H and H–D have very similar barriers for the rotation, there is no direct experimental evidence that the H–H rotation is also being practically blocked. From a theoretical point of view, the rotational barrier for the H–H rotation has been calculated for the simplified model complexes  $[Cl_2Ta(\eta^2-H_2)CO]^+$  and  $[Cl_2Ta(\eta^2-H_2)(PH_3)]^+$  to be 9.7 and 11.3 kcal mol $^{-1}$ , respectively [82]. It was necessary to obtain a new experimental example of a non-rotating coordinated H–D molecule in order to analyze its structure and the hydrogen exchange process. Such an example would provide theoretical evidence that in the corresponding H–H molecule this process is also classically blocked at the NMR time scale and allow the estimation of the order of magnitude of the exchange coupling. For these reasons new dihydrogen niobocene complexes were prepared [56]. In fact, low-temperature protonation of hydride isocyanide derivatives  $Cp'_2Nb(H)(CNR)$  ( $Cp' = \eta^5-C_5H_4SiMe_3$ ) with a slight excess of  $CF_3CO_2H$  led to the dihydrogen complexes  $[Cp'_2Nb(\eta^2-H_2)(CNR)]^+$ . The monodeuterated com-


plexes have also been prepared and coupling constants  $J_{\text{HD}}$  of ca. 30 Hz verify the presence of a H–D bond and provide effective proof of the presence of an unstretched dihydrogen complex. H–H distances between 1.01 and 1.06 Å were found assuming slow  $\text{H}_2$  spinning. The NMR spectra of these H–H complexes and their monodeuterated H–D isotopomers show a single high-field resonance at r.t. On lowering the temperature to 178 K, decoalescence of the signal was observed for the H–D complexes but not for the H–H ones. By combining DFT electronic structure calculations with a monodimensional rotation tunneling model, it has been shown that the absence of decoalescence of the H–H signal is due to the existence of a very large exchange coupling. Conversely, for the H–D isotopomers, the difference in zero point energy corresponding to two nonequivalent (H–D and D–H) positions leads to a slight asymmetry, which dramatically reduces the exchange coupling, allowing decoalescence to be observed. Therefore, the H–D classical rotation and the quantum exchange processes will not be practically observed for this complex, whereas only the classical case for the H–H species is quenched out on the NMR time scale.

#### 4.4. The hydride–olefin niobocene and tantalocene complexes

Olefin–hydride metal systems, including those corresponding to Group 5 metals, are of great interest in a host of catalytic processes. After the preparation of the first olefin–hydride bis-(cyclopentadienyl) complex of Group 5 metals,  $\text{Cp}_2\text{Nb}(\text{H})(\eta^2\text{-C}_2\text{H}_4)$  ( $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ ) by Tebbe and Parshall [7a], several olefin–hydride complexes of Nb and Ta were synthesized by employing two alternative processes, namely the reactions of  $\text{Cp}_2\text{MCl}_2$  and  $\text{Cp}_2\text{MH}_3$  ( $\text{M} = \text{Nb}, \text{Ta}$ ) with various alkylmagnesium halides and olefins, respectively.

##### 4.4.1. Preparations

The first niobocene and tantalocene olefin–hydride complexes have been prepared by the reaction of  $\text{Cp}_2\text{MH}_3$  with the corresponding olefin. This process takes place through thermal elimination of  $\text{H}_2$  followed by olefinic coordination on the  $\text{Cp}_2\text{MH}$  species Eq. (3).




$\text{M} = \text{Nb}, \text{Ta}$ .  $\text{R} = \text{H}$ ;  $\text{R}' = \text{H}$ . [7a]

$\text{M} = \text{Nb}$ .  $\text{R} = \text{Me}$ ;  $\text{R}' = \text{H}$ , Ph,  $p\text{-Me}_2\text{NC}_6\text{H}_4$ ,  $p\text{-MeOC}_6\text{H}_4$ ,  $p\text{-MeC}_6\text{H}_4$ ,  $p\text{-CF}_3\text{C}_6\text{H}_4$ . [83]

$\text{R}_5 = \text{H}_4\text{SiMe}_3$ ;  $\text{R}' = \text{Ph}$ ,  $p\text{-MeC}_6\text{H}_4$ ,  $p\text{-MeOC}_6\text{H}_4$ . [64, 84]

$\text{M} = \text{Ta}$ .  $\text{R} = \text{H}$ ;  $\text{R}' = \text{Me}$ . [85]

In contrast to the complexes  $(\text{C}_5\text{R}_5)_2\text{Nb}(\text{H})(\text{olefin})$  ( $\text{R}_5 = \text{H}_5$ ;  $\text{H}_4\text{SiMe}_3$ ), in which both *endo* and *exo* isomers are observed, when  $\text{R} = \text{Me}$  the product is obtained in a single isomeric form. Models indicate that the *endo* isomer is sterically favored over the *exo* isomer (Scheme 14).



Scheme 14. Adapted from Ref. [64].

A second, alternative process to obtain these complexes has been reported by Klazinga and Teuben [85]. The reactions of  $\text{Cp}_2\text{MCl}_2$  with Grignard reagents ( $\text{RMgCl}$ ) are reported to afford the complexes  $\text{Cp}_2\text{Nb}(\text{H})(\text{alkene})$ . These reactions presumably proceed via the Nb(III) intermediate ' $\text{Cp}_2\text{NbR}$ ', in which  $\beta$ -H elimination occurs stereospecifically to give the alkene hydrides (Eq. (4)). Two equivalents of the Grignard reagent are required for this reaction. Presumably, one equivalent acts as a reducing agent [ $\text{Nb}(\text{IV}) \rightarrow \text{Nb}(\text{III})$ ] and the other as the source of an alkyl ligand that  $\beta$ -H eliminates to generate the olefin-hydride complex.



$\text{M} = \text{Nb}$ .  $\text{R} = \text{Me}$ ;  $\text{R}' = \text{C}_2\text{H}_5$ ,  $n\text{-C}_3\text{H}_7$ ;  $\text{L} = \text{C}_2\text{H}_4$ ,  $\text{C}_3\text{H}_6$ . [83]


$\text{R}_5 = \text{H}_4\text{SiMe}_3$ ;  $\text{R}' = \text{PhC}_2\text{H}_4$ ;  $\text{L} = \text{styrene}$ . [64]

$\text{M} = \text{Ta}$ .  $\text{R} = \text{H}$ ;  $\text{R}' = n\text{-C}_3\text{H}_7$ ,  $i\text{-C}_3\text{H}_7$ ,  $n\text{-C}_4\text{H}_9$ ,  $s\text{-C}_4\text{H}_9$ ,  $n\text{-C}_5\text{H}_{11}$ . [85]

$\text{L} = \text{C}_3\text{H}_6$ ,  $\text{C}_4\text{H}_8$  (1-butene),  $\text{C}_5\text{H}_{10}$  (1-pentene)

This method leads mainly to the *endo* isomer. In this field there have been recent reports [84] describing an alternative to this synthetic method. In fact, the starting niobium complex  $[\text{Cp}'_2\text{NbCl}]_2$  ( $\text{Cp}' = \text{C}_5\text{H}_4\text{SiMe}_3$ ) reacts with one equivalent of  $\text{RMgX}$  ( $\text{R} = \text{Et}$ ,  $\text{CH}_2\text{CH}_2\text{C}_6\text{H}_5$ ) to generate, at r.t. and in high yields, the corresponding olefin-hydride complexes  $\text{Cp}'_2\text{Nb}(\text{H})(\eta^2\text{-RHC=CH}_2)$  ( $\text{R} = \text{H}$ ,  $\text{C}_6\text{H}_5$ ) (*endo* isomer) by a highly stereoselective  $\beta$ -elimination from the initially formed monoalkyl species  $\text{Cp}'_2\text{Nb}(\text{CH}_2\text{-CH}_2\text{R})$ , which has not been detected. In order to obtain a firm understanding of the  $\beta$ -hydrogen elimination reaction, Doherty and Bercaw [83] carefully examined its microscopic reverse, the migratory insertion that occurs upon reaction of  $\text{Cp}^*_2\text{Nb}(\text{H})(\text{CH}_2=\text{CHR})$  ( $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$ , with  $\text{L} = \text{CO}$  or  $\text{CNMe}$ ) to afford  $\text{Cp}^*_2\text{Nb}(\text{CH}_2=\text{CHR})\text{L}$ . Magnetization transfer experiments demonstrate a reversible migratory insertion  $\rightleftharpoons$   $\beta$ -H elimination equilibrium, although the  $\beta$ -elimination rate constant is orders of magnitude greater than the insertion rate constant. The migratory insertion reactions of the series of *para*-substituted styrene complexes  $\text{Cp}^*_2\text{Nb}(\text{H})(\text{CH}_2=\text{CHR})$  ( $\text{R} = \text{Ph}$ ,  $p\text{-Me}_2\text{NC}_6\text{H}_4$ ,  $p\text{-MeOC}_6\text{H}_4$ ,  $p\text{-MeC}_6\text{H}_4$ ,  $p\text{-CF}_3\text{C}_6\text{H}_4$  [83]) in the presence of  $\text{L}$  were examined in a detailed kinetic and mechanistic study, and were found to be consistent with a four-center transition state in which bond making and breaking are concerted (Scheme 15). A small degree of positive charge builds up on the alkene C- $\beta$  and negative charge on the migrating hydrogen, which is more consistent with its migration as a hydride  $\text{H}^-$ .

Closer examination of  $\text{Cp}^*_2\text{Nb}(\text{H})(\text{exo-CH}_2=\text{CHR})$  using spin-saturation transfer NMR experiments has provided compelling evidence for intermediates with agostic




Scheme 15. Adapted from Ref. [83].


H–C bonding in the migratory insertion  $\rightleftharpoons$   $\beta$ -H elimination reactions [86]. Initially, a mechanism for the exchange of NbH and the CH<sub>2</sub> hydrogens in *exo*-Cp<sub>2</sub><sup>\*</sup>Nb(H)(CH<sub>2</sub>=CHMe) was proposed based upon its rate being two orders of magnitude faster than the exchange of inequivalent Cp rings (Scheme 16) [87]. This mechanism involves a simple insertion followed by inversion about niobium and reinsertion to another *exo*-isomer.

However, in a later report by Green and co-workers [88], a new mechanism was proposed that involves an agostic alkyl ligand in the intermediate Cp<sub>2</sub>Nb(CHMe<sub>2</sub>), which is formed upon insertion (Scheme 17). Thus, the exchange of NbH and the CH<sub>2</sub> hydrogens occurs by in-place rotation of this agostic alkyl intermediate. For the exchange of Cp ligands, the niobium center becomes planar as the alkyl group ‘slides’ across the face of the metal to form a complex (a) with a molecular plane of symmetry in which two agostic interactions (one from each Me group) are now present.

The mechanism of the formation of Cp<sub>2</sub>'NbH(PhCH=CH<sub>2</sub>) (Cp' = C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>) has been discussed [64] in terms of a model that implies a consecutive reaction with a reversible step and a first order dependence with respect to the precursor Cp<sub>2</sub>'NbH<sub>3</sub>, see Scheme 18. A value of 20.6 kcal mol<sup>-1</sup> has been found for Ea. Likewise, studies on the insertion of styrene into the niobium–hydride bond on the mixture of *endo* and *exo* isomers of the aforementioned olefin–hydride niobocene complex were carried out by means of variable-temperature NMR studies using coalescence techniques.



Scheme 16. Adapted from Ref. [87].



Scheme 17. Adapted from Ref. [88].

#### 4.4.2. Reactivity studies

The reactions [89] of  $\text{Cp}'_2\text{Nb}(\text{H})(\eta^2\text{-CH}_2=\text{CH}_2)$  and *endo*- $\text{Cp}'_2\text{Nb}(\text{H})(\eta^2\text{-CH}_2=\text{CHPh})$  with  $\text{CS}_2$  in a 1:1 molar ratio were shown to be dependent on the experimental reaction conditions. Indeed, in an apolar solvent such as hexane or toluene, the alkyl/carbon disulfide complexes  $\text{Cp}'_2\text{Nb}(\text{Et})(\eta^2\text{-CS}_2\text{-C,S})$  and  $\text{Cp}'_2\text{Nb}(\text{CH}_2\text{CH}_2\text{Ph})(\eta^2\text{-CS}_2\text{-C,S})$  were formed exclusively. The carbon disulfide-promoted hydride–olefin insertion is the only process that occurs, and it gives rise exclusively to alkyl complexes. However, in polar solvents such as acetone, the dithioformato derivative  $\text{Cp}'_2\text{Nb}(\eta^2\text{-SC(H)S-S,S})$  and free olefin are generated along with small amounts of alkyl complexes (monitored by  $^1\text{H}$  NMR spectroscopy).


The reaction of niobocene olefin–hydride complexes with  $\text{CO}_2$  and  $\text{CO}$  affords the alkyl–carbon dioxide and alkyl–carbonyl niobocene complexes, respectively (Eq. (5)).



$\text{L} = \text{CO}_2$   $\text{R}_5 = \text{H}_4\text{SiMe}_3$ ;  $\text{R}' = \text{H}$ ,  $\text{C}_6\text{H}_5$ , *p*- $\text{MeC}_6\text{H}_4$ , *p*- $\text{MeOC}_6\text{H}_4$ . [84]

$\text{L} = \text{CO}$   $\text{R}_5 = \text{H}_4\text{SiMe}_3$ ;  $\text{R}' = \text{H}$ ,  $\text{C}_6\text{H}_5$ . [84]

$\text{R}_5 = \text{Me}$ ;  $\text{R}' = \text{C}_6\text{H}_5$ , *p*- $\text{MeNC}_6\text{H}_4$ , *p*- $\text{MeOC}_6\text{H}_4$ , *p*- $\text{MeC}_6\text{H}_4$ , *p*- $\text{CF}_3\text{C}_6\text{H}_4$ . [83]



Scheme 18. Adapted from Ref. [64].


The niobocene olefin–hydride complexes react with a large excess of isocyanide to form alkyl isocyanide derivatives analogous to the CO adducts [83]. The reaction of *endo*-Cp<sub>2</sub>Ta(H)L [L = C<sub>3</sub>H<sub>6</sub>, C<sub>4</sub>H<sub>8</sub> (1-butene)] with aryl and alkyl-isocyanides R'NC [R' = 2,6-(CH<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>, *cyclo*-C<sub>6</sub>H<sub>11</sub>, CH<sub>3</sub> and *t*-C<sub>4</sub>H<sub>9</sub>] gives the thermally stable isocyanide adducts of Cp<sub>2</sub>TaR (R = C<sub>3</sub>H<sub>7</sub>, C<sub>4</sub>H<sub>9</sub>) [90]. The extraordinarily strong metal–isocyanide interaction leads to a nonlinear ‘carbene-like’ structure for these complexes and gives rise to the possibility of reactions at the nitrogen lone-pair of electrons. Complexes of type Cp<sub>2</sub>Nb(H)(CH<sub>2</sub>=CHR) (R = H, CH<sub>3</sub>, C<sub>6</sub>H<sub>5</sub>) react with propyne, 1-butyne, 2-butyne and phenylacetylene to give Cp<sub>2</sub>NbCH<sub>2</sub>CH<sub>2</sub>R(alkyne) [91]. The resulting complexes prefer the *exo* geometry, which is in sharp contrast to the *endo* preference of the starting materials. Bulky 1-alkynes such as 3,3-dimethyl-1-butyne, (trimethylsilyl)acetylene and diphenylacetylene behave in a different manner and gave rise to the formation of hydridoniobocene complexes of the type Cp<sub>2</sub>Nb(H)(alkyne) with *exo* geometry. Cp<sub>2</sub>Ta(H)(alkene) complexes also react with alkynes in a similar manner. A recent report [92] has described the reactions of Cp<sub>2</sub>Ta(H)(C<sub>2</sub>H<sub>4</sub>) with perfluoro-*n*-butyl iodide or perfluorobenzyl iodide, which cleanly yield the iodo(ethylene) complex Cp<sub>2</sub>Ta(C<sub>2</sub>H<sub>4</sub>)I together with the hydrofluorocarbons CF<sub>3</sub>CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>H and C<sub>6</sub>F<sub>5</sub>CF<sub>2</sub>H, respectively. In sharp contrast, reactions of Cp<sub>2</sub>Ta(H)(C<sub>2</sub>H<sub>4</sub>) with perfluoro-*iso*-propyl iodide or pentafluorophenyl iodide cleanly afford the isomeric iodo(ethylidene) complex Cp<sub>2</sub>Ta(CHCH<sub>3</sub>)I.

## 5. Heterobimetallic complexes containing cyclopentadienyl–hydride niobium- and tantalum moieties

The first compound containing a cyclopentadienyl hydride niobium- or tantalum-transition metal bond, whose characteristic structure has been determined, is the niobium complex Cp<sub>2</sub>(CO)Nb(μ-H)Zn(BH<sub>4</sub>)<sub>2</sub> [93]. This compound was prepared by reduction of NbCl<sub>5</sub> with sodium cyclopentadienide and zinc powder in THF under a CO atmosphere at r.t., followed by treatment with NaBH<sub>4</sub>. The geometry of the molecule shows a mirror plane containing the Nb, Zn, H and C (CO) atoms. The plane relates the two C<sub>5</sub>H<sub>5</sub> rings and two BH<sub>4</sub> groups to each other. The large Zn...C(CO) (2.90 Å) and a practically linear structure of the Nb–CO fragment are indicative of the absence of a bond interaction between the carbonyl group and the Zn atom. The Nb–Zn bond (2.829 Å) should be described as a dative interaction Nb → Zn. There is another complex [{Cp<sub>2</sub>NbH<sub>2</sub>}<sub>2</sub>Zn], reported by Tebbe in 1973, obtained as the sole product of the reaction between Cp<sub>2</sub>NbH<sub>3</sub> and ZnEt<sub>2</sub> even when an excess of the latter reagent was employed [7b]. A structure with bridges of hydrogen atoms has been proposed for this compound. Although Cp<sub>2</sub>NbH<sub>3</sub> reacts with ZnEt<sub>2</sub> to give a well-defined product, the reaction of the tantalum analogs resulted only in the formation of a loose adduct. Boersma and coworkers have carried out several reactions between Cp<sub>2</sub>MH<sub>3</sub> (M = Nb, Ta) and Cp<sub>2</sub>Zn, which gives a convenient starting material for the preparation of cyclopentadienylzinc-

transition metal compounds. It reacts smoothly with transition metal hydrides, forming direct zinc–metal bonds, under conditions where dialkylzinc compounds either fail to react or react very slowly [94,95]. When the reaction of  $\text{Cp}_2\text{NbH}_3$  and  $\text{Cp}_2\text{Zn}$  is carried out in benzene, the use of an excess of the hydride niobocene resulted in the formation of the known compound reported by Tebbe, while with an excess of the Zn reagent,  $[\text{Cp}_2\text{NbH}\{\text{ZnCp}\}_2]$  was formed. In THF as the solvent,  $[\text{Cp}_2\text{NbH}_2\text{ZnCp}]$  was isolated as the sole product, whether an equimolar amount or an excess of  $\text{Cp}_2\text{Zn}$  was used. An X-ray diffraction study was carried out for this compound. The molecule resembles that of  $\text{Cp}_2\text{NbH}_3$  with the central hydrogen atom replaced by a  $\text{CpZn}$  group [94]. A similar structure has been assigned to the tantalum analogs [95]. Meanwhile, Tebbe obtained  $[\{\text{Cp}_2\text{NbH}_2\}\text{Zn}]$  as the sole reaction product, and evidence that the reactivity of  $\text{Cp}_2\text{Zn}$  is greater than that of  $\text{ZnEt}_2$  appears from the fact that it is able to effect the substitution of a second hydrogen atom to yield  $[\text{Cp}_2\text{NbH}\{\text{ZnCp}\}_2]$ . This formula was corroborated by the isolation and structural characterization of the tantalum analog, which was isolated in the *transoid* and *cisoid* isomeric forms. The *transoid* isomer was converted irreversibly and completely into the *cisoid* isomer when a benzene solution was heated at 50°C for 3 h [95]. Pasynskii and coworkers have also prepared  $[\{\text{Cp}_2(\text{CO})\text{Nb}(\mu\text{-H})_2\}_2\text{M}(\text{CO})_4]$  ( $\text{M} = \text{Mo}, \text{W}$ ) by the reaction of  $\text{Cp}_2\text{Nb}(\text{CO})\text{H}$  and  $\text{Ni}(\text{CO})_4$  and  $\text{M}(\text{CO})_6$  ( $\text{M} = \text{Mo}, \text{W}$ , respectively). In the molecular structures of these complexes the large  $\text{Nb}\dots\text{Ni}$  [3.218(2) Å],  $\text{Nb}\dots\text{Mo}$  [3.549(2) Å] and  $\text{Nb}\dots\text{W}$  [3.565(2) Å] distances show the weakening or even the absence of a direct niobium–transition metal bond interaction [96]. On the other hand,  $\text{Fe}(\text{CO})_5$  reacts rapidly with  $\text{Cp}_2\text{NbH}_3$  even at r.t., to give the complex  $[\text{Cp}_2(\text{CO})\text{Nb}(\mu\text{-H})\text{M}(\text{CO})_3]$  [97]. When a mixture in THF of  $\text{Cp}_2\text{NbH}_3$  and  $\text{M}(\text{CO})_6$  ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ) is irradiated using a mercury lamp for 4 h, the complexes  $[\text{Cp}_2(\text{CO})\text{Nb}(\mu\text{-H})\text{M}(\text{CO})_5]$  ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ) were formed [98]. The X-ray crystal study for the Cr derivative shows a bridging hydride between both metal centers, but the Nb–Cr distance [3.452(2) Å] indicates the absence of a direct metal–metal bond. These compounds can be viewed as containing three-center, two-electron Nb–H–M bonds, with the 18-electron moieties  $\text{Cp}_2(\text{CO})\text{NbH}$  effectively acting as a two-electron donor to the coordinatively unsaturated  $\text{M}(\text{CO})_5$  fragment. The tantalum analogs have also been reported by the reaction of  $\text{Cp}_2\text{TaH}(\text{CO})$ , as a two-electron donor, with the Lewis acid organometallic fragments  $\text{M}(\text{CO})_5$  ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ) or  $\text{CpMn}(\text{CO})_2$  [99]. Also, when a mixture in toluene of  $\text{Cp}_2\text{NbH}_3$  and  $[\text{Mn}_2(\text{CO})_{10}]$  is irradiated for 50 min. using a fluorescent lamp ( $\lambda_{\text{max}} = 3600$  Å), the ionic compound  $[\{\text{Cp}_2(\text{CO})\text{Nb}\}_2(\mu\text{-H})]^+ [\text{Mn}(\text{CO})_5]^-$  was obtained. The cation should be described as the result of the protonation reaction on the dimer of two 17-electron fragments  $[\text{Cp}_2(\text{CO})\text{Nb}\text{--}\text{Nb}(\text{CO})\text{Cp}_2]$  [98,100]. In contrast, when the same reaction is carried out with  $\text{Cp}_2\text{TaH}_3$ , the neutral complex  $[\text{Cp}_2(\text{CO})\text{Ta}(\mu\text{-H})\text{Mn}_2(\text{CO})_9]$  is isolated. The details about the sequence leading to one compound or the other are still unclear [98]. There is another cyclopentadienyl hydride niobium compound exhibiting an Nb–Nb bond whose crystal structure was determined by Guggenberger [101]. The complex was described as the result of the reduction reaction of  $\text{Cp}_2\text{NbCl}_2$  with Na amalgam in THF. The compound  $\text{Cp}_2\text{NbH}_3$  reacts with potassium hydride for





Scheme 19. Adapted from Ref. [105].

3 days in THF solution at r.t. to give an orange precipitate of  $[\text{K}][\text{Cp}_2\text{NbH}_2]$  and this compound is then treated with a suspension of ytterbium diiodide in 1,2-dimethoxyethane, which must contain a trace of diglyme, at r.t. for 12 h to yield the compound  $[\{\text{NbCp}_2\text{H}_2\}_2\text{Yb}\cdot\text{diglyme}]$  [102,103]. The X-ray crystal structure determination for this compound shows a central Nb–Yb–Nb core, with an Nb–Yb distance that lies between the sum of the Slater atomic radii and the sum of the metallic radii. In addition, there are two bridging hydrogens located symmetrically on each side of the Nb–Yb vector and these bridging hydrogens and the Nb and Yb atoms lie in a plane. Each Nb bears two Cp groups that are bent away from the Yb. The complete molecule has two-fold rotational symmetry about the Yb–O(CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>)<sub>2</sub> bond. Fajardo and coworkers reported, in a preliminary communication [104], the synthesis and structural characterization of the mixed metal cluster cation  $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{R})_2(\text{AuPPh}_3)_2]^+$  (R = H, SiMe<sub>3</sub>) as the PF<sub>6</sub><sup>−</sup> or ClO<sub>4</sub><sup>−</sup> salts. In that communication they report the first examples of triangular ‘NbAu<sub>2</sub>’ metal frameworks with X-ray evidence for the metal–metal bonds. The complexes represented a new class of heteronuclear niobium–gold clusters. The metal framework, located on the reflection plane of the two cyclopentadienyl groups, consists of a triangle with rather similar and large [2.9139(8) Å and 2.9098(8) Å] Nb–Au bond distances. In the X-ray diffraction study the two hydride ligands were not located in the final structure. Several years later, the cationic compound was reported as a model of a protonation reaction with  $[\text{Au}(\text{PPh}_3)]^+$  on  $\text{Cp}'_2\text{NbH}_3$  and in this case the hydrides were found [105]. Whatever the stoichiometry {molar ratio  $[\text{Au}(\text{PPh}_3)]^+:\text{complex}$ , 1:1 or 2:1} or reaction conditions employed only the compound  $[\{\text{Cp}_2\text{Nb}(\mu\text{-H})_2\{\text{Au}(\text{PPh}_3)_2\}]^+$  was obtained. The different steps involved in the formation of this cationic complex are proposed in Scheme 19.

It has also been reported [105] that when  $[\text{Au}\{\text{N}(\text{SiMe}_3)_2\}(\text{PPh}_3)]$  was used as an alternative gold reagent in order to avoid the final protonation process, the reaction with  $\text{Cp}'_2\text{NbH}_3$ , in a hydrocarbon solvent and at r.t. yielded a very air-sensitive product that was characterized as the raft cluster  $[\text{Au}_3\text{Nb}_3(\mu\text{-H})_6\{\eta^5\text{-C}_5\text{H}_4(\text{SiMe}_3)\}_6]$ . The structure contains a gold triangle surrounded by three  $\text{Cp}'_2\text{Nb}$  groups and six bridging hydrides. MO calculations indicate a striking electronic stabilization of this geometry [22].

The cationic complexes  $[\{\text{Nb}(\eta^5\text{-C}_5\text{H}_3\text{RR}')_2\text{H}_3\}_2\text{Au}]^+$  (R = R' = H; R = H, R' = SiMe<sub>3</sub>; R = R' = SiMe<sub>3</sub>) were synthesized, as mentioned previously, upon addition of  $[\text{Au}(\text{THT})][\text{PF}_6]$ , prepared in situ from  $[\text{AuCl}(\text{THT})]$  and TIPF<sub>6</sub>, to  $[\text{Nb}(\eta^5\text{-C}_5\text{H}_3\text{RR}')_2\text{H}_3]$  in THF [21b]. The crystal structure of one of these cationic complexes showed that the metal core consists of an almost linear arrangement of Nb, Au and Nb atoms [Nb–Au–Nb angle 175.92(2)°]. The coordination around the Au

atom is intermediate between tetrahedral and square planar. This geometry and the locations of the hydrides suggest a significant narrowing of the distance between one of the bridging hydrides and the terminal [17]. On the other hand, the reaction of  $\text{Cp}_2\text{NbHL}$  ( $\text{L} = \text{P}(\text{OMe})_3$ ,  $\text{P}(\text{OEt})_3$ ,  $\text{P}(\text{OPh})_3$ ) with  $[\text{Au}(\text{PPh}_3)][\text{PF}_6]$ , obtained in situ from  $[\text{Au}(\text{PPh}_3)\text{Cl}]$  and  $\text{TIPF}_4$ , in THF solution and at r.t. leads to the mixed-metal Nb–Au complexes  $[\text{Cp}_2\text{L}(\mu\text{-H})(\text{AuPPh}_3)][\text{PF}_6]$ , where the niobium–hydrogen–gold interaction can be considered to occur through a three-center two electron bond. The authors proposed an additional niobium–gold interaction [49]. The cationic complexes  $[\{\text{Cp}_2\text{Nb}(\text{CO})(\mu\text{-H})\}_2\text{M}]^+$  ( $\text{M} = \text{Cu}, \text{Ag}, \text{Au}$ ) are the result of the reaction of  $\text{Cp}_2\text{Nb}(\text{CO})\text{H}$  with  $[\text{Cu}(\text{MeCN})_4][\text{BF}_4]$ ,  $[\text{Cu}(\text{PPh}_3)\text{Cl}]/\text{TIPF}_6$ ,  $\text{AgBF}_4$ ,  $[\text{Ag}(\text{PPh}_3)\text{Cl}]/\text{TIPF}_6$  or  $[\text{Au}(\text{THT})\text{Cl}]/\text{TIPF}_6$ . In these complexes, the coinage cation is only linked to the hydride of each niobium center [23]. The most important result of this study is that the formation of the adducts essentially involves binding of the hydride of a niobium moiety to the Lewis acidic cation  $\text{M}(\text{I})$ . In addition, a weak interaction between the Lewis acid and CO is present, at least in the case of copper, with a pseudotetrahedral geometry around the copper. Nevertheless, the same reaction with  $[\text{AuPPh}_3\text{Cl}]/\text{TIPF}_6$  leads to the complex  $[\{\text{NbCp}_2(\text{CO})\}(\mu\text{-H})(\text{AuPPh}_3)][\text{PF}_6]$ , and this cationic complex contains one niobium atom linked to a ‘ $(\text{AuPPh}_3)$ ’ fragment. The nature of this bonding and the geometry around the gold atom are, however, unknown [23].

## Acknowledgements

The authors gratefully acknowledge financial support from the Dirección General de Enseñanza Superior e Investigación Científica (DGES) (Grant No. PB 95-0023-CO1-CO2) of Spain.

## References

- [1] (a) G.G. Hlatky, R.H. Crabtree, *Coord. Chem. Rev.* 65 (1985) 1. (b) B.R. James, *Homogeneous Hydrogenation*, Wiley, New York, 1973.
- [2] A recent review: E.W. Abel, G. Wilkinson, F.G.A. Stone (Eds.), *Comprehensive Organometallic Chemistry*, vol. 5, Pergamon Press, Oxford, 1995, p. 76.
- [3] (a) G.W. Parshall, *Homogeneous Catalysis*, Wiley, New York, 1980. (b) J.P. Collman, L.S. Hegedus, J.R. Norton, R.G. Finke, *Principles and Applications of Organotransition Metal Chemistry*, 2nd ed., University Science Books, Mill Valley, CA, 1987.
- [4] (a) G.J. Kubas, R.R. Ryan, B.I. Swanson, P.J. Vergamini, H.J. Wasserman, *J. Am. Chem. Soc.* 106 (1984) 451. (b) G.J. Kubas, *Acc. Chem. Res.* 21 (1988) 120. (c) G. Rattan, G.J. Kubas, C.J. Unkefer, L.S. Van Der Sluys, K.A. Kubat-Martin, *J. Am. Chem. Soc.* 112 (1990) 3855.
- [5] Recent reviews: (a) P.G. Jessop, R.H. Morris, *Coord. Chem. Rev.* 121 (1992) 155. (b) D.M. Heinekey, W.J. Oldman, *Chem. Rev.* 93 (1993) 913. (c) R.H. Crabtree, *Angew. Chem. Int. Ed. Engl.* 32 (1993) 789.
- [6] A. Antiñolo, B. Chaudret, G. Commenges, M. Fajardo, F. Jalón, R.H. Morris, A. Otero, C.T. Schweitzer, *J. Chem. Soc. Chem. Commun.* (1988) 1210.

- [7] (a) F.N. Tebbe, G. W. Parshall, *J. Am. Chem. Soc.* 93, (1971) 3793. (b) F.N. Tebbe, *J. Am. Chem. Soc.* 95 (1973) 5412.
- [8] J.A. Labinger, in: G. Wilkinson, F.G.A. Stone (Eds.), *Comprehensive Organometallic Chemistry*, vol. 3, Pergamon Press, Oxford, 1982, p. 770.
- [9] M.D. Curtis, L.G. Bell, W.M. Butler, *Organometallics* 4 (1985) 701.
- [10] (a) T. Arliguie, B. Chaudret, J. Devillers, R. Poilblanc, *C.R. Hebd. Séances Acad. Sci.* 305-II (1987) 1523. (b) B. Chaudret, G. Commenges, F. Jalón, A. Otero, *J. Chem. Soc. Chem. Commun.* (1989) 210. (c) T. Arliguie, C. Border, B. Chaudret, J. Devillers, R. Poilblanc, *Organometallics* 8 (1989) 1308. (d) T. Arliguie, B. Chaudret, F. Jalón, A. Otero, J.A. López, F.J. Lahoz, *Organometallics* 10 (1991) 1888. (e) D.M. Heinekey, T.G.P. Harper, *Organometallics* 10 (1991) 2891.
- [11] (a) D.M. Heinekey, N.G. Payne, G.K. Shulte, *J. Am. Chem. Soc.* 110 (1986) 2303. (b) D.M. Heinekey, J.M. Millar, T.F. Koetzle, N.G. Paine, K.W. Zilm, *J. Am. Chem. Soc.* 112 (1990) 909.
- [12] D.M. Heinekey, *J. Am. Chem. Soc.* 113 (1991) 6074.
- [13] D.G. Gusev, R. Kuhlman, G. Sini, O. Eisenstein, K.G. Caulton, *J. Am. Chem. Soc.* 116 (1994) 2685.
- [14] B. Chaudret, H.H. Limbach, C. Moise, *C.R. Hebd. Séances Acad. Sci.* 315-II (1992) 533.
- [15] (a) K.W. Zilm, D.M. Heinekey, J.M. Millar, N.G. Payne, P. Demou, *J. Am. Chem. Soc.* 111 (1989) 3088. (b) K.W. Zilm, D.M. Heinekey, J.M. Millar, N.G. Payne, S.P. Neshyba, J.C. Duchamp, J. Szczyrba, *J. Am. Chem. Soc.* 112 (1990) 92.
- [16] D. Jones, J.A. Labinger, D.P. Weitekamp, *J. Am. Chem. Soc.* 111 (1989) 3087.
- [17] A. Antiñolo, F. Carrillo-Hermosilla, B. Chaudret, M. Fajardo, J. Fernández-Baeza, M. Lanfranchi, H.-H. Limbach, M. Maurer, A. Otero, M.A. Pellinghelli, *Inorg. Chem.* 35 (1996) 7873.
- [18] H.-H. Limbach, M. Maurer, G. Scherer, B. Chaudret, *Angew. Chem.* 31 (1992) 1369.
- [19] J.C. Barthelat, B. Chaudret, J.P. Daudey, P. De Loth, R. Poilblanc, *J. Am. Chem. Soc.* 113 (1991) 9896.
- [20] S. Camanyes, F. Maseras, M. Moreno, A. Lledós, J.M. Lluch, *J. Am. Chem. Soc.* 118 (1996) 4617.
- [21] (a) A. Antiñolo, F. Carrillo, J. Fernández-Baeza, A. Otero, M. Fajardo, B. Chaudret, *Inorg. Chem.* 31 (1992) 5156. (b) A. Antiñolo, F. Carrillo, B. Chaudret, M. Fajardo, J. Fernández-Baeza, M. Lanfranchi, H.-H. Limbach, M. Maurer, A. Otero, M.A. Pellinghelli, *Inorg. Chem.* 33 (1994) 5163.
- [22] A. Antiñolo, J.K. Burdett, B. Chaudret, O. Eisenstein, M. Fajardo, F. Jalón, F. Lahoz, J.A. López, A. Otero, *J. Chem. Soc. Chem. Commun.* (1990) 17.
- [23] A. Antiñolo, F. Carrillo, B. Chaudret, M. Fajardo, S. García-Yuste, F.J. Lahoz, M. Lanfranchi, J.A. López, A. Otero, M.A. Pellinghelli, *Organometallics* 14 (1995) 1297.
- [24] S. Camanyes, F. Maseras, M. Moreno, A. Lledós, J.M. Lluch, J. Bertran, *Angew. Chem. Int. Ed. Engl.* 36 (1997) 265.
- [25] S. Camanyes, F. Maseras, M. Moreno, A. Lledós, J.M. Lluch, J. Bertran, *Inorg. Chem.* 37 (1998) 2334.
- [26] U. Drexler, R. Wiedenbruch, C. Scheurer, R. Meyer, R.R. Ernst, S. Chaloupka, L.M. Venanzi, *Mol. Phys.* 93 (1998) 471.
- [27] (a) R.H. Crabtree, *Chem. Rev.* 95 (1995) 987. (b) R.H. Crabtree, *Chem. Rev.* 85 (1985) 245. (c) B.A. Arndtsen, R.G. Bergman, T.A. Mobley, T.H. Peterson, *Acc. Chem. Res.* 28 (1995) 154. (d) E. Shilov, *Activation of Saturated Hydrocarbons by Transition Metal Complexes*, D. Reidel, Dordrecht, 1984.
- [28] G.W. Parshall, *Acc. Chem. Res.* 8 (1975) 113.
- [29] A. Antiñolo, F. Carrillo, M. Fajardo, A. Otero, M. Lanfranchi, M.A. Pellinghelli, *Organometallics* 14 (1995) 1518.
- [30] Q. Jiang, P.J. Carroll, D.H. Berry, *Organometallics* 10 (1991) 3648.
- [31] D.H. Berry, Q. Jiang, *J. Am. Chem. Soc.* 109 (1987) 6210.
- [32] A. Antiñolo, F. Carrillo-Hermosilla, A. Castel, M. Fajardo, J. Fernández-Baeza, M. Lanfranchi, A. Otero, M.A. Pellinghelli, G. Rima, J. Satgé, E. Villaseñor, *Organometallics* 17 (1998) 1523.
- [33] D.H. Harris, S.A. Keppie, M.F. Lappert, *J. Chem. Soc. Dalton Trans.* (1973) 1653.

- [34] (a) M.L.H. Green, A.K. Hughes, P. Mountford, *J. Chem. Soc. Dalton Trans.* (1991) 1407. (b) M.L.H. Green, A.K. Hughes, P. Mountford, *J. Chem. Soc. Dalton Trans.* (1991) 1699.
- [35] K. Colomer, R.J.P. Corriu, A. Vioux, *Inorg. Chem.* 21 (1982) 368.
- [36] (a) R.J.P. Corriu, *J. Organomet. Chem.* 400 (1990) 81. (b) M.-F. Fan, Z. Lin, *Organometallics* 17 (1998) 1092.
- [37] H.-G. Woo, W.P. Freeman, T.D. Tilley, *Organometallics* 11 (1992) 2198.
- [38] (a) G.I. Nikonov, E.V. Antomono, W. Massa, *Chem. Ber.* 130 (1997) 1629. (b) G.I. Nikonov, L.G. Kuzmina, P. Mountford, D.A. Lemenovskii, *Organometallics* 14 (1995) 3588. (c) G.I. Nikonov, D.A. Lemenovskii, J. Lorberth, *Organometallics* 13 (1994) 3127. (d) G.I. Nikonov, A.J. Blake, J. Lorberth, D.A. Lemenovskii, S. Wocadlo, *J. Organomet. Chem.* 547 (1997) 235.
- [39] (a) F. Carrillo-Hermosilla, F.A. Jalón, A. Otero, E. Villaseñor, *Anales de Química, Int. Ed.* 92 (1996) 339. (b) N. Etkin, M.T. Benson, S. Courtenay, M.J. McGlinchey, A.D. Bain, D.W. Stephan, *Organometallics* 16 (1997) 3504.
- [40] J.F. Hartwig, S.R. de Gala, *J. Am. Chem. Soc.* 116 (1994) 3661.
- [41] D.R. Lantero, D.L. Ward, M.R. Smith, *J. Am. Chem. Soc.* 119 (1997) 9699.
- [42] D.R. Lantero, D.H. Motry, D.L. Ward, M.R. Smith, *J. Am. Chem. Soc.* 116 (1994) 10811.
- [43] E.K. Barefield, G.W. Parshall, F.N. Tebbe, *J. Am. Chem. Soc.* 92 (1970) 5235.
- [44] J.F. Leboeuf, O. Lavastre, J.C. Leblanc, C. Moise, *J. Organomet. Chem.* 418 (1991) 359.
- [45] J.C. Leblanc, C. Moise, *J. Organomet. Chem.* 364 (1989) C3.
- [46] J.F. Leboeuf, J.C. Leblanc, C. Moise, *J. Organomet. Chem.* 364 (1989) C22–C24.
- [47] C.R. Lucas, M.L.H. Green, *J. Chem. Soc. Chem. Commun.* (1972) 1005.
- [48] A. Antiñolo, M. Fajardo, C. López-Mardomingo, A. Otero, C. Sanz-Bernabé, *J. Organomet. Chem.* 369 (1989) 187.
- [49] A. Antiñolo, F. Carrillo, M. Fajardo, S. García-Yuste, A. Otero, *J. Organomet. Chem.* 482 (1994) 93.
- [50] G.E. Herberich, I. Savupoulos, *J. Organomet. Chem.* 362 (1989) 345.
- [51] P.P. Deutsch, J.A. Maguire, W.D. Jones, R. Eisenberg, *Inorg. Chem.* 29 (1990) 35.
- [52] G.E. Herberich, U. Englert, H. Hoeveler, I. Savupoulos, *J. Organomet. Chem.* 399 (1990) 35.
- [53] J.A. Labinger, J. Schwartz, *J. Am. Chem. Soc.* 97 (1975) 1596.
- [54] G.E. Herberich, H. Mayer, *Organometallics* 9 (1990) 2655.
- [55] J. Sala-Pala, J. Amaudrut, J.E. Guerchais, R. Mercier, J. Douglade, J.G. Theobald, *J. Organomet. Chem.* 204 (1981) 347.
- [56] A. Antiñolo, F. Carrillo-Hermosilla, M. Fajardo, S. García-Yuste, A. Otero, S. Camanyes, F. Maseras, M. Moreno, A. Lledós, J.M. Lluch, *J. Am. Chem. Soc.* 119 (1997) 6107.
- [57] (a) J.R. Reynoud, J.C. Leblanc, C. Moise, *Organometallics* 4 (1985) 1059. (b) J.C. Leblanc, J.R. Reynoud, C. Moise, *C.R. Acad. Sci. Ser.* 2 295 (1982) 755. (c) C. Moise, J.R. Reynoud, J.C. Leblanc, R. Broussier, *J. Organomet. Chem.* 240 (1982) C15.
- [58] M.L.H. Green, B. Jousseume, *J. Organomet. Chem.* 193 (1980) 339.
- [59] R.A. Bell, S.A. Cohen, N.M. Doherty, R.S. Threnkel, J.E. Bercaw, *Organometallics* 1 (1982) 972.
- [60] C. McDade, V.C. Gibson, B.D. Santarsiero, J.E. Bercaw, *Organometallics* 7 (1988) 1.
- [61] V.C. Gibson, J.E. Bercaw, W.J. Burton, R.D. Sanner, *Organometallics* 5 (1986) 977.
- [62] A. Castro, M. Gómez, P. Gómez-Sal, A. Manzanero, P. Royo, *J. Organomet. Chem.* 518 (1996) 37.
- [63] See, for example: P.T. Wolczanski, J.E. Bercaw, *J. Am. Chem. Soc.* 101 (1979) 6450.
- [64] A. Antiñolo, F. Carrillo, S. García-Yuste, A. Otero, *Organometallics* 13 (1994) 2761.
- [65] D.F. Foust, R.D. Regers, M.D. Rausch, J.L. Atwood, *J. Am. Chem. Soc.* 102 (1982) 5446.
- [66] C.R. Lucas, *Inorg. Synth.* 16 (1976) 107.
- [67] J.F. Leboeuf, J.C. Leblanc, C. Moise, *J. Organomet. Chem.* 355 (1987) 331.
- [68] (a) G.E. Herberich, H. Mayer, *J. Organomet. Chem.* 322 (1987) C29. (b) V.C. Gibson, T.P. Kee, *J. Organomet. Chem.* 471 (1994) 105.
- [69] A. Antiñolo, F. Carrillo-Hermosilla, S. García-Yuste, M. Freitas, A. Otero, S. Prashar, E. Villaseñor, M. Fajardo, *Inorg. Chim. Acta* 259 (1997) 101.
- [70] (a) D.J. Darensbourg, C. Ovalles, *J. Am. Chem. Soc.* 109 (1987) 330. (b) E. Graf, W. Leitner, *J. Chem. Soc. Chem. Commun.* (1992) 623. (c) J. Tsai, K.M. Nicholas, *J. Am. Chem. Soc.* 114 (1992) 5117.

- [71] A. Antiñolo, M. Fajardo, S. García-Yuste, I. del Hierro, A. Otero, S. Elkrani, Y. Mourad, Y. Mugnier, *J. Chem. Soc. Dalton Trans.* (1995) 3409.
- [72] P.F. Fu, A.K. Fazlur-Rahman, K.M. Nicholas, *Organometallics* 12 (1994) 413.
- [73] Some representative examples: (a) H.C. Clark, W.S. Tsang, *J. Am. Chem. Soc.* 89 (1967) 533. (b) J.L. Davison, M. Green, F. Gordon, A. Stone, A.J. Welch, *J. Chem. Soc. Chem. Commun.* (1975) 286. (c) S. Otsuka, A. Nakamura, *Adv. Organomet. Chem.* 14 (1976) 245. (d) J.P. William, A. Wojcicki, *Inorg. Chem.* 16 (1977) 3116. (e) F.Y. Pétillon, F. de Floch-Pérennou, J.E. Guerchais, D.W.A. Sharp, L.J. Manojlovic-Muir, K.W. Muir, *J. Organomet. Chem.* 202 (1980) 23. (f) H. Scordia, R. Kergoat, M.M. Kubicki, J.E. Guerchais, *J. Organomet. Chem.* 249 (1983) 371. (g) F.Y. Pétillon, J.L. de Quéré, F. de Floch-Pérennou, J.E. Guerchais, M.B. Gomes de Lima, L.J. Manojlovic-Muir, K.W. Muir, D.W.A. Sharp, *J. Organomet. Chem.* 255 (1984) 231. (h) J. Amaudrut, J.C. Leblanc, C. Moise, J. Sala-Pala, *J. Organomet. Chem.* 295 (1985) 167. (i) H.C. Clark, G. Ferguson, A.B. Goel, E.G. Janzen, H. Ruegger, P.Y. Siew, C.S. Wong, *J. Am. Chem. Soc.* 108 (1986) 6961. (j) M. Cariou, M. Etienne, J.E. Guerchais, R. Kergoat, M.M. Kubicki, *J. Organomet. Chem.* 327 (1987) 393. (k) W.D. Jones, V.L. Chandler, E.J. Feher, *Organometallics* 9 (1990) 164.
- [74] G.E. Herberich, H. Mayer, *J. Organomet. Chem.* 347 (1988) 93.
- [75] (a) G.E. Herberich, W. Barlage, *Organometallics* 6 (1987) 1924. (b) G.E. Herberich, W. Barlage, *J. Organomet. Chem.* 331 (1987) 63.
- [76] A. Antiñolo, F. Carrillo-Hermosilla, M. Fajardo, S. García-Yuste, M. Lanfranchi, A. Otero, M.A. Pellinghelli, S. Prashar, E. Villaseñor, *Organometallics* 15 (1996) 5507.
- [77] R.S. Threlkel, J.E. Bercaw, *J. Am. Chem. Soc.* 103 (1981) 2650.
- [78] (a) L. Roullier, D. Lucas, Y. Mugnier, A. Antiñolo, M. Fajardo, A. Otero, *J. Organomet. Chem.* 412 (1991) 353. (b) J.F. Reynoud, J.F. Leboeuf, J.C. Leblanc, C. Moise, *Organometallics* 5 (1986) 1863. (c) F. Jalón, A. Otero, B. Manzano, E. Villaseñor, B. Chaudret, *J. Am. Chem. Soc.* 117 (1995) 10123. (d) J.F. Leboeuf, O. Lavastre, J.C. Leblanc, C. Moise, *J. Organomet. Chem.* 418 (1991) 359.
- [79] R.H. Crabtree, M. Lavin, *J. Chem. Soc. Chem. Commun.* (1985) 1661.
- [80] S. Sabo-Etienne, B. Chaudret, H.A. al Makarim, J.C. Barthelat, J.P. Daudey, C. Moise, J.C. Leblanc, *J. Am. Chem. Soc.* 116 (1994) 9335.
- [81] P.J. Desrosiers, L. Cai, Z. Lin, R. Richards, J. Halpern, *J. Am. Chem. Soc.* 113 (1991) 4173.
- [82] S. Sabo-Etienne, B. Chaudret, H.A. al Makarim, J.P. Daudey, S. Ulrich, H.H. Limbach, C. Moise, *J. Am. Chem. Soc.* 117 (1995) 11602.
- [83] N.M. Doherty, J.E. Bercaw, *J. Am. Chem. Soc.* 107 (1985) 2670.
- [84] A. Antiñolo, F. Carrillo-Hermosilla, I. del Hierro, A. Otero, *Organometallics* 16 (1997) 4161.
- [85] A.H. Klazinga, J.H. Teuben, *J. Organomet. Chem.* 157 (1978) 413.
- [86] J.E. Bercaw, B. Burger, M.H.L. Green, *J. Chem. Soc. Chem. Commun.* (1989) 734.
- [87] B.J. Burger, B.D. Santarsiero, M.S. Trimmer, J.E. Bercaw, *J. Am. Chem. Soc.* 110 (1988) 3134.
- [88] M.H.L. Green, A. Sella, L.-L. Wong, *Organometallics* 11 (1992) 2650.
- [89] A. Antiñolo, I. del Hierro, M. Fajardo, S. García-Yuste, S. Otero, O. Blacque, M.M. Kubicki, J. Amaudrut, *Organometallics* 15 (1996) 1966.
- [90] A.H. Klazinga, J.H. Teuben, *J. Organomet. Chem.* 192 (1980) 75.
- [91] H. Yasuda, H. Yamamoto, A. Takashi, A. Nakamura, J.Ch.Y. Kai, N. Kasai, *Organometallics* 10 (1991) 4058.
- [92] R.P. Hughes, S.M. Maddock, A.L. Rheingold, I.A. Guzei, *Polyhedron* 17 (1998) 1037.
- [93] M.A. Porai-Koshits, A.S. Antsyshkina, A.A. Pasynskii, G.G. Sadikow, Yu.V. Skripkin, V.N. Ostrikova, *Inorg. Chim. Acta* 34 (1979) 285.
- [94] P.H.M. Budzelaar, K.H. den Haan, J. Boersma, G.J.M. van der Kerk, A.L. Spek, *Organometallics* 2 (1994) 156.
- [95] P.H.M. Budzelaar, A.A.H. van der Zeijden, J. Boersma, G.J.M. van der Kerk, A.L. Spek, A.J.M. Duisenberg, *Organometallics* 3 (1984) 159.
- [96] Yu.V. Skripkin, A.A. Pasynskii, V.T. Kalinnikov, M.A. Porai-Koshits, L.K. Minacheva, A.S. Antsyshkina, V.N. Ostrikova, *J. Organomet. Chem.* 231 (1982) 205.
- [97] J.A. Labinger, K.S. Wang, W.R. Scheidt, *J. Am. Chem. Soc.* 100 (1978) 3254.

- [98] B. Balbach, S. Baral, H. Biersack, W.A. Herrmann, J.A. Labinger, W.R. Schidt, F.J Timmers, M.L. Ziegler, *Organometallics* 7 (1988) 325.
- [99] J.C. Leblanc, J.R. Reynoud, C. Moise, J. Organomet. Chem. 244 (1983) C26.
- [100] S. Baral, J.A. Labinger, W.R. Scheidt, F.J. Timmers, J. Organomet. Chem. 215 (1981) C53.
- [101] L.J. Guggenberger, *Inorg. Chem.* 12 (1973) 294.
- [102] M.L.H. Green, A.K. Hughes, D.M. Michaelidou, P. Moutford, J. Chem. Soc. Chem. Commun. (1993) 591.
- [103] D.M. Michaelidou, M.L.H. Green, A.K. Hughes, P. Mountford, A.N. Chernega, *Polyhedron* 14 (1995) 2663.
- [104] M. Fajardo, M.P. Gómez-Sal, P. Royo, S. Martínez-Carrera, S. García-Blanco, J. Organomet. Chem. 312 (1986) C44.
- [105] A. Antiñolo, F.A. Jalón, A. Otero, M. Fajardo, B. Chaudret, F. Lahoz, J.A. López, J. Chem. Soc. Dalton Trans. (1991) 1861.