

## HIGHER OXIDATION STATE CHEMISTRY OF MANGANESE

W LEVASON and C A. McAULIFFE

*Department of Chemistry, University of Manchester Institute of Science and Technology, Manchester 1 (Great Britain)*

(Received March 30th, 1971)

### CONTENTS

A Introduction	353
B. Manganese (VII)	353
C. Manganese (VI)	356
D Manganese (V)	358
E Electronic spectra and structure of the tetraoxoanions	359
F Manganese (IV)	359
G. Manganese (III)	367
H Spectra of manganese (III) compounds	377
References	378

### A. INTRODUCTION

In the higher oxidation states, manganese shows some resemblance to technetium and rhenium, and a more formal resemblance to chlorine, especially in the 7+ state. Manganese chemistry is dominated by the very stable manganese (II) cation ( $3d^5$ ), which is in marked contrast to technetium and rhenium where there is little evidence for any simple cations.

Apart from a review by Sidgwick over twenty years ago<sup>1</sup> and a recent review of the solution chemistry of manganese (III)<sup>2</sup>, no comprehensive collation of the higher oxidation state chemistry of this important metal has appeared.

Reference to Table 1 illustrates the multiplicity of oxidation states in which manganese exists, and it is the purpose of this review to describe the chemistry of  $Mn^{III}$ ,  $Mn^{IV}$ ,  $Mn^V$ ,  $Mn^{VI}$ , and  $Mn^{VII}$ .

### B MANGANESE (VII)

Manganese (VII) is found only in compounds containing oxide ligands. All  $Mn^{VII}$  compounds are very strongly oxidising, and with the exception of the permanganates, are thermally unstable, decomposing, often explosively, at or below room temperature.

TABLE 1

Redox potentials for manganese<sup>3</sup>

Reaction	Redox potential (v)
$\text{MnO}_2(\text{s}) + 4 \text{H}^+(\text{aq}) + 2 \text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 2 \text{H}_2\text{O}$	1.239
$\text{MnO}_2(\text{s}) + 4 \text{OH}^-(\text{aq}) \rightarrow \text{MnO}_4^{2-}(\text{aq}) + 2 \text{H}_2\text{O} + 3 \text{e}^-$	-0.588
$\text{MnO}_4^{3-}(\text{aq}) \rightarrow \text{MnO}_4^{2-}(\text{aq}) + \text{e}^-$	-0.3
$\text{Mn}^{2+}(\text{aq}) \rightarrow \text{Mn}^{3+}(\text{aq}) + \text{e}^-$	-1.5
$\text{Mn}^{2+}(\text{aq}) + 4 \text{H}_2\text{O} \rightarrow \text{MnO}_4^{2-}(\text{aq}) + 8 \text{H}^+(\text{aq}) + 5 \text{e}^-$	-1.51
$\text{MnO}_2(\text{s}) + 2 \text{H}_2\text{O} \rightarrow \text{MnO}_4^{2-}(\text{aq}) + 4 \text{H}^+(\text{aq}) + 3 \text{e}^-$	-1.69
$\text{MnO}_2(\text{s}) + 2 \text{H}_2\text{O} \rightarrow \text{MnO}_4^{2-}(\text{aq}) + 4 \text{H}^+(\text{aq}) + 2 \text{e}^-$	-2.26
$\text{Mn}^{3+}(\text{aq}) + 2 \text{H}_2\text{O} \rightarrow \text{MnO}_2(\text{s}) + 4 \text{H}^+(\text{aq}) + \text{e}^-$	-1.0
$\text{Mn}(\text{OH})_2(\text{s}) + 2 \text{OH}^-(\text{aq}) \rightarrow \text{MnO}_2(\text{s}) + 2 \text{H}_2\text{O} + 2 \text{e}^-$	0.03
$\text{Mn}(\text{OH})_2(\text{s}) + 6 \text{OH}^-(\text{aq}) \rightarrow \text{MnO}_4^{2-}(\text{aq}) + 4 \text{H}_2\text{O} + 5 \text{e}^-$	-0.34
$\text{MnO}_2(\text{s}) + 4 \text{OH}^-(\text{aq}) \rightarrow \text{MnO}_4^{3-}(\text{aq}) + 2 \text{H}_2\text{O} + \text{e}^-$	-0.9
$\text{MnO}_2(\text{s}) + 4 \text{OH}^-(\text{aq}) \rightarrow \text{MnO}_4^{2-}(\text{aq}) + 2 \text{H}_2\text{O} + 2 \text{e}^-$	-0.603
$\text{Mn}(\text{OH})_2(\text{s}) + \text{OH}^-(\text{aq}) \rightarrow \text{Mn}(\text{OH})_3(\text{s}) + \text{e}^-$	0.2
$\text{MnO}_4^{3-}(\text{aq}) \rightarrow \text{MnO}_4^{2-}(\text{aq}) + \text{e}^-$	-0.56

*(i) Oxide halides*

$\text{MnO}_3\text{X}$  ( $\text{X} = \text{F}, \text{Cl}$ ) are obtained from the reaction of  $\text{KMnO}_4$  with the appropriate halosulphonic acid<sup>4,5</sup> or <sup>4,6,7</sup> with anhydrous  $\text{HX}$ .  $\text{MnO}_3\text{F}$  is also formed<sup>8</sup> from  $\text{KMnO}_4$  and  $\text{IF}_5$ . Manganese trioxide fluoride forms dark green crystals, which melt at  $-38^\circ$  to a green liquid, which vaporises to a green gas (extrapolated b.p.  $+60^\circ$ ). Green-violet  $\text{MnO}_3\text{Cl}$  is similar but less stable.  $\text{MnO}_3\text{F}$  decomposes above  $0^\circ$ , often explosively, forming  $\text{MnO}_2$ ,  $\text{MnF}_2$ , and  $\text{O}_2$ , and hydrolyses instantly in water to  $\text{HMnO}_4$  and  $\text{HF}$ . Microwave spectroscopy<sup>9</sup> shows the structure to be tetrahedral, with  $\text{Mn}-\text{F} = 1.724 \text{ \AA}$ ,  $\text{Mn}-\text{O} = 1.586 \text{ \AA}$ , and  $\text{O}-\text{Mn}-\text{F} = 108.5^\circ$ . The IR and electronic spectra of both compounds have been reported<sup>5,10</sup>.

*(ii) Manganese heptoxide*

$\text{Mn}_2\text{O}_7$  is an oily liquid,  $D_4^{20} = 2.396$ , which appears red by transmitted, green by reflected light; the vapour is purple. It freezes at  $5.9^\circ$  and can be distilled at low pressure<sup>11</sup>. It separates on standing<sup>11-13</sup> from  $\text{KMnO}_4$  and cold, concentrated  $\text{H}_2\text{SO}_4$ .  $\text{Mn}_2\text{O}_7$  decomposes slowly above  $0^\circ$  into  $\text{MnO}_2$  and oxygen, at higher temperatures or in the presence of dust or organic matter, it explodes violently, forming  $\text{Mn}_2\text{O}_3$  and oxygen<sup>11</sup>. It dissolves in water to form permanganic acid of which it is the anhydride, and is soluble without reaction in freons and  $\text{SO}_2\text{Cl}_2$ , and only slowly reacts with  $\text{CCl}_4$ ; such solutions have been proposed as useful strong oxidising agents<sup>5</sup>. The structure is probably  $\text{O}_3\text{MnOMnO}_3$  (cf  $\text{Cl}_2\text{O}_7$ ), a proposal supported by the small dipole moment<sup>11</sup>.

(iii) *Permanganic acid and the permanganates*

A deep purple solution of  $\text{HMnO}_4$  is obtained by adding the calculated quantity of sulphuric acid to saturated barium permanganate solution<sup>14</sup>. The dilute solution is stable but decomposes on concentration. Dubois<sup>14</sup> obtained a 24.5% solution which froze at  $-11^\circ$  to a mass of ice and permanganic acid crystals, but these soon decomposed in diffuse daylight. By fractional vacuum sublimation of these crystals, anhydrous  $\text{HMnO}_4$  and the dihydrate  $\text{HMnO}_4 \cdot 2\text{H}_2\text{O}$  were obtained<sup>15</sup>.  $\text{HMnO}_4$  forms deep violet rectangular crystals, very soluble in water, and slightly soluble in  $\text{CCl}_3\text{CF}_3$  and perfluorodecalin. It explodes on contact with most organic materials, and on warming. The dihydrate is generally similar but reacts less violently.

$\text{KMnO}_4$  is extensively used as a strong oxidising agent, no attempt has been made to cover this aspect of its chemistry in this review. Dark purple, orthorhombic crystals of  $\text{KMnO}_4$  are normally obtained from  $\text{K}_2\text{MnO}_4$  by electrochemical oxidation, or by disproportionation with acids (usually  $\text{CO}_2$ ). Barium permanganate results from passing  $\text{CO}_2$  through an aqueous suspension of  $\text{BaMnO}_4$  (ref. 16), and other permanganates are obtained from this by metathesis with the appropriate sulphate<sup>17-19</sup>.  $\text{AgMnO}_4$ , which crystallises on mixing saturated solutions of  $\text{KMnO}_4$  and  $\text{AgNO}_3$ , can also be used with the appropriate chloride<sup>20-23</sup>. The sparingly soluble permanganates of  $\text{NH}_4^+$ , Rb, Cs, can be obtained directly from  $\text{KMnO}_4$  and  $\text{M}_2\text{SO}_4$  ( $\text{M} = \text{NH}_4, \text{Rb}, \text{Cs}$ )<sup>24</sup>.

The permanganates of K, Rb, Cs,  $\text{NH}_4$ , Ba, Ag, crystallise anhydrous, but many of the others form hydrates, e.g.  $\text{Li} (3\text{H}_2\text{O})$ ,  $\text{Na} (2 \cdot 1\text{H}_2\text{O})$  (ref. 25), Mg, Cd, Zn, Cu, Ni, (all  $6\text{H}_2\text{O}$ ), Sr ( $3\text{H}_2\text{O}$ ), Ca ( $4\text{H}_2\text{O}$ ). The hydrated alkali and alkaline earth permanganates can be dehydrated without decomposition, but the others decompose<sup>25</sup>.

The permanganate ion consists of a manganese atom surrounded by four oxygen atoms at the corners of a regular tetrahedron<sup>26</sup> with  $\text{Mn}-\text{O}$  (ave) =  $1.629 \pm 0.008 \text{ \AA}$ , and  $\text{O}-\text{Mn}-\text{O} = 109.4 \pm 0.7^\circ$ . The electronic spectrum is discussed in Sect. E and the vibrational frequencies are recorded in Table 2. The IR spectra of a number of individual permanganates have been recorded, and discussed in terms of the site symmetry of the anion<sup>25</sup>.


The thermal decomposition of  $\text{KMnO}_4$  has been the subject of a great deal of work, but the nature of some of the products is still uncertain. The water-soluble decomposition product is  $\text{K}_2\text{MnO}_4$ , but the nature of the water-insoluble part has been disputed. Thermogravimetric and differential thermal analyses have been used to study the decomposition

TABLE 2

Vibrational spectra of the tetraoxoanions

	Frequency ( $\text{cm}^{-1}$ )				Ref.
	$\nu_1(A_1)$	$\nu_2(E)$	$\nu_3(F_2)$	$\nu_4(F_2)$	
$\text{MnO}_4^-$	838	355	921	429	29
$\text{MnO}_4^{2-}$	810		862	328	30
$\text{MnO}_4^{3-}$	836		770	348	30

of the K and Rb salts<sup>27,28</sup>. Herbst et al<sup>28</sup> propose the idealised decomposition equation



at 250° in air. The phase in parentheses is of unknown structure but may be related to the  $\text{Mn}^{\text{IV}}-\text{Mn}^{\text{III}}$  complex oxides prepared by Scholder and Protzer<sup>51</sup>.

The structural parameters of some individual permanganates are given in Table 3


TABLE 3

Crystal structures of some permanganates

Compound	Type	Dimensions (Å)			Ref
		<i>a</i>	<i>b</i>	<i>c</i>	
$\text{KMnO}_4$	Orthorhombic	9.105	5.720	7.425	26
$\text{RbMnO}_4$	Orthorhombic	7.65	9.55	5.74	31
$\text{CsMnO}_4$	Orthorhombic	7.96	10.06	5.80	31
$\text{NH}_4\text{MnO}_4$	Orthorhombic	7.58	9.33	5.78	32
$\text{LiMnO}_4 \cdot 3\text{H}_2\text{O}$	Hexagonal	7.73		5.39	17,33
$\text{Ba}(\text{MnO}_4)_2$	Orthorhombic	14.71	11.86	7.39	34
$\text{Sr}(\text{MnO}_4)_2 \cdot 3\text{H}_2\text{O}$	Cubic	9.61			22
$\text{AgMnO}_4$	Monoclinic	5.66	8.27	7.13	35
$\text{Mg}(\text{MnO}_4)_2 \cdot 6\text{H}_2\text{O}$	Orthorhombic	7.81	13.56	5.28	31
$\text{Zn}(\text{MnO}_4)_2 \cdot 6\text{H}_2\text{O}$	Orthorhombic	7.81	13.56	5.23	31
$\text{Ni}(\text{MnO}_4)_2 \cdot 6\text{H}_2\text{O}$	Orthorhombic	7.75	13.45	5.20	31
$\text{Cd}(\text{MnO}_4)_2 \cdot 6\text{H}_2\text{O}$	Orthorhombic	8.04	13.91	5.34	19

(iv)  $\text{MnO}_3^+$  or  $\text{O}_3\text{MnOSO}_3\text{H}$

The green solution formed on dissolving  $\text{KMnO}_4$  in concentrated  $\text{H}_2\text{SO}_4$  contains  $\text{MnO}_3^+$  or  $\text{O}_3\text{MnOSO}_3\text{H}$ , or possibly both, depending upon concentration, and not  $(\text{MnO}_3)_2\text{SO}_4$  as was assumed in the older literature<sup>36,37</sup>. Cryoscopic, conductimetric, and electronic spectral studies were said to be consistent with the presence of a planar  $\text{MnO}_3^+$  cation, formed by the reaction<sup>36</sup>



In 100% sulphuric acid  $\text{O}_3\text{MnOSO}_3\text{H}$  is said to be present<sup>37</sup>

### C MANGANESE (VI)

$\text{Mn}^{\text{VI}}$  is confined to salts of the manganate (VI) ion, and  $\text{MnO}_2\text{Cl}_2$ .  $\text{Mn}^{\text{VI}}$  is stable only in basic solution, its most characteristic reaction being to disproportionate in acidic or neutral solution.

(i) *Manganese dioxide dichloride*

$\text{MnO}_2\text{Cl}_2$  is the only oxide halide of  $\text{Mn}^{\text{VI}}$  known, although presumably  $\text{MnO}_2\text{F}_2$

could be prepared. Briggs<sup>5</sup> obtained  $\text{MnO}_2\text{Cl}_2$  as a very unstable brown liquid, by reduction of  $\text{KMnO}_4$  in  $\text{HSO}_3\text{Cl}$  with  $\text{SO}_2$  at low temperatures. It begins to decompose at  $-30^\circ$ , and hydrolyses to  $\text{MnO}_4^-$ ,  $\text{MnO}_2$ , and  $\text{Cl}^-$


(ii) "Manganese trioxide"

The early claims<sup>38,39</sup> to have prepared  $\text{MnO}_3$  have not been substantiated by later work. Lankshear<sup>40</sup> showed that the red solid claimed as  $\text{MnO}_3$  is really a mixture of  $\text{HMnO}_4$  and some  $\text{MnO}_2$ , and it has since been demonstrated that  $\text{Mn}_2\text{O}_7$  loses oxygen to form  $\text{MnO}_2$  directly, with no evidence for any intermediate oxide<sup>12</sup>. Despite this, reports of a compound  $\text{MnO}_3$  still appear in textbooks from time to time<sup>1</sup>


(iii) Manganates (VI)

The parent acid is unknown, acidification of manganate (VI) solutions resulting in disproportionation. Manganates (VI) are obtained by fusing  $\text{MnO}_2$  with alkali metal hydroxides under oxidising conditions or by electrolytic oxidation of alkaline manganese (II) solutions<sup>41</sup>. Pure  $\text{K}_2\text{MnO}_4$  was obtained by boiling  $\text{KMnO}_4$  with concentrated  $\text{KOH}$  solution, and various hydrates of sodium manganate (VI) can be prepared similarly<sup>42</sup>. Alkali metal manganates (VI) are also formed by thermal decomposition of the corresponding permanganate<sup>43</sup>, or by heating  $\text{MnO}_2$  with the metal superoxide<sup>44</sup>. Barium manganate precipitates<sup>16</sup> on boiling  $\text{KMnO}_4$  with a saturated aqueous solution of  $\text{Ba}(\text{NO}_2)_2$  and  $\text{Ba}(\text{OH})_2$ . Lithium manganate does not appear to have been reported,  $\text{MnO}_2$  does not dissolve readily in fused  $\text{LiOH}$ , and  $\text{LiMnO}_4$  decomposes to  $\text{Li}_2\text{O}$  and  $\text{MnO}_2$  on heating<sup>43</sup>. A number of other manganates (VI) were reported in the older literature (see ref. 45), but these must be regarded as doubtful until reinvestigated.

Alkali metal manganates (VI) form deep green crystals, which often appear darker due to a superficial layer of permanganate. They dissolve in aqueous alkali to form green solutions, but in acidic, neutral, or even weakly basic solutions, they disproportionate



Thermal decomposition produces manganates (V) and other products, e.g., at  $620^\circ$  in nitrogen<sup>28</sup>



The electronic spectrum is discussed later (Section E) and the IR frequencies are shown in Table 2.  $\text{K}_2\text{MnO}_4(d^1)$  has a magnetic moment of 1.73 B.M., and follows the Curie-Weiss law with  $\theta \simeq 0^\circ$  (ref. 46). The structure of  $\text{K}_2\text{MnO}_4$  shows the  $\text{MnO}_4^{2-}$  ion to be tetrahedral with  $\text{Mn}-\text{O} = 1.659 \text{ \AA}$ , and  $\text{O}-\text{Mn}-\text{O} = 109.5 \pm 0.7^\circ$ , which means that the  $\text{Mn}-\text{O}$  bond length is 0.03 Å longer than in the permanganate<sup>47</sup>, in agreement with the predictions of molecular orbital theory. The structural parameters of the alkali manganates are reported in Table 4.

TABLE 4

## Manganate (VI) crystal structures

Compound	Type	Dimensions (Å)			Ref
		<i>a</i>	<i>b</i>	<i>c</i>	
K <sub>2</sub> MnO <sub>4</sub>	Orthorhombic	7 667	5 895	10 359	47
Rb <sub>2</sub> MnO <sub>4</sub>	Orthorhombic	7 997	10 670	6 044	44
Cs <sub>2</sub> MnO <sub>4</sub>	Orthorhombic	8 360	11 052	6 247	44

## D MANGANESE (V)

Manganese (V), the rarest of the higher oxidation states, occurs in the manganate (V) anion, and in MnOCl<sub>3</sub>. It is even more prone to disproportionation than manganese (VI), existing only in strongly basic solution.

*(i) Manganese oxide trichloride*

This is the only oxide halide, although as with MnO<sub>2</sub>Cl<sub>2</sub> the fluorine analogue should be capable of existence. MnOCl<sub>3</sub> is a mint-green liquid, with a deep yellow vapour, formed by reducing KMnO<sub>4</sub> dissolved in HSO<sub>3</sub>Cl with CHCl<sub>3</sub>, or better with sucrose<sup>5</sup>. The pure liquid decomposes above 0° to MnCl<sub>3</sub>, and hydrolyses readily, MnO<sub>4</sub><sup>3-</sup> being produced only if sufficient alkali is present to prevent its decomposition.

*(ii) Manganates (V)*

These were characterised by Lux<sup>48</sup> as recently as 1946, although other workers had previously obtained blue oxo-species of manganese without recognising their true nature. Deep blue sodium manganate (V) is obtained<sup>48-50</sup> on reduction of KMnO<sub>4</sub> in concentrated aqueous NaOH, with Na<sub>2</sub>SO<sub>3</sub>, KI, or HCOONa, or by oxidation of manganese (II) or MnO<sub>2</sub> in an alkaline melt<sup>48,49</sup>. A simple method for obtaining the Na, K, Rb compounds has recently been described<sup>51</sup>. Pure anhydrous alkali manganates (V) (including Li<sub>3</sub>MnO<sub>4</sub>) are formed<sup>50</sup> by heating the permanganate with MOH.

Lux<sup>48</sup> obtained a deep blue hydrate which he formulated as Na<sub>3</sub>MnO<sub>4</sub> · 10H<sub>2</sub>O, but Scholder et al.<sup>50</sup> suggested this was Na<sub>3</sub>MnO<sub>4</sub> · ½ NaOH · 12H<sub>2</sub>O, and found that on recrystallisation from cold concentrated NaOH, Na<sub>3</sub>MnO<sub>4</sub> · 7H<sub>2</sub>O is produced. A compound of limiting composition Na<sub>3</sub>MnO<sub>4</sub> · 0.25NaOH · 12H<sub>2</sub>O, analogous to the corresponding vanadate, phosphate, and arsenate has been obtained<sup>52</sup>. Klemm showed that Na<sub>3</sub>MnO<sub>4</sub> · 10H<sub>2</sub>O contains Mn<sup>V</sup> and not Mn<sup>VI</sup> and Mn<sup>IV</sup> by magnetic analysis<sup>53</sup>.

Ba<sub>3</sub>(MnO<sub>4</sub>)<sub>2</sub> (refs 53, 54), Sr<sub>2</sub>(MnO<sub>4</sub>)(OH) (refs 54, 55), Ba<sub>5</sub>(MnO<sub>4</sub>)<sub>3</sub>Cl (ref. 56), and Ba<sub>5</sub>(MnO<sub>4</sub>)<sub>3</sub>(OH) (ref. 53) have been reported.

Electronic (Sect. E) and vibrational spectra (Table 2), have been reported, but no structural data are available for the MnO<sub>4</sub><sup>3-</sup> ion. The magnetic moments of a number of Mn<sup>V</sup> compounds have been reported by Klemm et al.<sup>57</sup> as Na<sub>3</sub>MnO<sub>4</sub> · 10H<sub>2</sub>O 2.8B M,

$\text{Ba}_5(\text{MnO}_4)_3(\text{OH})$  2.83B M.,  $\text{Ba}_3(\text{MnO}_4)_2$  2.83B M.,  $\text{K}_3\text{MnO}_4$  2.68B M.\* ,  $\text{Rb}_3\text{MnO}_4$  2.61B M.\* and  $\text{Cs}_3\text{MnO}_4$  2.57B M.\*

In solution, manganates (V) disproportionate into  $\text{MnO}_4^{2-}$  and  $\text{MnO}_2$ , unless a large excess of alkali is present.  $\text{K}_3\text{MnO}_4$  is much more stable thermally than either  $\text{KMnO}_4$  or  $\text{K}_2\text{MnO}_4$ , in dry air it appears to be stable up to at least  $900^\circ$ , but is decomposed rapidly by moisture<sup>28</sup>.

A compound  $\text{Na}_5\text{MnO}_5$  was reported in Scholder's review<sup>54</sup> but no details were given.

## E ELECTRONIC SPECTRA AND STRUCTURE OF THE TETRAOXOANIONS

The tetraoxoanions of manganese have provided fruitful discussion for workers interested in the energy levels and electronic spectra and structures of these tetrahedral  $(\text{XO}_4)^{n-}$  species, which are characterised by strong visible and near-ultraviolet absorptions. There have been many interpretations, and these frequently differ from one another.

Wolfsberg and Helmholtz<sup>58</sup> (W-H) carried out some semi-empirical calculations on the permanganate ion, according to the molecular orbital scheme, and their most surprising result was the order of the first unfilled MO's ( $3t_2 < 2e$ ) which is not in agreement with crystal field (C F) theory. Subsequently, Ballhausen and Liehr<sup>59</sup> (B-L) proposed a different scheme which is in agreement with C F. theory and is justified by intensity calculations. The latter proposals were supported by the studies of Carrington and coworkers<sup>60-63</sup>, who showed on the basis of ESR measurements that the unpaired electron in the  $\text{MnO}_4^{2-}$  ion ( $d^1$ ), occupied an  $e$  level. More recently, Fenske and Sweeney<sup>64</sup> concluded that if the empirical parameters of the W-H calculations are substantially altered, any order of levels can be obtained, and, consequently, any transition assignment justified. A calculation of Viste and Gray<sup>65</sup> (V-G) has confirmed the B-L order of levels, but with a different transition assignment.

Oleary et al.<sup>66</sup> have carried out an S.C.F. -M O -L C A O. calculation for the  $\text{MnO}_4^-$  ion, and confirmed the order of levels previously assigned by B-L. A new spectral assignment was presented, attributing to each observed absorption band two electronic transitions occurring between one of the occupied MO's ( $t_1, 2t_2, 1t_2$ ) and the two lowest empty MO's ( $2e, 3t_2$ ). These workers subsequently extended their calculations to the  $\text{MnO}_4^{2-}$  and  $\text{MnO}_4^{3-}$  ions<sup>67</sup>.

The spectra of the three ions are shown in Fig. 1, and Table 5 contains the spectral assignments proposed by various workers.

## F MANGANESE (IV)

$\text{Mn}^{\text{IV}}$  has a more extensive chemistry than the higher oxidation states, but, in general, the compounds are not particularly stable. The apparent exception,  $\text{MnO}_2$ , owes its "stability" to its insolubility; other  $\text{Mn}^{\text{IV}}$  compounds are readily hydrolysed and reduced. There are few simple compounds, the majority of manganese (IV) compounds containing the element in a complex anion.

\*Low values may be due to the presence of some manganate (VI)

### Spectral assignments of tetraoxoanions

Anion	Band maxima (cm <sup>-1</sup> )	Wolfsberg-Helmholtz <sup>58</sup>	Ballhausen-Liehr <sup>59</sup>	Carrington-Jorgensen <sup>60</sup>	Carrington-Symons <sup>60</sup>	Viste-Gray <sup>61</sup>	Kingsley et al. <sup>62</sup>	Orgel <sup>63</sup>	Orlean et al. <sup>64,67</sup>
[MnO <sub>4</sub> ] <sup>-</sup>		$t_1 \rightarrow 3t_2$	$t_1 \rightarrow 2e$	$t_1 \rightarrow 2e$		$t_1 \rightarrow 2e$	$\begin{cases} t_1 \rightarrow 3t_2 \\ t_1 \rightarrow 2e \end{cases}$		
		$2t_2 \rightarrow 3t_2$	$t_1 \rightarrow 3t_2$	$t_1 \rightarrow 3t_2$			$2t_2 \rightarrow 2e$		
				$2t_2 \rightarrow 3t_2$			$2t_2 \rightarrow 3t_2$		
							$\begin{cases} 1t_2 \rightarrow 2e \\ 1t_2 \rightarrow 3t_2 \\ 1e \rightarrow 3t_2 \end{cases}$		
[MnO <sub>4</sub> ] <sup>2-</sup>	(12,000)				$2e \rightarrow 3t_2(^3T_2)$		$2e \rightarrow 3t_2(^3T_2)$		
	16,580	$2e \rightarrow 3t_2(^1)$		$2e \rightarrow 3t_2$	$t_1 \rightarrow 2e(^3T_2)$	$2e \rightarrow 3t_2$	$t_1 \rightarrow 2e(^3T_1)$		
	22,830	$t_1 \rightarrow 2e$		$t_1 \rightarrow 2e$	$t_1 \rightarrow 2e(^3T_1)$	$t_1 \rightarrow 2e$	$t_1 \rightarrow 2e(^3T_1)$		
	28,170	$2t_1 \rightarrow 2e(^1)$		$2t_1 \rightarrow 2e$	$t_1 \rightarrow 3t_2(^3T_2)$	$2t_2 \rightarrow 2e$	$\begin{cases} 2t_2 \rightarrow 2e(^3T_1) \\ t_1 \rightarrow 3t_2(^3T_2) \\ 2t_2 \rightarrow 2e(^3T_2) \end{cases}$		
					$t_1 \rightarrow 3t_2(^3T_1)$		$2t_2 \rightarrow 3t_2(^3T_2)$		
	33,200						$t_1 \rightarrow 3t_2(^3T_1)$		
[MnO <sub>4</sub> ] <sup>3-</sup>	11,000					$2e \rightarrow 3t_2(^3T_2)$	$2e \rightarrow 3t_2(^3T_2)$		
	14,810	$2e \rightarrow 3t_2$			$t_1 \rightarrow 2e(^3T_1)$	$2e \rightarrow 3t_2(^3T_1)$	$t_1 \rightarrow 2e(^3T_1)$		
	(16,700)					$2e \rightarrow 3t_2(^3T_1)$	$2e \rightarrow 3t_2(^3T_1)$		
	30,800		$t_1 \rightarrow 2e$		$t_1 \rightarrow 3t_2(^3T_1)$	$t_1 \rightarrow 2e(^3T_1)$	$t_1 \rightarrow 3t_2(^3T_1)$		






Fig 1 Electronic spectra of the tetraoxoanions (a), (b), Oleari et al <sup>67</sup>, (c) G van den Boef, H J van der Beek and T. Braaf *Rec. Trav. Chim.*, 77 (1958) 1064 Reproduced by permission

### (i) Halides

The only halide is the very reactive  $\text{MnF}_4$ , there is little evidence for  $\text{MnCl}_4$  despite a number of claims in the older literature<sup>70</sup>. Sharpe and Woolf<sup>71</sup> were unable to isolate  $\text{MnF}_4$  from the reaction of  $\text{Mn}(\text{IO}_3)_2$  with  $\text{BrF}_3$ , although the solution obtained gave  $\text{Mn}^{\text{IV}}$  fluorocomplexes upon addition of  $\text{KF}$ . The blue, very hygroscopic, tetrafluoride is produced by fluorination of manganese powder in a fluidised bed<sup>72</sup> or of  $\text{MnF}_2$ ,  $\text{MnF}_3$ ,  $\text{LiMnF}_5$ , or  $\text{Li}_2\text{MnF}_6$ , at  $550^\circ$  (ref 73) It obeys the Curie–Weiss law with  $\mu_{\text{eff}} = 3.48$  B.M. Fluorine is lost on standing, and it hydrolyses instantly in water. A small amount of red solid is produced from  $\text{ClF}$  and manganese powder in a fluidised bed<sup>72</sup>, this may be a mixed halide, e.g.  $\text{MnClF}_3$ . There appear to be no oxide halides known

### (ii) Complex halides

Hexahalomanganates (IV),  $\text{MnX}_6^{2-}$  ( $\text{X} = \text{Cl}, \text{F}$ ), and pentafluoromanganates (IV),  $\text{MnF}_5^-$ , are known. All attempts to prepare bromine analogues have failed.

Hexafluoromanganates (IV) are obtained by four basic methods.

(a) Direct fluorination of an intimate mixture of  $\text{MnSO}_4$ ,  $\text{MnCl}_2$ , or  $\text{NH}_4\text{MnF}_3$ , and the chloride, sulphate, or carbonate of the desired cation.

(b) Fluorination of the corresponding permanganate, manganate (VI), or manganate (IV)

(c) Electrolytic oxidation of  $\text{MnF}_2$  in 40% HF in the presence of the metal fluoride.

(d) Fluorination of the starting materials in (a) or (b) with  $\text{BrF}_3$  (ref. 89).

Some properties of these compounds are reported in Table 6, and their structures in Table 7

$(\text{NO})_2\text{MnF}_6$  and the heavy metal hexafluoromanganates are instantly decomposed by water, the alkali metal compounds hydrolyse only slowly in the cold. The  $\text{MnF}_6^{2-}$  ion ( $d^3$ ) is a regular octahedron<sup>77, 87</sup>, with  $\text{Mn}-\text{F} = 1.72-1.75 \text{ \AA}$ , and exhibits  $\nu(\text{Mn}-\text{F})$  at  $625 \text{ cm}^{-1}$  (ref. 81). Asprey et al.<sup>97a</sup> have recently studied the IR and Raman spectra of some hexafluoromanganates (IV).

The brick-red pentafluoromanganates (IV), formed on fluorination of  $\text{MMnF}_3$  ( $\text{M} = \text{K}, \text{Rb}, \text{Cs}$ ) at  $450-500^\circ$ , or of  $\text{LiF} + \text{MnF}_2$  at  $350^\circ$ , are instantly decomposed by water<sup>73, 74</sup>. They have magnetic moments in the range  $3.7-3.9 \text{ B.M.}$ , and the  $\text{Mn}-\text{F}$  stretching frequency in  $\text{KMnF}_5$  is at  $617 \text{ cm}^{-1}$  (ref. 81). Although the structures have not been determined, it seems probable that they contain condensed  $\text{MnF}_6$  units.

The very dark red hexachloromanganates (IV) are prepared<sup>88, 90</sup> by adding saturated alkali metal chloride solution to calcium permanganate in 40% hydrochloric acid at  $0^\circ$ . The potassium salt is also obtained from  $\text{KMnO}_4$  and saturated aqueous HCl (ref. 88), or by the reaction of potassium acetate and manganese (III) acetate in the presence of a large excess of acetyl chloride<sup>91</sup>. They readily lose chlorine on standing.  $\text{K}_2\text{MnCl}_6$  has  $\mu_{\text{eff}} = 3.9 \text{ B.M.}$ <sup>92</sup> and  $\nu(\text{Mn}-\text{Cl}) = 358 \text{ cm}^{-1}$  (ref. 93).

The diffuse reflectance spectra of the hexafluoromanganates have been studied by a number of workers<sup>94-96</sup>. Allen et al.<sup>94</sup> found five bands in the spectrum of  $\text{K}_2\text{MnF}_6$  at  $14.0 \text{ kK}$  ( ${}^4A_{2g} \rightarrow {}^2E_g, {}^2T_{1g}, t_{2g}^3$ ),  $29.3 \text{ kK}$  ( ${}^4A_{2g} \rightarrow {}^2T_{2g}, t_{2g}^3$ ),  $22.2 \text{ kK}$  ( ${}^4A_{2g} \rightarrow {}^2T_{2g}, t_{2g}^2e_g$ ),  $28.6 \text{ kK}$  ( ${}^4A_{2g} \rightarrow {}^2T_{1g}, t_{2g}^2e_g$ ),  $38.8 \text{ kK}$  ( $\pi \rightarrow t_{2g}$ ). Novotny and Sturgeon<sup>95</sup> reported the spectra of the K, Rb, Cs and Ba compounds, and examined the effect of the method of preparation on the spectrum of the K salt. Moews<sup>88</sup> reported bands at  $15.4$ ,  $16.6$ ,  $27.4$ ,  $33.3 \text{ kK}$  for  $\text{K}_2\text{MnCl}_6$ , but this was disputed by other workers, who reported absorption at  $17.8$ ,  $25.0$ ,  $43.5 \text{ kK}$ , which they assigned as  ${}^4t_2 \rightarrow {}^4t_5$ ,  ${}^4t_2 \rightarrow {}^4t_4$ , and charge transfer, respectively<sup>97</sup>.

### (iii) Manganese dioxide

There is an extensive literature on  $\text{MnO}_2$ , only a very small amount of which will be mentioned here<sup>98-110</sup>. The common form,  $\beta\text{-MnO}_2$  (pyrolusite), is made by heating manganese (II) nitrate at  $150-160^\circ$ , extracting with water, drying at  $200^\circ$ , and repeatedly extracting with boiling nitric acid to remove lower oxides<sup>105</sup>. Several modifications of the basic method have been described<sup>98</sup> and most of the other forms of  $\text{MnO}_2$  are said<sup>102</sup> to change into the  $\beta$  variety on heating in air at  $400^\circ$ . It has an ideal composition  $\text{MnO}_{2.00}$ , with the rutile structure  $a = 4.38$ ,  $c = 2.86 \text{ \AA}$ , but it is usually oxygen deficient<sup>98</sup>. Reducing agents attack it readily in acid solution, alkalis cause disproportionation, whilst concentrated  $\text{H}_2\text{SO}_4$  at  $110^\circ$  forms  $\text{Mn}^{\text{III}}$  sulphate and oxygen.

TABLE 6

## Hexafluoromanganates (IV)

Compound	Preparation*	Colour	Magnetic moment (B.M.)	Ref
$\text{Li}_2\text{MnF}_6$	a	Yellow	3.85	74
$\text{Na}_2\text{MnF}_6$	a, c	Yellow		75
$\text{K}_2\text{MnF}_6$	a, c, d	Yellow	3.86	71, 75, 79, 82-84
$\text{Rb}_2\text{MnF}_6$	a, c	Yellow		75, 79
$\text{Cs}_2\text{MnF}_6$	a, c	Yellow		75, 79
$(\text{NH}_4)_2\text{MnF}_6$	c	Yellow		85
$\text{CaMnF}_6$	a	Yellow	3.87	78, 79, 87
$\text{SrMnF}_6$	a, b	Yellow	3.82	78-80
$\text{BaMnF}_6$	a, b	Yellow	3.90	78-80
$\text{MgMnF}_6$	a	Orange-yellow	3.9	78, 79, 87
$(\text{NO})_2\text{MnF}_6$	d	Yellow		86
$\text{CdMnF}_6$	a	Yellow	3.78	87
$\text{NiMnF}_6$	a	Ochre-yellow	5.38	87
$\text{ZnMnF}_6$	a	Orange-yellow		87
$\text{HgMnF}_6$	a	Orange		87
$\text{CuMnF}_6$	a	Bright red	4.43	87
$\text{AgMnF}_6$	b	Dark brown	4.43	87

\*Letters refer to methods a, b, c and d discussed in text

TABLE 7

Structural parameters of  $\text{Mn}^{\text{IV}}$  complex halides

Compound	Type	Parameters (Å)		Ref
		a	c	
$\text{Li}_2\text{MnF}_6$	$\text{Na}_2\text{SiF}_6$ , hexagonal	8.42	4.59	74
$\text{Na}_2\text{MnF}_6$	$\text{Na}_2\text{SiF}_6$ , hexagonal	9.03	5.13	75, 76
$\text{K}_2\text{MnF}_6$	" $\text{Rb}_2\text{MnF}_6$ ", hexagonal	5.67	9.35	77
	trigonal	5.71	4.65	77
$\text{Rb}_2\text{MnF}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	8.28		77
	" $\text{Rb}_2\text{MnF}_6$ ", hexagonal	5.86	9.50	77
	$\text{K}_2\text{PtCl}_6$ , cubic	8.40		77
$\text{Cs}_2\text{MnF}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	8.92		77
$(\text{NH}_4)_2\text{MnF}_6$	" $\text{Rb}_2\text{MnF}_6$ ", hexagonal	5.91	9.55	85
$\text{MgMnF}_6$	$\text{LiSbF}_6$ , hexagonal	5.01	13.17	78, 79, 87
$\text{CaMnF}_6$	$\text{LiSbF}_6$ , hexagonal	5.21	14.17	78, 79, 87
$\text{SrMnF}_6$	$\text{BaGeF}_6$ , hexagonal	7.02	6.78	78, 79
$\text{BaMnF}_6$	$\text{BaGeF}_6$ , hexagonal	7.35	7.09	78, 79, 80
$\text{CdMnF}_6$	$\text{LiSbF}_6$ , hexagonal	5.08	14.00	87
$\text{NiMnF}_6$	$\text{VF}_3$ , hexagonal	4.91	13.16	87
$\text{ZnMnF}_6$	$\text{VF}_3$ , hexagonal	4.96	13.29	87
$\text{HgMnF}_6$	$\text{LiSbF}_6$ , hexagonal	5.08	14.12	87
$\text{K}_2\text{MnCl}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	$9.6445 \pm 0.002$		88
$\text{Rb}_2\text{MnCl}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	$9.82 \pm 0.02$		88
$\text{Cs}_2\text{MnCl}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	$10.17 \pm 0.02$		88
$(\text{NH}_4)_2\text{MnCl}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	$9.80 \pm 0.02$		88
$(\text{NMe}_4)_2\text{MnCl}_6$	$\text{K}_2\text{PtCl}_6$ , cubic	$12.70 \pm 0.02$		88


A large number of varieties of  $\text{MnO}_2$  have been reported, but despite a great deal of work, the nature of some of these is still in doubt. Many of them contain  $\text{Mn}^{\text{III}}$  ions, water, or other metal ions, and exist over a range of compositions. A brief discussion of some of these forms is given below (see refs 102–104 for references to earlier work).

Glemser and coworkers<sup>102–104</sup> described six other varieties,  $\alpha$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\eta$ , and the mineral ramsdellite. The latter is orthorhombic  $a = 4.53$ ,  $b = 9.27$ ,  $c = 2.87$  Å.  $\alpha$ - $\text{MnO}_2$  is related to the mineral cryptomelane, and always contains other large cations, it has been formulated as  $\text{A}_x\text{Mn}_8\text{O}_{16}$  ( $\text{A} = \text{K}, \text{Ba}, \text{etc.}, x < 1$ )<sup>101, 102</sup>. All forms lose oxygen on heating, eventually forming  $\text{Mn}_2\text{O}_3$ . It has since been shown that  $\delta$ - $\text{MnO}_2$  is a ternary oxide (see below). The phase  $\text{Mn}(\text{OH})_2 \cdot 2\text{MnO}_2$  is formed<sup>106</sup> upon wet oxidation of  $\gamma$ - $\text{Mn}_2\text{O}_3$  above  $\text{MnO}_{1.4}$ .

(iv) "Mixed-valence oxides"

The manganese–oxygen system, especially in the presence of other cations, and of water, is very complex, and a number of compounds have been discovered in addition to the simple oxides of  $\text{Mn}^{\text{II}}$ ,  $\text{Mn}^{\text{III}}$ , and  $\text{Mn}^{\text{IV}}$ . The older literature contains several reports of oxides containing manganese in more than one oxidation state<sup>45</sup>, many of these must be regarded as very doubtful and will not be discussed further. Compounds that do fit into the category of mixed-valence oxides are  $\text{Mn}_3\text{O}_4$ ,  $\text{Mn}_5\text{O}_8$ , and the recently discovered  $\text{Mn}_7\text{O}_{12} \cdot 6\text{H}_2\text{O}$ , and  $\text{Mn}_7\text{O}_{13} \cdot 5\text{H}_2\text{O}$ .

$\alpha$ - $\text{Mn}_3\text{O}_4$  occurs as the mineral hausmannite, and is the product<sup>98</sup> of heating any oxide, and many manganese salts, in air at  $1000^\circ$ . The purple-red powder has a distorted spinel structure (distortion is caused by the  $d^4$   $\text{Mn}^{\text{III}}$  ions),  $a = 8.13$  Å,  $c/a = 1.16$ . It is a normal spinel  $\text{Mn}^{\text{II}}\text{Mn}_2^{\text{III}}\text{O}_4$  (ref. 111) not  $\text{Mn}_2^{\text{II}}\text{Mn}^{\text{IV}}\text{O}_4$  as reported in some of the earlier work. It changes reversibly into a cubic modification  $\beta$ - $\text{Mn}_3\text{O}_4$  above  $1170^\circ$ . Concentrated  $\text{H}_2\text{SO}_4$  produces  $\text{Mn}^{\text{II}}$  and  $\text{Mn}^{\text{III}}$  sulphates, and  $\text{HNO}_3$  causes disproportionation.



$\text{Mn}_5\text{O}_8$  is produced on oxidation of  $\text{Mn}_3\text{O}_4$  in a nitrogen/oxygen stream at  $250$ – $500^\circ$ , or by heating  $\beta$ - $\text{MnO}(\text{OH})$  in air below  $500^\circ$  (ref. 113). It is a black powder, which loses oxygen above  $550^\circ$  to form  $\alpha$ - $\text{Mn}_2\text{O}_3$ . The structure determination supports the formulation  $\text{Mn}_2^{\text{II}}\text{Mn}_3^{\text{IV}}\text{O}_8$ , there is a distorted octahedral arrangement of oxygen atoms around  $\text{Mn}^{\text{IV}}$ , and a distorted trigonal prism around  $\text{Mn}^{\text{II}}$  (ref. 114).

Giovanoli et al.<sup>108, 109</sup> found that  $\text{Na}_4\text{Mn}_{14}\text{O}_{27} \cdot 9\text{H}_2\text{O}$ , reacts with dilute nitric acid to produce manganese (III) manganate (IV)  $\text{Mn}_7\text{O}_{13} \cdot 5\text{H}_2\text{O}$  (hexagonal  $a = 2.84$ ,  $c = 7.27$  Å), and with  $\text{Mn}(\text{NO}_3)_2$  solution to form manganese (II) manganate (IV)  $\text{Mn}_7\text{O}_{12} \cdot 6\text{H}_2\text{O}$ . Prolonged digestion of  $\text{Mn}_7\text{O}_{13} \cdot 5\text{H}_2\text{O}$  with nitric acid produces  $\gamma$ - $\text{MnO}_2$ , upon heating  $\text{Mn}_7\text{O}_{13} \cdot 5\text{H}_2\text{O}$ , the first product is  $\gamma$ - $\text{MnO}_2$  and finally a distorted  $\beta$ - $\text{MnO}_2$  results.  $\delta$ - $\text{MnO}_2$  seems to be related to these manganate (IV) species with some of the  $\text{Mn}^{\text{III}}$  replaced by  $\text{Mn}^{\text{IV}}$ , it also contains a fairly large amount of alkali metal ions<sup>107, 109</sup>. The structural relationships of these compounds, and the reduction of  $\text{Mn}_7\text{O}_{13} \cdot 5\text{H}_2\text{O}$  to  $\gamma$ - $\text{MnO}(\text{OH})$  with cinnamyl alcohol, have been described<sup>110</sup>.

*(v) Manganates (IV)*

A large number of mixed oxides containing  $\text{Mn}^{\text{IV}}$ , often described as manganates (IV) are known, but will not be described here.

On heating a mixture of  $\text{MnO}_2 + 2\text{KOH}$ , Scholder and Protzer<sup>51</sup> obtained a mixture of  $\text{K}_3\text{MnO}_4$ ,  $\text{KMnO}_2$ , and  $\text{K}_2\text{MnO}_3$ . At  $200^\circ$  this contained 75 mole % of  $\text{K}_2\text{MnO}_3$ , whilst at  $550^\circ$  only 18 mole % was present. Similar results were obtained with  $\text{MnO}_2 + 2\text{RbOH}$ . Some  $\text{Mn}^{\text{IV}}$  always disproportionates and the authors report that the isolation of pure  $\text{K}_2\text{MnO}_3$  is not possible.

Scholder<sup>54</sup> reported the compounds  $\text{Na}_4\text{MnO}_4$ ,  $\text{Ba}_2\text{MnO}_4$ ,  $\text{Sr}_2\text{MnO}_4$ , and  $\text{Li}_2\text{MnO}_3$ , but no details were given. The crystal structure of  $\text{Ba}_3\text{MnO}_5$  was determined by Mansmann<sup>115</sup>, who found it to be isostructural with  $\text{Cs}_3\text{CoCl}_5$ .  $\text{MMnO}_3$  ( $\text{M} = \text{Sr}, \text{Ba}$ ) and substances formulated  $\text{CaO}_x\text{MnO}_2$  ( $x = \frac{1}{2}, 1, 2, 3, 5$ ) have been reported<sup>45</sup>.

A number of "mixed valence" manganates exist, and will be dealt with here for convenience.  $\text{KMnO}_2$  and  $\text{RbMnO}_2$  oxidise readily in air to substances of empirical formulae  $\text{MMnO}_{2.25}$  ( $\text{M} = \text{Rb}, \text{K}$ ). These were formulated<sup>51</sup> as  $\text{M}_4\text{Mn}_2^{\text{III}}\text{Mn}_2^{\text{IV}}\text{O}_9$ . Giovanoli et al.<sup>108</sup> found that the oxidation of fresh  $\text{Mn}(\text{OH})_2$  in aqueous  $\text{NaOH}$  with molecular oxygen produced sodium manganese (II, III) manganate (IV),  $\text{Na}_4\text{Mn}_{14}\text{O}_{27} \cdot 9\text{H}_2\text{O}$ , and determined its structure. It has a double-layer structure resembling chalcophanite. A phase  $\text{K}_2\text{Mn}_4\text{O}_{7.8-8.0}$  was observed<sup>51</sup> from the reaction of  $\text{MnO}_2$  with  $2\text{KOH}$ , which is close to the composition observed by Delano<sup>116</sup> for the compound formed from a mixture of  $4\text{K}_2\text{CO}_3 + \text{MnO}_2$ . Recent studies of the decomposition products of  $\text{KMnO}_4$  have shown<sup>28</sup> that the water-insoluble product(s) have compositions approximating to " $\text{K}_4\text{Mn}_7\text{O}_{16}$ ", which may indicate that they are related to some of the compounds discussed above. Further work is required, however, before any conclusion can be reached.

*(vi) Peroxomanganates (IV)*

By reaction of  $\text{KMnO}_4$  with  $\text{H}_2\text{O}_2$  in 30%  $\text{KOH}$  at  $-18^\circ$ , Scholder obtained  $\text{K}_2\text{H}_2\text{Mn}(\text{O})(\text{O}_2)_3$  as a dark red crystalline solid, which decomposed to  $\text{MnO}_2$  and  $\text{O}_2$  in water, to  $\text{Mn}^{\text{II}}$  and  $\text{O}_2$  in acids, and tended to explode on warming<sup>117</sup>. The occurrence of  $\text{K}_3\text{H Mn}(\text{O})(\text{O}_2)_3$  and  $\text{K}_2\text{H}_2\text{Mn}(\text{O}_2)_4$  was also postulated.

*(vii) Manganese (IV) compounds of oxygen donor ligands*

The only simple compound appears to be the black crystalline  $\text{Mn}(\text{SO}_4)_2$  obtained on oxidising  $\text{MnSO}_4$  in hot concentrated  $\text{H}_2\text{SO}_4$  with permanganate<sup>119</sup>, which hydrolyses to  $\text{MnO}_2$  even in dilute sulphuric acid. Anodic dissolution of manganese in 18–22 *N* sulphuric acid produces<sup>118</sup> a dark-coloured solution containing  $\text{Mn}^{\text{IV}}$ .

Complexes with orthoperiodate, iodate, tellurate, and rather surprisingly, oxalate, are probably the best known. Reimer and Lister<sup>120</sup> showed that the dark red crystals obtained from  $\text{MnSO}_4$  and alkali periodate solutions, are  $\text{MMnIO}_6 \cdot \frac{1}{2}\text{H}_2\text{O}$  ( $\text{M} = \text{Na}, \text{K}$ ), not  $\text{M}_2\text{Mn}_2\text{I}_2\text{O}_{11}$  as originally reported<sup>121</sup>. They have  $\mu_{\text{eff}} = 4.17 \text{ B.M. (Na)}, 3.87 \text{ B.M. (K)}$ , and are remarkably stable, being insoluble in, and not hydrolysed by water. On oxidising the mixture of  $\text{MnSO}_4$  and periodate solutions with hypochlorite, two other compounds,

$\text{Na}_7\text{H}_4\text{Mn}(\text{IO}_6)_3 \cdot 17\text{H}_2\text{O}$  and  $\text{K}_7\text{H}_4\text{Mn}(\text{IO}_6)_3 \cdot 8\text{H}_2\text{O}$  result<sup>122</sup> They are also red, but hydrolyse slowly in water to  $\text{MnO}_4^-$ ,  $\text{IO}_3^-$ , and  $\text{H}_3\text{IO}_6^{2-}$

Alkaline hypochlorite oxidises  $\text{MnSO}_4$ , mixed with an alkali metal tellurate, to dark red complexes.  $\text{K}_6\text{H}_8\text{Mn}(\text{TeO}_6)_3 \cdot 5\text{H}_2\text{O}$  ( $\mu_{\text{eff}} = 3.30 \text{ B.M.}$ ) and  $\text{Na}_7\text{H}_7\text{Mn}(\text{TeO}_6)_3 \cdot 3\text{H}_2\text{O}$ , (refs 123, 124) which are less stable in solution than the periodate analogues<sup>125</sup> The spectra and structures of a number of periodato-, and tellurato-manganese (IV) compounds have been determined<sup>123,126,127</sup>

Complex iodates,  $\text{M}_2\text{Mn}(\text{IO}_3)_6$  ( $\text{M} = \text{NH}_4, \text{K}$ ) and  $\text{BaMn}(\text{IO}_3)_6$ , are produced as brown-violet solids on boiling  $\text{MnO}_2$ , iodic acid, and the metal iodate solutions<sup>128</sup> The K salt has  $\mu_{\text{eff}} = 3.82 \text{ B.M.}$ , and dissolves in water to give a brown solution, which deposits a brown precipitate ( $\text{MnO}_2?$ ), its IR spectrum has been reported<sup>129</sup>


Dark olive-green crystals of  $\text{K}_2\text{Mn}(\text{C}_2\text{O}_4)_2(\text{OH})_2 \cdot 2\text{H}_2\text{O}$ , which on microscopic examination proves to be a mixture of orange and green crystals, possibly the *cis* and *trans* isomers, are formed<sup>130</sup> by the reaction of oxalic acid, potassium oxalate, and  $\text{KMnO}_4$  in aqueous solution at  $0^\circ$ . The crystals decompose at room temperature, and the solutions slowly turn red-brown, indicating the formation of  $\text{Mn}^{\text{III}}$ .

Other complexes include the red-brown formaldoxime  $\text{Na}_2\text{Mn}(\text{CH}_2\text{NO})_6$ , obtained on air oxidation of manganese (II) solutions containing formaldoxime<sup>131</sup>, black crystalline  $(\text{NH}_4)_2\text{H}_2\text{MnE}_2\text{O}_9$  ( $\text{E} = \text{P, As}$ )<sup>132</sup>, and the curious yellow-red glycerylmanganates (IV), e.g.,  $\text{Na}_2\text{Mn}(\text{C}_3\text{H}_5\text{O}_3)_2$ , formed on heating freshly precipitated  $\text{MnO}_2$  with glycerol and aqueous alkali<sup>133</sup>

#### (viii) Manganese (IV) complexes of N-donor ligands

These seem to be confined to biguanidine and 2,2'-bipyridyl complexes On treatment of alkaline  $\text{KMnO}_4$  with biguanidine, or by oxidising  $\text{Mn}^{\text{II}}$  and biguanidine with alkaline persulphate, bright red crystals  $[(\text{OH})_2\text{Mn}(\text{BigH})_2](\text{OH})_2$  were formed, from which the  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{C}_2\text{O}_4^{2-}$ ,  $\text{CrO}_4^{2-}$ , and  $\text{IO}_3^-$ , can be obtained by metathesis. They have unusually low magnetic moments, in the range 2.0–2.5 B.M.<sup>134</sup>

$\text{Mn}(\text{bipy})\text{Cl}_4$  which is formed<sup>135</sup> as black crystals from the reaction of 2,2'-bipyridyl with  $\text{KMnO}_4$  in concentrated  $\text{HCl}$ , loses chlorine on standing, and obeys the Curie–Weiss law with  $\mu_{\text{eff}} = 3.82 \text{ B.M.}$  and  $\theta = 8^\circ$  (ref. 136) The *o*-phenanthroline analogue has not been obtained<sup>135,137</sup>, but red-brown  $[\text{MnO}(\text{phen})_2](\text{ClO}_4)_2 \cdot \frac{1}{2}\text{H}_2\text{O}$  forms on treating  $\text{MnCl}_3(\text{phen})\text{H}_2\text{O}$  mixed with *o*-phenanthroline, with  $\text{HClO}_4$ .<sup>136</sup> It is antiferromagnetic, and may be an oxygen-bridged dimer, viz.



although more highly condensed structures are also possible. Some N-donor complexes which contain both  $\text{Mn}^{\text{IV}}$  and  $\text{Mn}^{\text{III}}$ , are discussed under  $\text{Mn}^{\text{III}}$ .

*(ix)  $K_2Mn(CN)_6$* 

Nitrosyl chloride oxidises  $K_3Mn(CN)_6$  in DMF to canary yellow  $K_2Mn(CN)_6$ ,  $\mu_{\text{eff}} = 3.94 \text{ B.M.}$ , which is instantly decomposed by water<sup>138</sup>. Yakimach<sup>139</sup> claimed to have obtained  $K_4Mn(CN)_8$  by reaction of  $KMnO_4$  with alkaline KCN, but this was denied by Goldenberg<sup>140</sup>, who could not repeat the preparation, and obtained a manganese (III) cyanocomplex contaminated with KOH. The  $KMn^{II}Mn^{III}(CN)_6$  complex is similarly obtained<sup>138</sup> from NOCl and  $K_2Mn^{II}Mn^{II}(CN)_6$ .

*(x) Heteropolyanions*

Manganese (IV), like a number of other transition metal higher oxidation states, is stabilised by incorporation into a heteropolyanion. The 9-molybdomanganates (IV), e.g.  $(NH_4)_6MnMo_9O_{32} \cdot 8H_2O$  (ref. 141) have been known for some time, and recently 12-niobomanganates (IV)<sup>142,143</sup> have been obtained, and the structure of  $Na_{12}MnNb_{12}O_{38} \cdot 50H_2O$  determined<sup>143</sup>. Flynn and Stucky<sup>144,145</sup> have described three types of vanadomanganate (IV) —  $K_5MnV_{11}O_{32} \cdot 10H_2O$ ,  $K_5HMn_3V_{12}O_{39} \cdot 10H_2O$ , and  $K_7MnV_{13}O_{38} \cdot 16-18H_2O$ .

## G MANGANESE (III)

$Mn^{III}$  has a far more extensive chemistry than any of the other oxidation states discussed. There are relatively few simple compounds, but  $Mn^{III}$  is fairly stable in complexes, almost all of which are anionic, e.g. fluoro, chloro, cyano, or neutral  $\beta$ -diketonates; cationic species are rare. The complexes with oxygen donor ligands have been known for many years, but complexes with nitrogen donors have only recently been characterised. Sidgwick<sup>1</sup>, for example, stated in 1950 that "trivalent manganese has no tendency to coordinate with nitrogen to form amines, nitrocomplexes, or in other ways". There is an almost total absence of complexes of the heavier Group V donors.

*(i) Halides*

A red-purple trifluoride, and a very unstable trichloride are known.  $MnF_3$  is prepared<sup>146-149</sup> by fluorination of  $MnF_2$ ,  $MnCl_2$ , various manganese oxides, or, best,  $MnI_2$ , or by dissolving  $Mn(IO_3)_2$  in  $BrF_3$  and evaporating at  $140^\circ$  (ref. 71, 150). Ruby-red crystals of the hydrate  $MnF_3 \cdot 2H_2O$  are obtained on dissolving  $Mn_2O_3$  in aqueous HF, or by oxidising  $Mn^{II}$  in HF solution with  $KMnO_4$ , or electrolytically<sup>151,152</sup>. The anhydrous compound is moisture-sensitive, but the hydrate is not hydrolysed in water in the presence of HF (ref. 151).  $MnF_3$  liberates fluorine on heating, although the dissociation pressure is less than 0.1 atm at  $600^\circ$  (ref. 153) and has been extensively studied as a fluorinating agent, especially in organic chemistry<sup>147,154</sup>. The structure is most unusual in that there are three different Mn—F distances. The lattice is monoclinic,  $a = 8.509$ ,  $b = 5.037$ ,  $c = 13.448 \text{ \AA}$ , and the structure consists<sup>150</sup> of distorted  $MnF_6$  octahedra sharing corners with Mn—F 2.09, 1.92, 1.79  $\text{\AA}$  and Mn—F—Mn (ave.)  $146^\circ$ . The distortion has been discussed in terms of crystal field theory and the Jahn—Teller effect by Hepworth et al.<sup>155</sup>. Mass

spectra<sup>156</sup> show that  $\text{MnF}_3$  evaporates principally as the monomer, whilst the value for the heat of formation ( $-238 \text{ kcal. mole}^{-1}$ ) coupled with the heat of sublimation ( $68.0 \pm 3 \text{ kcal. mole}^{-1}$  at  $298^\circ \text{K}$ ) leads to a value for the average  $\text{Mn}-\text{F}$  bond energy of  $97.9 \pm 4 \text{ kcal. mole}^{-1}$ . The bond dissociation energies are  $D(\text{Mn}-\text{F}) = 101$ ,  $D(\text{FMn}-\text{F}) = 119$ ,  $D(\text{F}_2\text{Mn}-\text{F}) = 74 \text{ kcal. mole}^{-1}$ .  $\text{MnF}_3$  obeys the Curie-Weiss law with  $\mu_{\text{eff}} = 5.0 \text{ B.M.}$  and  $\theta = 8^\circ$ , and becomes antiferromagnetic at very low temperatures<sup>157</sup>. An adduct  $\text{MnF}_3 \cdot 4\text{XeF}_6$ , or possibly  $\text{MnF}_4 \cdot 4\text{XeF}_6$ , is formed on heating manganese with xenon and excess fluorine under pressure<sup>158</sup>.

A black solid, analysing as  $\text{MnCl}_3$  is formed<sup>70</sup> on suspending  $\text{MnO}_2$  in dry ether at  $-78^\circ$ , saturating with dry  $\text{HCl}$ , and precipitating with  $\text{CCl}_4$ . The action of liquid  $\text{HCl}$  on  $\text{Mn}(\text{OAc})_3$  at  $-100^\circ$  produces brown crystalline  $\text{MnCl}_3$ , soluble in organic solvents to green solutions, and readily losing chlorine above about  $-35^\circ$  (ref. 159).

#### (ii) Complex halides

$\text{Mn}^{\text{III}}$  forms a number of complex anions with the halogens, viz  $\text{MnF}_6^{3-}$ ,  $\text{MnF}_5^{2-}$ ,  $\text{MnF}_4^-$ ,  $\text{MnCl}_6^{3-}$ , and  $\text{MnCl}_5^{2-}$  but all attempts to prepare bromocomplexes have failed.

$\text{K}_3\text{MnF}_6$  and  $\text{K}_2\text{NaMnF}_6$  are formed on fusing  $\text{K}_2\text{MnF}_5 \cdot \text{H}_2\text{O}$  with  $\text{MHF}_2$  ( $\text{M} = \text{K}, \text{Na}$ )<sup>160,162</sup> and  $\text{Cs}_2\text{KMnF}_6$  has recently been obtained by direct fluorination<sup>161</sup>.  $\text{K}_3\text{MnF}_6$  dissolves in aqueous  $\text{HF}$  to reform the  $\text{MnF}_5 \cdot \text{H}_2\text{O}^{2-}$  ion, and is completely hydrolysed by water.  $[\text{M}(\text{NH}_3)_6]\text{MnF}_6$  ( $\text{M} = \text{Co}, \text{Cr}, \text{Rh}$ ) are obtained on oxidising  $\text{Mn}^{\text{II}}$  in 40%  $\text{HF}$  with  $\text{KMnO}_4$  in the presence of the cation and  $\text{NH}_4\text{F}$  (ref. 163). The IR spectra of these compounds have been examined, and discussed in terms of the distorted  $\text{MnF}_6^{3-}$  ion and the Jahn-Teller effect<sup>163</sup>.

Pink crystals of  $\text{K}_2\text{MnF}_5 \cdot \text{H}_2\text{O}$ , which are hydrolysed by water, form<sup>82,152</sup> on oxidising  $\text{MnF}_2$  in 40%  $\text{HF}$  followed by addition of  $\text{KHF}_2$ .  $\text{Cs}_2\text{MnF}_5 \cdot \text{H}_2\text{O}$  was crystallised from mixed solutions of  $\text{CsF}$  and  $\text{MnF}_3$  in aqueous  $\text{HF}$ , whilst  $\text{M}_2\text{MnF}_5$  ( $\text{M} = \text{Li}, \text{Na}, \text{NH}_4$ ) are formed<sup>166,167</sup> on adding  $\text{MHF}_2$  to  $\text{MnO}(\text{OH})$  in 20%  $\text{HF}$ . A substance which is probably  $\text{BaMnF}_5$  is formed<sup>79</sup> on hydrogen reduction of  $\text{BaMnF}_6$  at  $400^\circ$ . The " $\text{MnF}_5$ " group in  $(\text{NH}_4)_2\text{MnF}_5$  is actually a distorted  $\text{MnF}_6$  octahedron,  $\text{Mn}-\text{F} = 1.84, 1.85, 2.12 \text{ \AA}$ , the octahedra being linked into infinite chains<sup>167</sup>. (Note added in proof. a crystal structure of  $\text{K}_2\text{MnF}_5 \cdot \text{H}_2\text{O}$  has been determined by X-ray crystallography. The water molecule is not coordinated, and bridging fluorine atoms complete a distorted octahedron about the manganese. A. J. Edwards, *J. Chem. Soc. A*, (1971) 2653.)

The brown-violet  $\text{MMnF}_4$  ( $\text{M} = \text{Li}, \text{K}, \text{Rb}$ ) are formed<sup>73,74</sup> by hydrogen reduction of the corresponding  $\text{MMnF}_5$  at  $150-250^\circ$ .

Hexachloromanganates (III) are only found with large cations. Hatfield et al.<sup>168</sup> prepared  $[\text{M}(\text{pn})_3]\text{MnCl}_6$  ( $\text{M} = \text{Co}, \text{Rh}$ ) by oxidation of  $\text{MnSO}_4$  in concentrated hydrochloric acid with  $\text{NaClO}_3$  in the presence of  $[\text{M}(\text{pn})_3]\text{Cl}_3$ . The Co compound, and the less stable  $[\text{Co}(\text{en})_3]\text{MnCl}_6 \cdot 2\text{H}_2\text{O}$  may be obtained using  $\text{KMnO}_4$  as the oxidising agent<sup>169</sup>. These compounds are instantly hydrolysed in water, decompose on keeping, and show rather high  $\nu(\text{Mn}-\text{Cl})$  frequencies compared with other hexachlorometallates (III)<sup>93</sup>.

$(\text{NEt}_4)_2\text{MnCl}_5$  is formed by reaction of  $\text{MnO}_2$  with acetyl chloride in ether<sup>171</sup>, or by saturating  $\text{MnO}_2$  suspended in  $\text{CCl}_4$  with dry  $\text{HCl}$  and ether-extracting the black residue<sup>170</sup>. In both cases, addition of  $\text{NEt}_4\text{Cl}$  precipitates the dark green pentachloromanganate. Addition of the ligand to the dark brown solution of  $\text{KMnO}_4$  in concentrated  $\text{HCl}$  precipitates



TABLE 8

Complex halides of manganese (III)

Compound	Colour	Structural data	Magnetic moment (B M)	Ref
$K_3MnF_6$	Blue-violet	Tetragonal, $a = 8.75$ $c = 8.30 \text{ \AA}$	4.95	160
$K_2NaMnF_6$		Tetragonal, $a = 8.171$ $c = 8.577$		162
$Cs_2KMnF_6$	Violet	Tetragonal, $a = 8.93$ $c = 9.26$	4.88	161
$(NH_4)_3MnF_6$	Reddish			164
$[Co(NH_3)_6]MnF_6$	Orange-brown	Cubic, $a = 9.97$	4.94	163
$[Cr(NH_3)_6]MnF_6$	Golden-brown	Cubic, $a = 10.06$		163
$[Rh(NH_3)_6]MnF_6$	Violet	Cubic, $a = 10.05$		163
$K_2MnF_5 \cdot H_2O$	Pink		3.32	82, 152, 165
$Na_2MnF_5$	Pink	Orthorhombic, $a = 6.08$ $b = 7.86$ $c = 9.28$		167
$Li_2MnF_5$	Pink	Orthorhombic, $a = 5.46$ $b = 7.78$ $c = 8.81$		167
$(NH_4)_2MnF_5$	Pink	Orthorhombic, $a = 6.20$ $b = 7.94$ $c = 10.72$		167, 167a
$RbMnF_4$	Brown-violet		5.0	74
$KMnF_4$	Brown-violet		4.9	74
$LiMnF_4$	Brown		4.7	74
$[Co(pn)_3]MnCl_5$	Dark brown		4.91	168
$[Co(en)_3]MnCl_5 \cdot 2H_2O$	Dark brown		4.94	169
$(NEt_4)_2MnCl_5$	Dark green		5.0	170, 171
$(phen)_2MnCl_5$	Green-black		5.0	135, 136
$(bipy)_2MnCl_5$	Green-black	$a = 13.204$ $b = 7.103$ $c = 15.339$	4.88	135, 136, 173

the green-black  $(phen)_2MnCl_5$  and  $(bipy)_2MnCl_5$  (ref 135). The structure of the latter has been determined, it contains discrete  $MnCl_5^{2-}$  ions which are square pyramidal with a distortion towards trigonal bipyramidal,  $Mn-Cl = 2.53, 2.34 \text{ \AA}$  (ref 173). The alkali metal analogues have been described, but there is some doubt as to whether they are  $M_2MnCl_5$  or  $M_2MnCl_5 \cdot H_2O$ , and they would repay further study<sup>90,172</sup>

### (iii) Oxide

$\alpha-Mn_2O_3$  is best prepared by decomposing manganese (II) nitrate at  $150-200^\circ$ , and then heating the product to constant weight at  $600^\circ$ , above this temperature the product obtained is not completely stoichiometric<sup>98,105</sup>. It is the normal product of heating many manganese compounds in air between  $600-800^\circ$ , but above  $900^\circ$  oxygen is lost to form  $Mn_3O_4$ . The structure is body-centred cubic,  $a = 9.401 \text{ \AA}$  (c-sesquioxide type)<sup>174</sup>

$\gamma\text{-Mn}_2\text{O}_3$  was reported as a black powder obtained by careful dehydration of  $\gamma\text{-MnO(OH)}$  (ref 175), but recent work indicates that the  $\gamma\text{-Mn}_2\text{O}_3$  phase exists only in the range  $\text{MnO}_{1.33-1.40}$  and readily changes into  $\alpha\text{-Mn}_2\text{O}_3$  on heating<sup>106</sup>. The X-ray diffraction pattern is said to be similar to that of  $\text{Mn}_3\text{O}_4$  to which it is related in the same way as  $\gamma\text{-Fe}_2\text{O}_3$  is to  $\text{Fe}_3\text{O}_4$ .

There is much confusion in the literature about the nature of the various hydrated oxides of  $\text{Mn}^{\text{III}}$ . There is no good evidence for  $\text{Mn(OH)}_3$ , but two forms of  $\text{MnO(OH)}$  seem well-established  $\alpha\text{-MnO(OH)}$  (manganite) and  $\gamma\text{-MnO(OH)}$  (groutite) are both formed by the oxidation of  $\text{Mn(OH)}_2$  under carefully controlled conditions<sup>100</sup>. This reaction can also yield  $\text{MnO}_2$  (various forms), manganates (III), manganates (IV), and a number of other products, so the confusion in the literature is not surprising! A  $\beta\text{-MnO(OH)}$  was also reported to be formed in this reaction<sup>100,177</sup> but this is now in doubt<sup>106</sup>. Both  $\alpha$ - and  $\gamma\text{-MnO(OH)}$  contain  $\text{Mn}^{\text{III}}$  surrounded octahedrally by oxygen atoms, with all the latter involved in hydrogen bonding<sup>176</sup>. The difference in structures is in the extent of the distortion of the octahedron, in  $\alpha\text{-MnO(OH)}$  this is quite small with average  $\text{Mn-O} = 1.89 \text{ \AA}$ , whilst in  $\gamma\text{-MnO(OH)}$  there are four oxygens at  $1.88 \text{ \AA}$ , and two at  $2.30 \text{ \AA}$ , a much more distorted structure<sup>176</sup>.

#### (iv) Manganates (III)

The large number of complex oxides will not be discussed. Scholder and Kyri<sup>178</sup> found that  $\text{Mn(OH)}_2$  in 50%  $\text{NaOH}$  is oxidised to  $\text{NaMnO}_2$ , also obtained on heating  $\text{Mn}_2\text{O}_3$  and  $\text{Na}_2\text{CO}_3$  in air at  $1000^\circ$ , or along with the  $\text{K}$ ,  $\text{Rb}$  analogues by heating a mixture of  $\text{Mn}_2\text{O}_3$  and  $\text{MOH}$  in a 1:2 ratio in nitrogen<sup>51</sup>.  $\text{LiMnO}_2$  is formed by fusing the constituent oxides in argon<sup>179</sup>. Scholder and Kyri<sup>178</sup> also reported  $\text{Na}_3\text{Mn(OH)}_6 \cdot 2.4\text{H}_2\text{O}$ ,  $\text{Na}_4\text{Mn(OH)}_7 \cdot 5\text{H}_2\text{O}$ , and the  $\text{Ba}$  and  $\text{Sr}$  salts of the ions  $\text{Mn(OH)}_5^{2-}$ ,  $\text{Mn(OH)}_6^{3-}$ , and  $\text{Mn(OH)}_7^{4-}$ .

$\text{M}_2\text{Mn}_4\text{O}_7$  ( $\text{M} = \text{K}, \text{Rb}$ ) are formed on heating 1:1 mixtures of  $\text{Mn}_2\text{O}_3$  and  $\text{MOH}$ <sup>51</sup>.

#### (v) Manganese (III) compounds with oxygen donor ligands

Green  $\text{Mn}_2(\text{SO}_4)_3$  is formed on dissolving  $\text{KMnO}_4$ ,  $\text{MnO}_2$ , or  $\text{Mn}_2\text{O}_3$  in hot concentrated sulphuric acid, and drying the product at  $130^\circ$ . It dissolves in 70%  $\text{H}_2\text{SO}_4$  and on cooling red  $\text{HMn(SO}_4)_2 \cdot 2\text{H}_2\text{O}$  crystallises, but in more dilute acid, hydrolysis occurs<sup>180,181</sup>. Violet solutions of  $\text{Mn}^{\text{III}}$  are formed by electrolysis of  $\text{MnSO}_4$  in sulphuric acid solution<sup>182</sup>. Alums are formed with alkali sulphates, there are double sulphates with  $\text{Al}$ ,  $\text{Cr}^{\text{III}}$  and  $\text{Fe}^{\text{III}}$ .  $\text{CsMn(SO}_4)_2 \cdot 12\text{H}_2\text{O}$  is ruby-red, melts at  $40^\circ$ , and has  $\mu_{\text{eff}} = 4.9\text{ B.M.}$ , the  $\text{Rb}$ ,  $\text{K}$ ,  $\text{NH}_4$  analogues are known<sup>183,184</sup> but are less stable, and all are readily hydrolysed.


Grey-green  $\text{MnPO}_4 \cdot \text{H}_2\text{O}$  precipitates on oxidising  $\text{Mn}^{\text{II}}$  in phosphoric acid solution, but redissolves in concentrated  $\text{H}_3\text{PO}_4$  to form a violet solution, said by Meyer and Marek<sup>185</sup> to contain  $\text{H}_3\text{Mn(PO}_4)_2$ , from which double salts with alkali phosphates, e.g.  $\text{Na}_2\text{HMn(PO}_4)_2 \cdot 2\text{H}_2\text{O}$ , can be isolated. On dilution of the violet solution  $\text{MnPO}_4 \cdot \text{H}_2\text{O}$  is precipitated. Several  $\text{Mn}^{\text{III}}$  salts of the condensed phosphoric acids are known<sup>45</sup>.

Manganese (III) acetate is one of the easiest  $\text{Mn}^{\text{III}}$  compounds to obtain, and is a convenient starting material for the synthesis of several others. The cinnamon-brown anhydrous compound is produced in the violent reaction between  $\text{Mn(NO}_3)_6 \cdot 6\text{H}_2\text{O}$  and acetic

anhydride<sup>159</sup>, and the dihydrate by oxidation of the manganese (II) compound in glacial acetic acid with chlorine or  $\text{KMnO}_4$  (ref 186)


Cartledge and Eriks<sup>187</sup> prepared the trisoxalato and bisoxalato complexes of  $\text{Mn}^{\text{III}}$  by oxidation of the corresponding  $\text{Mn}^{\text{II}}$  complexes with  $\text{KMnO}_4$ . When care was taken to isolate a pure product, the trisoxalatomanganate (III) could be stored at  $20^\circ$  in the absence of light with little decomposition, but the bisoxalatomanganate (III) was less stable, although it kept fairly well at  $0^\circ$

Meyer and Schramm prepared several malonate complexes but it appears that these were impure in the majority of cases<sup>188</sup> Bullock et al.<sup>189</sup> have obtained and studied a large series of malonate complexes  $\text{M}[\text{MnL}_2(\text{H}_2\text{O})_2]$  ( $\text{M} = \text{Cs}, \text{Ti}, \text{NH}_4, \text{Rb}, \text{NMe}_4, \text{NEt}_4$ ),  $\text{M}[\text{MnL}_2(\text{H}_2\text{O})_2] \cdot 2\text{H}_2\text{O}$  ( $\text{M} = \text{Na}, \text{K}, \text{C}_5\text{H}_6\text{N}$ ),  $\text{M}[\text{MnL}_2(\text{H}_2\text{O})_2] \cdot 3\text{H}_2\text{O}$  ( $\text{M} = \text{Li}$ ),  $\text{M}[\text{MnL}_2]$  ( $\text{M} = \text{Na}, \text{K}$ ),  $\text{M}_3[\text{MnL}_3] \cdot 3\text{H}_2\text{O}$  ( $\text{M} = \text{K}$ ),  $\text{M}_3[\text{MnL}_3] \cdot \text{H}_2\text{O}$  ( $\text{M} = \text{K}$ ), and  $[\text{MnL}_2(\text{H}_2\text{O})_2]$ . The effective magnetic moment for these complexes lay between 4.90 and 5.03 B.M., and the visible spectra exhibited two absorption bands at  $22,000 \text{ cm}^{-1}$  and  $13,000 - 16,000 \text{ cm}^{-1}$ . Cartledge and Nichols<sup>190</sup> calculated the equilibrium constant for the reaction



and obtained an average value of  $K = 0.057$  at  $0^\circ$

The solid EDTA complex of  $\text{Mn}^{\text{III}}$  was characterised by Yoshino et al.<sup>191</sup> who found it was decomposed by light and heat, and was unstable at room temperature. They noted that in solution two forms existed depending on the pH of the solution, and suggested an equilibrium




which leads to the conclusion that either EDTA is acting as a quinque-dentate ligand, or that  $\text{Mn}^{\text{III}}$  is capable of forming seven-coordinate complexes. The idea of seven-coordination is reasonable in the light of the crystallographic establishment of the  $\text{Mn}^{\text{II}}-\text{EDTA}$  complex as seven-coordinate<sup>192</sup>

The complexes  $[\text{Mn}(\text{bipyO}_2)_3](\text{ClO}_4)_3 \cdot 3\text{H}_2\text{O}$ ,  $\mu_{\text{eff}} = 4.97 \text{ B.M.}$ <sup>193</sup>, and  $[\text{Mn}(\text{terpyO}_3)_2](\text{ClO}_4)_3$  (ref 194) have been obtained. The latter contains  $\text{terpyO}_3$  behaving as a tridentate ligand.

#### (vi) $\beta$ -ketoenolates

Fackler<sup>195</sup> has reviewed the literature up to 1965. Fernellus and Bryant<sup>196</sup> have described the preparation of  $\text{Mn}(\text{acac})_3$ , and the synthetic procedures have also been reviewed by Charles<sup>197</sup>. Brown-black  $\text{Mn}(\text{acac})_3$ , which is thermally less stable than the Cr, Ni, or Fe analogues, is most conveniently obtained by oxidising  $\text{Mn}^{\text{II}}$  with  $\text{KMnO}_4$  in the presence of acetylacetone<sup>198</sup>. The preparations of  $\text{Mn}^{\text{III}}$  complexes of diisobutylmethane<sup>199</sup>, dipivaloylmethane<sup>199</sup>, trifluoroacetylacetone<sup>200,201</sup>, 1-phenyl-1,3-butanedione<sup>201</sup>, hexafluoroacetylacetone<sup>202</sup>, and 3-cyano-2,4-pentanedione<sup>202</sup> have been described.

The standard heat of formation of  $\text{Mn}(\text{acac})_3$  at  $25^\circ$  has been determined by reaction calorimetry<sup>203</sup> to be  $-332.1 \text{ kcal mole}^{-1}$ . Enthalpy changes at  $25^\circ$  for the hypothetical gaseous reactions



have been derived and thus the heterolytic bond energy = 246 kcal. mole<sup>-1</sup>, and the homolytic bond energy = 44 kcal mole<sup>-1</sup> for the Mn—O bond in Mn(acac)<sub>3</sub>. The homolytic bond energy appears to provide the better basis for the comparison of M<sup>III</sup>—O bond energies for the first transition series acetylacetonates<sup>203</sup>.

Despite the high-spin *d*<sup>4</sup> electronic configuration of Mn<sup>III</sup> in Mn(acac)<sub>3</sub>, the room-temperature X-ray crystal structure indicates an almost regular octahedral arrangement about the metal<sup>204</sup>. The distortion from *D*<sub>3</sub> symmetry is slight, but real, with Mn—O = 1.88 Å (deviation 0.03 Å). Forman and Orgel<sup>205</sup> have suggested that, because of the unusual infrared spectrum<sup>206</sup>, Mn(acac)<sub>3</sub> is Jahn—Teller distorted. Morosin and Brathovde<sup>204</sup> suggest that, because there are two types of O—Mn—O bond angles (96.9° and 87.8°) in the complex, the bands being observed include some coupling to an Mn—O bond. On the other hand, the crystal structure of acetylacetonatobis(*N*-phenylaminotroponiminato)manganese (III) has revealed<sup>207</sup> the first case of a distorted octahedral complex of high-spin Mn<sup>III</sup>, where the distortion occurs because of different Mn—ligand bond lengths and not Mn—ligand bond angles. The Mn—O bond in the mixed-ligand complex, 1.96 Å, is significantly longer than in the acetylacetonate, this may be due to the presence of N donors in the complex. The distortion takes the form of four short bonds and two long bonds, and Bartlett and Palenik<sup>207</sup> have suggested an empirical rule for predicting the final configuration of octahedral Mn<sup>III</sup> complexes.

Barnum has carried out some Huckel LCAO—MO calculations on Mn(acac)<sub>3</sub> and discussed the metal—ligand  $\pi$ -bonding<sup>208</sup>. Piper and Carlin<sup>209</sup> have studied the polarised visible spectrum of Al(acac)<sub>3</sub> isomorphously replaced in part with Mn<sup>3+</sup>. Electronic energy levels were assigned and interpretations of the spectral intensities and estimates of octahedral and trigonal field strengths were presented. The trigonal field parameter, *K*, is 500 cm<sup>-1</sup>.

When Mn(acac)<sub>3</sub> is subjected to various acids in water, salts of type [Mn(acac)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>]<sup>+</sup> are formed<sup>210</sup>. Manganese can be extracted quantitatively from alkaline peroxide solutions with acetylacetone and chloroform<sup>211</sup>.


#### (*vu*) Schiff base complexes

There are not many reported investigations of Schiff base complexes of Mn<sup>III</sup>. Early work by Pfeiffer et al.<sup>212</sup> and Tsumaki<sup>213</sup> on hydroxy compounds of the type Mn(salen)(OH) and Mn(sal-NR)<sub>2</sub>OH, has recently been reinvestigated<sup>214,216</sup>. Lewis et al.<sup>216</sup> obtained two modifications of Mn(salen)(OH), one of which showed antiferromagnetic behaviour, possibly arising from an oxygen-bridged structure of the type [Mn<sub>2</sub>(salen)<sub>2</sub>O]·H<sub>2</sub>O. Derivatives of the type Mn(salen)X (X = halogen, acetate, etc.) have recently been studied and all display Curie—Weiss magnetic behaviour with small  $\theta$  values<sup>214</sup>. Mn(salen) is oxidised by nitric oxide in ethanol to an Mn<sup>III</sup> derivative<sup>214</sup>, and it has been shown that three different complexes containing Mn<sup>III</sup> and Mn<sup>IV</sup> can be formed on oxidation under various conditions<sup>215</sup>. The six-coordinate [Mn(TS<sub>2</sub>)]·H<sub>2</sub>O (TS<sub>2</sub> = bis-salicylaldehydetriethylenetetramine) was prepared<sup>217</sup> by reacting bis-salicylaldehyde-Mn<sup>II</sup> with triethylenetetramine,

concentrated HCl, and KI, the magnetic moment is normal (5.05 B.M.). The complexes  $\text{Mn}(\text{sal-NR})_3$  and  $\text{Mn}(\text{sal-NR})_2\text{X}$  ( $\text{X} = \text{Cl}, \text{Br}, \text{OAc}$ ) show high-spin magnetic behaviour ( $\mu_{\text{eff}} = 4.80\text{--}5.04$  B.M.) with small deviations from the expected susceptibilities according to the Curie law<sup>218</sup>. The Curie–Weiss behaviour of the  $\text{Mn}(\text{salen})\text{X}$  ( $\text{X} = \text{Br}, \text{I}$ ) has been explained in terms of very weak antiferromagnetic interactions<sup>218</sup>.

(viii) *Manganese (III) complexes with nitrogen-donor ligands*

Persulphate oxidation of  $[\text{Mn}(\text{bipy})_3]^{2+}$  produces a dark green complex<sup>219</sup>, characterised as  $[\text{Mn}(\text{bipy})_2\text{O}]_2(\text{S}_2\text{O}_8)_{1.5} \cdot 3\text{H}_2\text{O}$  by Turco and Nyholm<sup>220</sup> who also obtained  $[\text{Mn}(\text{bipy})_2\text{O}]_2(\text{ClO}_4)_3 \cdot 2\text{H}_2\text{O}$  by metathesis with  $\text{NaClO}_4$ . The effective magnetic moment is 1.7 B.M. indicating an average oxidation state of  $\text{Mn}_{3.5}$ , i.e. an  $\text{Mn}^{\text{III}}$ - and  $\text{Mn}^{\text{IV}}$ -containing molecule, for which the structure



was postulated


Goodwin and Sylva<sup>135</sup> found that *o*-phenanthroline and 2,2'-bipyridyl reacted with  $\text{KMnO}_4$  in concentrated HCl, to produce pentachloromanganates (III),  $[\text{bipyH}_2][\text{MnCl}_5]$  and  $[\text{phenH}_2][\text{MnCl}_5]$ . Solutions of lower acidity produced  $[\text{Mn}(\text{bipy})(\text{H}_2\text{O})\text{Cl}_3]$  ( $\mu_{\text{eff}} = 4.9$  B.M.) and  $[\text{Mn}(\text{phen})(\text{H}_2\text{O})\text{Cl}_3]$  ( $\mu_{\text{eff}} = 4.9$  B.M.).  $[\text{Mn}(\text{bipy})\text{Cl}_4]$  decomposes on heating to produce  $[\text{Mn}(\text{bipy})\text{Cl}_3]_2$ , a chlorine-bridged dimer, with the rather low magnetic moment of 3.9 B.M., which can be obtained directly, along with  $[\text{Mn}(\text{phen})\text{Cl}_3]_2$ , from the product of the interaction of  $\text{MnO}_2$  and dry HCl in  $\text{CCl}_4$  with either of these bidentate ligands<sup>135</sup>. The same workers<sup>136</sup> also prepared  $[\text{Mn}(\text{phen})(\text{H}_2\text{O})\text{F}_3]$  and  $[\text{Mn}(\text{terpy})\text{Cl}_3]$  and studied the magnetic properties of a number of these complexes over the range  $\sim 100\text{--}300^\circ\text{K}$ . The compounds  $[\text{MnL}(\text{H}_2\text{O})\text{X}_3]$  ( $\text{L} = \text{phen}, \text{bipy}, \text{X} = \text{F}, \text{Cl}$ ) are simple paramagnetics, but  $[\text{MnLCl}_3]$  are antiferromagnetic, in accordance with the dimeric, or possibly more highly polymeric, structures proposed. The tridentate ligand terpy forms  $[\text{MnLCl}_3]$  ( $\text{L} = \text{terpy}$ ) which is a normal, monomeric octahedral complex<sup>136</sup>.

Funk and Kreis<sup>221</sup> reacted  $\text{MnCl}_3$  with some simple amines and obtained mainly 1:3 adducts. The compounds were moisture-sensitive, but stable at room temperature, in contrast to the simple trichloride. With 2,2'-bipyridyl and *o*-phenanthroline, water-soluble complexes resulted in which the chloride could be substituted by other ions; similar addition compounds with ethers decomposed at room temperature. Reaction of  $\text{MnCl}_3$  with acetylacetone caused substitution of one or two chlorides by the diketone. Among the complexes isolated were  $\text{MnLCl}_3$  ( $\text{L} = \text{bipy}, \text{phen}, \text{py}$ ),  $\text{MnL}_2\text{Cl}_3$  ( $\text{L} = \text{en}, \text{dioxan}$ ), and  $\text{MnL}_3\text{Cl}_3$  ( $\text{L} = \text{NH}_3, \text{MeNH}_2, \text{EtNH}_2, \text{PrNH}_2, \text{py}$ ),  $\text{MnCl}_3 \cdot 4\text{THF}$ , and  $\text{Mn}(\text{acac})\text{Cl}_2$  and  $\text{Mn}(\text{acac})_2\text{Cl}$ . Unfortunately, no measurements were carried out on this interesting series of complexes.

Ray and coworkers have prepared a number of interesting  $\text{Mn}^{\text{III}}$  complexes containing the biguanide ligand. The compound  $[\text{Mn}_2(\text{big})_4(\text{OH})_2] \cdot \text{H}_2\text{O}$  becomes anhydrous on heating to  $90^\circ$  (ref. 222).  $\text{Mn}(\text{acac})_3$  reacts with biguanidine to yield<sup>223</sup>  $[\text{Mn}(\text{acac})(\text{big})(\text{OH})(\text{H}_2\text{O})]$ , which reacts further with  $\text{H}_2\text{O}$  to produce  $[\text{Mn}(\text{acac})(\text{bigH})(\text{H}_2\text{O})]\text{OH}$ . The mixed complex  $[\text{Mn}(\text{big})_2(\text{benzoylacac})]$  has also been produced<sup>224</sup>. These workers

have also prepared  $[\text{Mn}(\text{pic})_3]$  ( $\mu_{\text{eff}} = 4.9 \text{ B.M.}$ ) and  $[\text{Mn}(\text{oxin})_3]$  ( $\mu_{\text{eff}} = 4.8 \text{ B.M.}$ ) (pic = anion of picolinic acid, oxin = anion of  $\beta$ -hydroxyquinoline). The latter complex reacts with bromine<sup>225</sup> to form  $[\text{Mn}(\text{S},7\text{-dibromoxin})_3]$

Elvidge and Lever<sup>226</sup> found that the oxidation of pyridine solutions of phthalocyanato-manganese (II) with molecular oxygen resulted in precipitation of short, opaque crystals which they formulated as  $\text{Mn}^{\text{IV}}\text{Pc py.O}$  (Pc = phthalocyanato). An X-ray structure of this compound<sup>227</sup> showed it to be the more complicated phthalocyanatopyridinemanganese-(III)- $\mu$ -oxophthalocyanatopyridinemanganese (III). This novel molecule consists of two approximately flat and parallel manganese phthalocyanine ring systems, joined by an oxygen atom which is midway between the manganese atoms. Each Mn also has a pyridine



(Reproduced by permission of the American Chemical Society )

molecule coordinated opposite to the oxygen atom. The crystals also contain molecules of pyridine of crystallisation.

Manganese has been implicated in oxygen formation which occurs during green plant photosynthesis<sup>228</sup>. In fact, two Mn ions are found per photosynthetic unit in the chloroplast<sup>229</sup>. Several suggestions have been made as to the function of the Mn ion in the liberation of  $\text{O}_2$  by photosystem II. In these schemes the  $\text{Mn}^{\text{II}}-\text{Mn}^{\text{III}}-\text{Mn}^{\text{IV}}$  redox system is thought to take part in the electron-transfer chain from an electron donor to an electron acceptor<sup>230</sup>. Unfortunately, the metal atom binding sites and the ligands at these sites are not known. Calvin<sup>231</sup> has suggested that a relevant model for the Mn in the chloroplast would be Mn complexes of porphyrin-like ligands. The photochemical behaviour of phthalocyanine<sup>232</sup>, porphyrin<sup>233</sup>, and chlorophyll<sup>234</sup> complexes of Mn have been extensively investigated. In a detailed study of the photochemistry of phthalocyanine-


manganese, it has been shown that the stable oxidation level of manganese may be shifted among the (II), (III), and (IV) oxidation states, depending on the nature of the fifth and sixth coordinating groups<sup>235</sup>. Furthermore, photochemical oxidation as well as photochemical reduction of the phthalocyaninemanganese (III) has been observed, and photochemical reduction of the manganese (IV) compound demonstrated. Mn<sup>III</sup> etioporphyrins also are reduced to the (II)-oxidation state<sup>235</sup>. Some halide complexes of Mn<sup>III</sup> protoporphyrin IX dimethyl ester have been prepared and studied<sup>236</sup>. A manganese (III) porphyrin was first prepared by Zaleski<sup>237</sup>, and this mesoporphyrin IX was later studied by Taylor who examined its redox properties<sup>238</sup>. Also, the acetatoetioporphyrin II<sup>232</sup>, acetatohematoporphyrin IX dimethyl ester, and chlorohematoporphyrin IX dimethyl ester<sup>233</sup>, derivatives of Mn<sup>III</sup> have been prepared. The reaction of Mn<sup>III</sup> tetrapyrroldiporphines with various reducing agents has been studied<sup>239</sup>.

#### (ix) Hexacyanomanganates (III)

K<sub>3</sub>Mn(CN)<sub>6</sub> is prepared<sup>241</sup> by air oxidation of K<sub>4</sub>Mn(CN)<sub>6</sub>, or<sup>83, 240</sup> by the action of excess aqueous KCN on a manganese (III) complex such as MnPO<sub>4</sub>, K<sub>2</sub>MnF<sub>5</sub>·H<sub>2</sub>O or Mn(CH<sub>3</sub>COO)<sub>3</sub>. The Li, Na, Rb, NH<sub>4</sub> salts are also known. K<sub>3</sub>Mn(CN)<sub>6</sub> forms red crystals, isomorphous with K<sub>3</sub>Fe(CN)<sub>6</sub>, *a* = 13.59, *b* = 10.62, *c* = 8.52 Å (ref. 243). The magnetic moment is 3.50 B.M., considerably higher than the "spin-only" value for a low spin *d*<sup>4</sup> compound (2.83 B.M.). The Mn(CN)<sub>6</sub><sup>3-</sup> ion is one of the few low-spin Mn<sup>III</sup> compounds known (*t*<sub>2g</sub><sup>4</sup>).

The yellow, dilute aqueous solution hydrolyses slowly to MnO(OH). Exchange between Mn(CN)<sub>6</sub><sup>3-</sup> and CN<sup>-</sup> is very fast, first-order in Mn(CN)<sub>6</sub><sup>3-</sup> and zero-order in CN<sup>-</sup>.

The standard potential for the reaction<sup>242</sup>




is -0.24V

K<sub>3</sub>[Mn(CN)<sub>5</sub>OH] has been reported<sup>244, 245</sup>. The anion [Mn(CN)<sub>5</sub>NO]<sup>3-</sup>, although once regarded as an Mn<sup>III</sup> compound, exhibits an N—O stretching frequency of 1730 cm<sup>-1</sup> which indicates the presence of NO<sup>+</sup> and hence the oxidation state of the manganese is probably +1 (ref. 246).

Red-brown crystals of 1/2 [K<sub>4</sub>Mn(CN)<sub>6</sub>] · 0.7 [K<sub>3</sub>Mn(CN)<sub>6</sub>] · H<sub>2</sub>O, a mixed Mn<sup>II</sup>—Mn<sup>III</sup> cyanocomplex is formed<sup>247</sup> by reaction between K<sub>4</sub>Mn(CN)<sub>6</sub> and K<sub>3</sub>Mn(CN)<sub>6</sub> under carefully controlled conditions. It has a magnetic moment of 1.04 B.M.

#### (x) Other Mn<sup>III</sup> complexes

Nyholm and Sutton<sup>248</sup> were unable to oxidise [Mn(DAS)<sub>2</sub>X<sub>2</sub>] (X = Cl, Br, I) to Mn<sup>III</sup>, are found that DAS rapidly reduced Mn(OAc)<sub>3</sub> or Mn(acac)<sub>3</sub> to Mn<sup>II</sup> under anhydrous conditions. In the presence of small amounts of water, the red-purple [Mn(DAS)Cl<sub>2</sub>·H<sub>2</sub>O] (ClO<sub>4</sub>) was formed in low yield. The reaction can be represented




The complex has  $\mu_{\text{eff}} = 5.13$  B.M. A green complex, probably the bromo analogue was isolated using HBr in place of HCl in the above reaction.

*(xi) Manganese (III) in solution*

A detailed review<sup>2</sup> is available, in which the generation of  $\text{Mn}^{\text{III}}$  in solution, its estimation, and the equilibria and kinetics of a large number of reactions are discussed.


Manganese (III) is a strongly oxidising species and is unstable with respect to disproportionation. Solutions of  $\text{Mn}^{\text{III}}$  in perchlorate media generally produce solid hydrated  $\text{MnO}_2$  on standing, instability increasing with increasing  $[\text{Mn}^{\text{III}}]$  and decreasing acidity<sup>249,250</sup>.



Complexation of  $\text{Mn}^{\text{III}}$  with anions usually results in a reduction of the electrode potential of the  $\text{Mn}^{\text{III}}-\text{Mn}^{\text{II}}$  couple; for example, the electrode potential in the presence of EDTA is ca.  $-0.82 \text{ V}$  (ref. 251), while that in perchlorate media is ca.  $-1.58 \text{ V}$ . Moreover, the presence of a number of complexed  $\text{Mn}^{\text{III}}$  species complicates the interpretation of kinetic data. For example, Waters and Littler<sup>252</sup> have concluded that little information can be obtained from a study of the pH-dependence of reactions of manganese (III) pyrophosphate with organic reductants because of a lack of knowledge of oxidising species present in solution. An interesting observation regarding the magnitude of the effects of complexation on the oxidising power of  $\text{Mn}^{\text{III}}$  is that hydrogen peroxide is found as a product of the oxidation of oxalate by  $\text{Mn}^{\text{III}}\text{CyDTA}$  at low pH<sup>253</sup>.

In an investigation of the analytical applications of the EDTA complex of  $\text{Mn}^{\text{III}}$ , Pribil et al.<sup>254,255</sup> found that the complex slowly decomposed. Yoshino et al.<sup>256</sup> isolated the complex and found decomposition was enhanced by heat, light, and traces of  $\text{Mn}^{\text{II}}$  ion. Tanaka et al.<sup>257</sup> have measured the standard oxidation-reduction potential of the  $\text{Mn}^{\text{III}}\text{EDTA}-\text{Mn}^{\text{II}}\text{EDTA}$  complex. Hamm isolated the complexes of  $\text{Mn}^{\text{III}}$  with EDTA, *trans*-1,2-diaminocyclohexane-tetraacetic acid (CyDTA), and hydroxyethylethylenediamine-tetraacetic acid (HEDTA), and established the rates of decomposition in acidic solution. The standard potentials for the reaction  $\text{Mn}^{\text{III}}\text{Y} + \text{e}^- \rightleftharpoons \text{Mn}^{\text{II}}\text{Y}$  were determined, and all three complexes were equally good oxidising agents. Formation constants were calculated<sup>251</sup>. The products of the decomposition of the  $\text{Mn}^{\text{III}}$  complexes of CyDTA and EDTA were  $\text{Mn}^{\text{II}}$  complexes with oxidation of some of the ligand to formaldehyde, carbon dioxide, and the triacetate ligand<sup>258</sup>. The oxidation of oxalate with  $\text{Mn}^{\text{III}}\text{CyDTA}$  has been studied<sup>253</sup>.

Important information concerning the stabilities of several  $\text{Mn}^{\text{III}}$  complexes has been obtained by a study of the  $\text{Mn}^{\text{III}}$ -catalysed oxidation of oxalic acid by chlorine<sup>259</sup> and bromine<sup>260</sup>. The rate-determining step in this reaction in the first-order decomposition of the monoxalate complex, viz



Studies of the oxalate complexes of  $\text{Mn}^{\text{III}}$  showed<sup>259,261</sup> that the decomposition was dependent on the concentrations of oxalate and  $\text{H}^+$  ions. The decomposition of the tartrate<sup>262</sup> and glycolate<sup>263</sup> complexes was found to be first-order in the complex concentration in the presence of excess complexing agent, and to be inhibited by the addition of  $\text{Mn}^{\text{II}}$  ion.




Wells and coworkers have carried out spectrophotometric investigations of alcohol-<sup>264</sup> and aquo-manganese (III)<sup>265</sup> species in perchlorate media. The reactions of aquomanganese (III) ions with bromide ion<sup>266</sup>, hydrogen peroxide<sup>267,268</sup> hydrazoic acid<sup>269,270</sup> and hydroxylamine<sup>271</sup> in perchlorate media have been studied. The species  $\text{MnF}(\text{CH})^+_{\text{aq}}$  has been postulated by Davies and Kustin<sup>272</sup>.

Oxidation of organic compounds by  $\text{Mn}^{\text{III}}$  complexes in aqueous solution has been extensively studied<sup>252</sup>, e.g. the oxidation of toluene by  $\text{Mn}(\text{acetate})_3$  in refluxing acetic acid yields benzyl acetate, *o*-methylbenzylacetate, and tolylacetic acid<sup>273</sup>. Most of these results have been successfully interpreted in terms of inner-sphere one-electron-transfer processes<sup>274</sup>. The non-aqueous chemistry has been less extensively explored, and the oxidations reported have been interpreted as resulting from inner-<sup>275</sup> or outer-sphere<sup>276</sup> one-electron-transfer steps. Manganese (III) acetate oxidises olefins to  $\gamma$ -lactones<sup>277-280</sup>.

#### SPECTRA OF MANGANESE (III) COMPOUNDS

The electronic spectra of high-spin  $\text{Mn}^{\text{III}}$  compounds are of special interest because the ground electronic state anticipated in octahedral complex,  ${}^5E_g$ , is subject to strong Jahn–Teller forces<sup>265,281</sup>. In general, three bands are observed in the visible region, and satisfactory assignment has been made in terms of  $D_{4h}$  symmetry<sup>171,282</sup>. The two highest energy bands have been assigned to the transitions  ${}^5B_{1g} \rightarrow {}^5B_{2g} \rightarrow {}^5E_g$ .



However, despite intensive work on the preparation and optical properties of  $\text{Mn}^{\text{III}}$  complexes<sup>168,223,250,282-286</sup>, difficulties have arisen in assignment of the band in the near-infrared region. The relationship of the presence of this band to the structure of the complexes has not been clarified. The band (which appears between 8 and 13K) has been variously assigned to a spin-forbidden transition<sup>209,213</sup> from the ground state to  ${}^3T_{1g}$  (in  $O_h$  symmetry), as a low-energy charge-transfer transition<sup>284,286</sup> as a transition from the  ${}^5E_g$  ground state to the trigonally split  ${}^5T_{2g}$  excited state<sup>287</sup>, and as a spin-allowed

transition between components of the  ${}^5E_g$  (in  $O_h$  symmetry) ground state split by Jahn–Teller forces<sup>265,288</sup>. Davis et al. have convincingly shown that this band can be assigned to the  ${}^5B_{1g} \rightarrow {}^5A_{1g}$  transition in  $D_{4h}$  symmetry, and they formulated an interesting argument for this assignment<sup>171</sup>. The reflectance spectra of  $K_3MnF_6$  (ref. 285) and  $MnCl_6^{3-}$  (ref. 289) show the presence of a low-energy band at  $9,000\text{ cm}^{-1}$  and  $8,300\text{ cm}^{-1}$  respectively. The single-crystal spectrum of  $(NH_4)_2MnF_5$  shows a similar low-energy band at  $12,750\text{ cm}^{-1}$  which can be assigned to the  ${}^5B_{1g} \rightarrow {}^5A_{1g}$  transition on the basis of polarisation of the bands<sup>167a</sup>. All other halide complexes for which there are data also show a low-energy band, e.g.  $Na_2MnF_5$  ( $12,500\text{ cm}^{-1}$ ) (ref. 290),  $K_2MnCl_5$  ( $12,000\text{ cm}^{-1}$ ) (ref. 289),  $K_2MnF_5$  ( $12,100\text{ cm}^{-1}$ ) (ref. 289),  $[(C_2H_5)_4N]_2MnCl_5$  ( $11,300\text{ cm}^{-1}$ ) (refs. 135, 289),  $[(bipy)H_2][MnCl_5]$  ( $11,500\text{ cm}^{-1}$ ) (ref. 135),  $[(phen)H_2][MnCl_5]$  ( $12,500\text{ cm}^{-1}$ ) (ref. 135) and  $MnF_3$  ( $12,000\text{ cm}^{-1}$ ) (ref. 283). In the pentahalide species, the atom generally attains six-coordination by sharing axial ligands as in  $(NH_4)_2MnF_5$  (refs. 167, 291). In  $MnF_3$ , all fluorines are shared to complete the distorted octahedron about each  $Mn^{III}$  ion. The shift in position of the low-energy band in going from the hexahalides to the pentahalides and  $MnF_3$  is best understood on the basis of the Jahn–Teller effect. In the hexahalides none of the ligand ions is shared, hence dynamic motion (minima exchange) of the tetragonally displaced ligands about the manganese may occur. In the pentahalides and in  $MnF_3$ , where ligands are shared, the minima exchange should be decreased as stabilisation of the molecule in one tetragonally distorted form occurs. A shift in the low-energy ( ${}^5B_{1g} \rightarrow {}^5A_{1g}$ ) band to higher wave-numbers consistent with a statically distorted structure follows. A similar argument<sup>283</sup> was presented for aqueous  $MnF^{2+}$ .

#### (xiii) Low-spin compounds

The spectra of  $Mn(CN)_6^{3-}$  and of  $Mn(CN)_5(OH)^{3-}$  have been discussed by Chawla and Frank<sup>292</sup>. Earlier investigations<sup>293,294</sup> of the spectrum of the  $Mn(CN)_6^{3-}$  ion in aqueous solution are of doubtful value, since it has been shown that hydrolysis in such solutions is not negligible, as had been earlier assumed<sup>179,265</sup>.

The  $Mn(CN)_6^{3-}$  ion<sup>292</sup> shows charge-transfer absorptions at  $43,500$ ,  $21,800\text{ cm}^{-1}$ , with a spin-allowed transition at  $27,000\text{ cm}^{-1}$  ( ${}^3T_{1g} \rightarrow {}^3E_g$ ), and spin-forbidden transitions at  $12,000\text{ cm}^{-1}$  ( ${}^3T_{1g} \rightarrow {}^5E_g$ ),  $12,500\text{ cm}^{-1}$  ( ${}^3T_{1g} \rightarrow {}^1T_{2g}$ ),  $13,500\text{ cm}^{-1}$ , ( ${}^3T_{1g} \rightarrow {}^1E_g$ ), and  $24,500\text{ cm}^{-1}$  ( ${}^3T_{1g} \rightarrow {}^1A_{1g}$ ).

#### REFERENCES

- 1 N V Sidgwick, *Chemical Elements and Their Compounds*, Oxford University Press, London, 1950, p. 1262.
- 2 G Davies, *Coord Chem Rev.*, 4 (1969) 199.
- 3 T A Zordan and L G Hepler, *Chem Rev.*, 68 (1968) 739.
- 4 A Engelbrecht and A V Grosse, *J Amer Chem. Soc.* 76 (1954) 2042.
- 5 T S Briggs, *J Inorg Nucl. Chem.*, 30 (1968) 2866.
- 6 K. Wiechert, *Z Anorg Chem.*, 261 (1950) 310.
- 7 D Michel and A. Doiwa, *Naturwissenschaften*, 53 (1966) 129.
- 8 E.E Aynsley, *J Chem Soc.*, (1958) 2425.
- 9 A Javan and A Engelbrecht, *Phys Rev.*, 96 (1954) 649.

- 10 P J Aymonino, H. Schulze and A. Muller, *Z. Naturforsch.*, B 24 (1969) 1508
- 11 O. Glemser and H. Schroder, *Z. Anorg. Chem.*, 271 (1953) 293
- 12 A. Simon and F. Feher, *Z. Elektrochem.*, 38 (1932) 137
- 13 J M Loven, *Ber. Deut. Chem. Ges.*, 25 (1892) ref 620
- 14 P. Dubois, *Compt. Rend.*, 200 (1935) 1107.
- 15 N A Frigerio, *J. Amer. Chem. Soc.*, 91 (1969) 6200
- 16 G. Brauer, *Handbook of Preparative Inorganic Chemistry*, Vol. 2, Academic Press, New York, 1963, p. 1462.
- 17 A A Zinov'ev and K V Titova, *Russ. J. Inorg. Chem.*, 8 (1963) 823
- 18 A. Muller and B. Krebs, *Z. Naturforsch.*, B 21 (1966) 163
- 19 E J Baran and P J Aymonino, *Monatsh. Chem.*, 99 (1968) 606
- 20 E. Moles and M. Crespi, *Anal. Soc. Espan. Fis. Quim.*, 21 (1923) 305.
- 21 E. Moles and M. Crespi, *Anal. Soc. Espan. Fis. Quim.*, 20 (1922) 692
- 22 A. Ferrari, A. Braibanti, G. Bighardi and A M. Manotti Lanfredi, *Acta Cryst.*, 21 (1966) 681
- 23 E G. Prout and P J Herley, *J. Phys. Chem.*, 66 (1962) 961
- 24 P D Komissarova and S A Krestovnikova, *Chem. Abstr.*, 69 (1968) 56590
- 25 W P Doyle and I Kirkpatrick, *Spectrochim. Acta*, 24A (1968) 1495
- 26 G J Palenik, *Inorg. Chem.*, 6 (1967) 503
- 27 H Peters, K.H. Radeke and L. Till, *Z. Anorg. Chem.*, 346 (1966) 1
- 28 F H. Herstein, G. Ron and A. Weissman, *J. Chem. Soc. A*, (1971) 1821
- 29 P J Hendra, *Spectrochim. Acta*, 24A (1968) 125
- 30 W P Griffith, *J. Chem. Soc. A* (1966) 1467
- 31 A. Muller and B. Krebs, *Naturwissenschaften*, 52 (1965) 492
- 32 E J Baran and P J Aymonino, *Z. Anorg. Chem.*, 354 (1967) 85
- 33 J A. Ketelaar, *Z. Krist.*, 92 (1935) 155
- 34 A. Hardy, C. Piekarski and P. Hagen-Muller, *Compt. Rend.*, 249 (1959) 2579
- 35 K. Savari, *Z. Krist.*, 99 (1938) 9
- 36 J. Royer, *J. Inorg. Nucl. Chem.*, 17 (1961) 159
- 37 H C. Mishra and M C R. Symons, *J. Chem. Soc.*, (1962) 4411
- 38 B. Franke, *J. Prakt. Chem.*, 36 (1887) 31
- 39 T E. Thorpe and F J. Hambly, *J. Chem. Soc.*, 53 (1888) 175
- 40 F.R. Lankshear, *Z. Anorg. Chem.*, 82 (1913) 97
- 41 G M Damanskaya and R I Agladze, *Issled. v Obl. Elektrokhim. i Radiats. Khim. Akad. Nauk Gruz. SSR*, (1965) 156
- 42 R. Scholder and H. Waterstradt, *Z. Anorg. Chem.*, 277 (1954) 172
- 43 C. Rocchiccioli, *Compt. Rend. C*, 256 (1963) 1707
- 44 G. Duquenoy, *Compt. Rend. C*, 268 (1969) 1828.
- 45 J W. Mellor, *A Comprehensive Treatise of Inorganic and Theoretical Chemistry*, Vol. XII, Longmans, London, 1932
- 46 W. Klemm and K A. Jensen, *Z. Anorg. Chem.*, 237 (1938) 47
- 47 G J. Falenik, *Inorg. Chem.*, 6 (1967) 507
- 48 H. Lux, *Z. Naturforsch.*, 1 (1946) 281
- 49 B. Jezowska-Trzebiatowska, J. Nawojcka and M. Wroucka, *Roczniki Chem.*, 25 (1951) 405
- 50 R. Scholder, D. Fisher and H. Waterstradt, *Z. Anorg. Chem.*, 277 (1954) 234
- 51 R. Scholder and U. Protzer, *Z. Anorg. Chem.*, 369 (1969) 313.
- 52 H. Lux, E. Brodkorb, R. Mahr and E. Oeser, *Z. Anorg. Chem.*, 337 (1965) 200
- 53 W. Klemm, *Proc. Intern. Symp. Reactivity of Solids. Gottenberg 1952*, Pt 1, 1954, p. 173
- 54 R. Scholder, *Angew. Chem.*, 70 (1958) 583
- 55 E J. Baran and P J. Aymonino, *Monatsh. Chem.*, 100 (1969) 1674
- 56 L H. Brixner and J F. Weiher, *Inorg. Chem.*, 7 (1968) 1474
- 57 W. Klemm, C. Brendel and G. Wehrmeyer, *Chem. Ber.*, 93 (1960) 1506
- 58 M. Wolfsberg and L. Helmholtz, *J. Chem. Phys.*, 20 (1952) 837
- 59 C J. Ballhausen and A.D. Liehr, *J. Mol. Spectry*, 2 (1958) 342
- 60 A. Carrington and M.C.R. Symons, *J. Chem. Soc.*, (1960) 88

- 61 D S Schonland, *Proc Roy Soc (London), Ser A*, 254 (1960) 111  
62 A Carrington and D S Schonland, *Mol Phys*, 3 (1960) 331  
63 A Carrington and C K Jorgensen, *Mol Phys*, 4 (1961) 395  
64 R F Fenske and C Sweeney, *Inorg Chem*, 3 (1964) 1105  
65 A Viste and H B Grey, *Inorg Chem*, 3 (1964) 1113.  
66 L Oleari, G De Michelis and L Di Sipio, *Mol Phys*, 10 (1965) 111  
67 L Oleari, G De Michelis, L Di Sipio and E Tondello, *Coord Chem Revs*, 2 (1967) 53  
68 J D Kingsley, J S Prener and B Segall, *Phys Rev* 137A, (1965) 189  
69 L E Orgel, *Mol Phys*, 7 (1963) 397  
70 J Kubris and J H Krepelka, *Collection Czech Chem Commun* 7 (1935) 105  
71 A G Sharpe and A A Woolf, *J Chem Soc*, (1951) 798  
72 H W Roesky, O Glemser and K H Hellberg, *Chem Ber*, 98 (1965) 2046  
73 R Hoppe, W Dahne and W Klemm, *Ann Chem*, 658 (1962) 1  
74 R Hoppe, W Dahne and W Liebe, *Z Anorg Chem*, 307 (1961) 276  
75 B Cox, *J Chem Soc*, (1954) 3251  
76 D H Brown, K R Dixon, R W D Kemmit and D W A Sharp, *J Chem Soc*, (1965) 1559  
77 H Bode and W Wendt, *Z Anorg Chem* 269 (1952) 165  
78 R Hoppe, *J Inorg Nucl Chem*, 8 (1958) 435  
79 R Hoppe and K Blinne, *Z Anorg Chem*, 291 (1957) 269  
80 R Hoppe, *Rec Trav Chim*, 75 (1956) 569  
81 R D Peacock and D W A Sharp, *J Chem Soc*, (1959) 2762  
82 J T Grey, *J Amer Chem Soc*, 68 (1946) 605  
83 W G Palmer, *Experimental Inorganic Chemistry* Cambridge University Press, London, 1962  
84 H Bode, H Jensen and F Bandte, *Angew Chem* 65 (1953) 304  
85 A G Sharpe and B Cox, *J Chem Soc*, (1951) 798  
86 P Bouy, *Ann Chim (Paris)*, 4 (1959) 853  
87 R Hoppe and G Siebert, *Z Anorg Chem*, 376 (1970) 261  
88 P C Moews, Jr, *Inorg Chem* 5 (1966) 5  
89 R D Peacock, *J Chem Soc*, (1953) 3617  
90 R F Weinland and P Dinklacher, *Z Anorg Chem* 60 (1908) 173  
91 H D Hart and M Fleischer, *Z Anorg Chem* 357 (1968) 113  
92 N Elliot, *J Chem Phys*, 46 (1967) 1006  
93 D M Adams and D M Morris, *J Chem Soc A*, (1968) 694  
94 G C. Allen, G A M El-Sharkawy and K D Warren, *Inorg Nucl Chem Letters*, 5 (1969) 725  
95 D S Novotny and G D Sturgeon, *Inorg Nucl Chem Letters* 6 (1970) 455  
96 C K Jorgensen, *Acta Chem Scand*, 12 (1958) 1539  
97 B Jezowska-Trzebiatowska, S Wajda, M Baluka, L. Nathaniel and W Wojciechowski, *Inorg Chim Acta*, 1 (1967) 205  
97a L B Asprey, M J Reisfeld and N A Matwiyoff, *J Mol Spectry*, 34 (1970) 361  
98 T E Moore, M Ellis and P W Selwood, *J Amer Chem Soc*, 72 (1950) 856.  
99 A D Wadsley and A Walkley, *Rev Pure Appl Chem*, 1 (1951) 203  
100 W Feitknecht and A Marti, *Helv Chim Acta*, 28 (1945) 129  
101 K H Maxwell and H R Thurst, *J Chem Soc*, (1955) 4054  
102 G Gattow and O Glemser, *Z Anorg Chem*, 309 (1961) 121  
103 O Glemser, G Gattow and M Meisiek, *Z Anorg Chem*, 309 (1961) 1  
104 O Glemser and G Gattow, *Z Anorg Chem*, 309 (1961) 20  
105 S S Bhatnager, A Cameron, E H Habard, A King, P L Kapur and B Prakash, *J Chem Soc*, (1939) 1433  
106 A Schmier and G Sterr, *Z Anorg Chem*, 346 (1966) 181  
107 A. Schmier and G Sterr, *Z Anorg Chem*, 368 (1969) 265  
108 R Giovanoli, E Stahl and W Feitknecht, *Helv Chim Acta*, 53 (1970) 209  
109 R Giovanoli, E Stahl and W Feitknecht, *Helv Chim. Acta* 53 (1970) 453.  
110 R Giovanoli, W Feitknecht and F. Fischer, *Helv Chim Acta*, 54 (1970) 1112

- 11 N N Greenwood, *Ionic Crystals Lattice Defects, and Nonstoichiometry* Butterworths, London, (1968) p. 101.
- 12 H F McMurdie and E. Golovato, *J Res Natl Bur Std*, 41 (1948) 589
- 13 H R Oswald, W Feitknecht and M J Wampetich, *Nature*, 207 (1965) 72
- 14 H R Oswald and M J Wampetich, *Helv Chim Acta*, 70 (1967) 2023
- 15 M Mansmann, *Z Anorg Chem*, 339 (1965) 52
- 16 F H Delano, *Ind Eng Chem*, 42 (1950) 535
- 17 R Scholder, *Z Anorg Chem*, 260 (1949) 41, 231
- 18 N I Kharabadze, *Elektrokhim Margantsa Akad Nauk Gruz SSR*, 2 (1963) 255.
- 19 H Remy, *Treatise on Inorganic Chemistry*, Vol 2, Elsevier, Amsterdam, 1956
- 20 I Reimer and M.W Lister, *Can J Chem*, 39 (1961) 2431
- 21 W B Price, *J Amer Chem. Soc* 30 (1903) 182
- 22 M W Lister and Y Yoshino, *Can J Chem*, 38 (1960) 1291
- 23 Y Yoshino, T Takeuchi and H Kinoshita, *Nippon Kagaku Zasshi*, 86 (1965) 978
- 24 M W Lister and Y Yoshino, *Can J Chem*, 40 (1962) 1490
- 25 M W Lister, *Can J Chem*, 39 (1961) 2330
- 26 I D Brown, *Can J Chem*, 47 (1969) 3779
- 27 L Jenovsky, *Omagiu Raluca Ripon* (1966) 293
- 28 M A Berg, *Compt Rend*, 128 (1899) 673
- 29 W E Dasent and T C Waddington, *J Chem Soc* (1960) 2429
- 30 G H Cartledge and W P Ericks, *J Amer Chem Soc* 58 (1936) 2069
- 31 Z Marczenko, *Bull Soc Chim*, (1964) 939
- 32 V. Auger and A Yakimach, *Compt Rend*, 187 (1928) 603
- 33 P Schottlander, *Ann Chem*, 155 (1870) 230
- 34 M M Ray and P Ray, *J Ind Chem Soc*, 35 (1958) 595
- 35 H.A. Goodwin and R N Sylva, *Australian J Chem*, 18 (1965) 1743
- 36 H A Goodwin and R N Sylva, *Australian J Chem*, 20 (1967) 629
- 37 W Levason and C A McAuliffe, unpublished work
- 38 J R Fowler and J Kleinberg, *Inorg Chem.*, 9 (1970) 1005
- 39 A Yakimach, *Compt Rend*, 190 (1930) 681
- 40 N Goldenberg, *Trans Faraday Soc*, 37 (1940) 847
- 41 J T Waugh, *Acta Cryst*, 7 (1954) 438
- 42 B W Dale, J M Buckley and M T Pope, *J. Chem Soc A*, (1969) 301
- 43 C M Flynn and G D Stucky, *Inorg Chem*, 8 (1969) 333
- 44 C M Flynn and G D. Stucky, *Inorg Chem*, 9 (1970) 2009
- 45 C M Flynn and G D Stucky, *J Amer Chem Soc*, 92 (1970) 85
- 46 G Brauer, *Handbook of Preparative Inorganic Chemistry*, Vol 1, Academic Press, New York, 1963, p 263
- 47 R D Fowler, H C Anderson, J M Hamilton, W B Burford, A Spadette, S B Bitterlick and I Litant, *Ind Eng Chem*, 39 (1947) 343
- 48 H J Emeleus and G L Hunt, *J Chem Soc*, (1964) 396
- 49 E E Aynsley, R D Peacock and P L Robinson, *J Chem Soc*, (1950) 1622
- 50 M A Hepworth and K H Jack, *Acta Cryst*, 10 (1957) 345
- 51 J H Simons, *Fluorine Chemistry*, Vol 1, Academic Press, New York, 1950
- 52 B Cox and A G Sharpe, *J Chem Soc*, (1954) 1798
- 53 H V Wartenberg, *Z Anorg Chem*, 244 (1940) 337
- 54 R A Rausch, R A Davies and D W Osborne, *J Org Chem*, 28 (1963) 494
- 55 M A Hepworth, K H Jack and R S Nyholm, *Nature*, 179 (1957) 211
- 56 Z F Zimbov and J L Margrave, *J Inorg Nucl Chem*, 29 (1967) 673
- 57 R M Bozorth and J W Nielson, *Phys Rev*, 110 (1958) 879
- 58 J Aubert and G H. Cady, *Inorg Chem* 9 (1970) 2600
- 59 A Chretien and G Varga, *Bull Soc Chim*, 3 (1936) 2385
- 50 R D Peacock, *J Chem Soc.*, (1957) 4684

- 161 S Schneider and R Hoppe, *Z Anorg Chem* 376 (1970) 268  
162 K Knox, *Acta Cryst* 16 (1963) A45  
163 K Wieghardt and H Siebert, *Z Anorg Chem*, 381 (1971) 12  
164 J G Ryss and B S Vitukhnovskaya, *Chem Abstr*, 48 (1954) 13506.  
165 R S Nyholm and A G Sharpe, *J Chem Soc*, (1952) 3579  
166 J.G Ryss and B S Vitukhnovskaya, *Z Neorg Khim*, 3 (1958) 1185  
167 S Enori, M Inoue, M Kishita and M Kubo, *Inorg Chem*, 9 (1969) 1385  
167a R Dingle, *Inorg Chem*, 4 (1965) 1287  
168 W E Hatfield, R C Fay, C E Pfluger and T S Piper, *J Amer. Chem Soc*, 85 (1963) 265  
169 W Levason, C A McAuliffe and S G Murray, submitted for publication  
170 N S Gill, *Chem Ind (London)*, (1961) 989  
171 T S Davis, J P Fackler, Jr and M J Weeks, *Inorg Chem*, 7 (1968) 1994  
172 C E Rice, *J Chem Soc*, 73 (1898) 260  
173 I Bernal, N Elliot and R A Lalancette in M Cais (Ed), *Proc XI Intern Conf Coord Chem*, Haifa, Israel, 1968, Elsevier, Amsterdam, 1968, p 518  
174 L Pauling, *Z Krist*, 75 (1930) 128  
175 P Dubois, *Ann Chim*, 5 (1936) 401  
176 A F Wells, *Structural Inorganic Chemistry*, Oxford University Press, London, 3rd edn, 1962  
177 W Feitknecht, P Brunner and H R Oswald, *Z Anorg Chem*, 316 (1962) 154  
178 R Scholder and H Kyri, *Z Anorg Chem*, 270 (1952) 56  
179 R Hoppe, *Z Anorg Chem*, 368 (1969) 262  
180 L Domage, *Bull Soc Chim France* 4 (1937) 594  
181 L. Domage, *Bull Soc Chim France*, 6 (1939) 1452  
182 S V Gorbatshev and E E Schpitalski, *J Gen Chem. Russ*, 10 (1940) 1961  
183 O T Christensen, *Z Anorg Chem.*, 27 (1901) 329  
184 H Bommer, *Z Anorg Chem*, 246 (1941) 275  
185 J Meyer and J Marek, *Z Anorg Chem*, 133 (1924) 325  
186 O T Christensen, *Z Anorg Chem*, 27 (1901) 325  
187 G H Cartledge and W P Ericks, *J Amer. Chem Soc* 58 (1936) 2061  
188 J Meyer and W Schramm, *Z Anorg Chem*, 123 (1922) 56  
189 J I Bullock, M M Patel and J E Salmon, *J Inorg Nucl Chem*, 31 (1969) 415  
190 G H Cartledge and P.M Nichols, *J Amer Chem Soc*, 62 (1940) 3057  
191 Y Yoshino, A. Ouchi, T Tsunoda and M Kojima, *Can J Chem*, 40 (1962) 775  
192 S Richards, B Peterson, J V Silvertown and J L Hoard, *Inorg Chem*, 3 (1964) 27  
193 R S Nyholm and A Turco, *J Chem Soc*, (1962) 1121  
194 W M. Reiff and W.A Baker, *Inorg Chem*, 9 (1970) 570  
195 J P Fackler, Jr, *Progr Inorg Chem.*, 7 (1966) 361.  
196 W C Fernelius and B E Bryant, *Inorg Syn*, 6 (1957) 105  
197 R G Charles, *Inorg Syn*, 6 (1960) 164  
198 G H Cartledge, *J Amer Chem Soc* 73 (1951) 4416, *US Pat* 2,556,316  
199 G S Hammond, W.G Bordman and G A Guter, *J Amer Chem Soc*, 81 (1959) 4682  
200 A E Martell and M Calvin, *Chemistry of the Metal Chelate Compounds*, Prentice-Hall, New York, 1952  
201 R C. Fay and T S Piper, *J Amer Chem Soc*, 84 (1952) 2303, 85 (1964) 500  
202 T S Davis, J P Fackler, Jr and M J Weeks, *Inorg Chem*, 7 (1968) 1994  
203 J O Hill and R J Irving, *J Chem Soc A*, (1968) 3116  
204 B Morosin and J R Brathovde, *Acta Cryst*, 17 (1964) 705  
205 A Forman and L.E. Orgel, *Mol Phys* 2 (1959) 362  
206 K Nakamoto, *Infrared Spectra of Inorganic and Coordination Compounds*, Wiley, New York, 1963, 1st edn, p 216  
207 M Bartlett and G J Palenik, *Chem Commun*, (1970) 416  
208 D.W Barnum, *J Inorg Nucl Chem*, 21 (1961) 221, 22 (1961) 183  
209 T S Piper and R L. Carlin, *Inorg Chem*, 2 (1963) 260

- 210 G H Cartledge, *J Amer Chem Soc* 74 (1952) 6015.
- 211 T Shigematsu and M. Tabustin, *J. Chem Soc. Japan Pure Chem Sect.*, 83 (1962) 814
- 212 P Pfeiffer, E Breith, E Lubbe and T Tsumaki, *Ann Chem*, 503 (1933) 84.
- 213 T Tsumaki, *Bull Chem Soc Japan*, 13 (1938) 579
- 214 A Earnshaw, E A. King and L F. Larkworthy, *J Chem Soc A*, (1968) 1048.
- 215 T Yarino, T Matsuchita, I Masuda and K. Shinra, *Chem Commun*, (1970) 1317.
- 216 J Lewis, F E Mabbs and H Weigold, *J Chem Soc. A*, (1968) 1699.
- 217 B Das Sarma, R Ray, R E Sievers and J C Bailor, Jr, *J Amer Chem Soc*, 86 (1964) 14
- 218 A Van den Bergen, K S Murray, M J O'Connor and B O West, *Australian J Chem*, 22 (1969) 39.
- 219 F H Burstall and R S Nyholm, *J. Chem Soc*, (1952) 3570
- 220 A Turco and R S Nyholm, *Chem Ind (London)*, (1960) 74
- 221 H Funk and H Kreis, *Z Anorg Chem*, 349 (1967) 45
- 222 M M. Ray and P Ray, *J Ind Chem Soc*, 35 (1955) 595
- 223 M M Ray, J N Adhya, D. Biswas and S N Podder, *Australian J Chem*, 19 (1966) 881
- 224 M M Ray, J N Adhya, D Biswas and S N Podder, *Australian J Chem*, 21 (1968) 801
- 225 M M Ray, J N Adhya, D Biswas and S N Podder, *Australian J Chem*, 19 (1966) 1737
- 226 J A Elvidge and A B P Lever, *Proc Chem Soc*, (1959) 195
- 227 L H Vogt, Jr, A Zalkin and D H Templeton, *Inorg Chem*, 6 (1967) 1725, *Science*, 151 (1966) 569
- 228 E Kessler, W Arther and I E Brugger, *Arch Biochem Biophys* 71 (1957) 326
- 229 R D Park and N G Pon, *J Mol Biol*, 6 (1963) 105
- 230 A D Swensen and L P Vernon, *Biochim Biophys Acta*, 102 (1965) 349.
- 231 M Calvin, *Rev Pure Appl Chem*, 15 (1965) 1
- 232 A Yamamoto, L K. Phillips and M Calvin, *Inorg Chem*, 7 (1968) 847
- 233 P A Loach and M Calvin, *Biochemistry*, 2 (1963) 361
- 234 P.A Loach and M Calvin, *Nature*, 202 (1964) 343
- 235 G Engelsma, A Yamamoto, E Markham and M Calvin, *J Phys Chem*, 66 (1962) 2517
- 236 J L Boucher, *J Amer Chem Soc*, 90 (1968) 6640
- 237 J Zaleski, *J Physiol Chem*, 43 (1904) 11
- 238 J F Taylor, *J Biol Chem.*, 135 (1940) 569
- 239 P Hambright and E.B Fleischer, *Inorg Chem*, 4 (1965) 912
- 240 J A Lowther and W.C Fernelius, *Inorg Syn*, 2 (1946) 213
- 241 J Meyer, *Z Anorg Chem*, 81 (1913) 385
- 242 W D Treadwell and W E Rath, *Helv Chim Acta*, 35 (1952) 2275
- 243 C Gotfried and J G Nagelschmitt, *Z Krist*, 73 (1930) 357
- 244 H K Saha, *Chem. Abstr.*, 57 (1962) 1833
- 245 I D Chawla and M J Frank, *J Inorg Nucl Chem.*, 32 (1970) 555.
- 246 B M Chadwick and A G Sharpe, *Advan Inorg Chem Radiochem*, 8 (1966) 83.
- 247 A E. McCarthy, *J Chem Soc A*, (1970) 1379
- 248 R S Nyholm and G J Sutton, *J Chem Soc*, (1958) 564
- 249 D R Rosseinsky, *J Chem Soc*, (1963) 1181
- 250 H Diebler and N Sutn, *J Phys Chem*, 68 (1964) 174
- 251 R E Hamm and M A Suwyn, *Inorg Chem*, 6 (1967) 139
- 252 W A Waters and J S Littler, in K B Wiberg (ed) *Oxidation in Organic Chemistry*, Academic Press, New York, 1965.
- 253 M A Suwyn and R E Hamm, *Inorg Chem*, 6 (1967) 142
- 254 R Pribil and J Horacek, *Collection Czech Chem Commun.*, 14 (1949) 454
- 255 R Pribil and E Hornychova, *Collection Czech. Chem Commun*, 15 (1950) 456
- 256 Y Yoshino, Y Tsumodu and A Ouski, *Bull Chem. Soc Japan*, 34 (1961) 1194
- 257 N Tanaka, T Shirakashi and H Ogino, *Bull Chem Soc Japan*, 38 (1965) 1515
- 258 K A Schroder and R E Hamm, *Inorg Chem*, 3 (1964) 391
- 259 H Taube, *J Amer Chem Soc*, 70 (1948) 1216
- 260 H Taube, *J Amer Chem Soc*, 70 (1948) 3928

- 261 F R Duke, *J Amer Chem Soc*, 69 (1947) 2885  
262 T N Srivastava, *Z Phys Chem* 209 (1958) 22  
263 T N Srivastava, *Z Phys Chem*, 211 (1959) 251  
264 C F Wells and C Barnes, *J Chem Soc A*, (1968) 1626  
265 C F Wells and G Davies, *J Chem Soc A*, (1967) 1858  
266 C F Wells and D Mays, *J Chem Soc A*, (1968) 577  
267 C F Wells and D Mays, *J Chem. Soc A*, (1968) 665  
268 G Davies, L J Kirschenbaum and K Kustin, *Inorg Chem*, 7 (1968) 146  
269 C F Wells and D Mays, *J Chem. Soc. A* (1968) 1622  
270 G Davies, L J. Kirschenbaum and K Kustin, *Inorg Chem* 8 (1969) 663  
271 G Davies and K. Kustin, *Inorg Chem*, 8 (1969) 484  
272 G Davies and K Kustin, *Inorg Chem*, 8 (1969) 1196  
273 E I Heiba, R M Dessau and W J Koehl, Jr, *J Amer Chem Soc*, 91 (1969) 138  
274 G Brauer, *Handbook of Preparative Inorganic Chemistry*, Vol 2, Academic Press, New York, 1963, p 186.  
275 R Van Helden and E C Kodyman, *Rec Trav Chim*, 80 (1961) 57  
276 P J Andrulis, M J S Dewar, R Deitz and R L Hunt, *J Amer Chem Soc* 88 (1966) 5473  
277 C Furlani and A Ciani, *Ann Chim (Rome)*, 48 (1958) 286  
278 O Holmes and D S McClure, *J Chem Phys*, 26 (1957) 1686  
279 E I Heiba, R M. Dessau and W J Koehl, Jr, *J Amer Chem Soc*, 90 (1968) 5905  
280 J B Bush, Jr and H Finkbeimer, *J Amer Chem. Soc* 90 (1968) 5903  
281 A D Liehr, *Prog Inorg Chem*, 3 (1962) 281  
282 J P Fackler and I D Chawla, *Inorg Chem*, 3 (1964) 1130  
283 R J H Clark, *J Chem Soc*, (1964) 417  
284 J P Fackler, T S Davis and I D Chawla, *Inorg Chem*, 4 (1965) 130  
285 W E Hatfield and W E Parker, *Inorg Nucl Chem Letters*, 1 (1965) 7  
286 R Dingle, *Acta Chem Scand*, 20 (1966) 33  
287 C K Jorgensen, *Acta Chem Scand*, 16 (1962) 2406  
288 R Dingle, *J Mol Spectry*, 9 (1962) 426  
289 T S Davis, *Ph D Thesis* Case Inst of Techn, 1967  
290 D Oelkrug, *Angew Chem Intern Ed Engl*, 5 (1966) 744  
291 D R Sears, *Ph D Thesis* Cornell Univ, 1958  
292 I D Chawla and M J Frank, *J Inorg Nucl Chem*, 32 (1970) 555  
293 R Samuel and A R R Desparde, *Z Phys*, 80 (1933) 395  
294 J J Alexander and H B Gray, *J Amer Chem Soc*, 90 (1968) 4260