


HIGH-SPIN BEHAVIOUR AND CHEMICAL PROPERTIES OF ORGANOMETALLIC DERIVATIVES OF IRON(III)

C. FLORIANI and F. CALDERAZZO

Istituto di Chimica Generale ed Inorganica, University of Pisa 56100 Pisa (Italy)

Relatively few examples of organometallic compounds of iron(III) have been reported in the literature. Iron—carbon σ -bonds have been prepared recently¹ by a Grignard reaction on iron(III)-halo derivatives with an etioporphyrin type of structure. The compounds isolated were the *p*-tolyl and the ethyl derivatives as dark blue and purple solids, respectively, sensitive to light. Besides these compounds, no other reports concerning the isolation of iron(III) organometallic compounds have appeared, to the best of our knowledge. The thermal instability of compounds of this type was further suggested by the



recently reported² failure to isolate iron(III) organometallic compounds, in spite of their suggested formation in the course of the reaction between anhydrous FeCl_3 and lithium alkyls. Since the metal—etioporphyrin and the metal-*N,N'*-ethylenebis(salicylideneiminato) skeletons (see structures I and II, respectively) are rather closely related either sterically or electronically, it was considered of interest to see whether it was possible to obtain stable organometallic derivatives of *N,N'*-ethylenebis(salicylideneiminato) iron.

In this paper a metal complex containing the *N,N'*-ethylenebis(salicylideneiminato) ligand will be abbreviated as [MSaEn], such as [FeSaEn] or [CoSaEn].

In the past years we have reported the occurrence of reductions taking place in non-aqueous solvents with several bis(salicylideneiminato)metal complexes^{3–5}. In particular, chemical reductions by sodium metal in tetrahydrofuran were carried out and the corresponding sodium derivatives of formula $\text{Na}[\text{MSaEn}]$ were isolated in an analytically pure state ($\text{M} = \text{Fe}, \text{Co}$). It is worth mentioning that we had also previously reported⁵ the preparation of the $[\text{CoSaEn}]^{2-}$ anion, which was prepared by treating CoSaEn with two




Fig. 1 Temperature dependence of magnetic susceptibility for $[\text{FeSaEn}]^-$

equivalents of sodium metal. We had demonstrated the existence of the anion by the reaction



Recently, Russian workers⁶ have confirmed the existence of the anion by electron transfer reactions to cyclooctatetraene to give $\text{C}_8\text{H}_8^{2-}$ and $[\text{CoSaEn}]^-$.

The most interesting properties of these compounds are (a) the bulk magnetic susceptibilities, and (b) the chemical reactivity, especially toward electrophilic reagents such as alkyl halides. As far as magnetic susceptibility measurements are concerned, the $[\text{FeSaEn}]^-$ and the $[\text{CoSaEn}]^-$ anions can be regarded as d^7 and d^8 systems of, respectively, high-spin and low-spin behaviour. We have, in fact, established that our previous results⁵ on the magnetic properties of $\text{Na}[\text{CoSaEn}]$, suggesting the presence of a residual small paramagnetism, were in error because of partial oxidation of the samples. Under the best operative conditions, the sodium derivative of cobalt is diamagnetic.

The reduction of FeSaEn takes place in tetrahydrofuran as solvent by using the stoichiometric amount of sodium



The sodium derivative is then precipitated from the resulting green solution by partial evaporation of the solvent and addition of heptane. The deep blue solid is extremely sensitive to air and has to be handled with air strictly excluded. The magnetic susceptibility of this compound was measured by the Gouy method and found to follow the Curie–Weiss law reasonably well from room temperature down to liquid nitrogen temperature (Fig. 1). The value of the magnetic moment at room temperature is 3.71 BM, quite close to the theoretical spin-only value (3.87 BM) for three unpaired electrons. The Weiss con-




Fig. 2 Temperature dependence of magnetic susceptibility for $\text{FeSaEn-CH}_2\text{C}_6\text{H}_5$


stant is -45°

It has previously been observed^{4,5,7} that $[\text{CoSaEn}]^-$ reacts with organic halides to give the corresponding cobalt(III) organometallic compounds



In view of the presently reported magnetic properties of $[\text{CoSaEn}]^-$, reaction (3) can be rationalised in terms of an oxidative addition to a planar d^8 system such as those usually encountered with cobalt(I), rhodium(I) and iridium(I)^{8,9}. Reaction (3) can, in fact, be interpreted as the intermediate formation of an anionic cobalt(III) organometallic derivative, $[\text{CoSaEn(R)X}]^-$. The latter would then be unstable with respect to X^- elimination and formation of the neutral cobalt(III) organometallic compound. Consistent with this view is the fact under the initial conditions of reaction (3), i.e. addition of the alkyl halide to the solution of $[\text{CoSaEn}]^-$ at ca. -80°C , no precipitation of sodium chloride was observed, in spite of the fact that the change of colour from green to orange takes place practically within the time of mixing. Sodium chloride elimination usually takes place at room temperature and is completed by addition of water in the case of the cobalt complexes⁵.


In view of past experience with cobalt complexes, a reaction similar to (3) was also carried out with $[\text{FeSaEn}]^-$



Reaction (4) was carried out under conditions similar to those previously reported for the cobalt complex. After addition of water, the benzyl derivative was isolated as red needle-like crystals in about 90% yield. Recrystallisation of the iron(III) organometallic compound was carried out by dissolution in tetrahydrofuran and precipitation with heptane.


The compound is very sensitive to air oxidation (see below) and is light-sensitive, as expected for a compound containing metal carbon σ -bonds.

Besides the benzyl derivative, the phenyl derivative, $\text{FeSaEn}-\text{C}_6\text{H}_5$ was also prepared and characterised*, by reaction of the phenyl Grignard reagent with $\text{FeSaEn}-\text{I}$ in tetrahydrofuran as solvent



The phenyl compound was isolated from the reaction mixture by a procedure similar to that previously mentioned for the benzyl derivative.

Both the phenyl and the benzyl derivatives of iron(III) are thermally unstable and reactive towards oxygen. The results obtained from the investigation of the behaviour of the organometallic derivatives towards heat and oxygen represent an additional piece of evidence for the presence of iron-carbon bonds in these compounds. The reactions



have been established. Although the reaction of the benzyl derivative with oxygen was somewhat more complicated, with the phenyl derivative the dimeric μ -oxo-bis [ethylenebis(salicylideneiminato)iron(III)]¹⁰ was the only isolable compound. We have never obtained substantial evidence for the formation of peroxo-alkyl compounds of the type found^{11,12} in certain alkyl-cobaloximes. We believe that in reaction (7) the determining factor is the labilisation of the metal-carbon bond, prior to attack by oxygen.

Both reactions (6) and (7) are typical of metal-carbon bonds and they therefore exclude the possibility that the alkyl group might be attached to some other point of the molecule. Further evidence of iron-carbon bonds in these compounds came from the magnetic susceptibility data, the $\text{FeSaEn}-\text{R}$ compounds are thermally much less stable than, for example, the corresponding low-spin $\text{CoSaEn}-\text{R}$ derivatives^{4,5,15}. However, a satisfactory Curie-Weiss plot was obtained (see Fig. 2), the Weiss constant was found to be about -9° and the magnetic moment at room temperature was 5.87 BM, very close to the theoretical value (5.92 BM) for five unpaired electrons. Details concerning the magnetic susceptibility measurements have been published elsewhere¹³.

The high-spin behaviour of $\text{FeSaEn}-\text{CH}_2\text{C}_6\text{H}_5$ is very unusual in organometallic chemistry. Only the σ -mesityl and the σ -naphthyl derivatives of chromium(II)¹⁴ recently prepared by Stolze and Hahle behave similarly. The iron(III) organometallics with the etio-porphyrin type of structure¹ might also be high-spin systems but unfortunately no detailed magnetic susceptibility data were reported to establish this point.

In accordance with the occupancy of the highest molecular orbitals, as suggested by magnetic susceptibility data, the $\text{FeSaEn}-\text{R}$ compounds are thermally much less stable than for example, the corresponding low-spin $\text{CoSaEn}-\text{R}$ derivatives^{4,5,15}. However,

* Unless otherwise stated, all the new compounds reported in this paper analysed correctly for the formulations given.

the iron(III) compounds were isolated by normal operating procedures at room temperature and should be regarded as stable at least in the solid state, provided air and light are excluded.

We have also attempted the preparation of the corresponding methyl and acetyl derivatives. In both cases reactions between the $[\text{FeSaEn}]^-$ anion and the proper organic halide were observed but no analytically pure product could be obtained.

By analogy with the magnetic results obtained¹⁶ for the chloro derivatives FeSaEn-Cl and $[\text{FeSaEn-Cl}]_2$, the magnetic susceptibility data of the present work might be interpreted in terms of a mononuclear pentacoordinate structure for the benzyl derivative, $\text{FeSaEn-CH}_2\text{C}_6\text{H}_5$. This is indicated by the value of the magnetic moment at room temperature and by the relatively small value of the Weiss constant, suggesting that no important interactions take place between different iron centres, as one would expect for a dimeric, oxygen-bridged structure. Of course, the results of an X-ray investigation would be required to confirm this suggestion.

ACKNOWLEDGEMENT

The authors thank the American Cyanamid Co. for a gift of laboratory equipment to the University of Pisa and the National Research Council (CNR, Rome) for financial support.

REFERENCES

- 1 D A Clarke, D Dolphin, R Grigg, A W Johnson and H A Pinnock, *J Chem Soc C* (1968) 881
- 2 G Kobrich and H Buttner, *J Organometal Chem* 18 (1969) 117
- 3 F Calderazzo and C Floriani, *Chem. Commun.* (1968) 417
- 4 F Calderazzo and C Floriani, *Chem. Commun.*, (1967) 139
- 5 C Floriani, M Puppis and F Calderazzo, *J Organometal Chem* 12 (1968) 209
- 6 I Ya Levitkin, Z V Todrez, M E Vol'pin and D N Kursanov, *Zh Obshch Khim*, 41 (1971) 193
- 7 G Costa and G Mestroni, *Tetrahedron Lett*, (1967) 1783
- 8 P B Chock and J Halpern, *J Amer Chem Soc*, 88 (1966) 3511
- 9 A J Hart-Davis and W A G Graham, *Inorg Chem*, 9 (1970) 2658
- 10 J Lewis, F E Mabbs and A Richards, *J Chem Soc A*, (1967) 1014
- 11 C Giannotti, A Gaudemer and C Fontaine, *Tetrahedron Lett* (1970) 3209
- 12 K N V Duong, C Fontaine, C Giannotti and A Gaudemer, *Tetrahedron Lett*, (1967) 1187
- 13 C Floriani and F Calderazzo, *J Chem Soc A*, (1971) 3665
- 14 G Stolze and J Hahle, *J Organometal Chem*, 7 (1967) 301
- 15 G Costa, G Mestroni and G Pellizer, *J Organometal Chem*, 11 (1968) 333
- 16 M Gerloch, J Lewis, F E Mabbs and A Richards, *J Chem Soc A* (1968) 117