

## 1. CHROMIUM<sup>†</sup>

R. COLTON

### CONTENTS

Introduction . . . . .	1
1.1 Chromium(VI) . . . . .	2
1.2 Chromium(V) . . . . .	3
1.3 Chromium(IV) . . . . .	5
1.4 Chromium(III) . . . . .	5
1.4.1 Complexes of simple ligands . . . . .	5
1.4.2 Complexes of bidentate ligands . . . . .	8
1.4.3 Complexes of polydentate ligands . . . . .	12
1.4.4 Dinuclear complexes . . . . .	17
1.4.5 Polynuclear complexes . . . . .	18
1.4.6 Photochemistry . . . . .	20
1.5 Chromium(II) . . . . .	21
1.5.1 Simple compounds of chromium(II) . . . . .	21
1.5.2 Metal-metal bonded species . . . . .	23
1.6 Chromium(I) and chromium(0) . . . . .	24
Acknowledgement . . . . .	24
References . . . . .	24

### INTRODUCTION

The review on chromium chemistry this year deals mainly with the inorganic and coordination chemistry of the element which was cited in Chemical Abstracts, Volumes 102 and 103. Thus most of the work published in 1985 and some published towards the end of 1984 will be reviewed. Chromium organometallic chemistry, including carbonyl chemistry, is not formally included in this review but reference is made to some papers of general interest.



The organization of the review is similar to that of previous years. The various oxidation states are dealt with in turn with individual systems ordered in terms of the nature of the donor atoms. Chromium chemistry is dominated by the trivalent state so, for convenience, this large section of the review is divided into sections according to the nature of the ligands.

---

<sup>†</sup> NO REPRINTS AVAILABLE

## 1.1 CHROMIUM(VI)

The oxidation of lactic acid by chromium(VI) has been studied in detail and found to proceed in two separate and measurable steps, both of which give pyruvic acid



The buildup and decay of chromium(V) intermediates accompanied the disappearance of Cr(VI). Mechanisms were deduced for each step and it was suggested that the intervention of chromium(V) may be widespread in chromium(VI) oxidations [1].

A study has been made of the interaction at physiological pH's of chromium(VI) with a series of low molecular weight cellular reductants and other model compounds since it is known that reduction to chromium(III) is involved in the carcinogenic properties of chromium(VI). It was found that only ascorbate and compounds containing a thiol group reduced Cr(VI) at an appreciable rate. Several mechanisms of interaction were observed, but with glutathione, which is the most abundant intracellular thiol, clear evidence was obtained for the formation of a chromium(VI) thioester in a rapid pre-equilibrium step. This was followed by a slower reduction step involving a further molecule of glutathione [2].

$\text{CrOF}_4$  has been prepared by the reaction of  $\text{CrO}_3$  and fluorine at  $140^\circ\text{C}$  and 4 atm. pressure in a Monel autoclave with a watercooled lid.  $\text{CrOF}_4$  is a powerful fluorinating agent and is rapidly hydrolyzed by water. A sample isolated in a nitrogen matrix showed a strong  $\text{Cr}=\text{O}$  stretch at  $1028\text{cm}^{-1}$  and a doublet at  $746, 742\text{cm}^{-1}$  due to the chromium-fluorine stretches. An analysis of the spectrum and comparison with similar data for  $\text{MoOF}_4$  and  $\text{WOF}_4$  suggested a square pyramidal structure with a  $\text{O}-\text{Cr}-\text{F}$  angle of  $106^\circ$ . Reaction between  $\text{CrOF}_4$  and  $\text{CsF}$  under nitrogen gave  $\text{Cs}[\text{CrOF}_5]$  which has a  $\text{Cr}=\text{O}$  stretch at  $955\text{cm}^{-1}$  and a broad band at  $650-720\text{cm}^{-1}$  due to  $\text{Cr}-\text{F}$  stretches [3].

$\text{CrF}_6$  has been prepared under conditions considerably less extreme than those in the original report of this compound. Reaction between  $\text{CrO}_3$  and fluorine in a Monel reactor fitted with a watercooled lid at  $170^\circ\text{C}$  and 25 atm. pressure followed by cooling to room temperature and then to  $-63^\circ\text{C}$  led to the formation of  $\text{CrF}_6$ . Excess  $\text{F}_2$  and  $\text{CrO}_2\text{F}_2$  were pumped off to yield a mixture of red  $\text{CrF}_5$  and yellow  $\text{CrF}_6$ . Matrix isolation IR studies showed the  $\text{Cr}-\text{F}$  stretch to occur at  $759\text{cm}^{-1}$  [4].


Force fields and mean amplitudes of vibration have been calculated for  $\text{MF}_6$

and  $\text{MO}_2\text{F}_2$  (M Cr, Mo, W). The agreement between calculated and observed spectra was good [5].

## 1.2 CHROMIUM(V)

A review has appeared on the coordination chemistry of chromium(V) [6].  $\text{CrN(TTP)}$  is known to have a distorted porphyrin core while  $\text{MnN(TTP)}$  has a regular core. These structural differences are reflected in their resonance Raman spectra. Whereas the  $\text{Mn}\equiv\text{N}$  stretch at  $1052\text{cm}^{-1}$  is greatly enhanced in intensity, no such enhancement is observed for  $\text{CrN(TTP)}$  [7].

It has been shown that the most likely mechanism of the interaction between iodosylbenzene and  $\text{ClCr(TPP)}$  involves transfer of an oxygen atom to give  $\text{Cr(V)}$ , which then reacts with further  $\text{Cr(III)}$  to give  $\text{Cr(IV)}$



but in the presence of excess of  $\text{PhI=O}$  the final product is  $\text{Cr(V)}$  [8]. The kinetics of the interactions of  $\text{ClCr(TPP)}$  with a number of percarboxylic acids and hydroperoxides were also investigated. The data indicated that all the reactions proceeded with similar mechanisms involving heterolytic fission of the O-O bond [8].

1-phenyl-1,2-ethanediol is oxidized by *p*-cyano-*N,N*-dimethylaniline *N*-oxide in the presence of  $\text{ClCr(TPP)}$  as catalyst. It was shown that photo-initiation of the reaction is necessary and the likely catalytic intermediate is  $\text{ClCr(O)(TPP)}$  [9].

One-electron oxidation of  $\text{Cr(O)(TPP)}$  by electrochemical techniques gave a  $\text{Cr(V)}$  oxoporphyrin which is stable on the coulometric timescale. It is reactive towards norbornadiene to give the epoxide in a 1:1 stoichiometric reaction. A 2:1 mixture of  $\text{ClCr(TPP)}$  and iodosylbenzene gave a product which was thought to be the  $\text{Cr(IV)}$  dimer  $[\text{ClCr(TPP)}]_2\text{O}$  which disproportionates to  $\text{Cr(III)}$  and  $\text{Cr(V)}$  [10].

A number of substituted  $\text{sal}_2\text{en}$  complexes of oxochromium(V) have been isolated. The synthetic procedure involved reaction of  $\text{Cr(III)}$  triflate with the ligand and subsequent air oxidation to give  $[(\text{sal}_2\text{en})\text{Cr}(\text{H}_2\text{O})_2]\text{O}_3\text{SCF}_3$ . Oxidation with iodosylbenzene in MeCN yielded the air-stable  $[(\text{sal}_2\text{en})\text{CrO}]^+$  species. The crystal structure of  $[(7,7'\text{-Me}_2\text{sal}_2\text{en})\text{CrO}]^+$  revealed a square pyramidal structure with the oxo group apical and the Cr atom  $0.53\text{\AA}$  above the mean  $\text{N}_2\text{O}_2$  plane. Bond distances are  $\text{Cr=O} = 1.545(2)$ ,  $\text{Cr-O} = 1.809(\text{av})$  and  $\text{Cr-N} = 1.98\text{\AA}(\text{av})$ . All the complexes reacted with a variety of donor ligands

and the crystal structure of pyNO adduct showed that the additional ligand completed the octahedral coordination about the metal. EPR and magnetic data confirm the  $d^1$  configuration of these complexes [11].

The catalytic epoxidation of various olefins with iodosylbenzene may be carried out in the presence of a series of Cr(III) cations  $[\text{Cr}(\text{sal}_2\text{en})]^+$  with or without the presence of pyNO. Kinetic studies showed that the Cr(V) species  $[(\text{sal}_2\text{en})\text{CrO}]^+$  and  $[(\text{sal}_2\text{en})\text{Cr}(\text{O})(\text{pyNO})]^+$ , described above, are the active intermediates. The mechanism of the oxygen transfer involves the rate determining attack by the Cr(V) oxocation on the olefin [12].

It has been shown that the EPR signals for paramagnetic  $[\text{CrO}_3]^-$  and  $[\text{CrO}_4]^{3-}$  may be distinguished by a combination of  $g$ - and  $A$ -tensor analysis and the temperature dependence of the line widths. This arises because the electronic spin-lattice relaxation time for a species is dependent upon its coordination and point symmetry. The results indicate that several previous assignments distinguishing between these species need to be reviewed [13].

The relatively stable Cr(V) complex sodium bis(2-ethyl-2-hydroxybutanato)-oxochromate(V) has been the subject of several investigations. The chromium(V) complex disproportionates



and is most stable at pH 3-4 in the presence of a small amount of free 2-ethyl-2-hydroxybutanoic acid. Oxidation of organic substrates with this Cr(V) reagent have been studied and compared with similar oxidations using Cr(VI). Generally, oxidations with Cr(V) are faster than those with Cr(VI) (hence Cr(V) is not usually observed as an intermediate in Cr(VI) oxidations) except for the oxidation of pinacol (faster with Cr(VI)) and ethylene glycol which is oxidized at about the same rate by the two reagents. The organic substrates investigated included oxalic acid, primary and secondary hydroxy acids, alcohols etc [14].

Reduction of sodium bis(2-ethyl-2-hydroxybutanato)oxochromate(V) by  $[\text{VO}]^{2+}$  has been investigated by spectrophotometric and EPR techniques [15]. In reactions with  $[\text{VO}]^{2+}$  in less than 1:1 stoichiometry, the chromium appeared to be reduced to Cr(IV) and only after addition of further  $[\text{VO}]^{2+}$  was subsequent reduction to Cr(III) observed. The EPR signal of Cr(V) decreased as  $[\text{VO}]^{2+}$  was added until at 1:1 stoichiometry no signal was observed. The eight line spectrum of  $[\text{VO}]^{2+}$  only appeared after 2 moles had been added. Kinetic measurements also indicated the intervention of Cr(IV) in the overall reaction scheme [15]. In a similar reduction by Fe(II), spectrophotometric evidence suggested transient formation of Cr(IV), but the observation was not so clear cut as in the vanadium system [16]. In the reduction of the Cr(V) species by

Ti(III), Cr(IV) was not detected even though the reduction must pass through this stage. Hence it was deduced that Cr(IV) must react with Ti(III) at least 20 times faster than Cr(V) [17].

Matrix isolation studies show that  $\text{CrF}_5$  rapidly disproportionates in the gas phase to  $\text{CrF}_4$  and  $\text{CrF}_6$  [4]. This result therefore casts some doubt on the interpretation of electron diffraction data for  $\text{CrF}_5$  reported in the review for 1984.  $\text{CrF}_5$  reacts with  $\text{NF}_4 \cdot \text{HF}_2$  to give  $(\text{NF}_4)[\text{CrF}_6]$  and with NOF to yield  $(\text{NO})[\text{CrF}_6]$ . The hexafluorochromate(V) anion was identified by analysis and IR and Raman spectroscopies [18].

### 1.3 CHROMIUM(IV)

The compound  $\text{CrH}_4(\text{dmpe})_2$  has been prepared in two ways; by the interaction of  $\text{CrCl}_2(\text{dmpe})_2$  with lithium butyl or by the photolysis of  $\text{Cr}(\text{N}_2)_2(\text{dmpe})_2$  in hexane. This is the first hydride of Cr(IV) and the first eight coordinate complex of Cr(IV). The structure is dodecahedral with bond distances  $\text{Cr-P} = 2.255(3)$  and  $\text{Cr-H} = 1.53(3)\text{--}1.60(3)\text{\AA}$  [19].

The compounds  $\text{Sr}_2\text{CrO}_4$ ,  $\text{Ba}_2\text{CrO}_4$  and  $\text{Ba}_3\text{CrO}_5$  have been prepared by high temperature reactions between the appropriate chromate and  $\text{Cr}_2\text{O}_3$ . All contain the tetrahedral  $[\text{CrO}_4]^{4-}$  ion. Magnetic moments are close to spin-only values with small Weiss constants and weak antiferromagnetic ordering was indicated [20].

Chromyl chloride,  $\text{CrO}_2\text{Cl}_2$ , and chromyl acetate react with various alkylaromatics (toluene, ethylbenzene etc) to yield solids which, on the basis of IR and magnetic studies, contain Cr(IV) [21].

Chromium tetrafluoride has been isolated in a dilute matrix and displays bands at  $784$  and  $303\text{cm}^{-1}$  which were shown to be due to monomeric  $\text{CrF}_4$ . On warming, changes occurred in the spectrum which are consistent with a monomeric to polymeric rearrangement [4].

### 1.4 CHROMIUM(III)


#### 1.4.1 Complexes of simple ligands

X-ray diffraction and structural refinement on  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{O})][\text{Cr}(\text{CN})_6]$  at 295 and 120 K showed it to be isomorphous with  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$ . The data for both compounds were refined using an aspherical valence electron population model. The metal configuration, the observed metal-ligand overlap populations and the ammonia lone pair populations could all be analyzed in terms of a single two parameter per bond  $\sigma$ -bonding LF model, using only metal  $d$  orbitals [22]. The spin distribution in the anion of  $\text{Cs}_2\text{K}[\text{Cr}(\text{CN})_6]$  has been


determined by polarized neutron diffraction and analyzed in terms of atomic orbital populations [23].

There has been controversy in the literature as to whether the hydrolysis of aquo Cr(III)- $\alpha$ -hydroxyalkyl complexes proceeds through intervention of a water molecule *cis* to the alkyl group, or whether a solvent water molecule is involved. The hydrolysis of number of such complexes have been studied including  $[(\text{edta})\text{Cr}(\text{CR}_1\text{R}_2\text{OH})]^{2-}$  (no coordinated water), and complexes of two quadridentate ligands (nta and [15]aneN<sub>4</sub>) which have a coordinated water molecule and the  $\alpha$ -hydroxyalkyl group *cis* and *trans* respectively. The results indicate that a water molecule *cis* to the alkyl group has no major role in the hydrolysis mechanism [24]. In addition, the specific rates of reaction of the free radicals  $\cdot\text{CH}_2\text{OH}$ ,  $\cdot\text{CH}(\text{Me})\text{OH}$  and  $\cdot\text{C}(\text{Me})_2\text{OH}$  with  $[\text{Cr}^{\text{II}}(\text{edta})]^{2-}$  and  $[\text{Cr}^{\text{II}}(\text{nta})]^-$  are reported [24].

The reaction



appears to give the expected products, but the Cr(III) complex has some unusual properties which are best explained by chelation of the organo group. For example, reaction with Hg(II) does not give  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  and an organomercurial as usual [25]. It is suggested that the reaction is




Substitution of  $\text{K}_3[\text{Cr}(\text{NCS})_6]$  with  $\text{PET}_2\text{Ph}$  or  $\text{PET}_2(p\text{-tol})$  gives the new series of complexes  $[\text{Cr}(\text{NCS})_4(\text{PET}_2\text{R})_2]^-$  [26]. The solvation of the  $[\text{Cr}(\text{NCS})_4(\text{morpholine})_2]^-$  anion has been studied in water/methanol mixtures varying from 9.7 to 87.3 volume % MeOH at several temperatures. Mechanisms of substitution are discussed [27].

It has been shown that the coordinated trifluoromethane sulphonate in  $[\text{Cr}(\text{NH}_3)_5(\text{OSO}_2\text{CF}_3)]^{2+}$  is readily replaced by a large number of O and N donor ligands including  $\text{H}_2\text{O}$ , MeOH, dmsO,  $\text{OP}(\text{OMe})_3$ , dma, MeCN, urea etc thus opening the way to the synthesis of many such derivatives [28].

The displacement of water by dmsO in the second coordination sphere of  $[\text{Cr}(\text{NH}_3)_5(\text{dmsO})]^{3+}$  has been investigated in dmsO/ $\text{H}_2\text{O}$  mixtures by line broadening effects in  $^1\text{H}$  NMR spectroscopy. The number of molecules in the second coordination sphere was found to be 10 and stepwise equilibrium constants for the replacement of  $\text{H}_2\text{O}$  by dmsO were determined. The free energy increase per replacement is about  $1.42 \text{ kJmol}^{-1}$  [29].

The determination of the equilibrium constant K for the reaction



several years ago by a kinetic method and by the method of mixtures gave two values which differed by an order of magnitude. The actual values have now been verified by new determinations [30]. The discrepancy arises because medium effects and/or other specific effects seriously influence the interpretation of the kinetic data and therefore the value from the method of mixtures is preferred.

The kinetics of the stepwise aquation of *trans*- $[\text{Cr}(\text{NH}_3)_4(\text{CN})_2]^+$  to *trans*- $[\text{Cr}(\text{NH}_3)_4(\text{CN})(\text{H}_2\text{O})]^{2+}$  and finally to *trans*- $[\text{Cr}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{3+}$  have been studied spectrophotometrically in acid solution. Both aquation steps can be described in terms of an acid independent pathway ( $k_o$ ) and an acid dependent one ( $k_H$ ), the latter consequent to instantaneous and reversible (K) protonation of some of the reactant. Then

$$k_{\text{obs}} = (k_o + k_H K[\text{H}^+]) / (1 + K[\text{H}^+])$$

and although the pattern for dependence for the two steps are different for the two species, both favour the acid assisted pathway. Comparisons with the kinetics of aquation of other cyano complexes of Cr(III) are discussed in detail [31].

The emission spectra of some  $\text{CrN}_4\text{XY}$  complexes ( $\text{N}_4 = 4\text{py}, 2\text{en}$  etc) have been examined as a function of glassy solvent at 77 K. When X or Y is F or OH, the splitting of the  ${}^2\text{T}_1$  level is reduced in hydrogen bonding solvents. The solvent effect on the spectra depends upon whether the lowest component of  ${}^2\text{T}_1$  ( ${}^2\text{E}_Q$ ) is above or below  ${}^2\text{E}$  in all solvents. If it is above, the spectra are sharp and solvent independent while if it is below the spectra are broader and solvent dependent [32].

The crystal structures of *trans*- $[\text{Cr}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}_2$  and *trans*- $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{I}$  have been determined [33]. Important bond distances are

	$[\text{Cr}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]^{2+}$	$[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$
Cr-Cl	2.298(1)	2.325(1)
Cr-N	2.064(3)-2.086(3)	2.071(2)
Cr-O	2.023(2) Å	

The water exchange between solvent and  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  and  $[\text{Cr}(\text{H}_2\text{O})_5(\text{OH})]^{2+}$  has been investigated by  ${}^{18}\text{O}$  tracer studies at various pH's as a function of pressure. For  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ ,  $\ln k$  is a linear function of pressure to 400MPa implying no solvational change in the activation process [34]. The rate of aquation of  $[\text{Cr}(\text{H}_2\text{O})_5\text{F}]^{2+}$  has been determined in relatively concentrated

(1-10M)  $\text{HClO}_4$  solution and compared with the corresponding data for  $[\text{Cr}(\text{H}_2\text{O})_5(\text{N}_3)]^{2+}$  [35].

A series of complexes  $\text{CrCl}_3(\text{py} \cdot 3\text{COOR})_3$  ( $\text{R} = \text{Me, Et, Pr, Bu}$ ) have been prepared with the esters of nicotinic acid being used to prevent coordination of nicotinic acid through its carboxylate group. IR studies indicate that all the compounds have the ligand coordinated through nitrogen in a *meridional* configuration [36].

#### 1.4.2 Complexes of bidentate ligands

Reactions of substituted biguanidines ( $\text{Rbiq}$ ) and  $[\text{Cr}(\text{NCS})_6]^{3-}$  in 3:1 molar proportions in refluxing  $\text{EtOH}$  gave  $[\text{Cr}(\text{Rbiq})_3](\text{CNS})_3$  [37].

The kinetics have been determined for the quenching of the long lived excited states of  $[\text{Cr}(\text{L-L})_3]^{3+}$  ( $\text{L-L} = \text{bipy, phen}$ ), and the corresponding ruthenium complexes, by the copper proteins plastocyanin, azurin and stellacyanin. The kinetic behaviour of the reaction with plastocyanin was discussed in terms of model in which the chromium complex binds to the protein at a site remote from the copper atom [38].

The reaction of  $[\text{Cr}(\text{en})_3]\text{Cl}_3$  with  $\text{KCN}$  in  $\text{dmsO}$  at  $80^\circ\text{C}$  gave good yields of *cis*- $[\text{Cr}(\text{en})_2(\text{CN})_2]\text{Cl}$ . The stereochemistry of the cation was elucidated by ion exchange studies, vibrational spectroscopy and by resolution [39].


Irradiation of solid  $[\text{Cr}(\text{en})_3]_2(\text{C}_2\text{O}_4)_3 \cdot 3\text{H}_2\text{O}$  in the region of the LF  $4\text{A}_{2g} \rightarrow 4\text{T}_{2g}$  bands gave conversion to  $[\text{Cr}(\text{en})_2(\text{ox})][\text{Cr}(\text{en})(\text{ox})_2]$ . The yield was high and the kinetics of the reaction were studied. The products were isolated by ion exchange techniques and fully characterized [40]. The photoaquation of  $[\text{Cr}(1,3\text{-pn})_3]^{3+}$  has been studied in both acid and alkali solution by absorption spectroscopy, pH measurements and phosphorescence intensity measurements. It was concluded that two main products,  $[\text{Cr}(\text{pn})_2(\text{pnH})(\text{H}_2\text{O})]^{4+}$  and  $[\text{Cr}(\text{pn})_2(\text{H}_2\text{O})_2]^{3+}$ , were formed [41].

Room temperature circular dichroism (CD) spectra have been measured in the spin forbidden  $2\text{E}_g, 2\text{T}_{1g} \leftarrow 4\text{A}_{2g}$  transition for diastereoisomers of a number of complexes of  $\text{Cr}(\text{III})$  containing (R)- or (S)-propylenediamine or (1R,2R)- or (1S,2S)-1,2-*trans*-cyclohexanediamine. The compounds studied were of the types  $[\text{Cr}(\text{en})_x(\text{diamine})_{3-x}]^{3+}$  ( $x = 0, 2$ ), *cis*- $[\text{Cr}(\text{NH}_3)_2(\text{diamine})_2]^{3+}$  and  $[\text{Cr}(\text{NH}_3)_4(\text{diamine})]^{3+}$ . The differences in the observed CD spectra between each pair of diastereoisomers were accounted for by the separability and additivity of the configurational and vicinal CD effects [42].

*Cis*- $[\text{Cr}(\text{en})_2(\text{CN})_2]^+$  has been shown to interact in solution with other cations such as  $\text{Hg}(\text{I})$ ,  $\text{Ag}(\text{I})$  and  $\text{Cd}(\text{II})$  and the association constants have been measured. Isolation of the silver salt formed by addition of  $\text{AgCN}$  and determination of its structure revealed the interesting stereochemistry shown



in (1) which shows that linkage isomerism has occurred at the chromium atom. Bond distances were not reported [43].




A series of complexes of the type  $[\text{Cr}(\text{Rbiq})_2(\text{NCS})_2](\text{CNS})$  (Rbiq = substituted biguanidine) have been prepared by the interaction of the ligands with  $[\text{Cr}(\text{NCS})_6]^{3-}$  in 2:1 proportions in refluxing EtOH [44].

Hydrolysis of  $[\text{Cr}(\text{en})_2(\text{S-XCOO})]^+$  ( $\text{X} = \text{CH}_2, \text{CH}_2\text{CH}_2, \text{CMe}_2$ ) involves an acid catalyzed Cr-S bond cleavage as the first step. The product,  $[\text{Cr}(\text{en})_2(\text{H}_2\text{O})(\text{OOC-X-SH})]^{2+}$ , can then react in three ways:- (i) the pendant thiol can coordinate to regenerate the starting material (ii) the pendant thiol can anchimerically assist Cr-O bond cleavage to release the complete ligand or (iii) the pendant thiol can assist Cr-N bond cleavage to give further reactive products. The relative rates of these reactions are influenced by the ring size of the chelated ligand and steric effects in the comparisons between  $\text{X} = \text{CH}_2$  and  $\text{CMe}_2$  [45].

A number of mixed chelate complexes of Cr(III) containing en, 1,3-pn and fluoride have been prepared from *trans*- $[\text{Cr}(\text{en})\text{F}_2(\text{H}_2\text{O})_2]\text{Cl}$  in refluxing EtOH by the following reaction scheme (2). Heating *trans*- $[\text{Cr}(\text{en})(\text{pn})\text{FCl}]\text{Cl}$  or *cis*- $[\text{Cr}(\text{en})(\text{pn})\text{F}(\text{H}_2\text{O})]\text{Br}_2$  gave a mixture of products including *cis*- $[\text{Cr}(\text{en})(\text{pn})\text{FX}]\text{X}$  which on reaction with  $\text{NH}_4\text{F}$  gave *cis*- $[\text{Cr}(\text{en})(\text{pn})\text{F}_2]^+$  which was resolved [46]. The complete series of fluoro complexes of the type  $[\text{Cr}(\text{aa})(\text{bb})\text{F}_2]^+$  (where aa, bb are diamines such as en, pn, chxn) have been synthesized using the general reaction



The compounds were all *trans* except for *cis*- $[\text{Cr}(\text{en})(\text{pn})\text{F}_2]\text{Br}$  [47]. The



different stereochemistries for  $[\text{Cr}(\text{en})(\text{pn})\text{F}_2]^+$  reported in these two papers may well be correct and due to the different conditions of preparation.

The solid phase thermal dehydration of  $\text{trans}-[\text{Cr}(\text{en})_2\text{F}(\text{H}_2\text{O})][\text{M}(\text{CN})_4]$  ( $\text{M} = \text{Ni}, \text{Pd}, \text{Pt}$ ) to give  $\text{trans}-[\text{Cr}(\text{en})_2(\text{F})\text{NCM}(\text{CN})_3]$  has been investigated by non-thermal DSC together with isothermal and non-isothermal TG methods. The proposed mechanism is  $\text{S}_{\text{N}}1$  involving a square pyramidal activated complex and elimination of water as Frankel defects [48].

The crystal structure of  $\text{trans}-[\text{Cr}(\text{en})_2\text{F}(\text{NH}_3)](\text{ClO}_4)_2$  has been determined. As expected, the stereochemistry is octahedral with bond distances  $\text{Cr}-\text{F} = 1.862(3)$ ,  $\text{Cr}-\text{NH}_3 = 2.093(6)$  and  $\text{Cr}-\text{N} = 2.057(8)-2.088(6)\text{\AA}$  [49].

It has been found that  $l\text{-cis}-[\text{Cr}(\text{en})_2\text{Cl}_2]\text{Cl}$  undergoes thermal racemization at  $158^\circ\text{C}$  without any *cis* to *trans* conversion. This result is in accord only with the rhombic twist mechanism originally proposed for  $[\text{M}(\text{aa})_3]^{3+}$  complexes, since the alternative trigonal twist would give both racemization and isomerization [50].

$\text{Trans}-[\text{Cr}(\text{en})_2\text{Br}_2]\text{Br}\cdot\text{H}_2\text{O}$  undergoes efficient photoaquation in the solid state to give  $\text{cis}-[\text{Cr}(\text{en})_2\text{Br}(\text{H}_2\text{O})]\text{Br}_2$ . The chloride behaves similarly, but no other complex of the type  $\text{trans}-[\text{Cr}(\text{en})_2\text{X}_2]^+$  was found to be photosensitive [51].

The compounds  $(\text{aa}-\text{H}_2)[\text{Cr}(\text{aa})\text{F}_4]\text{Cl}$  ( $\text{aa} = \text{en}, \text{chxn}$ ) have been prepared by the interaction of  $\text{CrCl}_3\cdot 6\text{H}_2\text{O}$ , HF and the ligand. The crystal structure of the en complex showed an octahedral stereochemistry for the anion with  $\text{Cr}-\text{F} = 1.885(10)-1.930(4)$  and  $\text{Cr}-\text{N} = 2.079(6)\text{\AA}(\text{av})$ . The compounds hydrolysed in HBr solution to give  $[\text{Cr}(\text{aa})\text{F}_2(\text{H}_2\text{O})_2]$  [52].

$\text{CrCl}_3$  reacts with guanine (LH) in refluxing EtOH/triethylformate to give  $\text{CrCl}_3(\text{LH})_2$ . IR evidence suggested that the ligand was not bonded through the

exocyclic groups (CO and  $\text{NH}_2$ ) and that all the chlorines were terminal. A polymeric octahedral structure with one chelated and one bridging LH ligand was postulated [53].

The kinetics of the reaction between  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  and L-cysteine have been investigated. A simple first order process was observed which suggests that ring closure is rapid. The results are strikingly similar to those of an earlier study of the glycine/Cr(III) system which suggests that a  $N,O$  complex is formed initially [54]. The stability of glycine (HG) complexes of Cr(III) have been determined by paper electrophoresis to be  $[\text{CrG}]^{2+} 10^{8.07}$ ,  $[\text{CrG}_2]^+ 10^{14.32}$  and  $\text{CrG}_3 10^{19.23}$  [55].

The solubility of  $\text{Cr}(\text{acac})_3$  has been measured in  $\text{H}_2\text{O}/^t\text{BuOH}$  and  $\text{H}_2\text{O}/\text{MeOH}$  mixtures as a function of temperature and the thermodynamic parameters of solution calculated [56]. New complexes of the types  $[\text{Cr}(\text{acac})_{3-x}(\text{L-L})_x]^{n+}$  ( $\text{L-L} = \text{bipyO}_2$ ,  $3,3'\text{-Me}_2\text{bipyO}_2$ ) have been prepared. The number of conformational isomers differed in the two series as a result of the steric interactions of the methyl groups in the latter ligand. The isomers were resolved with visible and CD spectra being recorded. Isomerization occurred via an intermolecular conformation inversion of the  $\text{bipyO}_2$  ring and racemization occurred via an intermolecular twist mechanism [57].

$\text{CrO}_3$  reacts with  $\text{SO}_2$  in dmsO to give  $\text{Cr}_2(\text{S}_2\text{O}_7)_3 \cdot 24\text{dmsO}$  which is the first disulphate of Cr(III) [58].

The chiral octahedral complex  $\text{tris}((\text{-})\text{cyclic } O,O'\text{-l(R)2R dimethylethylene-dithiophosphato})\text{chromium(III)}$  has been prepared in EtOH solution. CD spectra in various solvents show a configurational inversion occurs which gives the diastereoisomeric equilibrium [59]



The sodium salt of N-benzoylglycine (hippuric acid, hipH) reacts with  $\text{CrCl}_3(\text{thf})_3$  in refluxing MeOH to give  $\text{Cr}(\text{hip})_3$  with the ligand acting as a bidentate. Mixed carboxylates  $\text{Cr}(\text{hip})_{3-x}(\text{carboxylate})_x$  were also synthesized together with some adducts  $\text{Cr}(\text{hip})_3 \cdot 2\text{L}$  ( $\text{L} = \text{py}$ ,  $\text{thf}$  etc) in which one hippuric acid group is displaced as free anion [60].

It has been found that  $\text{Pb}^{2+}$  strongly catalyzes the aquation of  $[\text{Cr}(\text{ox})_3]^{3-}$  to  $\text{cis-}[\text{Cr}(\text{ox})_2(\text{H}_2\text{O})_2]^-$ . The catalysis is greater than would be expected on the basis of a previously determined relationship (for some transition metals) between the rate of catalysis and  $K_{\text{M-ox}}$ , the formation constant for the first oxalato complex [61].

The electronic structure of  $\text{trans-}[\text{Cr}(\text{ox})_2(\text{py})_2]^-$  has been calculated by the spin polarization SCF- $X\alpha$ -SW method. The ground state for  $\text{Cr}^{3+}$  is  $^4\text{B}_{1g}$  and the calculated electronic spectrum associated with  $d-d$  and pyridine  $\pi \rightarrow \pi^*$

transitions agrees well with the experimentally observed spectrum. However, the calculated energies of ligand to metal charge transfer transitions show poor agreement [62].

The complexes *cis* and *trans*-H[Cr(mal)<sub>2</sub>(nic-N)<sub>2</sub>] were synthesized starting from Cr(II) solutions and their stereochemistries determined using <sup>2</sup>D NMR studies [63]. Complexes of sulphosalicylic acid (SsalH<sub>3</sub>) have been prepared for a number of M(III) ions including Cr(III). The species [Cr(Ssal)(H<sub>2</sub>O)<sub>4</sub>] and Cs<sub>3</sub>[Cr(Ssal)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>] were isolated and characterized by electrophoresis, IR and ion exchange studies. It was concluded that the ligand bonds through the carboxylate and phenolate oxygen atoms [64].

The 1:1 charge transfer adducts between Cr(dtc)<sub>3</sub> compounds and iodine have been investigated and compared with those formed between thiuramdisulphides and iodine. The latter form slightly stronger complexes, reflecting the lower basicity of the sulphur atoms on coordination [65].

Complexes of dibenzoyldisulphide (L) with a number of transition metal halides have been reported and CrCl<sub>3</sub> reacts to give CrCl<sub>3</sub>L. IR spectra indicate chelation of the ligand through its two oxygen atoms with sulphur taking no part in the coordination. A polymeric structure with chlorine bridges was proposed to give an octahedral arrangement about chromium [66].


#### 1.4.3 Complexes of polydentate ligands

CrCl<sub>3</sub>(thf)<sub>3</sub> reacts with K[HB(py<sub>3</sub>)<sub>3</sub>] in dichloromethane to give [CrCl<sub>3</sub>(HB{py<sub>3</sub>)<sub>3</sub>)]<sup>-</sup>, isolated as the (AsPh<sub>4</sub>)<sup>+</sup> salt. This anion reacts with nitrogen bases L (such as py, py<sub>3</sub>) to easily replace one halogen to give CrCl<sub>2</sub>(HB{py<sub>3</sub>)<sub>3</sub>)L and a crystal structure determination on the py compound showed the donor atoms of the tridentate ligand to be arranged in the *facial* arrangement. Bond distances are Cr-Cl = 2.311(1), Cr-N(py) = 2.108(3), Cr-N = 2.043(3)-2.065(3) Å [67].

The complex (L-alanine-N-acetato)(L-histidinato)chromium(III) monohydrate has been synthesized and its crystal structure determined by X-ray diffraction.

The L-alanine ligand coordinates as the dianion through two oxygens and one nitrogen while the histidine ligand coordinates through two nitrogens and one oxygen. Each ligand coordinates in a *facial* geometry, but the arrangement of donor atoms about the metal is *meridional* as shown in (3) [68].


The crystal structure of bis(salicylaldehydethiosemicarbazonato(1-))chromium(III) perchlorate, [Cr(C<sub>8</sub>H<sub>8</sub>N<sub>3</sub>OS)<sub>2</sub>](ClO<sub>4</sub>).3H<sub>2</sub>O, has been determined. The cation has a bis tridentate (S,N,O) *meridional* ligand arrangement with the S and O donor atoms of the ligands *cis* to each other and the nitrogen atoms *trans*. Bond lengths are Cr-O = 1.937, Cr-S = 2.416 and Cr-N = 2.038 Å [69].



Bond distances are

(alanine ligand)	(histidine ligand)
Cr-O(acetyl) = 1.960(3)	Cr-N(amine) = 2.062(4)
Cr-O = 1.946(4)	Cr-N(imid) = 2.036(5)
Cr-N = 2.078(5)	Cr-O = 1.957(4) Å

The compounds  $K_2[Cr(H_3L)(H_2L)] \cdot 3H_2O$  and  $K_2[Cr(H_2L)(A)] \cdot nH_2O$  have been prepared (where  $H_5L$  = glutathione (4) and A = anion of amino acids such as



L-cysteine, L-glutamic acid and L-aspartic acid). All show intense charge transfer bands associated with Cr-S bonds. Comparison with the IR spectra of other compounds suggests that glutathione is coordinated through a deprotonated sulphur and the terminal glycine (N,O) group with the glutamic acid residue also interacting with chromium [70].

The magnetic properties of the linear chain molecules  $[Cr(Pc)F]_n$  and  $[Cr(Pc)(CN)]_n$  have been measured over the temperature range 4.2-225 K. The fluoro compound shows antiferromagnetic exchange with  $J = -12.2\text{cm}^{-1}$  and  $g = 2.00$  whilst the data for  $[Cr(Pc)(CN)]_n$  indicates ferromagnetic exchange with

$J = 0.2\text{cm}^{-1}$  and  $g = 1.98$ . This is the first example of ferromagnetic coupling in a linear chain Cr(III) compound. The differences between the systems were discussed in terms of the electronegativities of the bridges and the orbitals which transmit the superexchange interactions [71].

The interactions of Cr(TPP)Cl with various nitrogen bases in dichloromethane have been investigated spectrophotometrically. It was found that axial ligation is rapid and that a correlation exists between  $\ln K$  for the equilibrium



and  $\text{pK}_a$  for the ligand, thus suggesting a predominantly  $\sigma$ -bonding interaction in the adducts [72].

Reaction between Cr(TPP)Cl with various thioglycollate esters in chloroform in the presence of  $\text{NMe}_4\text{OH}$  gave intensely coloured green solutions which are thought to contain bis(thiolate)Cr(III)(TPP) species [73]. The Soret band in the near UV spectrum was split in a manner similar to that of cytochrome  $\text{P}_{450}$  in the reduced state in the presence of carbon monoxide.

It has been found that oxidative cleavage of 1-phenyl-1,2-ethanediol to give benzaldehyde stoichiometrically occurs in the presence of Cr(TPP)Cl and the exogeneous oxidant 4-cyano-N,N-dimethylaniline-N-oxide. This carbon-carbon lyase reaction is analogous to one step in the removal of the side chain of cholesterol by cytochrome  $\text{P-450}_{\text{SCC}}$ . The process involves transfer of an oxygen atom from the N-oxide to chromium to give Cr(TPP)O followed by reaction of this species with the diol, regenerating Cr(TPP)Cl [74].

Cr(TPP) reacts with oxygen to give Cr(TPP)O and these species interact to give (TPP)Cr-O-Cr(TPP) as a blue, very air sensitive material which reacts with excess oxygen to give Cr(TPP)O. The magnetic moment of the dimer at room temperature is 1.61 BM per chromium atom and its variation with temperature indicates antiferromagnetic coupling. An IR band at  $860\text{cm}^{-1}$  is assigned to the Cr-O-Cr linkage [75]. Similar hetero dinuclear complexes (P)Cr-O-Fe(P') were prepared by the interaction of  $\text{Cr}^{\text{IV}}(\text{P})\text{O}$  and  $\text{Fe}^{\text{II}}(\text{P}')$  where P and P' are the dianions of different porphyrins. These dinuclear complexes react with only one mole of heterocyclic bases and a partial structural determination of the py adduct showed the base to be coordinated to chromium. The magnetic properties of the complexes were interpreted in terms of strong antiferromagnetic coupling between Cr(III) ( $S = 3/2$ ) and Fe(III) ( $S = 5/2$ ) centres with  $J = -130$  to  $-150\text{cm}^{-1}$  [76].

Complexes of 1,4,7,10-tetraazacyclotridecane,  $[\text{13}] \text{aneN}_4$ , (L), have been prepared for a number of first row transition metals including Cr(III). Reaction of  $\text{CrCl}_3 \cdot 3\text{H}_2\text{O}$  and L in a mixture of dmf and 2,2-dimethoxypropane gave

violet crystals of *cis*-[CrCl<sub>2</sub>L]Cl·H<sub>2</sub>O, the *cis* configuration being confirmed by the electronic spectrum [77]. There is a trend in the stereochemistries of Cr(III) complexes from [12]aneN<sub>4</sub> to [15]aneN<sub>4</sub> with the [12] and [13] ligands giving *cis* complexes exclusively, [14] gives a mixture of *cis* and *trans* while [15]aneN<sub>4</sub> gives only *trans* complexes. This trend is related to the size of the cavity in the ligand.

*Cis*-[Cr(NCS)<sub>2</sub>(tetb)]CNS has been resolved using potassium antimonyl(+)tartrate and then converted back to the thiocyanate salt (tetb = (RR);(SS)-C(5,12)-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane. It was found that the chiroptical properties show extensive solvent dependence. This was attributed to the fact that the 5 and 6 membered rings are of opposite helicity and the solvent properties depend upon which ring pair is dominant in a particular solvent [78].




The crystal structure of *trans*-[Cr(L)(H<sub>2</sub>O)Cl](NO<sub>3</sub>)<sub>2</sub> (L = 5,5,7e,12,12,14e-hexamethyl-1,4,8,11-tetraazacyclotetradecane) has been determined. The macrocyclic ligand is planar and the *sec* NH protons adopt the *meso* R(SSR) configuration and the Me groups at the 7 and 14 positions adopt an equatorial orientation. Bond distances are Cr-Cl = 2.307(2), Cr-O = 2.090(6) and Cr-N = 2.074(11)-2.102(7)Å [79].

The crystal structure determination of *trans*-[Cr(L)(NCS)(H<sub>2</sub>O)]CNS)<sub>2</sub> (L = [15]aneN<sub>4</sub>) shows that the ligand forms an equatorial belt about the metal atom with all four N-H bonds orientated to one side of the metal, away from the water ligand and towards the thiocyanato ligand. Bond distances are Cr-O = 1.997(4), Cr-NCS = 1.976(4) and Cr-N = 2.050(7)-2.108(6)Å [80].


In anhydrous solvents Cr(III) reacts with 1,5,9,13-tetraazacyclohexadecane (L) to give species containing the CrL unit which appear to exist in one *cis* and at least two *trans* forms [81].

Linear aliphatic tetraamines NH<sub>2</sub>(CH<sub>2</sub>)<sub>n</sub>NH(CH<sub>2</sub>)<sub>m</sub>NH(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub> (denoted as n,m,n-tet) have been reacted with [Cr(NH<sub>3</sub>)<sub>6</sub>]Br<sub>3</sub> and the products identified as [Cr(tet)Br<sub>2</sub>]Br. The ligands used were 1,4,7,10-tetraazadecane (2,2,2-tet), 1,4,8,11-tetraazaundecane (3,2,3-tet) and 1,5,8,12-tetraazadodecane (3,2,3-tet). With 2,3,2-tet all three possible isomers of [Cr(tet)Br<sub>2</sub>]Br shown in (5) were isolated although *cis*-α is the thermodynamically stable isomer. In the solid state *cis*-β converts to *cis*-α at 20°C and the *trans* isomer converts to *cis*-α at 200°C [82].

Reaction of Cr(III) salts with *m*-phenylenediamine and 2,3-butanedione in refluxing MeOH gave greyish green compounds which, on the basis of IR, magnetic and electronic spectral data, were formulated as containing the macrocyclic ligand (6) bound to chromium with the anions (Cl;Br;NCS) completing the octahedral geometry about the metal [83].

*cis-β**cis-α*

(5)



(6)

The decomposition reaction of  $\text{H}_2\text{O}_2$  in the presence of  $[\text{Cr}(\text{edta})(\text{H}_2\text{O})]^-$  proceeds via two pathways in which both the aquo and the hydroxy-edta complexes act as catalysts. Substitution controlled mechanisms were proposed for the reactions [84].

The studies of  $^2\text{D}$  NMR of  $\text{Cr}(\text{III})$  complexes reported in previous reviews have been extended to include fully and partially deuterated edta and related



ligands. The hexadentate coordination of cdta (= 1,2-cyclohexanediamine-N,N,N',N'-tetraacetate) and pdta (the 1,2-propanediamine analogue) was demonstrated by  $^2\text{D}$  NMR spectroscopy. For the first time assignments of the contact shifted  $^2\text{D}$  resonances were made whereas in previous studies structures had been deduced simply from the number of resonances and the possible molecular symmetries [85].

#### 1.4.4 Dinuclear complexes

New dinuclear complexes of the types  $[(\text{NH}_3)_5\text{MNCr}(\text{CN})_5]$  ( $\text{M} = \text{Rh}, \text{Ir}$ ) and  $[(\text{NH}_3)_5\text{MNCr}(\text{NO})(\text{CN})_4]$  ( $\text{M} = \text{Co}, \text{Rh}, \text{Ir}$ ) have been prepared by heating in the solid state the double salts  $[\text{M}(\text{NH}_3)_5(\text{H}_2\text{O})][\text{Cr}(\text{CN})_6]$  and  $[\text{M}(\text{NH}_3)_5(\text{H}_2\text{O})][\text{Cr}(\text{NO})(\text{CN})_5]$  respectively. The solid state deaquation-anation reactions were followed by TG measurements. Activation energies were in the range  $(101.4 \pm 5 - 127.8 \pm 5) \text{ kJ mol}^{-1}$  and these low values were taken as an indication that an  $\text{S}_{\text{N}}1$  dissociative mechanism with a square pyramidal intermediate was operating [86].

Single crystal EPR spectra have been observed over a wide temperature range for the complexes  $[\text{LCr}(\text{OH})_3\text{CrL}]\text{X}_3 \cdot n\text{H}_2\text{O}$  ( $\text{X} = \text{I}$ ,  $n = 3$ ;  $\text{X} = \text{ClO}_4$ ,  $n = 1$ ;  $\text{L} = 1,4,7$ -trimethyl-1,4,7-triazacyclononane). The zero field splittings of each excited state ( $S = 1, 2, 3$ ) in the exchange coupled system ( $-2J = 128 \text{ cm}^{-1}$ ) were derived [87].


The magnetic susceptibility of di- $\mu$ -hydroxobis[ $\{(S)\text{alaninato}\}\text{chromium(III)}$ ] trihydrate has been measured over the temperature range 1.8–290 K. The molecule is known to contain a highly distorted  $\text{Cr}_2\text{O}_2$  unit and so provides a stringent test of the proposed correlation between  $J$  and the geometry of the bridging groups. The experimental data show that a simple averaging of parameters for the two bridges is not sufficient and further theoretical work is required to fully understand the interactions in this type of system [88].

It has been found that perchlorate catalyses the oxo-bridge cleavage in  $[(\text{H}_2\text{O})_5\text{Cr}-\text{O}-\text{Cr}(\text{H}_2\text{O})_5]^{4+}$  almost as well as chloride and bromide ions. Furthermore, anation of  $\text{Cr(III)}$  is not the operative mechanism since only  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  is observed as the product in all cases and none of the inert  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]^{2+}$  was observed even under conditions when the chloride catalysed pathway predominates. The proposed mechanism involves bending of the originally linear  $\text{Cr}-\text{O}-\text{Cr}$  unit followed by protonation of the emergent lone pair on the  $\mu$ -oxo group [89]. It has been found that both strong and weak reductants, such as  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$  and ascorbic acid, also catalyse the acid hydrolysis of  $[(\text{H}_2\text{O})_5\text{Cr}-\text{O}-\text{Cr}(\text{H}_2\text{O})_5]^{4+}$  to give only  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ . Electrochemical studies on the aquo dimer show an irreversible one-electron reduction ( $E_{1/2} = +0.52 \text{ V vs SHE}$ ) which is about 0.93V more positive than for  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ . This may be

understood in terms of electron acceptance into the essentially non-bonding  $e_g(\pi)$  MO of the linear Cr-O-Cr system which disrupts  $2p_\pi(O) - 3d_\pi(Cr)$  overlap, leading to bridge rupture [90].

Chromium(III) catalyses the oxidation of EtOH by chloranilic acid (2,5-dihydroxy-3,6-dichlorobenzoquinone =  $H_2L$ ). Kinetic and spectrophotometric studies indicated formation of a binuclear complex  $Cr_2L$  as the active catalytic species [91].

The condensation of the triketone 2,2-dimethyl-3,5,7-octatrione,  $H_2PAA$ , with en gave  $(H_2PAA)_2en$ . Reaction of this ligand with  $Cr_2(OAc)_4$  and recrystallization of the product from pyridine gave  $Cr_2[(HPAA)_2en]_3 \cdot 6py \cdot H_2O$  which was examined by X-ray crystallography. Remarkably, the ligand is in the unfolded form with the two 1,3-diketone moieties bonded to different chromium atoms as shown in (7) [92].



Three such ligands are coordinated in this manner to give a Cr-Cr separation of 10.3 Å.

The structure of monoclinic  $NaBaCr_2F_9$  has been determined. The lattice is built of tilted double *cis* chains of  $[Cr_2F_9]_n^{3n-}$ , containing octahedrally coordinated chromium, linked by the  $Na^+$  and  $Ba^{2+}$  ions. Bond distances are  $Cr-F(t) = 1.86$  and  $Cr-F(b) = 1.92-1.95 \text{ Å}$  [93].

The energy dispersion of the singlet-triplet dimer exciton in  $Cs_3[Cr_2Cl_9]$  has been studied by inelastic neutron scattering at temperatures down to 1.3 K. Three exchange parameters were determined; intradimer, interdimer-intra-sublattice and interdimer-intersublattice. The systematic variation of the exchange parameters with substitution of chloride by bromide and iodide were discussed [94].

#### 1.4.5 Polynuclear complexes


It has been suggested previously that dynamic rather than static distortions best explain the magnetic susceptibility data for  $[Cr_3O(OAc)_6(H_2O)_3]^+$ . However, it has now been shown [95] that the complex

undergoes a phase transition at 211.4 K and optical spectral data clearly show the presence of two types of trimer in the solid state at low temperatures. Hence the distortions at low temperature must be static.

The complexes  $[\text{Cr}_n\text{Fe}_{3-n}\text{O}(\text{glycinate})_6(\text{H}_2\text{O})_3]^{3+}$  ( $n = 1, 2, 3$ ) have been prepared in  $\text{EtOH}/\text{H}_2\text{O}$  media at pH 2. The composition of the  $\text{CrFe}_2$  complex was confirmed by a structure determination. At low pH's glycinate can be made to coordinate through the two oxygens of the carboxylate because the amino acid is in the Zwitterion form [96].

Reaction of  $\text{Cr}_2(\text{OAc})_4$  with metal acetates in pyridine solution in air gives the complexes  $[\text{Cr}_2\text{MO}(\text{OAc})_6(\text{py})_3]$  ( $\text{M} = \text{Mg}, \text{Co}, \text{Ni}$ ) which have basically the same geometrical structure as the  $[\text{Cr}_3\text{O}(\text{OAc})_6(\text{H}_2\text{O})_3]^+$  cations. The trimetallic complexes  $[\text{CrFeMO}(\text{OAc})_6(\text{py})_3]$  ( $\text{M} = \text{Mn}, \text{Fe}$ ) were also prepared and the magnetic interactions between the metal ions were investigated in detail. They exhibit antiferromagnetic exchange and the strength of the  $\text{Cr(III)}-\text{Cr(III)}$  and the  $\text{Cr(III)}-\text{Fe(III)}$  interactions are largely independent of the nature of  $\text{M(II)}$ , but approximately double those in the  $[\text{Cr}_3\text{O}]^+$  cation. This suggests that the central oxygen provides the main superexchange pathway in these compounds [97].

Reaction of  $\text{CrCl}_3(\text{thf})_3$  with 2-hydroxyethanediol ( $\text{hetH}_2$ ) in the presence of base yielded  $\text{Cr}_3(\text{het})_6$ . A crystal structure determination revealed the interesting structure of a linear trimer with ligands chelating to the outer chromium atoms and the alkoxide groups acting as  $\mu_2$ -bridges to give an octahedral array of  $\text{O}_3\text{S}_3$  about the outer chromiums and an  $\text{O}_6$  array about the inner chromium as shown in (8) [98].



There has been a re-appraisal of the structures of  $[\text{Cr}_3(\text{OH})_4]^{5+}$  and  $[\text{Cr}_4(\text{OH})_6]^{4+}$  which are early members of the series leading to the complete polymerisation of  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  solutions as the pH is raised. It is now suggested that the unusual formulations for the aquo species (see Review for 1983) are unnecessary, as the acidic character of the complexes can be explained in terms of internal hydrogen bonding. A series of structures

involving more usual  $\mu_2$ -OH bridges and hydrogen bonding is now proposed [99].

$\text{Cr}_2(\text{mhp})_4$  (mhp = anion of 6-methyl-2-hydroxypyridine) in warm thf or diglyme gave a dark green solution from which was isolated  $\text{Cr}_4(\text{OH})_4(\text{mhp})_8$ . A crystal structure determination revealed a distorted cubane-type  $\text{Cr}_4(\text{OH})_4$  core, arranged such that there are two long Cr-Cr diagonals of  $2.974(1)\text{\AA}$  across two  $\text{Cr}_2\text{O}_2$  faces and four short Cr-Cr diagonals of  $2.829(1)\text{\AA}$ . Four of the mhp ligands bridge the short diagonals between chromium atoms, whilst the remaining four bond to chromium atoms through oxygen and H-bond to a  $\mu_3$ -OH group through nitrogen [100].

#### 1.4.6 Photochemistry

Phosphorescence lifetimes and their temperature dependencies have been determined for  $[\text{Cr}(\text{NH}_3)_5(\text{CN})]^{2+}$  and *cis* and *trans*- $[\text{Cr}(\text{NH}_3)_4(\text{CN})_2]^+$  in a variety of solvents. The lifetimes are in the 10-100 $\mu\text{s}$  range, which is relatively long for Cr(III) acidoammines. Considerable solvent effects were observed and the results are discussed in terms of the various processes effecting the lifetime of the doublet state [101].

The emission lifetimes of a number of Cr(III) complexes such as  $[\text{Cr}(\text{NH}_3)_6]^{3+}$ ,  $[\text{Cr}(\text{ND}_3)_6]^{3+}$ ,  $[\text{Cr}(\text{en})_3]^{3+}$  and *trans*- $[\text{Cr}(\text{en})_2\text{F}_2]^+$  etc have been examined in aqueous solution as a function of pressure up to 207MPa. The apparent volumes of activation range from  $0.1\text{mLmol}^{-1}$  for *trans*- $[\text{Cr}(\text{cyclam})(\text{CN})_2]^+$  to  $+6.9\text{mLmol}^{-1}$  for  $[\text{Cr}(\text{NH}_3)_5(\text{NCS})]^{2+}$ . The effects of applied pressure on the LF absorption bands of some compounds and the doublet state emission spectrum of  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  were also examined [102].

The compounds *cis* and *trans*- $[\text{Cr}(\text{cyclam})(\text{NH}_3)_2]^{3+}$  have been prepared from the corresponding dinitrato complexes in liquid  $\text{NH}_3$ . Both isomers exhibit strong phosphorescence in solution at room temperature. The emission of the *trans* isomer is very intense and long lived (136 $\mu\text{s}$  in dmsO at  $20^\circ\text{C}$ ) and shows a 12-fold enhancement on N-H deuteration. The *cis* isomer has a much shorter emission lifetime and shows only a weak deuterium isotope effect. Possible rationalizations of these effects are discussed [103].

Electronic absorption and emission spectra, lifetimes and phosphorescence yields of the lowest lying doublet states have been measured for the hexamine Cr(III) cage complexes  $[\text{Cr}(\text{sar})]^{3+}$  and  $[\text{Cr}(\text{diasar})]^{3+}$  (sar = 1,4,8,11,15,18-hexaazabicyclo[666]icosane; diasar = 3,6,10,13,16,19-hexaazabicyclo[666]-icosane). The photophysics were found to be markedly different to those of  $[\text{Cr}(\text{en})_3]^{3+}$ . This has led to a questioning of the mechanism of relaxation of the excited states in this type of molecule. The earlier work with  $[\text{Cr}(\text{sep})]^{3+}$  was interpreted as favouring surface crossing to a reactive ground state intermediate with higher coordination number, but this work favours back

intersystem crossing from the doublet state to an excited quartet state [104].


## 1.5 CHROMIUM(II)

### 1.5.1 Simple compounds of chromium(II)

The reaction of Cr(II) with the fumaronitrile complexes of Co(III) and with fumaronitrile itself have been investigated in detail. The reaction between  $[(\text{NH}_3)_5\text{Co}(\text{NCCH}=\text{CHCN})]^{3+}$  (A) and Cr(II) follows the rate law

$$-\text{dln}[A]/\text{dt} = k[\text{Cr}^{2+}]$$


with  $k = (1.9 \pm 0.2) \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$  ( $25^\circ\text{C}$ ,  $0.50 \text{M LiClO}_4/\text{HClO}_4$ ). The products of the reaction after separation by ion exchange were  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ ,  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  and fumaronitrile in equimolar proportions. When the reaction was carried out with a slight deficiency of Cr(II) an intermediate was detected spectrophotometrically which, based on its absorption spectrum and its rate of aquation to  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  was thought to be (9).



Cr(II) reacts with fumaronitrile with a complex series of colour changes from blue to red (20 mins) to blue green (200 mins) so that the reaction is not simply




although the final product was shown to be succinamide. It was shown that 2mol of  $\text{Cr}^{2+}$  per mole of fumaronitrile are involved in the first stage of the reaction and that no further  $\text{Cr}^{2+}$  is involved in the latter part of the reaction. Several intermediates were separated by ion exchange techniques and all were thought to contain Cr-C linkages, one example being (10) [105].



The reductions of Co(III) cysteinesulphonato and cysteinesulphinato derivatives containing en by Cr(II) have been investigated by stopped flow spectrophotometry [106].

When the green complex  $[\text{Cr}\{\text{OC}^t\text{Bu}_3\}_2 \cdot \text{LiCl} \cdot (\text{thf})_2]$  (described in the Review for 1983) is dissolved in warm hexane LiCl is precipitated and a blue green complex is obtained. An X-ray structural analysis showed it to be  $[\text{Cr}(\mu\text{-OCH}^t\text{Bu}_2)(\text{OC}^t\text{Bu}_3)]_2$  containing roughly trigonal three coordinate Cr(II) as shown in (11). The bridging ligands are derived from tri-*t*-butylmethoxide by elimination of  $\text{C}_4\text{H}_9$  which was detected by CG/MS analysis [107].



Bond distances are

$$\text{Cr}-\text{Cr} = 3.075(1)$$

$$\text{Cr}-\mu\text{O} = 1.974(\text{av})$$

$$\text{Cr}-\text{O} = 1.838(3)\text{\AA}$$

Electron diffraction studies on  $\text{CrCl}_2$  at  $895^\circ\text{C}$  suggested the compound exists as a bent monomeric species with about 30% of a dimeric form with two

bridging chlorine atoms also present [108]. The geometrical parameters derived from the analysis are

	monomer	dimer
Cr-Cl(t)	2.207(10)	2.208(8)
Cr-Cl(b)		2.353(10)Å
Cl(t)-Cr-Cl(t)	72°	
Cl(b)-Cr-Cl(b)		82°

There has been a detailed study of the optical spectrum of single crystals of the two dimensional ferromagnet  $(\text{MeNH}_3)_2[\text{CrCl}_4]$  in the region of the quartet to triplet LF transitions at 5330 and 6270Å from 1.6-9 K. Below 2 K there is evidence for cold band intensity arising from a small amount of spin canting [109].

The thf adduct of  $\text{CrCl}_2$  reacts with dmpe to give *trans*- $\text{CrCl}_2(\text{dmpe})_2$ . Bond distances are Cr-Cl = 2.348(av) and Cr-P = 2.368Å(av). It reacts with LiMe to give *trans*- $\text{CrMe}_2(\text{dmpe})_2$  which has bond distances of Cr-C = 2.168(4) and Cr-P = 2.345Å(av) [110].

$(\text{NMe}_4)[\text{CrBr}_3]$  has been prepared by the addition of  $\text{NMe}_4\text{Br}$  to a solution obtained by the reaction of chromium metal with HBr gas in glacial acetic acid. X-ray powder diffraction patterns are similar to those of the fully characterized chloro analogue. The magnetic susceptibility data at low temperatures was analyzed in terms of the linear chain model and it shows the compound to be antiferromagnetic [111].

### 1.5.2 Metal-metal bonded species

Generalized molecular orbital and CI calculations on  $\text{Cr}_2(\text{O}_2\text{CH})_4$  and  $\text{Cr}_2(\{\text{NH}_2\text{CH}\})_4$  at various Cr-Cr bond lengths have been reported. The calculations show that the Cr-Cr bond length in the aminoiminato complex is 0.48Å shorter than in the formate complex. This is due to the stronger donor  $(\text{NH})_2\text{CH}$  causing the chromium orbitals to expand, thus giving better Cr-Cr overlap. Although axial ligation effects the Cr-Cr distance, these theoretical studies show that the nature of the bridging group does effect it also [112]. Hartree-Fock-Slater calculations have been performed on  $\text{M}_2(\text{O}_2\text{CH})_4$  and  $[\text{M}_2(\text{O}_2\text{CH})_4]^+$  (M = Cr, Mo, W) to yield the M-M bond distances and force constants. For the cations, the three alternatives of the electron being removed from  $\sigma$ ,  $\pi$  and  $\delta$  orbitals were considered [113].

The kinetics of the reaction between  $\text{Cr}_2(\text{O}_2\text{CMe})_4$  and  $[\text{I}_3]^-$  have been studied in HOAc in the presence of NaI. The reaction is rapid and the spectrum of the final Cr(III) product closely resembles that of  $[\text{Cr}_3\text{O}(\text{OAc})_6(\text{H}_2\text{O})_3]^+$ . The kinetics support a mechanism of diiodide adduct formation (presumably axial)

before dissociation into monmeric species and subsequent oxidation to Cr(III) [114].

## 1.6 CHROMIUM(I) AND CHROMIUM(0)

The electronic and geometric structures  $\text{CrH}^+$ ,  $\text{CrCH}_3^+$ ,  $\text{CrCH}_2^+$  and  $\text{CrCH}^+$  have been investigated by *ab-initio* MCSCF and CI methods. The ligands  $\text{CH}_3$ ,  $\text{CH}_2$  and  $\text{CH}$  form single, double and triple bonds respectively with  $\text{Cr}^+$  [115].

Reduction of *trans*- $\text{CrCl}_2(\text{dmpe})_2$  under  $\text{N}_2$  by Na/Hg in thf gave red *trans*- $\text{Cr}(\text{N}_2)_2(\text{dmpe})_2$ , although if the time taken for the reaction was limited some brown crystals were isolated which are thought to be of the *cis* isomer. A crystal structure determination of *trans*- $\text{Cr}(\text{N}_2)_2(\text{dmpe})_2$  gave the bond distances  $\text{Cr-N} = 1.874(3)$ ,  $\text{Cr-P} = 2.2970(5)$  and  $\text{N-N} = 1.122(3)\text{\AA}$ . The shortness of the Cr-P bonds compared with those in *trans*- $\text{CrMe}_2(\text{dmpe})_2$  suggests that some back bonding to phosphorus occurs in the dinitrogen complex [115].

Cr atoms were co-condensed with water in an argon matrix at 15 K. An adduct was formed resulting in a decrease of the  $\nu_2$  bending mode of water by  $13.1\text{cm}^{-1}$ . Photolysis of the adduct caused insertion of chromium to give  $\text{HCr}(\text{OH})$  which is non-linear [116].

## ACKNOWLEDGEMENT

I thank Penny Panagiotidou for carefully proof reading the manuscript.

## REFERENCES

1. G.P. Haight, G.M. Jursich, M.T. Kelso and P.J. Merrill, *Inorg. Chem.* 24 (1985) 2740
2. P.H. Connert and K.E. Wetterhahn, *J. Am. Chem. Soc.* 107 (1985) 4282
3. E.G. Hope, P.J. Jones, W. Levason, J.S. Ogden, M. Tajik and J.W. Turff, *J. Chem. Soc. Dalton Trans.* (1985) 529
4. E.G. Hope, P.J. Jones, W. Levason, J.S. Ogden, M. Tajik and J.W. Turff, *J. Chem. Soc. Dalton Trans.* (1985) 1443
5. I.V. Kochikov, A.G. Yagola, G.M. Kuramshina, V.M. Kovba and Yu.A. Pentin, *Spectrochim. Acta* 41A (1985) 185
6. M.Miteva and P. Bonchev, *Coord. Chem. Revs.* 61 (1985) 241
7. C. Campochiaro, J.A. Hofmann Jr. and D.F. Bocian, *Inorg. Chem.* 24 (1985) 449
8. L-C. Yuan and T.C. Bruice, *J. Am. Chem. Soc.* 107 (1985) 512
9. L-C. Yuan, T.S. Calderwood and T.C. Bruice, *J. Am. Chem. Soc.* 107 (1985) 8273
10. S.E. Creager and R.W. Murray, *Inorg. Chem.* 24 (1985) 3824
11. K. Srinivasan and J.K. Kochi, *Inorg. Chem.* 24 (1985) 4671
12. E.G. Samsel, K. Srinivasan and J.K. Kochi, *J. Am. Chem. Soc.* 107 (1985) 7606
13. S.R. Maple and N.S. Dalal, *J. Am. Chem. Soc.* 107 (1985) 4082
14. M. Krumpolc and J. Rocek, *Inorg. Chem.* 24 (1985) 617



15. Y-T. Fanchiang, R.N. Bose, E. Gelerinter and E.S. Gould, *Inorg. Chem.* 24 (1985) 4679
16. R.N. Bose and E.S. Gould, *Inorg. Chem.* 24 (1985) 2832
17. R.N. Bose and E.S. Gould, *Inorg. Chem.* 24 (1985) 2645
18. R. Bougon, W.W. Wilson and K.O. Christe, *Inorg. Chem.* 24 (1985) 2286
19. J.E. Salt, G.S. Girolami, G. Wilkinson, M. Motavalli, M. Thornton-Pett and M.B. Hursthouse, *J. Chem. Soc. Dalton Trans.* (1985) 685
20. B.L. Chamberland, M.P. Herrero-Fernandez and T.A. Hewston, *J. Solid State Chem.* 59 (1985) 111
21. M. Sowinska and A. Bartecki, *Transition Met. Chem.* 10 (1985) 63
22. B.N. Figgis and P.A. Reynolds, *Inorg. Chem.* 24 (1985) 1864
23. B.N. Figgis, J.B. Forsyth, R. Mason and P.A. Reynolds, *Chem. Phys. Lett.* 115 (1985) 454
24. A. Rotman, H. Cohen and D. Meyerstein, *Inorg. Chem.* 24 (1985) 4158
25. P. Kita and R.B. Jordan, *Inorg. Chem.* 24 (1985) 2701
26. I. Ganescu, Cs. Varhelyi, A. Hantz and I. Papa, *Z. anorg. allg. Chem.* 520 (1985) 196
27. J. Zsako, I. Ganescu, Cs. Varhelyi and Gh. Brinzan, *Z. phys. Chemie Leipzig* 265 (1984) 1001
28. N.J. Curtis and G.A. Lawrance, *Inorg. Chim. Acta* 100 (1985) 275
29. W.L. Reynolds, L. Reichley-Yinger and Y. Yuan, *J. Chem. Soc. Chem. Commun.* (1985) 526
30. L. Monsted, T. Ramasami and A.G. Sykes, *Acta Chem. Scand.* A39 (1985) 437
31. P. Ricciari and E. Zinato, *Inorg. Chem.* 24 (1985) 441
32. A.F. Fucaloro, L.S. Forster, S.G. Glover and A.D. Kirk, *Inorg. Chem.* 24 (1985) 4242
33. J.V. Brencic, I. Leban and J. Zule, *Z. anorg. allg. Chem.* 521 (1985) 199
34. F-C. Xu, H.R. Krouse and T.W. Swaddle, *Inorg. Chem.* 24 (1985) 267
35. J.W. Vaughn and E.L. King, *Inorg. Chem.* 24 (1985) 4221
36. J. Barrett, M.K. Kormoh and P. O'Brien, *Inorg. Chim. Acta* 107 (1985) 269
37. C. Gheorghiu, C. Guran and M. Muntean, *Rev. Roum. Chim.* 29 (1984) 741
38. B.S. Brunschwig, P.J. DeLaive, A.M. English, M. Goldberg, H.B. Gray, S.L. Mayo and N. Sutin, *Inorg. Chem.* 24 (1985) 3743
39. Y. Sakabe and Y. Matsumoto, *Inorg. Chem.* 24 (1985) 2119
40. F. Benda and B. Hajek, *Z. anorg. allg. Chem.* 517 (1984) 231
41. E. Gowin and F. Wasgestian, *Inorg. Chem.* 24 (1985) 3106
42. S. Kaizaki, M. Ito, N. Nishimura and Y. Matsushita, *Inorg. Chem.* 24 (1985) 2080
43. U. Sakaguchi, K. Tomioka and H. Yoneda, *Inorg. Chim. Acta* 101 (1985) 23
44. C. Gheorghiu, C. Guran, A. Meghea and I. Jitaru, *Rev. Roum. Chim.* 29 (1984) 475
45. I.K. Adzamlı and E. Deutsch, *Inorg. Chem.* 24 (1985) 4086
46. J.W. Vaughn and J.S. Frigerio, *Inorg. Chem.* 24 (1985) 2110
47. J. Casabo, A. Solans, C. Diaz, J. Ribas, A. Segui and M. Corbella, *Transition Met. Chem.* 10 (1985) 128
48. J. Ribas, M. Serra, A. Escuer and M.D. Baro, *Thermochim Acta* 80 (1984) 103
49. K.A. Beveridge, G.W. Bushnell and A.D. Kirk, *Acta Cryst.* C41 (1985) 899
50. D. Bannerjee and J.C. Bailar Jr., *Transition Met. Chem.* 10 (1985) 331
51. A.D. Kirk and P.A. Warren, *Inorg. Chem.* 24 (1985) 720
52. C. Diaz, A. Segui, J. Ribas, X. Solans, M. Font-Altaba, A. Solans and J. Casabo, *Transition Met. Chem.* 9 (1984) 469
53. C.M. Mikulski, L. Mattucci, L. Weiss and N.M. Karayannis, *Inorg. Chim. Acta* 92 (1984) 275
54. M. Abdullah, J. Barrett and P. O'Brien, *Inorg. Chim. Acta* 96 (1985) L35
55. H.L. Yadava, S. Singh, P. Prasad, R.K.P. Singh, P.C. Yadava and K.L. Yadava, *Bull. Chim. Soc. Fr.* (1984) I-314
56. M. Yamamoto, *Bull. Chem. Soc. Jpn.* 58 (1985) 1505
57. H. Kanno, S. Utsuno and J. Fujita, *Bull. Chem. Soc. Jpn.* 57 (1984) 3443
58. B. Jeffreys, J.B. Gill and D.C. Goodall, *J. Chem. Soc. Dalton Trans.* (1985) 99
59. P. Biscarini, *Inorg. Chim. Acta* 99 (1985) 183

60. S.C. Tripathi, B.P. Baranwal and A.K. Shukla, *Synth. React. Inorg. Met-Org. Chem.* 15 (1985) 641
61. R. Banerjee, *Transition Met. Chem.* 10 (1985) 147
62. W.D. Wheeler and R.D. Poshusta, *Inorg. Chem.* 24 (1985) 3100
63. W.E. Broderick and J.I. Legg, *Inorg. Chem.* 24 (1985) 3725
64. F. Zonnevillie and G. Brunisholz, *Inorg. Chim. Acta* 102 (1985) 205
65. G. Crisponi, P. Deplano, V. Nurchi and E.F. Trogu, *Polyhedron* 3 (1984) 1241
66. I.K. Puri and J.M. Miller, *Inorg. Chim. Acta* 97 (1985) 179
67. M.J. Abrams, R. Faggiani and C.J.L. Lock, *Inorg. Chim. Acta*, 106 (1985) 69
68. M. Sato, M. Kosaka and M. Watabe, *Bull. Chem. Soc. Jpn.* 58 (1985) 874
69. M. Soriano-Garcia, J. Valdes-Martinez, R.A. Toscano and J. Gomez-Lara, *Acta Cryst.* C41 (1985) 500
70. M. Abdullah, J. Barrett and P. O'Brien, *J. Chem. Soc. Dalton Trans.* (1985) 2085
71. M. Schwartz, W.E. Hatfield, M. Joesten, M. Hanack and A. Datz, *Inorg. Chem.* 24 (1985) 4198
72. T. Ozawa and A. Hanaki, *Inorg. Chim. Acta* 102 (1985) 169
73. H. Sakurai, J. Tamura and T. Yoshimura, *Inorg. Chem.* 24 (1985) 4227
74. R.I. Murray and S.G. Sligar, *J. Am. Chem. Soc.* 107 (1985) 2186
75. D.J. Liston and B.O. West, *Inorg. Chem.* 24 (1985) 1568
76. D.J. Liston, B.J. Kennedy, K.S. Murray and B.O. West, *Inorg. Chem.* 24 (1985) 1561
77. R.W. Hay and M.A. Ali, *Inorg. Chim. Acta* 103 (1985) 23
78. A. Watson and D.A. House, *Inorg. Chim. Acta* 97 (1985) L45
79. R. Temple, D.A. House and W.T. Robinson, *Acta Cryst.* C40 (1984) 1789
80. W. Clegg, P. Leupin, D.T. Richens, A.G. Sykes and E.S. Raper, *Acta Cryst.* C41 (1985) 530
81. J. Eriksen and O. Monsted, *Acta Chem. Scand.* A38 (1984) 775
82. E.A. Pisarev and Yu.N. Shevchenko, *Russ. J. Inorg. Chem.* 30 (1985) 671
83. V.K. Chauhan, S.K. Agarwal, P. Singh and V.B. Rana, *Acta Chim. Hung.* 118 (1985) 315
84. E.N. Rizkalla, S.S. Anis and M.S. Antonious, *Inorg. Chim. Acta* 97 (1985) 165
85. W.D. Wheeler and J.I. Clegg, *Inorg. Chem.* 24 (1985) 1292
86. J. Ribas, A. Escuer and M. Monfort, *Inorg. Chem.* 24 (1985) 1874
87. S. Kremer, *Inorg. Chem.* 24 (1985) 887
88. D.E. Bolster, W.E. Hatfield and L.W. ter Haar, *Transition Met. Chem.* 10 (1985) 161
89. R.F. Johnston and R.A. Holwerda, *Inorg. Chem.* 24 (1985) 3181
90. R.F. Johnston and R.A. Holwerda, *Inorg. Chem.* 24 (1985) 3176
91. R.F. Johnston and R.A. Holwerda, *Inorg. Chem.* 24 (1985) 153
92. R.L. Lintvedt, G. Ranger and C. Ceccarelli, *Inorg. Chem.* 24 (1985) 1819
93. G. Ferey, M. Leblanc, A. de Kozak, M. Samouel and J. Pannetier, *J. Solid State Chem.* 56 (1985) 288
94. B. Leuenberger, H.U. Gudel, J.K. Kjems and D. Petitgrand, *Inorg. Chem.* 24 (1985) 1035
95. H.U. Gudel, *J. Chem. Phys.* 82 (1985) 2510
96. B.P. Straughan and O.M. Lam, *Inorg. Chim. Acta* 98 (1985) 7
97. A.B. Blake, A. Yavari, W.E. Hatfield and C.N. Sethulekshmi, *J. Chem. Soc. Dalton Trans.* (1985) 2509
98. J.R. Nicholson, R.-J. Wang, J.C. Huffman, G. Christou, H.-R. Chang and D.N. Hendrickson, *J. Chem. Soc. Chem. Commun.* (1985) 1781
99. L. Monsted, O. Monsted and J. Springborg, *Inorg. Chem.* 24 (1985) 3496
100. L. Akhtar, W. Clegg, D. Collison and C.D. Garner, *Inorg. Chem.* 24 (1985) 1725
101. E. Zinato, A.W. Adamson and P. Ricciari, *J. Phys. Chem.* 89 (1985) 839
102. S.H. Lee, W.L. Waltz, D.R. Demmer and R.T. Walters, *Inorg. Chem.* 24 (1985) 1531
103. N.A.P. Kane-Maguire, K.C. Wallace and D.B. Miller, *Inorg. Chem.* 24 (1985) 597

104. P. Comba, A.W.H. Mau and A.M. Sargeson, *J. Phys. Chem.* 89 (1985) 394
105. W.C. Kupferschmidt and R.B. Jordan, *Inorg. Chem.* 24 (1985) 3357
106. N.A. Lewis and A.M. Ray, *Inorg. Chem.* 24 (1985) 340
107. B.D. Murray, H. Hope and P.P. Power, *J. Am. Chem. Soc.* 107 (1985) 169
108. M. Hargittai, O.V. Dorofeeva and J. Tremmel, *Inorg. Chem.* 24 (1985) 3963
109. C. Bellitto, T.E. Wood and P. Day, *Inorg. Chem.* 24 (1985) 558
110. G.S. Girolami, G. Wilkinson, A.M.R. Galas, M. Thornton-Pett and M.B. Hursthouse, *J. Chem. Soc. Dalton Trans.* (1985) 1339
111. C. Bellitto, D. Fiuorani and S. Viticoli, *Inorg. Chem.* 24 (1985) 1939
112. R.A. Kok and M.B. Hall, *Inorg. Chem.* 24 (1985) 1542
113. T. Ziegler, *J. Am. Chem. Soc.* 107 (1985) 4453
114. L.M. Wilson and R.D. Cannon, *Inorg. Chem.* 24 (1985) 4366
115. A.E. Alvarado-Swaigood, J. Allison and J.F. Harrison, *J. Phys. Chem.* 89 (1985) 2517
116. J.W. Kauffman, R.H. Hauge and J.L. Margrave, *J. Phys. Chem.* 89 (1985) 3541