

# Thio-oxalates: their ligand properties and coordination chemistry

Wolfgang Dietzsch, Peter Strauch and Eberhard Hoyer

Department of Chemistry, University of Leipzig, D/O-7010 Leipzig (FRG)

(Received 20 December 1991)

## CONTENTS

A. Introduction	43
B. Topology, syntheses and structures of non-coordinated thio-oxalates	44
C. Modes of ligation in thio-oxalate complexes	47
D. Systematics of thio-oxalate complexes	48
(i) Monothio-oxalate, $[\text{O}_2\text{C}-\text{COS}]^{2-}$ , mto	51
(ii) 1,1-Dithio-oxalate, $[\text{O}_2\text{C}-\text{CS}_2]^{2-}$ , i-dto	51
(iii) 1,2-Dithio-oxalate, $[\text{SOC}-\text{COS}]^{2-}$ , dto	56
(a) Solid state structure characteristics of dto complexes	58
(b) Syntheses of dto complexes	71
(c) Magnetic properties; EPR and Mössbauer data	87
(d) Electronic spectra	93
(e) IR, Raman and resonance Raman spectra	96
(f) ESCA spectra	98
(g) Redox behaviour	101
(h) Photo- and thermochemical behaviour	103
(i) Ligand exchange reactions and mixed ligand complexes	104
(j) Analytical use of dto complexes	109
(iv) Trithio-oxalate, $[\text{SOC}-\text{CS}_2]^{2-}$ , trto	109
(v) Tetrathio-oxalate, $[\text{S}_2\text{C}-\text{CS}_2]^{2-}$ , tto	112
(a) Metal-promoted head-to-head dimerizations of $\text{CS}_2$	114
(b) Dimethyl tetrathio-oxalate, $\text{ttoMe}_2$ , and dimethyl ethenetetrathiolate, $\text{ettMe}_2$ complexes	116
(c) Authentic tetrathio-oxalate — synthesis and complex behaviour	118
(d) Related ligand systems containing the $\text{C}_2\text{S}_4$ core	121
Acknowledgements	123
References	123

## A. INTRODUCTION

Oxalates occurring in nature, as well as man made, belong to the classics of chemistry and especially of coordination chemistry. As early as 1868 the German

Correspondence to: E. Hoyer, Department of Chemistry, University of Leipzig, D/O-7010 Leipzig, FRG.

chemist Hermann Kolbe succeeded in the condensation of two  $C_1$  building blocks to form  $C_2O_4^{2-}$  by passing  $CO_2$  over molten sodium [1] according to the reaction



Of course, he extended the experiment also to  $CS_2$  in an effort to get tetrathio-oxalate, but failed.

The first paper on a thio-oxalate, namely 1,2-dithio-oxalate (dto) (and diphenyl-dithio-oxalate), had already appeared in 1909, dealing with its ligand properties towards Ni(II) and also other metal ions [2]. The authors isolated a Ni(II) bis-complex as a dipotassium salt and proposed the ligand for the analytical determination of nickel, cobalt and iron (the X-ray structure of  $K_2[Ni(dto)_2]$  had already been published in 1935; Ni(II) is square-planar  $S_4$ -chelated, and the corresponding Pd(II) and Pt(II) compounds are isomorphous [3]).

Oxalate and its thio-homologues, as compounds with a  $C_2$  backbone, belong principally to organic chemistry and are thus included in comprehensive works such as Beilstein (*Handbook of Organic Chemistry*) or Houben-Weyl (*Methoden der Organischen Chemie*). The Gmelin (*Handbook of Inorganic Chemistry*), however, also records oxalate and its coordination chemistry but not thio-oxalates. Lacking a single hydrogen atom directly connected to carbon, all thio-oxalates are strictly speaking inorganic as well. In this review we exclude most of the organic chemistry aspects of thio-oxalates, unless they are of key importance for synthesis or for structure and ligand properties in connection with coordination compounds. We direct the reader's attention to two reviews in which most of organic chemistry of thio-oxalate derivatives is summarized [4,5].

This review is divided into three main parts. After an overview on the topology and synthesis of thio-oxalates and the structures of the isolated alkali and onium salts (Sect. B), facts on the modes of ligation follow as Sect. C. Section D is devoted to the five individual thio-oxalates, and contains most knowledge up to the beginning of 1991. Papers on thio-oxalates which are mainly physically oriented are also included here, e.g. spectroscopy of all kinds and wavelengths, magnetic behaviour, electrical conductivity in the solid state, crystal chemistry, and kinetics of ligand substitution. As well as the references, a second bibliography is arranged by metal (heterobimetallic chelates are indexed twice; see Tables 4, 6 and 7).

## B. TOPOLOGY, SYNTHESIS AND STRUCTURES OF NON-COORDINATED THIO-OXALATES

Oxalate and its sulphur analogues are the smallest (composed of only six atoms) four-ligand atom ligands which provide possibilities for both low-strained five-membered (side-on) or four-membered (end-on) chelate rings for purely topological reasons. The possible alternatives (bridging or non-bridging) are obvious. The number of possible coordination modes is dependent on the number of sulphur atoms introduced instead of oxygen. In addition, further variation is introduced by the

different torsion angles of the two carboxylate halves of the ligands (between  $0^\circ$  and  $90^\circ$  for mto, i-dto, trto and tto or between  $0^\circ$  and  $180^\circ$  for dto). The principal topologies of the five thio-oxalate dianions are depicted in Fig. 1 together with the common nomenclature and abbreviations used.

There are X-ray crystal and molecular structure analyses from all but one of the dianionic representatives isolated as alkali or “onium” salts; three monomethylated monoanions and diseleno-oxalate are incorporated in Table 1. The data are self-explanatory.

The C–C bond length has the expected value for carbon–carbon single bonds with an apparently slight multiple bond contribution related more or less systematically to the number of sulphur atoms in the thio-oxalate dianion. The series oxalate  $\rightarrow$  1,2-dithio-oxalate  $\rightarrow$  tetrathio-oxalate shows a stepwise shortening of the C–C bond length:  $1.574 \rightarrow 1.516 \rightarrow 1.461$  Å.

As in oxalates, the dihedral distortion of the molecule along the C–C bond in ionic thio-oxalates is obviously connected with packing effects, although values near  $90^\circ$  are favoured, thereby minimizing the charge repulsion, maximizing electron delocalization and, in a way, “non-bond” sulphur–sulphur interaction.

Unfortunately, the precision of determination of the carbon–chalcogen bond lengths (C–Y) is normally not sufficient for answering the question whether there are two different C–Y bonds in a  $CY_2$  moiety (as found for free oxalic acid dihydrate [14] with symmetric H-bridges;) i.e. different “carbonyl” C=O and “hydroxy” C–O bond lengths.

With the exception of tetrathio-oxalate, the syntheses of the different thio-




Fig. 1. Topology of thio-oxalates (also oxalate) and abbreviations used.

TABLE 1

Structural data of oxalate, thio-oxalates and seleno-oxalates

Compound	C–C (Å)	Torsion angle about C–C (°)	Ref.
$K_2[O_2C-CO_2] \cdot H_2O^a$	1.574(2)	0	6
$Cs_2[S_2C-CO_2] \cdot CsCl \cdot H_2O$	1.50(2)	90	7
$K_2[SOC-COS]$	1.516(3)	103.5 <sup>b</sup>	8
$K_2[S_2C-COS] \cdot KCl^c$	1.52(3)	85.7	9
	1.51(3)	89.7	
$(Ph_4P)_2[S_2C-CS_2] \cdot 6H_2O$	1.461(19)	79.5(1.0)	10
$K[O_2C-COOCH_3]$	1.546(6)	1.2	11
$K[O_2C-COSCH_3]$	1.572(6)	12.5	11
$K[SOC-COSCH_3]$	1.562(9)	9.9 <sup>b</sup>	12
$K_2[SeOC-COSe]$	1.59(2)	<i>trans-planar</i>	13

<sup>a</sup>“Trans-planar”; hydrogen bonds to only two oxygens of  $C_2O_4^{2-}$ .<sup>b</sup>Referred to a planar *cis*-configuration.<sup>c</sup>Two independent  $C_2S_3O^{2-}$  anions are present.

oxalates (and diseleno-oxalate) are standard organic chemistry work starting with synthons in which the  $C_2$  backbone is already present and ready for nucleophilic attack by  $HS^-$  or  $S(e)^{2-}$  (as alkali salts), namely oxalic acid diethyl ester for monothio-oxalate, trichloroacetic acid for 1,1-dithio-oxalate, oxalyl chloride (via dialkyl- or diphenyl-dithio-oxalic acid ester) for 1,2-dithio-oxalate (diphenyl oxalic ester for diseleno-oxalate) and trichloroacetic acid phenyl ester for trithio-oxalate. The long-sought tetrathio-oxalate can be made by reductive electrodimmerization of carbon disulphide in acetonitrile in the presence of a supporting electrolyte with a cation which precipitates the tetrathio-oxalate, thus preventing follow-up reactions. Table 2 summarizes all preparative opportunities.

TABLE 2

Synthetic ways to thio-oxalate dianions

Isolated compound	Starting compound	S-Nucleophile	Ref.
$K_2[SOC-CO_2] \cdot H_2O$	$HOOC-COOC_6H_5$	KHS	15
	$H_5C_2OOC-COOC_2H_5$	KHS	16
$K_2[S_2C-CO_2]$	$Cl_3C-COOH$	$K_2S$	15
$K_2[SOC-COS]$	$ClOC-COCl$	KHS	2,17
	(via $RSOC-COSR$ )		
$K_2[S_2C-COS]$	$Cl_3C-COOC_6H_5$	KHS	15
$K_2[S_2C-COS] \cdot KCl \cdot H_2O$		$K_2S/KHS$	9
$A_2[S_2C-CS_2]^a$	$CS_2$ electroreduction		10,18
$K_2[SeOC-COSe]$	$H_5C_6OOC-COOC_6H_5$	$K_2Se$	13

<sup>a</sup>A = K,  $Me_4N$ ,  $Et_4N$ .

## C. MODES OF LIGATION IN THIO-OXALATE COMPLEXES

For purely topological reasons the modes of coordination shown in Fig. 2 have to be taken into consideration.

Taking into account the chalcogeno atom make-up of the ligands, the variation in number and geometry is remarkably large. It turns out, however, that side-on coordination is preferred no matter whether the chalcogen is oxygen or sulphur. The only authentic example of an end-on coordination was reported recently in  $[(\text{Ph}_3\text{P})_2\text{Ag}(\text{trto})\text{Ag}(\text{PPh}_3)_2]$ , where trithio-oxalate links a side-on bound silver ion with an end-on bound silver ion [19(c)]. The end-on coordination of oxalate to  $\text{Co}^{2+}$  (also end-on bridging oxalate between two  $\text{Co}^{2+}$ ) within the cavity of the macrocyclic ligand 1,4,10,13,16,22-hexaaza-7,19-dioxacyclotetracosane, proposed by Martell and Motekaitis recently [20], has not been confirmed up to now by X-ray structure analysis. Further apparent exceptions belong either to “ethenetetrachalcogenolates”  $[\text{Fe}_4(\text{CO})_{12}(\text{C}_2\text{S}_4)]$ , see below) or oxalates where four metal atoms are bound by just one bridging ligand ( $\text{Cs}_6[\text{Mo}_4\text{O}_6\text{S}_2(\text{C}_2\text{O}_4)_5] \cdot \text{H}_2\text{O}$  [21];  $\text{K}_6[\text{Mo}_4\text{O}_6\text{S}_2(\text{C}_2\text{O}_4)_5] \cdot 10\text{H}_2\text{O}$  [22]). It should be added, however, that an end-on mode is found with oxalate in two binuclear complexes containing tin(IV) or VO(II), but it turns out that the “carboxylate” side is only monodentate in both these cases [23,24] (see Fig. 3).

The same coordination type was found for bridging dto in a binuclear manga-




Fig. 2. Principal ligating functions of thio-oxalates.




Fig. 3. Monodentate oxalate bridging  $[\text{Sn}(\text{ox})_3]^{2-}$  units [23].




Fig. 4. Monodentate 1,2-dithio-oxalate bridging  $[\text{Mn}(\text{CO})_5]^+$  units [25].

nese(I) compound [25] (see Fig. 4) and is postulated also for  $\text{Cs}_4[\text{Re}_2(\text{dto})_3(\text{CO})_6]$  [26].

For stoichiometric reasons of charge, mononuclear chelates are anionic. In these cases the choice of the donor atom set is governed by “hard/soft” relationships between metal ion and ligator. This holds also for the linking of two different metal ions by (side-on bound) thio-oxalates. In addition, because the bridging ligand 1,2-dithio-oxalate is usually planar, linkage isomers can be observed [27]. Some possible cases of ligation are depicted for dto in Fig. 5. Besides X-ray evidence there are other spectroscopic means from which the coordination mode can be concluded, mainly infrared data. A representative IR treatment is given by Coucouvanis and his group in [16] for mto and in [27–29] for dto on the basis of normal coordinate analyses carried out by Nakamoto and co-workers, who also published Raman and resonance Raman spectra of dto complexes [30,31] (see also Sect. D.(iii)(e)).

Ligand “flipping” from *S,S*-chelation to *O,O*-chelation is observed with dto in  $\text{K}_3[\text{Fe}(\text{S}_2\text{C}_2\text{O}_2)_3]$  (dissolved in  $\text{CH}_2\text{Cl}_2$ ) during the reaction of the latter with  $[\text{Ag}(\text{PPh}_3)_2]^+$  (see Fig. 6). The C–O stretching frequency drops from  $1561\text{ cm}^{-1}$  in the starting material to  $1380\text{ cm}^{-1}$  in the resulting bimetallic tetranuclear chelate  $\{\text{Fe}[(\text{O}_2\text{C}_2\text{S}_2)\text{Ag}(\text{PPh}_3)_2]_3\}$  [27]; for an X-ray structure of this compound, see [32].





A similar change in only one of three ligands of the compound  $\text{K}_2[\text{Sn}(\text{S}_2\text{C}_2\text{O}_2)_3]$  (C–O stretching frequency  $1598\text{ cm}^{-1}$  [29]) was found after reaction with  $[\text{Cu}(\text{PTol}_3)_2]^+$  (Tol = tolyl,  $\text{C}_7\text{H}_7$ ) giving  $\text{K}\{(\text{dto})_2\text{Sn}[(\text{O}_2\text{C}_2\text{S}_2)\text{Cu}(\text{PTol}_3)_2]\} \cdot (\text{acetone})_2$  (C–O stretching frequencies  $1627$  and  $1371\text{ cm}^{-1}$ ; for X-ray structure see [33]); see Fig. 7.

In some cases also, changes in coordination modes of mto [16] or dto [27] in solution can be followed by IR spectrometry: the complexes  $\{\text{Cr}[(\text{S}_2\text{C}_2\text{O}_2)\text{M}(\text{PPh}_3)_2]_3\}$  (M = Cu(I) or Ag(I)), for instance, slowly undergo a kinetically controlled linkage isomerization in  $\text{CH}_2\text{Cl}_2$  yielding  $\{\text{Cr}[(\text{O}_2\text{C}_2\text{S}_2)\text{M}(\text{PPh}_3)_2]_3\}$  via  $\{\text{Cr}[(\text{SOC}_2\text{SO})\text{M}(\text{PPh}_3)_2]_3\}$ , as shown in Fig. 8.

#### D. SYSTEMATICS OF THIO-OXALATE COMPLEXES

In the following section, all reported thio-oxalate complexes (mostly chelates) up to the beginning of 1991 are collected under the heading of the individual thio-oxalates. After discussion of interesting properties, the complexes are grouped roughly following the Periodic Table, starting with the representative elements.

## a) terminal



## b) bridging












Fig. 5. Some ligation possibilities depicted for dto (outlined cases have been confirmed by X-ray structures).




Fig. 6. Ligand "flipping" from S,S'- to O,O'-chelation [27].




Fig. 7. Structure of the anion in  $\text{K}\{(\text{dto})_2\text{Sn}[(\text{O}_2\text{C}_2\text{S}_2)\text{Cu}(\text{PTol}_3)_2]\} \cdot (\text{acetone})_2$  [33].




Fig. 8. Linkage isomerization in  $\{\text{Cr}[(\text{dto})\text{Ag}(\text{PPh}_3)_2]_3\}$  [27].



(i) Monothio-oxalate  $[O_2C-COS]^{2-}$ , mto

There are as many as 27 published complexes containing monothio-oxalate as ligand. They are described in 10 papers which deal mainly with the synthesis, properties and structure of mto and its complexes. Besides introducing the ligand as the alkali salt there are two special ways of preparation. In the complex  $[Cr(mto)(en)_2]^+$  or  $[Co(mto)(en)_2]^+$  the ligand is formed by selective oxidation of coordinated mercaptoacetate (thioglycolate) through Ce(IV), Np(VI), *N*-chlorosuccinimide in *N,N*-dimethylformamide or excess acetic anhydride in dimethyl sulphoxide [34–37]



The other method was found by chance: treating  $Cs_2[Mo_2O_4(C_2O_4)_2(H_2O)_2] \cdot 2H_2O$  with  $Cs_2dto$  in the presence of  $O_2$  forms the monothio-oxalate compound  $Cs_2[MoO(S_2)_2(SOC_2O_2)]$  besides other products [21,38,39].

$[Co(en)_2(mto)]Cl \cdot H_2O$  has been found to undergo photoredox decomposition in concentrated HCl,  $HClO_4$  or water when irradiated in the near UV region (Xe arc lamp,  $NiSO_4$  or  $CoSO_4$  solution filter) giving Co(II),  $H_2S$ ,  $S_8$  and oxalate  $C_2O_4^{2-}$  as products [40].

Like other thio-oxalates mto complexes are mainly concentrated round the middle and late members of the first transition metal series. Because of the presence of two different side-on positions (*O,S* or *O,O*), mto can act both as a semi-soft and as a hard sided ligand. Examples of the latter (*O,O*) action are mono- or tetranuclear complexes of aluminium and gallium. Evidence for the *O,O*-chelate structure of these compounds was given by IR spectra [16,41]. Discussions on the IR assignments can be found under the heading “1,2-dithio-oxalate”. Very recent work deals with electron paramagnetic resonance (EPR) investigations of aqueous solutions containing different vanadyl/mto species [42]. There are three X-ray structures, of two compounds containing the complex unit  $[(en)_2Co(mto)]^+$  and of the Mo(VI) compound, mentioned above. Besides the unusually short C–O bond length involving the oxygen of the  $CO_2$  part which is not bound to the molybdenum (1.09(7) Å, compared with 1.230(3) and 1.228(7) Å in the Co(III) compounds) there are no other interesting details in the structures. It should be mentioned that mto is almost planar in the cobalt complex, the dihedral angle being 15.8° (torsion about the C–C bond) in  $Cs_2[MoO(S_2)_2(mto)]$  (see Table 3). Table 4 collects all the individual monothio-oxalates.

(ii) 1,1-Dithio-oxalate  $[O_2C-CS_2]^{2-}$ , i-dto

1,1-Dithio-oxalate is the thio-oxalate with the unique constitution of carbon-coupled  $CS_2$  and  $CO_2$  entities. Interestingly, its coordination chemistry is very scanty.

TABLE 3  
Structural data of the monothio-oxalate ligand in its complexes

Compound	Distances in Å							Ref.
	M-S	M-O	C-S (ligator atoms)	C-O	C-C	C-O (-COS)	C-O (-CO <sub>2</sub> )	
[Co(mto)(en) <sub>2</sub> ][Cl·H <sub>2</sub> O] <sup>a</sup>	2.241(1)	1.920(2)	1.723(3)	1.281(3)	1.546(4)	1.211(4)	1.230(3)	37
[Co(mto)(en) <sub>2</sub> ] <sub>2</sub> (S <sub>2</sub> O <sub>6</sub> )·2H <sub>2</sub> O <sup>a</sup>	2.237(2)	1.904(4)	1.702(6)	1.269(7)	1.560(11)	1.217(8)	1.228(7)	36
Cs <sub>2</sub> [MoO(S <sub>2</sub> ) <sub>2</sub> (mto)] <sup>b</sup>	2.53(2)	2.32(4)	1.61(6)	1.29(7)	1.53(8)	1.22(7)	1.09(7)	39

<sup>a</sup>Torsion angle in mto near 0°.

<sup>b</sup>Torsion angle in mto 15.8°.

TABLE 4  
Monothio-oxalate complexes

M	No.	Compound	M.p. (°C)	Colour (solid state)	Magnetism ( $\mu_B$ )	IR ( $\bar{\nu}_{C-O}$ )	Electronic spectra ( $\bar{\nu}_{max} \times 10^{-3} \text{ cm}^{-1} (\epsilon_0)$ )	Ref.
K		see compound 19						
Cs		see compound 18						
Ca		see compound 19						
Ba	1	Ba(mto) <sup>a</sup>						
Al	2	(Ph <sub>4</sub> P) <sub>3</sub> [Al(mto) <sub>3</sub> ]·CH <sub>2</sub> Cl <sub>2</sub>	183–5	Yellow	Diam.	1671s 1422s 1319s		16
Al	3	{Al[(mto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	191–3	Yellow	Diam.	1668s 1460s	27.8 (9700)	16
Al	4	{Al[(mto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }·CH <sub>2</sub> Cl <sub>2</sub>	177–9	Yellow	Diam.	1671s 1456s 1353s		16
Ga	5	(Ph <sub>4</sub> As) <sub>3</sub> [Ga(mto) <sub>3</sub> ] <sup>b</sup>			Diam.	1670s 1420 1315		41
Ga	6	{Ga[(mto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	173dec	Light green	Diam.	1660s 1460s 1330s		41
In	7	(Ph <sub>4</sub> As) <sub>4</sub> [In <sub>2</sub> (mto) <sub>5</sub> ]	158dec	Colourless	Diam.	1660s 1620s 1560m 1520m 1380m 1320m		41
Pb	8	Pb(mto) <sup>a</sup>						43
Cu	9	(BzPh <sub>3</sub> P) <sub>2</sub> [Cu(mto) <sub>2</sub> ]	117–8	Green	1.89(2)	1631s 1588s 1325s	16.7 (233) 29.4 (7600)	16
Cu	10	(BzPh <sub>3</sub> P) <sub>2</sub> [Cu(mto) <sub>2</sub> ]·2H <sub>2</sub> O	105–6	Violet	1.80(2)	1645s 1590s	16.7 (265) 18.3 (236)	16

TABLE 4 (continued)

M No.	Compound	M.p. (°C)	Colour (solid state)	Magnetism ( $\mu_B$ )	IR ( $\bar{\nu}_{C-O}$ )	Electronic spectra ( $\bar{\nu}_{max} \times 10^{-3} \text{ cm}^{-1} (\epsilon_0)$ )	Ref.
Cu	see also compounds 3,14,15,20,23				1316s	26.3 (6800)	
Ag	see compounds 4,6,21						
Zn 11	(BzPh <sub>3</sub> P) <sub>2</sub> [Zn(mto) <sub>2</sub> ]	181	White	Diam.	1640s 1586s 1330s		16
V 12	(Ph <sub>4</sub> As) <sub>2</sub> [VO(mto) <sub>2</sub> ]	163–7	Grey	EPR <sup>c</sup> : A = 170.7 g = 1.996	1675s 1640s 1310s	15.3sh(140) 17.3 (160) 20.8 (160) 23.1sh(180) 34.4 (5440) 16.4 (204)	44
Cr 13	(BzPh <sub>3</sub> P) <sub>3</sub> [Cr(mto) <sub>3</sub> ]·CH <sub>2</sub> Cl <sub>2</sub>	144–5	Green	3.90(2)	1640s 1583s 1320s		16
Cr 14	{Cr[(mto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }·CH <sub>2</sub> Cl <sub>2</sub> (Cr–SO isomer)	151–2	Green	3.84(2)	1613s 1548s	16.7(265)	16
Cr 15	{Cr[(mto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }·CH <sub>2</sub> Cl <sub>2</sub> (Cr–OO isomer)	166–7	Yellow	Diam. <sup>d</sup>	1364s 1632s 1435s 1342s	17.3 (134) 28.6 (11000)	16
Cr 16	[(en) <sub>2</sub> Cr(mto)]Cl <sup>e</sup>				1670 1630 1340		36
Cr 17	[(en) <sub>2</sub> Cr(mto)]I						34
Mo 18	Cs <sub>2</sub> [MoO(S <sub>2</sub> ) <sub>2</sub> (mto)]		Deep red to black		1635vs 1590s 1570s		39

Fe 19	KCa[Fe(mto) <sub>3</sub> ]·3H <sub>2</sub> O	350	Red	5.74(2)	1338s 1639m 1592s	16
Fe 20	{Fe[(mto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	147	Violet	5.69(2)	1302s 1632s 1421s	16
Fe 21	{Fe[(mto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	121–3	Orange	4.71(2)	1333s 1629s 1428s	16
Co 22	(BzPh <sub>3</sub> P) <sub>3</sub> [Co(mto) <sub>3</sub> ]·CH <sub>2</sub> Cl <sub>2</sub>	126–8	Green	Diam.	1317s 1632s 1578s	16
Co 23	{Co[(mto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }·CH <sub>2</sub> Cl <sub>2</sub>	154–6	Green	Diam.	1332s 1615s 1555s	16
Co 24	[(en) <sub>2</sub> Co(mto)] <sub>2</sub> (S <sub>2</sub> O <sub>6</sub> )·2H <sub>2</sub> O				1380s	36
Co 25	[(en) <sub>2</sub> Co(mto)]Cl·H <sub>2</sub> O				1660	37
Co 26	[(en) <sub>2</sub> Co(mto)]ClO <sub>4</sub>				1625	37
Ni 27	(BzPh <sub>3</sub> P) <sub>2</sub> [Ni(mto) <sub>2</sub> ]	181–2	Brown	Diam.	1340 1651s 1610s 1302s	16

<sup>a</sup> Mentioned in ref. 43 without details.

<sup>b</sup> No pure sample.

<sup>c</sup> In frozen pyridine solution.

<sup>d</sup> Seems likely to be erroneous.

<sup>e</sup> Only mentioned in ref. 36.

The first authentic example exhibiting a composition  $\{[(\text{Ph}_3\text{P})_2\text{Ag}]_2(i\text{-dto})\}$  is mentioned in ref. 45. Very recently, the structure of the analogous binuclear Cu(I) compound could be confirmed by X-ray analysis [19(d)]. Figure 9 shows the arrangement of the bridging 1,1-dithio-oxalate ligand linking two Cu(I) ions (see also Table 15).

A series of heterobimetallic *i*-dto complexes has also been isolated and characterized [19(f),(g)] (see Table 5). The EPR spectrum of  $[(\text{Ph}_3\text{P})_2\text{Cu}^{\text{I}}(i\text{-dto})\text{Cu}^{\text{II}}(i\text{-dto})\text{Cu}^{\text{I}}(\text{PPh}_3)_2]$  confirms the  $\text{Cu}^{\text{II}}\text{S}_2\text{O}_2$  central entity. Preliminary EPR results for vanadyl *i*-dto species in solution have been published [42].

The title of a paper on “(1,1-dithio-oxalato-*S,S'*)bis-( $\mu_3$ -sulphido)-2,2,3,3-tetrakis(trimethylphosphine)-triangulo-tripalladium(II)” is misleading because the starting ligand, as well as the ligand found in the complex, is in fact 1,2-dithio-oxalate [46(b)].

(iii) 1,2-Dithio-oxalate  $[\text{SOC}-\text{COS}]^{2-}$ , *dto*

Dto is the thio-oxalate with the most extended coordination chemistry reported. There are more than 70 papers on its transition metal as well as its non-transition metal complexes starting as early as 1909 [2] with Ni(II), Co(III) and Fe(III). The ligand can even be purchased commercially as the dipotassium salt  $\text{K}_2\text{dto}$  (CAS registry number 20267–56–5) at a reasonable price (1986), about 50 times more expensive than potassium oxalate but only half the price of toluene-3,4-dithiole, another important 1,2-dithio ligand. Its synthesis runs from oxalyl chloride via the corresponding dithio-oxalic ester and the solvolytic cleavage with KHS. The standard




Fig. 9. Molecular structure of  $[(\text{Ph}_3\text{P})_2\text{Cu}(i\text{-dto})\text{Cu}(\text{PPh}_3)_2]$  (phenyl rings omitted) [19].

TABLE 5  
1,1-Dithio-oxalate complexes

Compound	M.p. (°C)	Colour	Electronic spectra ( $\bar{\nu}_{\max} \times 10^3 \text{ cm}^{-1} (\epsilon_0)$ )	Ref.
$\{[(\text{Ph}_3\text{P})_2\text{Cu}]_2(i\text{-dto})\}$	193–4	Brown	22.4 (11700)	19(e), (f), (g)
$\{[(\text{Ph}_3\text{P})_2\text{Ag}]_2(i\text{-dto})\}$	140–3	Light brown	26.2 (4100) 21.5 (2900)	19(e), (f), (g)
$\{\text{Cu}[(i\text{-dto})\text{Cu}(\text{PPh}_3)_2]_2\}$	145–8(dec.)	Dark brown	22.0 (13500)	19(f), (g)
$\{\text{Ni}[(i\text{-dto})\text{Cu}(\text{PPh}_3)_2]_2\}$	168–71(dec.)	Dark brown	27.2 (10300) 23.6 (13000) 16.7 (2900)	19(f), (g)
$\{\text{Zn}[(i\text{-dto})\text{Cu}(\text{PPh}_3)_2]_2\}$	181–3(dec.)	Brown	15.3 (3200) 26.6 (10300)	19(f), (g)
$\{\text{Cu}[(i\text{-dto})\text{Ag}(\text{PPh}_3)_2]_2\}$	167–9(dec.)	Dark brown	22.3 (15300) 22.7 (4100)	19(f), (g)

method of preparing complexes starts with the alkali salts of dto, which are water soluble and fairly stable (see below).

The common feature of thio-oxalates, namely the ability to extend the coordination of binary complexes to further metal-containing species yielding definite multinuclear arrangements, is especially evident, and has indeed been widely investigated with dto itself. Important contributions to this subject were given by Coucouvanis and his group in a series of papers and summarized by himself in a conference volume [47].

*(a) Solid-state structure characteristics of dto complexes*

Table 6 allows a comparison of selected bond distances and angles in complexes containing terminal or bridging dithio-oxalate ligands as well as a few non-complex dithio-oxalate compounds.

**Mononuclear complexes.** For mononuclear chelates of dto, the structural characteristics of  $K_2[Ni(dto)_2]$  are representative in several respects. Its structure has been examined three times by different authors, Cox et al. in 1935 [3], Coucouvanis et al. in 1973 [28], and Gleizes et al. in 1979 [60]. Curiously enough, in the last structural study the authors found a second polymorphic “black” form. By treating the normal “red” form with  $K_2Cr_2O_7$  in aqueous solution, and after evaporation, a blend of “red” and “black” crystals is obtained with similar but different unit cells. In the “red” form (C2/c; “black” form  $P2_1/n$ ) the  $S_2C_2O_2$  moieties and the  $KO_2$  units are nearly coplanar; however, both dto planes are twisted, causing a dihedral angle of  $9^\circ$ . In contrast to the “red” form, the “black” one exhibits a linear stacking of planar  $[Ni(dto)_2]^{2-}$  anions, which are tilted with respect to the stacking axis (see Fig. 10). Similarly, this holds also for  $Li_2[Ni(dto)_2] \cdot 2H_2O$  and  $Na_2[Ni(dto)_2] \cdot 2H_2O$  [59]. Alkali and alkaline-earth counter ions influence the molecular structures of the negatively charged dto complex ions. This is because these cations are coordinated by the dto oxygens (compare also  $Ba[Ni(dto)_2] \cdot 6H_2O$  [62]). Strictly speaking, dto acts as bridging ligand in the former complexes. For quantitative reasons, however, we record the compounds with alkali and alkaline-earth cations under the heading “mononuclear”. In corresponding complexes with organic “onium” cations (e.g. pyridinium) [73,74(a),76] or ethylenediammonium [75] this interaction is lacking.

There are two copper bis-complexes differing only in the oxidation number (and of course in the number of cations) [51,52]. Whereas the Cu(II) bis-dto chelate was investigated as the  $[K(18-crown-6)]^+$  salt, the Cu(III) bis-dto chelate could be isolated and crystallographically characterized as the  $[(Ph_3P)_2N]^+$  salt. In the case of  $(TTF^+)[Cu(dto)_2]$  (TTF=tetrathiafulvalene), decomposition occurs when the complex is exposed to X-rays, preventing the resolution of the crystal structure [77(a)]. Both  $[Cu(dto)_2]^{2-}$  and  $[Cu(dto)_2]^-$  anions are planar and located on the centres of symmetry. Crown ether chelated  $K^+$  ions are bound further to the alpha-diketone sides of the  $[Cu(dto)_2]^{2-}$  complex. The major differences between the





TABLE 6 (continued)

M	No.	Compound	M-S	M-O	C-S	C-O	C-C	Intriligand		Torsion angle along S-C-C-S	Ref.
								S-M-S	O-M-O		
Cu	8	K(18-crown-6)] <sub>2</sub> [Cu(dto) <sub>2</sub> ]-DMF	2.268(2) 2.253(2)	K-O 2.777(4)	1.703(5)	1.222(6)	1.545(7)	88.85(7)		Planar	51,52
Cu	9	[(Ph <sub>3</sub> P) <sub>2</sub> N][Cu(dto) <sub>2</sub> ]	2.164(1) 2.178(1)		1.730(5)	1.217(6) 1.186(6)	1.524(7)	87.48(5)		Planar	51,52
Cu	10	[CuMn(dto) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ]-4.5H <sub>2</sub> O <sup>c</sup>	2.274(6)	2.31(1)	1.70(2)	1.24(2)	1.51(2)	91.7(2)	69.2(5)	Nearly planar	53
Cu	see also compound 7										
Ag	11	{[(Ph <sub>3</sub> P) <sub>2</sub> Ag] <sub>2</sub> (dto)}	2.513(2)	2.510(6)	1.717(8)	1.213(9)	1.561(11)			Trans-planar	45
Ag	see also compounds 5,19										
Zn	12	(Ph <sub>4</sub> As) <sub>2</sub> [Zn(dto) <sub>2</sub> ] <sup>a</sup>	2.311(2) 2.322(2)		1.728(7) 1.741(7)	1.203(9) 1.218(8)	1.566(11) 1.531(10)	95.6(1) 95.2(1)		31.7(2) 43.3(2)	54
Zn	see also compound 50										
Y, La, Ce, Nd, Sm, Eu, Gd, Dy, Er, Yb	see compounds 31–46										
Th	see compound 47										
Mo	13	Cs <sub>2</sub> [MoO(dto)] <sub>2</sub> S <sub>2</sub>	2.42(3)	Mo=O 1.70(4)	1.65(5)	1.28(7)	1.62(8)	79.0(9)		Planar	21
Mo	14	Cs <sub>2</sub> [MoO(dto) <sub>2</sub> ]-H <sub>2</sub> O	2.407(2) 2.379	Mo=O 1.660(8)	1.725(10) 1.737	1.22(1) 1.209	1.54(1) 1.555	85.3(1)		2.0, 15.1 Trans-planar	39 25
Mn	15	{[(CO) <sub>5</sub> Mn] <sub>2</sub> (dto)}		Tc=N							55
Mn	see also compounds 10,48,61,65										
Tc	16a	(Ph <sub>4</sub> As) <sub>2</sub> [TcN(dto) <sub>2</sub> ] <sup>c</sup>	2.387(2) 2.394(2)	Tc=N 1.613(4) 1.606(7)	1.727(5) 1.740(6)	1.213(10) 1.211(8)	1.549(12) 1.558(9)	86.2(1) 86.2(1)			56
Tc	16b										55
Tc	17	Ph <sub>4</sub> As[TcO(dto) <sub>2</sub> ]	2.329(1)	Tc=O 1.646(4)	1.758(6)	1.199(7)	1.528(10)	86.8(1)			



TABLE 6 (continued)

M	No.	Compound	M-S	M-O	C-S	C-O	C-C	Intraligand		Torsion angle along S-C-C-S	Ref.
								S-M-S	O-M-O		
Ni	27	$[\text{Ca}_4(\text{H}_2\text{O})_8(\text{C}_2\text{O}_4)][\text{Ni}(\text{dto})_2]_2 \cdot 6\text{H}_2\text{O}^j$	2.14(1) 2.174(4) 2.174(4)	2.48(3) 2.468(9) 2.500(9)	1.70(5) 1.68(1) 1.70(1)	1.26(5) 1.26(1) 1.25(1)	1.52(6) 1.51(2)	94.4(5) 92.2(1) 87.8(1)	64.6(3) 78.4(4)		61
Ni	28	$\text{Ba}[\text{Ni}(\text{dto})_2] \cdot 6\text{H}_2\text{O}^k$	2.177(3)	Ba-O 3.045(6)	1.713(9)	1.226(10)	1.548(12)	92.0(2)			62
Ni	29	$(\text{BzPh}_3\text{P})_2[(\text{dto})\text{Ni}(\text{dtoSnCl}_4)]$	dto 2.156(9) 2.136(7)		1.76(2) 1.78(2)	1.17(3) 1.13(3)	1.61(3)	91.7(3)		Planar	28
			$\mu\text{-dto}$	Sn-O	1.77(3)	1.19(3)	1.48(3)	91.2(3)			
			2.171(8) 2.159(8)	2.168(16) 2.154(13)	1.78(2)	1.20(3)					
Ni	30	$(\text{BzPh}_3\text{P})_2[\text{Ni}(\text{dtoSnCl}_4)_2]$	2.183(2)	2.212(5)	1.670(8)	1.264(9)	1.513(10)	92.6(1)		Planar	28
Ni	31	$[(\text{RE})_2(\text{H}_2\text{O})_{2n}][\text{Ni}(\text{dto})_2]_3 \cdot x\text{H}_2\text{O}$	Ni(1)-S 2.185(2) 2.174(2)	Eu-O 2.544(4) 2.484(5)	1.719(7) 1.706(7) 1.697(6)	1.206(9) 1.216(8) 1.250(7)	"terminal dto's" 1.498(9) 1.524(1)	O-Eu-O 91.69(7) 91.93(7)		Nearly planar	63,64
	-40	RE = Y, La, Ce, Nd, Sm, Eu, Gd, Dy, Er, Yb <sup>l</sup>	2.174(2) Ni(2)-S 2.182(2) 2.174(2)	2.482(5) 2.533(4) 2.533(4)	1.697(6) 1.686(6)	1.239(8) 1.259(8) 1.267(7)		62.66 63.47			
			Ni(1)-S 2.177(2)	Yb-O 2.170(2)	1.709(6) 1.703(6)	"terminal dto's" 1.234(7) 1.223(7)		O-Yb-O 91.75(6)			
			2.196(2) 2.189(2)	2.359(4) 2.329(4)	1.674(6) 1.684(6)	1.251(7) 1.248(7)	1.528(7)	92.93(7)	67.33 66.30		
			Ni(2)-S 2.188(2) 2.182(2)	2.341(4) 2.395(4)	1.679(6) 1.690(5)	1.256(7) 1.238(6)	1.521(8)	92.79(6)			

[illegible]

TABLE 6 (continued)

M	No.	Compound	M-S	M-O	C-S	C-O	C-C	Intraligand	Torsion angle along		Ref.
									S-M-S	O-M-O S-C-C-S	
Pt	62	$K_2[Pt(dto)_2]^{b,a}$									3
Pt	63	$(TTF)_2[Pt(dto)_2]^{c,z}$									77
Pt	64	$(TTF)_2[Pt(dto)_2]^*$	2.31(12)		1.735(8)	1.232(10)	1.550(9)	89.57(6)		Planar	77
Pt	65	$[PtMn(dto)_2(H_2O)_3] \cdot 4.5H_2O^e$		Mn-O 2.295(11)	1.70(4)	1.30(4)	1.45(5)	90.3(4)	O-Mn-O 68.6(10)	Planar	53

<sup>a</sup>Compounds  $\{M[(dto)Ag(PPh_3)_2]_3\}$  (M = Al, Fe) are isomorphous.

<sup>b</sup>"Bridging" dto ligand between Sn and K, S,S'-bound to Sn; two different kinds of dto ligands: one chelates the potassium, while the other one acts additionally as a monodentate ligand for another K.

<sup>c</sup>Compounds  $[AMn(dto)_2(H_2O)_3] \cdot 4.5H_2O$  (A = Cu, Ni, Pd, Pt) are isostructural. Bimetallic one-dimensional compounds with dto; infinite zigzag chain molecules  $\{[Mn(H_2O)_3](O_2C_2S_2)A(S_2C_2O_2)\}$  stacked chains; each layer of stacked chains is separated from the next one by intervening water molecules.

<sup>d</sup>Two different dto ligands, tetrahedral angle = 86.8°.

<sup>e</sup>Two different modifications (unit cells) are reported.

<sup>f</sup>Three different ligands (1), (2), (3).

<sup>g</sup>Compounds  $M_2[Ni(dto)_2] \cdot 2H_2O$  (M = Li, Na) are isostructural.

<sup>h</sup>Compounds  $K_2[M(dto)_2]$  (M = Ni, Pd, Pt) are isomorphous; in  $K_2[Ni(dto)_2]$  there is twisting of the ligands by 9° relative to each other.

<sup>i</sup>Formed from the "normal red form" by reaction with  $K_2Cr_2O_7$ . Totally planar  $Ni(dto)_2$  units stack along the *b* direction. There is a straightforward relationship between the structures of "red" and "black" form.

<sup>j</sup>" $[Ca_3Ni_2]$ " results from " $[CaNi]$ " in aqueous medium, where the dto is partly transformed into oxalate; both structures are built of zigzag chains of nickel atoms and either tetrahydrated or dihydrated calcium atoms with dto as bridging ligands. These chains are stacked into layers: in " $[CaNi]$ " the layers are weakly bonded to each other by a few intervening  $H_2O$  molecules. In contrast, in " $[Ca_3Ni_2]$ " the layers are firmly bridged by oxalate anions at the Ca atoms.

<sup>k</sup>It is of one-dimensional type, consisting of flat  $[Ni(dto)_2]^{2-}$  anions stacked in a zigzag way with  $Ba^{2+}$  and  $H_2O$  filling the tunnels between the anionic columns. The anionic stacking differs from the "roof-tile-like" type observed in the one-dimensional alkali derivatives of  $[Ni(dto)_2]^{2-}$ .

<sup>1</sup>The former part of the series (up to Dy) crystallizes in the monoclinic system ( $P2_1/c$ ) and the latter part crystallizes in the triclinic system ( $P1$ ). The monoclinic network exhibits channels running parallel to the direction  $[100]$  in which some water molecules are inserted. The triclinic structure can be described as a pseudo-lamellar structure, water molecules being inserted between the layers. There are two different Ni atoms in each heteropentanuclear entity.

<sup>m</sup>Infinite chains  $\cdots La-O_2C_2O_2-La \cdots$ , grafted on each La atom by a  $[Ni(dto)_2]^{2-}$  group.

<sup>n</sup>No detailed structural data reported.

<sup>o</sup>Two La(III) atoms are bridged, in an infinite way, by one glycine with contacts  $La-La$  respectively equal to 6.204(1) and 6.248(1) Å.

<sup>p</sup>Cyclic tetramers  $(YbO_2C_2O_2)_4$  are linked together by  $[Ni(dto)_2]^{2-}$  forming infinite chains.

<sup>q</sup>Two Yb(III) atoms are twofold bridged, in an infinite way, by two glycine ligands thus forming a dinuclear entity with a short Yb–Yb distance of 4.734 Å.

<sup>r</sup>Bidimensional structure with layers of  $[Ce_2(C_4O_4)]_\infty$  parallel to plane (001) and  $[Ni(dto)_2]^{2-}$  anions branched on the cerium atoms.

<sup>s</sup> $\beta$ -Alanine and oxalate bridge several positions of the cerium atom, and the  $[Ni(dto)_2]^{2-}$  is branched on the cerium atom.

<sup>t</sup>Heterotrinuclear entity  $Th(H_2O)_6Ni_2(dto)_4$ .

<sup>u</sup>Binuclear chain, ribbons  $\cdots Ni(S_2C_2O_2)Zn(O_2C_2S_2)Ni \cdots$ , Zn is octahedrally surrounded by six oxygen atoms (four from dto, two from water molecules; the remaining water molecules are bound to only 4% of all Ni atoms, giving non-stoichiometry).

<sup>v</sup> $C_{12}N_2H_{12}^{2+} = 6,7$ -dihydrodipyrido[1,2-*a*:2',1'-*c'*]pyrazinium.

<sup>w</sup> $C_5H_6N^+$  = pyridinium (some crystallographic data reported also for the 3-methylpyridinium, 3-ethylpyridinium, 4-methylpyridinium and 4-ethylpyridinium salts).

<sup>x</sup>The crystal structure consists of mixed stacks of donor  $(TTF^+)_2$  dimers and  $[Pd(dto)_2]^{2-}$  acceptors.

<sup>y</sup>The anion possesses a “chair” conformation. The dihedral angle between the  $PdS_4$  plane and the  $S_2C_2O_2$  plane is ca.  $8^\circ$ .

<sup>z</sup>Compounds  $(TTF)_2[M(dto)_2]$  ( $M = Pd, Pt$ ) are isomorphous.

<sup>\*</sup>Two different TTF species are present:  $[(TTF^+)_2(TTF^+)] [Pt(dto)_2]$ .




Fig. 10. View down the stacking axis of the unit cell of "black"  $K_2[Ni(dto)_2]$  [60].

molecular structures of the  $[Cu(dto)_2]^{-,2-}$  anions lie in the different mean Cu–S bond lengths: Cu(II)–S is 2.260 Å and Cu(III)–S 2.171 Å [52]. The latter value agrees very well with analogous distances in other Cu(III) dithiolates [79,80]. Shorter bonds in the "higher valent" metal complexes compared with longer ones for metal(II) are generally expected and are found also for other dithiolates [81,82]. Zn(II) forms a tetrahedral dto bis-complex  $[(C_6H_5)_4As]_2[Zn(dto)_2]$  (dihedral angle 86.8°). In addition there are two different torsion angles between the two COS entities along the C–C axis in the two ligands: 31.7° and 43.3° [54].

In  $KCa[Co(dto)_3] \cdot 4H_2O$  (the only "mononuclear" tris-chelate of dto for which the X-ray structure is known [58]), potassium and calcium ions influence the actual structure by coordinating the dto oxygens as in bis-chelates (see above).

The mononuclear mixed-ligand complex  $[(PMe_3)_2Pd(dto)]$  contains one sulphur-bound dto and two phosphorus atoms in square-planar coordination geometry around the palladium [46(b)]. A significant lengthening of the Pd–P distances in this compound (2.301 Å) compared with 2.253 Å in the oxalate compound  $[(PEt_3)_2Pd(C_2O_4)]$  suggests that the thio-oxalate ligand is a better *trans*-labilizing group than oxalate.

*Mononuclear five-coordinated oxometal chelates of dto.* As expected in the three mononuclear  $MO(dto)_2$  complexes (M=Mo [39], Tc [55], Re [26] and in  $[TcN(dto)_2]^{2-}$  [55,56], the coordination polyhedrons have roughly  $C_{4v}$  symmetry with the metals considerably above the  $S_4$  base plane (see Fig. 11). However, this is due mainly to the mutual inclination of the two ligands. In the molybdenyl complex






Fig. 11. Structure of the anion in  $(\text{Ph}_4\text{As})_2[\text{TcN}(\text{dto})_2]$  [56].

the “bite” of the dto ligand is 3.260 Å, which is, according to the authors, the largest one ever reported.

**Binuclear homometal dto complexes.** At present, two symmetric binuclear dto chelates are known which differ in constitution (except the central bridging unit dto):  $[(\text{Ph}_3\text{P})_2\text{Ag}(\mu\text{-dto})\text{Ag}(\text{PPh}_3)_2]$  [45] and  $(\text{Ph}_4\text{As})_4[(\text{dto})_2\text{In}(\mu\text{-dto})\text{In}(\text{dto})_2]$  [50]. Thus, according to the X-ray structures, dto links two tetrahedrally coordinated silver(I) ions (Fig. 12) or two octahedrally coordinated indium(III) ions (Fig. 13). In both




Fig. 12. Molecular structure of  $[(\text{Ph}_3\text{P})_2\text{Ag}(\text{dto})\text{Ag}(\text{PPh}_3)_2]$  (phenyl rings omitted) [45].




Fig. 13. Structure of the anion in  $(\text{Ph}_4\text{As})_4[(\text{dto})_2\text{In}(\text{dto})\text{In}(\text{dto})_2]$  [50].

compounds the bridging ligand turns out to be planar, whereas in the indium compound the terminal dtos are twisted with different torsion angles:  $15^\circ$  or  $45^\circ$  (the anion possesses a centre of symmetry, space group  $P\bar{1}$ ).

In a third binuclear ("homometal") complex  $[(\text{en})_2\text{Ni}(\mu\text{-dto})\text{Ni}(\text{dto})]$  the bridging dto links an octahedral  $\text{Ni}_4\text{NiO}_2$  core with a planar  $\text{NiS}_4$  core as shown in Fig. 14 [71].

$\text{Cs}_2[\text{Mo}_2\text{O}_2\text{S}_2(\text{dto})_2]$  [21] is a homobinuclear compound without dto bridges but possessing  $\mu$ -sulphido bridges (see Fig. 15).

As already mentioned, dto indeed links two  $\text{Mn}(\text{CO})_5$  entities in  $[(\text{CO})_5\text{Mn}(\mu\text{-dto})\text{Mn}(\text{CO})_5]$ . The bridging ligand acts as monodentate, however, via sulphur to each manganese atom [25].

**Binuclear heterometal dto complexes.** The only genuine heterobimetallic dto complex is the binuclear  $(\text{BzPh}_3\text{P})_2[(\text{dto})\text{Ni}(\text{S}_2\text{C}_2\text{O}_2)\text{SnCl}_4]$  [28], a typical "Couvouranis complex", with dto linking a planar  $\text{NiS}_4$  unit with an octahedral  $\text{Sn}(\text{IV})$  (see Fig. 16). Another heterometallic dto chelate is strictly speaking trimetallic because the potassium is also coordinated by dto oxygens (see above under "mononuclear"):  $\text{K}\{(\text{dto})_2\text{-Sn}[(\text{O}_2\text{C}_2\text{S}_2)\text{Cu}(\text{PTol}_3)_2]\} \cdot (\text{acetone})_2$  (see Fig. 7) [33]. In the latter case dto links octahedral  $\text{Sn}(\text{IV})$  with tetrahedral  $\text{Cu}(\text{I})$ .

**Oligo- and polynuclear dto complexes.** In  $(\text{BzPh}_3\text{P})_2[\text{Ni}(\text{S}_2\text{C}_2\text{O}_2\text{SnCl}_4)_2]$  the  $\text{Ni}(\text{II})$  ions, square planar coordinated by four sulphur donor atoms, form the centres of




Fig. 14. Molecular structure of  $[(en)_2Ni(dto)Ni(dto)]$  [71].




Fig. 15. Structure of the anion in  $Cs_2[(dto)MoO(\mu-S)_2MoO(dto)]$  [21].




Fig. 16. Structure of the anion in  $(BzPh_3P)_2[(dto)Ni(dtoSnCl_4)]$  [28].

trinuclear complexes and are linked via two dto ligands with two hexa-coordinated tin atoms which themselves are in an  $O_2Cl_4$  environment [28]. Infrared data and bond length changes reflect strong differences between the “parent” complexes and the “ $SnX_4$  adducts” suggesting a predominance of resonance form B (see Fig. 17) in the valence bond (VB) description of these “adducts”.

The Sn–O distances (mean 2.212 Å) are markedly longer than in cases with only one dto- $SnCl_4$  unit in the Ni, Sn binuclear complex mentioned above (2.161 Å). The Ni–S bonds are also slightly lengthened. Coucouvanis attributed this to different  $\pi$ -back-bonding effects.

In the other reported trinuclear dto complex, three Pd atoms are triangularly connected by two central  $\mu_3$ -sulphides [46(b)]. There is, however, only one dto ligand terminally S,S-bound to one of the three palladium atoms, as can be seen from Fig. 18.

Iron(III) and aluminium(III) form tetranuclear heterobimetallic complexes with phosphine shielded Ag(I) in  $\{M[(O_2C_2S_2)Ag(PPh_3)_2]_3\}$  [32].




Fig. 17. Resonance forms of the Ni(dto)Sn cores [28].




Fig. 18. Molecular structure of  $\{(dto)Pd(\mu_3-S)_2[Pd(PMe_3)_2]_2\}$  (methyl groups omitted) [46(b)].

There is a great variety of polynuclear dto complexes which all consist of  $\cdots(\text{O}_2\text{C}_2\text{S}_2)\text{M}(\text{S}_2\text{C}_2\text{O}_2)\cdots$  ( $\text{M}=\text{Ni}$ ,  $\text{Pd}$  or  $\text{Pt}$ ) units interconnected by “hard–hard” building blocks containing rare-earth [63–68],  $\text{Ca}$  [61],  $\text{Ba}$  [62],  $\text{Zn}$  [72],  $\text{Th}$  [69] or  $\text{Mn}$  [53,70,78] ions and water [53,61–70, 72,78], oxalate [65,68], squarate [67], glycine [66] or  $\beta$ -alanine [68] ligands. In the case of  $\text{Mn}$ , complexes with  $\text{Ni}$ ,  $\text{Pd}$ ,  $\text{Pt}$ ,  $\text{Cu}$  or  $\text{Mn(II)}$  could be formed [53,70]. The general structural feature of that group of polynuclear complexes is the presence of infinite networks of chains, ribbons and layers giving rise to various stacks, columns or channels. The complexes can be obtained as crystals by co-precipitation of corresponding mixtures of the starting materials from aqueous solution.

*Solid state conductivities and magnetism.* Some compounds were studied with regard to their solid state conductivity behaviour. Although the conductivity behaviour is anisotropic the values do not reach the typical “organic metal” level as found in the classic TTF–TCNQ [62,69,77]. The solid state conductivity increases in the series  $\text{Ba}[\text{Ni(dto)}_2] \cdot 6\text{H}_2\text{O}$  ( $10^{-12} \Omega^{-1} \text{cm}^{-1}$ ) < non-1-D “red”  $\text{K}_2[\text{Ni(dto)}_2]$  ( $4 \times 10^{-11}$ ) < 1-D “black”  $\text{K}_2[\text{Ni(dto)}_2]$  ( $2 \times 10^{-8} - 5 \times 10^{-9}$ ) <  $\text{Na}_2[\text{Ni(dto)}_2] \cdot 2\text{H}_2\text{O}$  ( $5 \times 10^{-7} - 5 \times 10^{-8}$ ) <  $\text{Li}_2[\text{Ni(dto)}_2] \cdot 2\text{H}_2\text{O}$  ( $10^{-6} - 10^{-7}$ ). For a single crystal of the 1-D “black”  $\text{K}_2[\text{Ni(dto)}_2]$   $\Lambda$  is significantly higher ( $10^{-6}$ ) along the stacking direction than is found for powder conditions. Thus, for a  $\text{Li}_2[\text{Ni(dto)}_2]$  single crystal, values of about  $10^{-4} \Omega^{-1} \text{cm}^{-1}$  could be expected in this direction. The temperature dependence of “black”  $\text{K}_2[\text{Ni(dto)}_2]$  indicates a semiconductor behaviour.

The complexes  $(\text{TTF})_2[\text{M(dto)}_2]$  ( $\text{M}=\text{Pd}$ ,  $\text{Pt}$ ) have been found to be insulators, whereas the metal(III) compounds  $\text{TTF}[\text{M(dto)}_2]$  ( $\text{M}=\text{Ni}$ ,  $\text{Cu}$ ) are semiconductors showing conductivities of  $1.2 \times 10^{-4}$  and  $1.1 \times 10^{-6} \Omega^{-1} \text{cm}^{-1}$ , respectively [77].

The compounds  $[(\text{RE})_2(\text{H}_2\text{O})_{2n}][\text{Ni(dto)}_2]_3 \cdot x\text{H}_2\text{O}$  ( $\text{RE}=\text{La}$ ,  $\text{Ce}$ ,  $\text{Nd}$ ,  $\text{Sm}$ ,  $\text{Eu}$ ,  $\text{Gd}$ ,  $\text{Y}$ ,  $\text{Dy}$ ,  $\text{Er}$ ,  $\text{Yb}$ ) [63] and  $[\text{Th}(\text{H}_2\text{O})_6][\text{Ni(dto)}_2]_2 \cdot 6.5\text{H}_2\text{O}$  [69] exhibit conductivities in the range  $10^{-8} - 2 \times 10^{-7} \Omega^{-1} \text{cm}^{-1}$ ; a value of  $10^{-6} \Omega^{-1} \text{cm}^{-1}$  was measured for the one-dimensional compound  $[\text{NH}_3(\text{CH}_2)_2\text{NH}_3][\text{Ni(dto)}_2]$  [75].

The compound  $\{[\text{Mn}(\text{H}_2\text{O})_3](\text{O}_2\text{C}_2\text{S}_2)\text{Cu}(\text{S}_2\text{C}_2\text{O}_2)\} \cdot 4.5\text{H}_2\text{O}_n$  reveals another remarkable property. It is the first molecular “one-dimensional ferrimagnetic compound” to be reported in the literature, containing alternating  $\text{Cu(II)}$  1/2 spins and  $\text{Mn(II)}$  5/2 spins (see Fig. 19) antiferromagnetically coupled [53,70,78,83] (see also Sect. D(iii)(c)).

### (b) Syntheses of dto complexes

In general, the complexes are formed by direct metathetical reaction between an alkali salt (mainly potassium or caesium) of the ligand, i.e.  $\text{K}_2\text{dto}$  or  $\text{Cs}_2\text{dto}$ , dissolved in a suitable solvent, generally water, and an appropriate metal compound [2]. Sometimes the complexes can be separated from the starting “parent” complex or the aqueous ligand solution by extraction into a halocarbon solvent, e.g.  $\text{CH}_2\text{Cl}_2$  or  $\text{CHCl}_3$ , containing coordinatively unsaturated species (or large “onium” ions, like




Fig. 19. Molecular structure of  $\{[\text{Mn}(\text{H}_2\text{O})_3](\text{dto})\text{M}(\text{dto}) \cdot 4.5\text{H}_2\text{O}\}_n$  ( $\text{M} = \text{Cu}, \text{Ni}, \text{Pd}, \text{Pt}$ ) [53].

$\text{Ph}_4\text{P}^+$  or  $\text{Ph}_4\text{As}^+$ ). Representative examples are  $\{\text{M}[(\text{dto})\text{M}'(\text{PR}_3)_2]_3\}$  [27]. Large counter ions or special combinations of alkali with alkaline-earth or large ions can be used for precipitation of the complexes directly from the aqueous phase, e.g.  $(\text{BzPh}_3\text{P})_2[\text{M}(\text{dto})_2]$  [28],  $\text{KBa}[\text{Co}(\text{dto})_3] \cdot 4\text{H}_2\text{O}$  [84] or  $\text{K}(\text{Me}_3\text{PhN})_2[\text{Cr}(\text{dto})_3] \cdot \text{H}_2\text{O}$  [27]. All known dto compounds are listed in Table 7.

Several special syntheses should be emphasized:

$\text{SnX}_4$  "adducts" of some dto bis-chelates (e.g.  $(\text{BzPh}_3\text{P})_2[\text{Ni}(\text{dtoSnCl}_4)_2]$  [108]) have been prepared by reaction of the parent bis-dithio-oxalates with the tin halides in acetone or  $\text{CH}_2\text{Cl}_2$  [28] under anhydrous conditions.

Refluxing of violet  $(\text{Ph}_4\text{As})_3[\text{Fe}(\text{dto})_3] \cdot 3\text{CH}_3\text{NO}_2$  in  $\text{CH}_2\text{Cl}_2$  over 12 hours resulted in the formation of a brown product analyzed as  $(\text{Ph}_4\text{As})_4[\text{Fe}_2(\text{dto})_5]$  [96] with EPR and IR spectra similar to those of the five-coordinate  $(\text{Ph}_4\text{P})_2[\text{Fe}(\text{dto})_2\text{X}]$  complexes which are prepared by reaction of  $\text{I}_2$ ,  $\text{Br}_2$  or  $\text{FeCl}_3$  with  $[\text{Fe}(\text{dto})_3]^{3-}$  in nitromethane, acetone or  $\text{CH}_2\text{Cl}_2$  [96,98].

$(\text{BzPh}_3\text{P})_2[\text{Fe}(\text{dto})_2\text{NO}]$  has been isolated from a green solution as a crystalline product formed upon heating of violet  $[\text{Fe}(\text{dto})_3]^{3-}$  solution in the presence of either aqueous  $\text{NaNO}_2$  or nitroalkanes ( $\text{CH}_3\text{NO}_2$  or  $\text{C}_2\text{H}_5\text{NO}_2$ ) [96].

Addition of polar solvents (e.g. DMF,  $\text{CH}_3\text{CN}$ ,  $\text{H}_2\text{O}$ ) or nucleophiles (e.g.  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ ,  $\text{CN}^-$  or  $\text{SCN}^-$ ) turns the green colour of the  $\text{CH}_2\text{Cl}_2$  solution of  $[\text{Cu}(\text{dtoSnCl}_4)_2]^{2-}$  to violet, but only one compound could be isolated pure and crystallized, i.e.  $(\text{Ph}_4\text{As})_3\text{P}[\text{Cu}(\text{dtoSnCl}_4)_2\text{Cl}]$  [88]. Oxidation of  $(\text{BzPh}_3\text{P})_2[\text{Cu}(\text{dto})_2]$  in  $\text{CH}_2\text{Cl}_2$  in the dark with  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$  as oxidant, dissolved in acetone, yielded the crystalline Cu(III) complex  $\text{BzPh}_3\text{P}[\text{Cu}(\text{dto})_2] \cdot \text{CH}_2\text{Cl}_2$ , which can be transformed into the mixed ligand Cu(I) compound  $\text{BzPh}_3\text{P}[\text{Cu}(\text{dto})(\text{PPh}_3)_2]$  by  $\text{Ph}_3\text{P}$  in  $\text{CH}_2\text{Cl}_2$  with COS being split off [51]. This complex is formed also via a compound described as " $(\text{Ph}_3\text{P})_2\text{Cu}(\text{S}_2\text{C}_2\text{O}_2\text{H})$ " which was prepared by simultaneous reaction of  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  and  $\text{Ph}_3\text{P}$  with  $\text{K}_2\text{dto}$  in water-ethanol [29].

**TABLE 7**  
**1,2-Dithio-oxalate complexes**

M	No.	Compound	M.p. (°C)	Colour	Magnetism (μ <sub>B</sub> )	IR (cm <sup>-1</sup> )	$\bar{\nu}_{C-C+Cl-S}$	$\bar{\nu}_{M-SO_2}$	Others <sup>a</sup>	Electronic spectra ( $\bar{\nu}_{max} \times 10^{-3} \text{ cm}^{-1} (\epsilon_0)$ )	Ref.
Li		see compound 121									
Na		see compounds 122, 123, 168									
K		see compounds 2, 7-9, 21, 27, 43, 48-50, 56, 69-72, 78, 83, 91-93, 96, 97, 102, 111, 113-115, 118, 124-126, 187, 210									
Cs		see compounds 57, 60, 65, 66									
Ca		see compounds 49, 51, 92-94, 111, 112, 144, 145									
Sr		see compound 72									
Ba		see compounds 50, 69-71, 95, 96, 113-115, 146, 147, 188									
Al	1	(Ph <sub>4</sub> As) <sub>3</sub> [Al(dto) <sub>3</sub> ]	110dec	Light brown	Diam.	1360s	1026s	370m			27
Al	2	K <sub>3</sub> [Al(dto) <sub>3</sub> ] <sup>a</sup>									27
Al	3	[Al]([dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub>	175dec	Brown	Diam.	1402s	1030s	607m	667sh	22.5 (11900)	27
Al	4	[Al]([dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> <sup>b</sup>	195dec	Light brown	Diam.	1395s	1025s	609m	667sh	25.0 (3480)	27
Ga	5	[Ga]([dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub>	192dec	Olive green	Diam.	1390s				28.5 (10300)	41
In	6	(Ph <sub>4</sub> As) <sub>3</sub> [In <sub>3</sub> (dto) <sub>3</sub> ]	165dec	Yellow	Diam.	1610s					41
Sn	7	K <sub>2</sub> [Sn(dto) <sub>3</sub> ]		Yellow		1520s					
Sn	8	K([Sn(dto) <sub>2</sub> ](dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> )]		Orange		1598		Sn-S			29
Sn	9	K([dto) <sub>2</sub> Sn([dto)Cu(PToI <sub>3</sub> ) <sub>2</sub> ]) <sub>3</sub> ·(acetone) <sub>2</sub>				1371s		332, 323			29
Sn	10-12	(X <sub>2</sub> Sn([dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> )] <sub>3</sub> <sup>c,d</sup> X = Cl, Br, Me		Brown		1627					33
Sn						1371					
Sn						1368		Sn-Cl			29
Sn								350			
Sn								Sn-Br			
Sn								284			
Sn	13-15	(X <sub>2</sub> Sn([dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> )] <sub>3</sub> <sup>c,d</sup> X-Cl and/or Me						Sn-Cl			29
Sn								350			

TABLE 7 (continued)

M	No.	Compound	M.p. (°C)	Colour (solid state)	Magnetism ( $\mu_B$ )	IR (cm <sup>-1</sup> )			Electronic spectra ( $\bar{\nu}_{max} \times 10^{-3} \text{ cm}^{-1} (\epsilon_0)$ )			Ref.
						$\bar{\nu}_{C=O}$	$\bar{\nu}_{C-C+Cl-S}$	$\bar{\nu}_{C-S}$	$\bar{\nu}_{M-SO_2}$	Others <sup>y</sup>		
Sn	16-18	BzPh <sub>3</sub> P(X <sub>2</sub> Cl <sub>3</sub> Sn[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ]) <sup>a,d</sup> X = Cl and/or Me				1402 1421 1447						29
Sn	19	{Sn <sub>2</sub> Cl <sub>3</sub> [(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ]} <sub>∞</sub> <sup>e</sup>				1331			Sn-Cl 268			29
Sn	20	{Sn[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ]} <sub>3</sub> <sup>e</sup>										29
Sn	see also compounds 38,39,148-153,197-201,217,218											
Pb	see compound 154											
Cu	21	K <sub>2</sub> [Cu(dto) <sub>2</sub> ]		Black	EPR							85,86, 87
Cu	22	(BzPh <sub>3</sub> P) <sub>2</sub> [Cu(dto) <sub>2</sub> ]	169-70	Brown	1.81(5)	1620s 1585s				24.6	(9130)	51,88
Cu	23	(Ph <sub>4</sub> As) <sub>2</sub> [Cu(dto) <sub>2</sub> ] <sup>e</sup>			1.81							77(a),88
Cu	24	(Ph <sub>4</sub> P) <sub>2</sub> [Cu(dto) <sub>2</sub> ] <sup>e</sup>								26.1	(11000)	51,87
Cu	25	(EtPh <sub>3</sub> P) <sub>2</sub> [Cu(dto) <sub>2</sub> ] <sup>e</sup>								24.7	(8490)	51
Cu	26	(R <sub>4</sub> N) <sub>2</sub> [Cu(dto) <sub>2</sub> ] (R = Et, Bu) <sup>e</sup>								25.0	(11400)	51,77(b)
Cu	27	[K(18-crown-6)] <sub>2</sub> [Cu(dto) <sub>2</sub> ] · DMF <sup>a,e</sup>		Dark green								51,52
Cu	28	[Mn(H <sub>2</sub> O) <sub>3</sub> ][Cu(dto) <sub>2</sub> ] · 4.5H <sub>2</sub> O		Brown								53,78
Cu	29	[CuMn(dto) <sub>2</sub> (H <sub>2</sub> O)] <sup>e</sup>		Black								53
Cu	30	[Cu(dto)(mnt)] <sup>e</sup> · *										86
Cu	31	BzPh <sub>3</sub> P[Cu(dto) <sub>2</sub> ] · CH <sub>2</sub> Cl <sub>2</sub> O <sup>a</sup>		Red	Diam.	1640 1590				16.3	(738)	51
Cu	32	Bu <sub>4</sub> N[Cu(dto) <sub>2</sub> ]		Red						26.2	(22300)	51
Cu	33	[(Ph <sub>3</sub> P) <sub>2</sub> N][Cu(dto) <sub>2</sub> ] <sup>e</sup>		Red						16.3	(614)	51
Cu	34	TTF[Cu(dto) <sub>2</sub> ]		Dark green	EPR <sup>a</sup>					26.3	(20000)	52
										12.4, 19.2 24.1 <sup>1</sup>		77



Cu	35	{[(Ph <sub>3</sub> P) <sub>2</sub> Cu] <sub>2</sub> (dto)}	205–7	Orange	Diam.	1528s			27,88
Cu	36	BzPh <sub>3</sub> P[Cu(dto)(PPh <sub>3</sub> ) <sub>2</sub> ] <sup>+</sup>		Orange					29,51
Cu	37	[(Ph <sub>3</sub> P) <sub>2</sub> Cu(dtoH)]		Orange	Diam.	1536			29
Cu	38	[(Ph <sub>3</sub> P) <sub>2</sub> N] <sub>2</sub> [Cu(dtoSnCl <sub>4</sub> ) <sub>2</sub> ]	169–72	Green	1.83(5)	1475s			88
Cu	39	(Ph <sub>4</sub> As) <sub>3</sub> [Cu(dtoSnCl <sub>4</sub> ) <sub>2</sub> Cl]	161–2	Violet	Diam.				88
Cu	see also compounds 3,8–20,44,52,53,87,99,119,155								
Ag	40	{[(Ph <sub>3</sub> P) <sub>2</sub> Ag] <sub>2</sub> (dto)}	168–70	Yellow brown	Diam.	1545s			27,45, 88
Ag	see also compounds 4,5,54,55,88,100,101,156,157								
Au	41	K[Au(dto) <sub>2</sub> ]					31.0sh 36.5 41.5sh	(16000)	89
Au	42	{[(Ph <sub>3</sub> P)Au] <sub>2</sub> (dto)}	151	Light yellow					45
Zn	43	(Ph <sub>4</sub> As) <sub>2</sub> [Zn(dto) <sub>2</sub> ]	195–198	Light yellow	Diam.	1610s	327m		27,54
Zn	44	{Zn[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> }	150dec	Brown	Diam.	1475s		(7500)	27,59
Zn	45	K <sub>2</sub> [Zn(dto) <sub>2</sub> ] <sup>+</sup>							27
Zn	see also compound 177								
Y, La, Ce, Nd, Sm, Eu, Gd, Dy, Er, Yb	see compounds 158–173								
Th	see compound 174								
Zr	46	[Zr(dto) <sub>4</sub> ] <sup>4+</sup> a,c							27
V	47	(Ph <sub>4</sub> As) <sub>2</sub> [VO(dto) <sub>2</sub> ]	112–114	Dark green	EPR	1630s 1580s	1070s 915m	V–O 1010s	42,90
Cr	48	K(Me <sub>3</sub> PhN)[Cr(dto) <sub>3</sub> ]·H <sub>2</sub> O	180dec	Dark green	3.8x(3)	1570s	325s		27
Cr	49	d,l-K <sub>2</sub> [Cr(dto) <sub>3</sub> ]·6H <sub>2</sub> O		Dark green	3.74				91,92, 93
Cr	50	KBa[Cr(dto) <sub>3</sub> ]·4H <sub>2</sub> O			3.89				91,92
Cr	51	Ca[Co(en) <sub>2</sub> (NO <sub>2</sub> ) <sub>2</sub> ][Cr(dto) <sub>3</sub> ]·2H <sub>2</sub> O <sup>a,t</sup>		Green					92
Cr	52	{Cr[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> } <sup>+</sup>	175dec	Brown	3.8x(3)	1508s	339s	(14200)	27
Cr	53	{Cr[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> } <sup>m</sup>	190dec	Brown	3.79(3)	1372s	370s	(14700)	27

TABLE 7 (continued)

M	No.	Compound	M.p. (°C)	Colour (solid state)	Magnetism ( $\mu_B$ )	IR (cm <sup>-1</sup> )	Electronic spectra ( $\bar{\nu}_{max} \times 10^{-3} \text{ cm}^{-1} (\epsilon_0)$ )				Ref.
							$\bar{\nu}_{C=O}$	$\bar{\nu}_{C-C+O-S}$	$\bar{\nu}_{C-S}$	Others <sup>a</sup>	
Cr	54	$\{Cr[(dioxo)Ag(PPh_3)_2]_3\}^+$	131dec	Light brown	3.86(5)	1622m 1440s 1572s			325s 370s	16.7 (456) 20.8 (1910)	27
Cr	55	$\{Cr[(dioxo)Ag(PPh_3)_2]_3\}^{3+}$	174–177 dec	Light brown	3.95(5)	1370s 1120w			663m 375s	31.3 (39900) 17.2 (365) 22.2 (3500)	27
Cr	56	$K_3[Cr(dto)_3]^+$								32.8 (47800)	27
Mo	57	$Cs_3[Mo_2O_7S_2(dto)_2]$		Olive green		1638vs 1620vs	1070s			Mo–O <sub>i</sub> 952 919	21
Mo	58	$(Ph_4P)_2[Mo_2O_7S_2(dto)_2]$				1631s 1637s			Mo=O 960s Mo–S <sub>b</sub> 452w	330sh 290sh 266sh	94
Mo	59	$(Ph_4P)_2[Mo_2O_7S_2(dto)]$				1630s			Mo=O 942m 956s Mo–S <sub>b</sub> 466	462br 330sh 290sh 265sh	94
Mo	60	$Cs_3[MoO(dto)_2] \cdot H_2O$		Red <sup>a</sup>		1612 1590	1045 920			Mo O <sub>i</sub> 949	39
Mn	61	$\{[Mn(CO)_3(dto)]^+\}$									25
Mn	see also compounds 28,29,175,202,219 <sup>a</sup>										
Tc	62a	$(Ph_4As)_2[TcN(dto)_2]$	215dec	Pale purple		1612	1079			Tc–N 1071m	55
Tc	62b	different modification	142–144	Violet		1620				1045	56

Tc	63	$\text{Bu}_4\text{N}[\text{TcO}(\text{dto})_2]$	126–127	Burgundy Maroon	0.34/0.17 <sup>p</sup>		Tc—O <sub>1</sub> 972	20.0sh 20.8 24.4sh 26.7 33.6	(275) (3400) (14700)	95
Tc	64	$\text{Ph}_4\text{As}[\text{TcO}(\text{dto})_2]$		Red brown			Tc—O <sub>1</sub> 980	19.6sh 20.8 24.4sh 26.7 33.7	(275) (3310) (14600)	55,95
Re	65	$\text{Cs}_4[\text{Re}(\text{CO})_3(\text{dto})_3(\text{dto})]$		Red		1590s 1559vs 1665vs 1640vs	935w, 542m 818s, 512m 548m	1065m		26
Re	66	$\text{Cs}[\text{ReO}(\text{dto})_2]$		Brown red		1665vs 1640vs	1060vs			26
Re	67	$\text{Bu}_4\text{N}[\text{ReO}(\text{dto})_2]$	137–138	Red brown	1.46/0.99 <sup>p</sup>		Re—O <sub>1</sub> 915m Re—O <sub>1</sub> 990	19.2sh 21.1 22.3 23.9 29.4sh 34.3	(340) (330) (480) (18460)	95
Re	68	$\text{Ph}_3\text{MeAs}[\text{ReO}(\text{dto})_2]$					Re—O <sub>1</sub> 991	19.2sh 21.0 23.9 34.0	(300) (430) (18300)	95
Fe	69	$\text{KBa}[\text{Fe}(\text{dto})_3] \cdot 3\text{H}_2\text{O}$	180dec		2.35(5)	1550s 1516s				96
Fe	70	$\text{KBa}[\text{Fe}(\text{dto})_3] \cdot 4\text{H}_2\text{O}$		Dark brown	2.95					92
Fe	71	$\text{KBa}[\text{Fe}(\text{dto})_3] \cdot 6\text{H}_2\text{O}$			2.28			13.6sh 18.3 20.9	(124) (2605) (2455)	91
Fe	72	$\text{KSr}[\text{Fe}(\text{dto})_3] \cdot \text{H}_2\text{O}$			2.36	1553 1520 1562 1520				97

TABLE 7 (continued)

M	No.	Compound	M.p. (°C)	Colour (solid state)	Magnetism ( $\mu_B$ )	IR (cm <sup>-1</sup> )	Electronic spectra ( $\bar{\nu}_{\max} \times 10^{-3}$ cm <sup>-1</sup> (sol))				Ref.
							$\bar{\nu}_{C=O}$	$\bar{\nu}_{C-C+O-S}$	$\bar{\nu}_{C-S}$	Others <sup>a</sup>	
Fe	73	(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> ] <sup>2-</sup>									87,98
Fe	74	(Ph <sub>4</sub> As) <sub>2</sub> [Fe(dto) <sub>2</sub> ]·3CH <sub>3</sub> NO <sub>2</sub>	104-107	Violet	2.30(5)	1622sh 1561s			1046	300	27,96
					2.93	1630					97
						1560					
Fe	75	(PhNH <sub>2</sub> ) <sub>2</sub> [Fe(dto) <sub>2</sub> ]·H <sub>2</sub> O		Black	2.61	1580 1510					97
Fe	76	[Co(en) <sub>2</sub> ][Fe(dto) <sub>2</sub> ]			2.35						92
Fe	77	[Co(en) <sub>2</sub> ][Fe(dto) <sub>2</sub> ]·H <sub>2</sub> O			2.34	1575 1518					97
Fe	78	K <sub>2</sub> [Fe(dto) <sub>2</sub> ] <sup>2-</sup>									27,87
Fe	79	(Ph <sub>4</sub> As) <sub>4</sub> [Fe <sub>2</sub> (dto) <sub>4</sub> ]	127-130	Brown	3.5(2)	1615s, br					96
Fe	80	(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> ]	230dec		4.03	1615s					96,98
Fe	81	(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> X]			EPR						98
	82	X = Cl, Br									
Fe	83	{K <sub>2</sub> [Fe(dto) <sub>2</sub> NO]·H <sub>2</sub> O} <sub>2</sub>		Black						N-O 1665	57
Fe	84	(BrPh <sub>3</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]	158-161	Green	2.22(5)	1585s, br				N-O 1685	96,99, 100
Fe	85	(Et <sub>4</sub> N) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]			2.24	1558				N-O 1688	97
Fe	86	(Bu <sub>4</sub> N) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]			2.30	1590					97
Fe	87	{Fe(dto) <sub>2</sub> Cu(PPPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> }	135-136	Black	5.93(5)	1382s			604m 650m	312s	27,96, 97,101
Fe	88	{Fe(dto) <sub>2</sub> Ag(PPPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> }	145-147	Black	5.81(5)	1380s			603m 653m	314s, br	27,96 97,101
Fe	89	(PhNH <sub>2</sub> ) <sub>2</sub> [Fe(dto) <sub>2</sub> ]·H <sub>2</sub> O <sup>a</sup>		Brown red							84

Ru	90	(Ph <sub>4</sub> As) <sub>2</sub> [Ru(dto) <sub>2</sub> ] <sup>+</sup>	EPR		102
Co	91	K <sub>3</sub> [Co(dto) <sub>3</sub> ]·2H <sub>2</sub> O <sup>c</sup>	Dark brown		27,84
Co	92	KCa[Co(dto) <sub>3</sub> ]·6H <sub>2</sub> O	Dark brown	Diam.	91,103 (1263) (4725)
Co	93	(+) <sub>iso</sub> -KCa[Co(dto) <sub>3</sub> ] <sup>a</sup>	Burgundy		58
Co	94	Ca[Co(en) <sub>2</sub> (NO <sub>2</sub> ) <sub>2</sub> ][Co(dto) <sub>3</sub> ]·H <sub>2</sub> O <sup>a,±</sup>			92
Co	95	KBa[Co(dto) <sub>3</sub> ]·4H <sub>2</sub> O or 2H <sub>2</sub> O <sup>c</sup>	Reddish brown		84
Co	96	d- and l-KBa[Co(dto) <sub>3</sub> ]·5H <sub>2</sub> O	Dark brown		92
Co	97	(+) <sub>iso</sub> -KBa[Co(dto) <sub>3</sub> ]·5H <sub>2</sub> O			93
Co	97	K(Me <sub>3</sub> PhN) <sub>2</sub> [Co(dto) <sub>3</sub> ]	245dec Brown	Diam.	27 (680) (2950)
Co	98	(PhNH <sub>2</sub> ) <sub>3</sub> [Co(dto) <sub>3</sub> ] <sup>c</sup>	Dark brown		84
Co	99	{Co[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	161–164 Brown	Diam.	27 (7500)
Co	100	{Co[(dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	dec		27 (25500)
Co	100	{Co[(dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	130dec Light brown	Diam.	27 (912) (2870)
Co	101	{Co[(dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	159dec Dark brown	Diam.	27
Co	102	A <sub>2</sub> [Co(dto) <sub>2</sub> ] <sup>b</sup> A = K, Ph <sub>4</sub> P		1498s	87
Co	103	[Co(en) <sub>2</sub> (dto)]Cl·2H <sub>2</sub> O		1600	93
Co	104	[Co(en) <sub>2</sub> (dto)]Cl	Brown	1569	36,93 (224) (1340)
Co	105–110	[Co(phen) <sub>2</sub> (dto)]X·nH <sub>2</sub> O X = Cl(n = 2), Br(1), I(1), NO <sub>3</sub> (1), ClO <sub>4</sub> (0), CH <sub>3</sub> COO(3)		1070	104 (16400) (18500)
Co	see also compounds 51,76,77,94,112		> 190dec Orange, red or brown	≈ 1600	≈ 19.0 ≈ 27.8 ≈ 33.3sb
Rh	111	KCa[Rh(dto) <sub>3</sub> ]·4H <sub>2</sub> O	Yellow		92
Rh	112	Ca[Co(en) <sub>2</sub> (NO <sub>2</sub> ) <sub>2</sub> ][Rh(dto) <sub>3</sub> ]·H <sub>2</sub> O <sup>a,±</sup>	Yellow		92
Rh	113	KBa[Rh(dto) <sub>3</sub> ]·4H <sub>2</sub> O <sup>c</sup>	Intense yellow		84
Rh	114	d- and l-KBa[Rh(dto) <sub>3</sub> ]·6H <sub>2</sub> O <sup>c</sup>			92
Rh	115	(+) <sub>iso</sub> -KBa[Rh(dto) <sub>3</sub> ]·6H <sub>2</sub> O <sup>c</sup>			93

TABLE 7 (continued)

M	No.	Compound	M.p. (°C)	Colour - (solid state)	Magnetism ( $\mu_B$ )	IR ( $\text{cm}^{-1}$ )		$\bar{\nu}_{\text{C}-\text{O}}$	$\bar{\nu}_{\text{C}-\text{C}+\text{C}-\text{S}}$	$\bar{\nu}_{\text{C}-\text{S}}$	$\bar{\nu}_{\text{M}-\text{SO}_2}$	Others <sup>y</sup>	Electronic spectra ( $\bar{\nu}_{\text{max}} \times 10^{-3} \text{ cm}^{-1} (\epsilon_{\text{M}})$ )	Ref.
						$\bar{\nu}_{\text{C}=\text{O}}$								
Rh	116	(PhNH <sub>3</sub> ) <sub>2</sub> [Rh(dto) <sub>2</sub> ]·H <sub>2</sub> O <sup>a</sup>												84
Rh	117	(Ph <sub>4</sub> As) <sub>2</sub> [Rh(dto) <sub>2</sub> ]·3CH <sub>3</sub> NO <sub>2</sub>	104dec	Brown	Diam.	1558s			1043m		307m			27
											290m			
Rh	118	K <sub>2</sub> [Rh(dto) <sub>2</sub> ] <sup>a</sup>												27
Rh	119	[Rh[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> ]												27
Ni	120	"H <sub>2</sub> [Ni(dto) <sub>2</sub> ]" <sup>bc</sup>	170dec	Brown	Diam.	1503s			1098s		321m			84
Ni	121	Li <sub>2</sub> [Ni(dto) <sub>2</sub> ]·2H <sub>2</sub> O <sup>a</sup>												59
Ni	122	Na <sub>2</sub> [Ni(dto) <sub>2</sub> ]·2H <sub>2</sub> O <sup>a</sup>												59
Ni	123	Na <sub>2</sub> [Ni(dto) <sub>2</sub> ]·2.5H <sub>2</sub> O <sup>a</sup>		Black										84
Ni	124	K <sub>2</sub> [Ni(dto) <sub>2</sub> ] "red form"		Dark red to black	Diam.	1602 1585			1084		362		≈17.7 ≈19.8	23, 31,60, 84,86, 105,106 31,60
Ni	125	K <sub>2</sub> [Ni(dto) <sub>2</sub> ] "black form"		Black		1505 1490			1144		374			
Ni	126	K <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>a</sup>		(Blue)									≈16.5	60
Ni	127	(NH <sub>4</sub> ) <sub>2</sub> [Ni(dto) <sub>2</sub> ]·4H <sub>2</sub> O <sup>a</sup>		Black										84
Ni	128	(Et <sub>4</sub> N) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>a</sup>												77b
Ni	129	(Bu <sub>4</sub> N) <sub>2</sub> [Ni(dto) <sub>2</sub> ]											17.7sh (1300)	89
													19.9 (3700)	
													33.4 (24000)	
													38.6 (18800)	
													44.4 (16700)	
Ni	130	(PhNH <sub>3</sub> ) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>a</sup>		Reddish brown										84
Ni	131	(NC <sub>4</sub> H <sub>4</sub> NH) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>a</sup>		Black										73
Ni	132	(C <sub>3</sub> H <sub>5</sub> NH) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>a</sup>		Black		1575vs			1090vs		368s			74a
											410m			

No.	Element	Chemical formula	Color	Diam.	$\lambda_{\text{max}}$	$\nu_{\text{max}}$	$\tau_{\text{max}}$	$\phi_{\text{max}}$
Ni 133	Ni	(3-Me-C <sub>3</sub> H <sub>7</sub> NH) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>a</sup>	Black		530w	1600s	1090vs	365s 415m
Ni 134	Ni	(3-Et-C <sub>3</sub> H <sub>7</sub> NH) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>b</sup>	Black		575w	1575vs	1087s	360s 415m
Ni 135	Ni	(4-Me-C <sub>3</sub> H <sub>7</sub> NH) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>c,a</sup>	Black		525m	1590vs	1090vs	365vs 409m
Ni 136	Ni	(4-Et-C <sub>3</sub> H <sub>7</sub> NH) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>b</sup>	Black		535s	1595vs	1090s	360vs 414m
Ni 137	Ni	[H <sub>3</sub> N(CH <sub>2</sub> ) <sub>2</sub> NH <sub>3</sub> ][Ni(dto) <sub>2</sub> ] <sup>c,a</sup>	Brown-red					
Ni 138	Ni	[C <sub>12</sub> N <sub>2</sub> H <sub>12</sub> ][Ni(dto) <sub>2</sub> ] <sup>c,a</sup>	Dark red					
Ni 139	Ni	(Ph <sub>4</sub> P) <sub>2</sub> [Ni(dto) <sub>2</sub> ] <sup>c</sup>						
Ni 140	Ni	(Ph <sub>4</sub> P) <sub>2</sub> [Ni(dto) <sub>2</sub> ] · CH <sub>3</sub> NO <sub>2</sub>						
Ni 141	Ni	(BzPh <sub>3</sub> P) <sub>2</sub> [Ni(dto) <sub>2</sub> ]						
Ni 142	Ni	(Ph <sub>4</sub> As) <sub>2</sub> [Ni(dto) <sub>2</sub> ]						
Ni 143	Ni	TTF[Ni(dto) <sub>2</sub> ] <sup>c,a</sup>	Dark blue					
Ni 144	Ni	[Ca(H <sub>2</sub> O) <sub>6</sub> ][Ni(dto) <sub>2</sub> ] · 2H <sub>2</sub> O*	Red					
Ni 145	Ni	[Cs <sub>9</sub> (H <sub>2</sub> O) <sub>8</sub> (C <sub>2</sub> O <sub>4</sub> )] [Ni(dto) <sub>2</sub> ] · 6H <sub>2</sub> O*	Black					
Ni 146	Ni	Ba[Ni(dto) <sub>2</sub> ] · 4H <sub>2</sub> O*	Black					
Ni 147	Ni	Ba[Ni(dto) <sub>2</sub> ] · 6H <sub>2</sub> O*	Black					
Ni 148	Ni	(BzPh <sub>3</sub> P) <sub>2</sub> [Ni(dto)(dtoSnF <sub>4</sub> )] <sup>d</sup>	Blue	Diam.	649w 615sh	1640s 1625s 1460s	1050m 1025w	378m 365m
Ni 149	Ni	(BzPh <sub>3</sub> P) <sub>2</sub> [Ni(dto)(dtoSnCl <sub>4</sub> )] · H <sub>2</sub> O**	Blue	Diam.	645w 614m	1640s 1613s 1445s	1052m 1025w	385m 363s 328br 221s 184s 160s







TABLE 7 (continued)

M	No.	Compound	M.p. (°C)	Colour (solid state)	Magnetism ( $\mu_B$ )	IR (cm <sup>-1</sup> )				$\bar{\nu}_{M-SO_2}$	Others <sup>c</sup>	Electronic spectra ( $\bar{\nu}_{max} \times 10^{-3} \text{ cm}^{-1} (e_0)$ )	Ref.
						$\bar{\nu}_{C=O}$	$\bar{\nu}_{C-C} \cdot C-S$	$\bar{\nu}_{C-S}$					
Pd	193	(Ph <sub>4</sub> As) <sub>2</sub> [Pd(dto) <sub>2</sub> ]		Yellow								41.3 (17000)	74b
Pd	194	(Ph <sub>4</sub> P) <sub>2</sub> [Pd(dto) <sub>2</sub> ] <sup>+</sup>											87
Pd	195	(Ph <sub>4</sub> P) <sub>2</sub> [Pd(dto) <sub>2</sub> ] <sup>+</sup> ·CH <sub>3</sub> NO <sub>2</sub>	227										107
Pd	196	(TTF) <sub>2</sub> [Pd(dto) <sub>2</sub> ] <sup>+</sup>		Dark brown	EPR <sup>a</sup>							12.4, 19.2, 25.6 <sup>d</sup>	77
Pd	197	(BzPh <sub>3</sub> P) <sub>2</sub> [Pd(dto)(dtoSnCl <sub>4</sub> )] <sup>+</sup>	168–169	Orange	Diam.	1635s 1612s 1448s	1050m	642m 628m	261m 267m	Sr–X 355s 331s 219s 168s 155s		22.5 (5300) 23.5 (5200) 27.0 (3600)	28, 108
Pd	198	(Ph <sub>4</sub> As) <sub>2</sub> [Pd(dto)(dtoSnCl <sub>4</sub> )]	201–202		Diam.	1623s 1580m 1480s 1450s						22.5 (5300) 23.5 (5200) 27.0 (3600)	108
Pd	199	(BzPh <sub>3</sub> P) <sub>2</sub> [Pd(dtoSnCl <sub>4</sub> )] <sup>+</sup>	219–221	Orange	Diam.	1465s	1150s	628m		355s 327s 216s 169s 156s		22.2sh 24.1 (12800) 27.4sh (8200)	28, 108
Pd	200	(BzPh <sub>3</sub> P) <sub>2</sub> [Pd(dtoSnBr <sub>4</sub> )]	209–210	Orange-red	Diam.	1460s	1143m	626w	331s	248s 234s 178s 131s 98s		22.2sh (8600) 23.8 (13200) 26.2sh (9100)	28

Pd	201	(BzPh <sub>3</sub> P) <sub>2</sub> [Pd(dtoSnL) <sub>2</sub> ]	152–153	Red	Diam.	1455s	1144m	62.5w	329s	194s 188s 180s 131s 104s	22.9 27.6	(2700) (5100)	28
Pd	202	[Mn(H <sub>2</sub> O) <sub>3</sub> ][Pd(dto) <sub>2</sub> ] <sub>2</sub> ·4.5H <sub>2</sub> O		Orange	Diam.								53.78
Pd	203	[Pd(dto)(Et <sub>2</sub> PCH <sub>2</sub> CH <sub>2</sub> PEt <sub>2</sub> )] <sup>+</sup>		Yellow		1632							46b
Pd	204	[Pd(dto)(Ph <sub>2</sub> PCH <sub>2</sub> CH <sub>2</sub> PPh <sub>2</sub> )]				1632							46b
Pd	205	[Pd(dto)(PMe <sub>3</sub> ) <sub>2</sub> ]		Yellow		1636							46b
Pd	206	[Pd <sub>3</sub> (μ <sub>3</sub> -S) <sub>2</sub> (dto)(PMe <sub>3</sub> ) <sub>4</sub> ] <sup>+</sup>		Orange									46b
Pd	207–208	[Pd(dto)L] <sub>2</sub> <sup>2+</sup> L = ned, i-nm <sup>4,4a</sup>											112
Pd	209	[Pd(dto)(CN) <sub>2</sub> ] <sub>2</sub> <sup>2+</sup> · <sup>a</sup>											112
Pt	210	K <sub>2</sub> [Pt(dto) <sub>2</sub> ] <sup>c</sup>		Dark red									3.53, 87.105, 106
Pt	211	(Et <sub>4</sub> N) <sub>2</sub> [Pt(dto) <sub>2</sub> ] <sup>c</sup>											77
Pt	212	(Bu <sub>4</sub> N) <sub>2</sub> [Pt(dto) <sub>2</sub> ]											89
											21.6	(2500)	
											23.1sh		
											23.4	(8500)	
											30.0	(1000)	
											41.7	(26000)	
											45.5	(25000)	
Pt	213	(Ph <sub>4</sub> P) <sub>2</sub> [Pt(dto) <sub>2</sub> ] <sup>c</sup>											87
Pt	214	(Ph <sub>4</sub> P) <sub>2</sub> [Pt(dto) <sub>2</sub> ] <sub>2</sub> ·2CH <sub>3</sub> NO <sub>2</sub>	219	Dark brown	EPR <sup>b</sup>						12.4, 19.2,		107
Pt	215	(TTF) <sub>2</sub> [Pt(dto) <sub>2</sub> ] <sup>+</sup>									24.1 <sup>a</sup>		77
Pt	216	(TTF) <sub>2</sub> [Pt(dto) <sub>2</sub> ] <sup>+</sup>		Black	EPR <sup>b</sup>						12.4, 19.2,		77
											24.1 <sup>a</sup>		
Pt	217	[Pt(dto)(dtoSnCl <sub>4</sub> )] <sub>2</sub> <sup>2+</sup> · <sup>a,1</sup>											107
Pt	218	[Pt(dtoSnCl <sub>4</sub> )] <sub>2</sub> <sup>2+</sup> · <sup>a,1</sup>											107
Pt	219	[Mn(H <sub>2</sub> O) <sub>3</sub> ][Pt(dto) <sub>2</sub> ] <sub>2</sub> ·4.5H <sub>2</sub> O		Black									53
Pt	220	[Pt(dto)(Et <sub>2</sub> PCH <sub>2</sub> CH <sub>2</sub> PEt <sub>2</sub> )] <sup>+</sup>											46b
Pt	221	[Pt(dto)(Ph <sub>2</sub> PCH <sub>2</sub> CH <sub>2</sub> PPh <sub>2</sub> )] <sup>+</sup>				1635							46b
Pt	222	[Pt(dto)(PMe <sub>3</sub> ) <sub>2</sub> ] <sup>+</sup>				1638							46b
						1639							46b

TABLE 7 (continued)

<sup>a</sup> Not isolated in the solid state.	<sup>f</sup> Detailed susceptibility and magnetization studies: quasi-one-dimensional ferrimagnetic chain; $M^T$ vs. $T$ presents a minimum at 130 K and a maximum at 7.5 K. Magnetization data at 1.3 and 4.2 K in the field range 0–5 T are consistent with an antiferromagnetic interaction between Cu(II) and Mn(II) through the dto bridge [83].
<sup>b</sup> Reported as also containing two molecules of $\text{CHCl}_3$ .	<sup>g</sup> First by mistake reported as $\text{BzPh}_3\text{P}[\text{Cu}(\text{dto})(\text{COS})_2] \cdot \text{CH}_2\text{Cl}_2$ ; also as the $\text{Ph}_4\text{As}^+$ salt [88].
<sup>c</sup> No further data reported.	<sup>h</sup> Weak narrow signal with a $g$ value of ca. 2.003, typical of TTF radical cations.
<sup>d</sup> Some IR absorptions are given.	<sup>i</sup> The IR spectrum is presented as a picture.
<sup>e</sup> Only X-ray structure.	<sup>j</sup> Diffuse reflectance.
	<sup>k</sup> Diastereo-isomers were isolated.
	<sup>l</sup> Cr–S, M–O (M = Cu, Ag) isomer.
	<sup>m</sup> Cr–O, M–S (M = Cu, Ag) isomer.
	<sup>n</sup> Red needles from water; blue powder from water/ethanol–ether.
	<sup>o</sup> In [70] a compound $[\text{Mn}(\text{H}_2\text{O})_3][\text{Mn}(\text{dto})_2] \cdot 4.5\text{H}_2\text{O}$ (?) is mentioned in a footnote.
	<sup>p</sup> Measured at 7250 or 14 000 G, respectively.
	<sup>q</sup> Published composition seems to be incorrect; most likely it concerns a metal(III) tris-complex.
	<sup>r</sup> Very unstable.
	<sup>s</sup> $\text{NC}_4\text{H}_4\text{NH}^+ = \text{pyrazinium}$ , $\text{C}_5\text{H}_5\text{NH}^+ = \text{pyridinium}$ , $3\text{-Me-C}_5\text{H}_4\text{NH}^+ = 3\text{-methylpyridinium}$ , $3\text{-Et-C}_5\text{H}_4\text{NH}^+ = 3\text{-ethylpyridinium}$ , $4\text{-Me-C}_5\text{H}_4\text{NH}^+ = 4\text{-methylpyridinium}$ , $4\text{-Et-C}_5\text{H}_4\text{NH}^+ = 4\text{-ethylpyridinium}$ , $\text{C}_{12}\text{N}_2\text{H}_{12}^{2+} = 6,7\text{-dihydrodipyrido}[1,2\text{-}a:2'1'\text{-}c]\text{pyrazinium}$ , $\text{TTF}^+$ -tetrafulvalenium.
	<sup>t</sup> The paramagnetic metal(I) species $[\text{Ni}(\text{dto})(\text{dtoSnF}_4)]^{3-}$ , $[\text{M}(\text{dto})(\text{dtoSnCl}_4)]^{3-}$ and $[\text{M}(\text{dtoSnCl}_4)_2]^{3-}$ could be generated electrochemically in solution [28,107].
	<sup>u</sup> In [108] reported without water.
	<sup>v</sup> The first report [108] contains somewhat different spectroscopic data.
	<sup>w</sup> The paramagnetic metal(I) species $[\text{M}(\text{dto})(\text{diphosphine})]^-$ or $[\text{M}(\text{dto})(\text{PR}_3)_2]^-$ could be generated electrochemically in solution [46b,109].
	<sup>x</sup> cdc = <i>N</i> -cyanodithiocarbamate, <i>i</i> -mnt = 1,1-dicyanoethene-2,2-dithiolate, cpd = 1-cyano-1-phenylethene-2,2-dithiolate, ned = 1-nitroethene-2,2-dithiolate, mnt = maleonitrile dithiolate.
	<sup>y</sup> Subscripts: b = bridging ligand; t = terminal ligand.

In acetone,  $K_2[Sn(dto)_3]$  reacts with  $(Ph_3P)_3CuCl$  forming the complex  $K\{Sn(dto)_2[(dto)Cu(PPh_3)_2]\}$ , which can be transformed in  $CH_2Cl_2$  by  $(Ph_3P)_3CuX$  into  $\{XSn[(dto)Cu(PPh_3)_2]_3\}$  ( $X = Cl, Br$ ) [29]. A series  $\{X_2Sn[(dto)Cu(PPh_3)_2]_2\}$  ( $X = Cl, Me$ ) has been prepared starting with either  $Me_nSnCl_{4-n}$ ,  $K_2dto$  and  $(Ph_3P)_3CuCl$  in THF, water and  $CH_2Cl_2$  or  $\{Zn[(dto)Cu(PPh_3)_2]_2\}$  and  $Me_nSnCl_{4-n}$  in acetone. Obviously  $\{MeSn[(dto)Cu(PPh_3)_2]_3\}$  can be prepared via  $\{Me_2Sn[(dto)Cu(PPh_3)_2]_2\}$  using  $[(PPh_3)_2Cu(dto)]^-$  [29].

A useful method for preparing binuclear compounds  $\{(dto)[M(PPh_3)_2]_2\}$  ( $M = Cu, Ag$ ) starts with a pyridine solution of  $(Ph_3P)_3MCl$  in which  $K_2dto$  is suspended and subsequently dissolved by careful addition of water [45].

$Li_2[Ni(dto)_2] \cdot 2H_2O$  and  $Na_2[Ni(dto)_2] \cdot 2H_2O$  are prepared from  $K_2[Ni(dto)_2]$  using  $LiClO_4$  or  $NaClO_4$  in concentrated aqueous solutions [59]. The "black" form of  $K_2[Ni(dto)_2]$  results from the "normal red" form by use of  $K_2Cr_2O_7$  in aqueous solution [60] (see above).

Metal compounds  $MCl_2L_2$  ( $M = Ni, Pd, Pt$ ;  $L_2$  = diphosphine or two phosphine ligands) react in suitable solvents, e.g. water, methanol or ethanol, to form mixed ligand species  $[M(dto)L_2]$ . It should be noted, however, that in the case of  $[Ni(dto)(PMe_3)_2]$  this route is not successful because  $[Ni(dto)_2]^{2-}$  is formed. Therefore the desired compound was synthesized via  $[(CH_3OCH_2CH_2OCH_3)Ni(dto)]$  by ligand exchange using  $PMe_3$ . If the DMF solution of  $[(PMe_3)_2Pd(dto)]$  is subjected to several heating and cooling cycles, thermolysis of the compound occurs and orange needles of  $[Pd_3(\mu_3-S)_2(dto)(PMe_3)_4]$  can be separated [46(b)].

Usually, coordination polymers containing the nickel bis-dithio-oxalate unit  $[Ni(dto)_2]^{2-}$  on the one hand and more or less "hard" coordination entities on the other crystallize when highly concentrated solutions of the water-soluble components are mixed; examples are:  $[Ca(H_2O)_4]^{2+}$  [61],  $[Ba(H_2O)_5]^{2+}$  [62],  $[(RE)_2(H_2O)_{2n}]^{6+}$  with  $n=5$  for Y, La, Ce, Nd, Sm, Eu and Gd and  $n=4$  for Er and Yb [63,64],  $[La(H_2O)_4(C_2O_4)]^+$  [65],  $[La(H_2O)_4(-O_2CCH_2NH_3^+)]^{3+}$  [66],  $[Yb(H_2O)_2(-O_2CCH_2NH_3^+)_2]^{3+}$  [66],  $[Ce_2(H_2O)_{10}(C_4O_4)]^{2+}$  [67],  $\{[Ce(H_2O)_3(-O_2CCH_2CH_2NH_3^+)]_2(C_2O_4)\}^{4+}$  (oxalate is formed by decomposition of dithio-oxalate during the reaction) [68],  $[Yb(C_2O_4)]^+$  (also here oxalate is formed by decomposition of dithio-oxalate during preparation of the heterobimetallic compound in the presence of sodium maleate or maleic acid) [65],  $[Th(H_2O)_6]^{4+}$  [69],  $[Mn(H_2O)_3]^{2+}$  [53,70] (in this case the hard units can be linked also by the soft partner  $[Pd(dto)_2]^{2-}$ ,  $[Pt(dto)_2]^{2-}$  or  $[Cu(dto)_2]^{2-}$  and  $[Zn(H_2O)_2]^{2+}$  [72]).

The compounds  $(TTF)_m[M(dto)_2]$  ( $m = 1, M = Cu, Ni$ ;  $m = 2, M = Pd, Pt$ ;  $m = 3, M = Pt$ ), which are insoluble in common solvents, have been prepared by metathesis or slow interdiffusion of saturated acetonitrile solutions of  $(TTF)_3(BF_4)_2$  and  $A_2[M(dto)_2]$  ( $A = Ph_4As^+$  or  $Et_4N^+$ ) under inert conditions [77].

### (c) Magnetic properties; EPR and Mössbauer data

The dto complexes behave as magnetically normal, i.e. they can be viewed as composed of innocent dto dianions and the metal ion in the corresponding electron

configuration. Paramagnetic compounds are reported with V(IV) (as VO(II)), Cr(III), Tc(V) (as TcO(III) or TcN(II)), Re(V) (as ReO(III)), Fe(II) (as Fe(NO)(II)), Fe(III), Ru(III), Ni(II), Ni(I), Pd(I), Pt(I), Cu(II) and Cu(II)/Mn(II).

All chromium complexes show magnetic moments from 3.74 to 3.95  $\mu_B$  because of their three unpaired electrons, as expected [27,91].

Bu<sub>4</sub>N[TcO(dto)<sub>2</sub>], (Ph<sub>4</sub>As)<sub>2</sub>[TcN(dto)<sub>2</sub>] and Bu<sub>4</sub>N[ReO(dto)<sub>2</sub>] are weakly paramagnetic. The effective magnetic moments are field strength dependent (0.17–1.46  $\mu_B$ ) and have been suggested to be due to temperature-independent paramagnetism, as found for some other d<sup>2</sup> configurations with local C<sub>4v</sub> symmetry [56,95].

For iron(III) the question arises whether the compounds are high-spin, low-spin, medium-spin or spin-crossover systems. The magnetic moments of all Fe(III) tris-dithio-oxalate compounds containing different cations or solvate molecules fit more or less well the value expected for a low-spin complex (2.0–2.5  $\mu_B$ ) [27,91,92,96,97,113] (presumably some workers had difficulties in obtaining pure samples of the complex salts [91]). The large negative Weiss constant for KBa[Fe(dto)<sub>3</sub>]·6H<sub>2</sub>O (compare also (Ph<sub>4</sub>As)<sub>3</sub>[Fe(dto)<sub>3</sub>]·CH<sub>3</sub>NO<sub>2</sub> [96]) first published by Carlin and Canziani [91] was confirmed later by other authors [97]. There is no evidence for spin-crossover behaviour. If, however, [Fe(dto)<sub>3</sub>]<sup>3-</sup> was used as a “ligand” for the [M(PPh<sub>3</sub>)<sub>2</sub>]<sup>+</sup> cations (M = Cu, Ag), the high-spin compounds {Fe[(dto)M(PPh<sub>3</sub>)<sub>2</sub>]<sub>3</sub>} were formed. This is expected by analogy with the iron(III) oxalate complexes (which are high-spin) because the dto ligands flip from Fe–S,S chelation to Fe–O,O chelation (see above).

(BzPh<sub>3</sub>P)<sub>2</sub>[Fe(dto)<sub>2</sub>NO] exhibits a magnetic moment of 2.22  $\mu_B$  [96]. This suggests a structure similar to that known for the iron dithiocarbamate nitrosyl complexes having also one unpaired electron in the [Fe(NO)] core [114]. Analogies exist also between [Fe(R<sub>2</sub>dtc)<sub>2</sub>X] compounds (X = halide ion) and [Fe(dto)<sub>2</sub>X]<sup>2-</sup> ions [96,98]. The five-coordinated dithio-oxalate complex ions are paramagnetic down to 1.4 K, showing an intermediate spin of  $S = 3/2$  as expected.

In contrast, the loosely bound dimer {K<sub>2</sub>[Fe(dto)NO]·H<sub>2</sub>O}<sub>2</sub> exhibits strong intramolecular antiferromagnetic interactions via an unsymmetrical Fe<sub>2</sub>S<sub>2</sub> unit (with two short and two rather long iron–sulphur distances) [57].  $J$  was found to be  $-23.8 \text{ cm}^{-1}$ .

The magnetism of (Ph<sub>4</sub>As)<sub>4</sub>[Fe<sub>2</sub>(dto)<sub>5</sub>] (3.5  $\mu_B$ ) is rather surprising. One expects a strongly reduced amount because of antiferromagnetic interactions which should appear in the proposed bridging structure [96] (found for the indium compound (Ph<sub>4</sub>As)<sub>4</sub>[In<sub>2</sub>(dto)<sub>5</sub>] [50]).

The polymeric mixed-metal copper–manganese complex [CuMn(dto)<sub>2</sub>·(H<sub>2</sub>O)<sub>3</sub>]·4.5H<sub>2</sub>O, a quasi-one-dimensional compound, consists of chains ...Cu–(S<sub>2</sub>C<sub>2</sub>O<sub>2</sub>)Mn(H<sub>2</sub>O)<sub>3</sub>(O<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)..., criss-crossing glide planes and stacking along these planes. Each layer of stacked chains is separated from the next one by intervening water molecules. The magnetic susceptibility of the compound shows a minimum at 130 K and a maximum at 7.5 K. The antiferromagnetic interaction between Cu(II)

and Mn(II) through the dithio-oxalate bridges leads to “one-dimensional ferrimagnetism” ( $J = -30.3 \text{ cm}^{-1}$ ) [53,70,78,83].

The EPR spin Hamiltonian parameters of dithio-oxalate complexes are summarized in Table 8. The only single-crystal EPR studies on dto systems were carried out for  $[\text{Cu}(\text{dto})_2]^{2-}$  in different host lattices and for undiluted  $\{\text{Fe}[(\text{dto})\text{-Cu}(\text{PPh}_3)_2]_3\}$ . White and Belford studied the system  $\text{K}_2[\text{Cu}/\text{Ni}(\text{dto})_2]$  in the Q-band and analyzed the secondary ( $\Delta m_1 = 1$ ) transitions for the quadrupole coupling parameter which is  $\text{QD} = 0.7 \times 10^{-4} \text{ cm}^{-1}$  [85]. The authors assign this small value (also found for other planar  $\text{Cu-S}_4$  systems with the ligands isomaleonitriledithiolate and dithiocarbamate) to an effectively spherical symmetry caused by strong covalent nature of the  $\text{Cu-S}(\sigma)$  bond. In [117] the spin Hamiltonian parameters of  $\text{K}_2[\text{Cu}/\text{Ni}(\text{dto})_2]$  are compared with those of some other planar and tetrahedral  $\text{CuS}_4$  species. As expected, the behaviour of  $[\text{Cu}(\text{dto})_2]^{2-}$  is markedly different in a  $(\text{Ph}_4\text{As})_2[\text{Zn}(\text{dto})_2]$  single crystal with a tetrahedral  $\text{ZnS}_4$  coordination sphere [54]. The copper guest, however, does not accept the structure of the host lattice completely [115]. The dihedral angle is found to be only about  $10^\circ$  ( $(\text{Ph}_4\text{As})_2[\text{Zn}(\text{dto})_2]$ :  $86.8^\circ$  [54]).

The EPR powder spectra of tetrathiafulvalenium salts  $(\text{TTF})_{1.0}[\text{Cu}(\text{dto})_2]$ ,  $(\text{TTF})_2[\text{M}(\text{dto})_2]$  ( $\text{M} = \text{Pd}, \text{Pt}$ ) and  $(\text{TTF})_3[\text{Pt}(\text{dto})_2]$  at room temperature show a weak, narrow signal with  $g$  values of about 2.003, typical for  $\text{TTF}^+$  radical cations [77].

For  $(\text{Ph}_4\text{As})_3[\text{Fe}(\text{dto})_3] \cdot 3\text{CH}_3\text{NO}_2$  no EPR spectra could be obtained, either at 300 or 78 K. However, at 1 K a strong signal has been observed for the  $^2\text{T}_2$  ground state [96]. Later on the EPR spectrum could be obtained for  $\text{KBa}[\text{Fe}(\text{dto})_3] \cdot 3\text{H}_2\text{O}$  also at higher temperature [113]. The situation is similar for  $[\text{Ru}(\text{dto})_3]^{3-}$  [102]. Calculated resonance line positions and intensities are compared with the experimental EPR spectra of both isomorphous high-spin ( $S = 5/2$ ) compounds  $\{\text{Fe}[(\text{dto})\text{M}(\text{PPh}_3)_2]_3\}$  ( $\text{M} = \text{Cu}$  as single crystal,  $\text{M} = \text{Ag}$  as powder) [101].

Five-coordinate iron(III) dithio-oxalato halides  $(\text{Ph}_4\text{P})_2[\text{Fe}(\text{dto})_2\text{X}]$  show a behaviour expected for  $S = 3/2$  with an axial character of the electronic spin Hamiltonian for  $\text{X} = \text{I}$  and increasing rhombicity for  $\text{X} = \text{Br}$  and  $\text{Cl}$  [98].

Compounds containing the anion  $[\text{Fe}(\text{dto})_2\text{NO}]^{2-}$  have been investigated in detail by EPR in liquid and rigid solution and as powder. The magnetic symmetry is axial. The spin Hamiltonian parameters are temperature- and solvent-dependent, indicating a vacant sixth site solvent coordination [99,100]. Addition of  $\text{SnCl}_4$  under anhydrous conditions gives a ligand perturbation which shows that the unpaired electron is in an  $a_1$  or  $a_2$  type orbital (in  $\text{C}_{2v}$  symmetry,  $d_{z^2}$  or  $d_{x^2-y^2}$ ).

Another five-coordinate complex containing one unpaired electron is  $(\text{Ph}_4\text{As})_2[\text{VO}(\text{dto})_2]$  [90]. Its EPR parameters differ remarkably from those of other five-membered ring sulphur ligand (dithiolene) chelates. The differences could be caused by the nonrigidity of the dto ligands.

EPR is well suited to detect mixed-ligand species formed in solution [118] if

TABLE 8

EPR investigations of dithio-oxalate complexes (in the Q band)

Compound (system)	Solvent/state	Temperature (K)	Spin Hamiltonian parameters		Ref.
			$g$	$A^a$	
$K_2[Cu/Ni(dto)_2]^b$	Single-crystal (and powder)	Ambient	$g_z$	$A_z^{Cu}$	85
			$g_x$	$A_x$	
			$g_y$	$A_y$	
$(Ph_4As)_2[^{63}Cu/Zn(dto)_2]$	Single-crystal	Ambient	$g_1$	$A_1^{Cu}$	115
			$g_2$	$A_2$	
			$g_3$	$A_3$	
$(Bu_4N)_2[Cu(dto)_2]$			$g_{iso}$	$A_{iso}^{Cu}$	
$[Cu(dto)(n-bu_2btu)]^{c-e}$			$g_0$	$a_0^{Cu}$	116
$[Cu(dto)(t-bu_2bsu)]^{c-e}$			$g_0$	$a_0^{Cu}$	116
$(Ph_4As)_2[VO(dto)_2]$	Acetone		$g_0$	$a_0^{Cu}$	116
		295	$g_0$	$a_0^V$	90
		133	$g_z$	$A_z$	
			$g_y$	$A_y$	
			$g_x$	$A_x$	
$[Ni(Ph_2PCH_2CH_2PPh_2)(dto)]^{d-f}$	$CH_2Cl_2$	233	$g_0$	$a_0^P$	109
		113	$g_1$	$A_1$	
			$g_2$	$A_2$	
			$g_3$	$A_3$	
$[Ni(dto)(dtoSnCl_4)]^{3-d}$	$CH_2Cl_2$	293	$g_0$	$a_0^{Sn}$	107
$[Ni(dtoSnCl_4)_2]^{3-d}$	$CH_2Cl_2$	293	$g_0$	$a_0^{Sn}$	107
$[Pd(dto)(dtoSnCl_4)]^{3-d}$	$CH_2Cl_2$	293	$g_0$	$a_0^{Sn}$	107
		113	$g_1$	$A_1^{Sn}$	
			$g_2$	$A_2$	
			$g_3$	$A_3$	
$[Pd(dtoSnCl_4)_2]^{3-d}$	$CH_2Cl_2$	293	$g_0$	$a_0^{Sn}$	107
$[Pt(dto)(dtoSnCl_4)]^{3-d}$	$CH_2Cl_2$	293	$g_0$	$a_0^e$	107
		113	$g_1$	$A_1$	
			$g_2$	$A_2$	
			$g_3$	$A_3$	
$[Pt(dtoSnCl_4)_2]^{3-d}$	$CH_2Cl_2$	293	$g_0$	$a_0^{Sn}$	107
$(Ph_4As)_3[Fe(dto)_3] \cdot 3CH_3NO_2$	Powder	1	$g_0$	$a_0^e$	107
$KBa[Fe(dto)_3] \cdot H_2O$	Powder		one line (no data)		96
			$g_{  }$		113



	Single-crystal	Line shape analysis		
		Line shape analysis	g	
{Fe[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	Powder	2.00(2)	g	101
{Fe[(dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	Powder	2.06(1)	g <sup>f</sup>	101
{Fe[(dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	Powder	2.04(1)	g <sup>f</sup>	96
(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> Cl]	Powder	2.04(1)	g <sup>f</sup>	98
(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> Br]	Powder	2.032(5)	g <sup>f</sup>	98
(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> I]	Powder	2.0329–	g <sup>f</sup>	98
(BzPh <sub>3</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]	Frozen solution	2.0336	g	96
(BzPh <sub>3</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]	g	2.0333(10)	g <sub>0</sub>	100
[Fe(dto) <sub>2</sub> NO] <sup>2-</sup>	Methanol–glycerol (1:1)	2.0162(5)	g <sub>0</sub>	99
		2.0345(15)	g <sub>  </sub>	12.6(1)
		2.0284(10)	g <sub>⊥</sub>	15.6(1)
			g <sub>iso</sub>	12.7(1)
				13.7(1)
ccr121874L/14-10-92 13:11:03				
	CH <sub>2</sub> Cl <sub>2</sub>	2.0337(10)	g <sub>0</sub>	12.8(1)
		2.0322(10)	g <sub>0</sub> <sup>N</sup>	13.2(1)
		2.061	g <sup>h</sup>	57
{K <sub>2</sub> [Fe(dto) <sub>2</sub> NO]·H <sub>2</sub> O} <sub>2</sub>	Powder	2.0359(10)	g <sub>0</sub>	99
(BzPh <sub>3</sub> P) <sub>2</sub> [Fe(dto)SnCl <sub>4</sub> NO]	CH <sub>2</sub> Cl <sub>2</sub>	2.0150(5)	g <sub>  </sub>	11.6(1)
		2.0383(15)	g <sub>⊥</sub>	14.0(1)
		2.0305(10)	g <sub>iso</sub>	10.9(1)
		2.037(3)	g <sub>  </sub>	11.9(1)
[TcN(dto) <sub>2</sub> ] <sup>-1</sup>	CHCl <sub>3</sub>	2.04	g <sub>1</sub>	206.0(1.5)
(Ph <sub>4</sub> As) <sub>3</sub> [Ru(dto) <sub>3</sub> ]	CHCl <sub>3</sub> –toluene	2.04	g <sub>2</sub>	56
		1.98	g <sub>3</sub>	102

<sup>a</sup> A values in 10<sup>-4</sup> cm<sup>-1</sup>.<sup>b</sup> Quadrupole coupling constant QD = 0.7 × 10<sup>-4</sup> cm<sup>-1</sup>.<sup>c</sup> *n*-bu, *btu* = di-*n*-butylbenzoylthiourea; *i*-bu, *bsu* = di-isobutylbenzoylselenourea.<sup>d</sup> Electrochemically generated.<sup>e</sup> Probably <sup>195</sup>Pt coupling.<sup>f</sup> Determined by susceptibility data.<sup>g</sup> 16 different solvents: methanol, HCONHMe, ethanol, *n*-propanol, *n*-butanol, tetrahydrothiophen-1,1-dioxide, MeNO<sub>2</sub>, Me<sub>2</sub>SO, propylene carbonate, MeCN, DMF, PhNO<sub>2</sub>, acetone, pyridine, CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub>.<sup>h</sup> Broad line.<sup>i</sup> Generated by bromine oxidation; paramagnetic mixed-ligand (dto/bromide) compounds showing S<sub>3</sub>Br, S<sub>2</sub>Br<sub>2</sub> and SBr<sub>3</sub> coordination spheres are also detected.

the spin Hamiltonian parameters are changed strongly on moving from the starting complexes to mixed-ligand systems.  $[\text{Cu}(\text{dto})_2]^{2-}$ , for example, reacts with copper(II) benzoylthioureas (e.g. 1,1-di-*n*-butyl-3-benzoylthiourea (*n*-bu)<sub>2</sub>btu) or benzoylsele-noureas (e.g. (*i*-bu)<sub>2</sub>bsu) giving mixed-ligand chelates containing a five-membered (dto) and a six-membered ring [116].

Oxidation of  $[\text{TcN}(\text{dto})_2]^{2-}$  with  $\text{Cl}_2$  or  $\text{Br}_2$  yields paramagnetic (Tc(VI)) mixed ligand compounds which contain dto as well as the corresponding halide, detected by EPR [56].

A reversible one-step reduction (−1.19 V) of  $[(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)\text{Ni}(\text{dto})]$  in  $\text{CH}_2\text{Cl}_2$  gives the corresponding paramagnetic anion with a  $d^9$  configuration, giving rise to an EPR spectrum with the expected  $^{31}\text{P}$  superhyperfine structure [109].

Unlike dithio-oxalato complexes, the  $\text{SnCl}_4$  “adducts” also undergo well-defined reversible electrochemical reduction to  $d^9$  species [107]. EPR spectra could be taken from the products obtained by in situ electrolysis. They consist of intense central lines with  $g$  values close to the free electron value and satellite lines due to hyperfine coupling with magnetic isotopes of tin. The closeness of the  $g$  values to the free electron value and the small degree of anisotropy in the  $g$  values for the frozen solutions are indicative of an essentially ligand-based orbital for the unpaired electron. The reason for the stabilization effect of  $\text{SnCl}_4$  on the one-electron reduction products is a lowering of the energy of a  $\pi$ -antibonding orbital on the dto ligand. In  $[(\text{dto})\text{M}(\text{dtoSnCl}_4)]^{3-}$  the unpaired electron could be expected to reside mainly on the dto which is coordinated to the  $\text{SnCl}_4$  unit. In  $[\text{M}(\text{dtoSnCl}_4)_2]^{3-}$  (the  $\pi^*$  orbitals of both dto ligands are lowered! See Fig. 20 [28,108]), on the other hand, the electron should be delocalized over both dto ligands. Thus the spin density near the Sn nuclei is expected to be higher in the 1:1 than in the 1:2 complexes. This fact is reflected by a decrease in the Sn coupling constants on going from  $[\text{M}(\text{dto})(\text{dtoSnCl}_4)]^{3-}$  to  $[\text{M}(\text{dtoSnCl}_4)_2]^{3-}$  (see Table 8).

As can be seen from Table 9, the shift values in the Mössbauer spectra of all Fe(III) dithio-oxalates lie in the expected range found also for 1,2-dithiolenes or dithiocarbamates, indicating that the extent of delocalization is similar in these systems [97]. Lower shift values are normally associated with low-spin configurations.  $\{\text{Fe}[(\text{dto})\text{M}(\text{PPh}_3)_2]_3\}$  complexes (*O,O*-chelation), however, formed by ligand flips from  $[\text{Fe}(\text{dto})_3]^{3-}$  (*S,S*-chelation) have the lowest values for Fe(III) dithio-oxalates despite their high-spin character. On the other hand, the expected lower isomer shift values for  $[\text{Fe}(\text{dto})_2\text{NO}]^{2-}$  could be verified, reflecting the powerful  $\pi$ -bonding character of the  $\text{NO}^+$  group. The relatively large quadrupole splitting ( $\approx 1 \text{ mm s}^{-1}$ ) observed for  $\{\text{Fe}[(\text{dto})\text{M}(\text{PPh}_3)_2]_3\}$  (unexpected for spherically symmetric  $^6\text{A}_1$  systems) confirms a considerable distortion from  $O_h$  symmetry [32].

Because the Fe(III) tris-dithio-oxalates exhibit no spin-crossover behaviour different quadrupole splittings (varying from 0.3 to  $1.7 \text{ mm s}^{-1}$ ) should be the result of different degrees of distortion from octahedral symmetry in all cases.

The isomer shifts found for the complexes  $(\text{Ph}_4\text{P})_2[\text{Fe}(\text{dto})_2\text{X}]$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ )




Fig. 20. Energy level diagram for  $[M(dto)_2]^{2-}$  and  $[M(dto)_2(SnCl_4)_x]^{2-}$  ( $x=1,2$ ) [108].

lie between 0.25 and 0.30  $\text{mm s}^{-1}$  (relative to metallic iron). They agree very well with those published for analogous five-membered chelate ring Fe(III) dithiolenes having  $S=3/2$  (four-membered chelate ring compounds, e.g.  $\text{Fe}(\text{R}_2\text{dtc})_2\text{X}$ , where  $\text{R}_2\text{dtc}$  signifies dithiocarbamate, show isomer shifts in the region of 0.5  $\text{mm s}^{-1}$ ), indicating substantial d electron delocalization in the ligand systems. The observed quadrupole splittings between 3.25 and 3.6  $\text{mm s}^{-1}$  are the highest reported for intermediate spin complexes [98].

A mixture of ferric nitrate and potassium dithio-oxalate in aqueous solution at 77 K gives more than one iron species, the relative intensities of which vary with the  $\text{Fe}^{3+}/\text{dto}^{2-}$  ratio. The Mössbauer signals have been attributed to low-spin Fe(III) and also high-spin Fe(II) [97].

#### (d) Electronic spectra

Qualitative spectroscopic data of  $[M(dto)_2]^{2-}$  chelates were first published by Cox et al. in 1935 [3]. The next paper did not follow until 1951. Its authors studied

TABLE 9

Mössbauer data of dithio-oxalate complexes

Compound	Temperature (K)	$\delta$ (mm s <sup>-1</sup> ) (vs. metallic iron)	$\Delta E_Q$ (mm s <sup>-1</sup> )	Ref.
KBa[Fe(dto) <sub>3</sub> ]·6H <sub>2</sub> O	77	0.37	0.42	97
	196	0.33	0.35	
	298	0.27	0.31	
KSr[Fe(dto) <sub>3</sub> ]·H <sub>2</sub> O	4	0.37	1.06	97
	77	0.36	1.09	
	298	0.26	0.68	
[Co(en) <sub>3</sub> ][Fe(dto) <sub>3</sub> ]·H <sub>2</sub> O	77	0.40	1.68	97
	298	0.32	1.36	
	330	0.28	1.05	
(PhNH <sub>3</sub> ) <sub>3</sub> [Fe(dto) <sub>3</sub> ]·H <sub>2</sub> O	4	0.40	1.27	97
	77	0.38	1.12	
	123	0.40	1.01	
	173	0.38	0.93	
	223	0.36	0.81	
(PhNH <sub>3</sub> ) <sub>3</sub> [Fe(dto) <sub>3</sub> ]·H <sub>2</sub> O <sup>a</sup>	298	0.29	0.64	97
	77	0.38	1.14	
	298	0.29	0.64	
(Ph <sub>4</sub> As) <sub>3</sub> [Fe(dto) <sub>3</sub> ]·3CH <sub>3</sub> NO <sub>2</sub>	77	0.53	0.90	97
	298	0.43	0.95	
{Fe[(dto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	77	0.42	0.99	97
	298	0.24	0.87	
{Fe[(dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>3</sub> }	77	0.33	1.18	97
	298	0.24	1.17	
(Et <sub>4</sub> N) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]	77	0.29	1.04	97
(Bu <sub>4</sub> N) <sub>2</sub> [Fe(dto) <sub>2</sub> NO]	4	0.29	1.05	97
	196	0.26	1.04	
	298	0.36	0.67	
(Bu <sub>4</sub> N) <sub>2</sub> [Fe(dto) <sub>2</sub> NO] <sup>a</sup>	77	0.28	1.03	97
	298	0.35	0.68	
(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> Cl]	4.2–77	0.25(5)	3.60(1)	98
(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> Br]	4.2–77	0.29(2)	3.25(2)	98
(Ph <sub>4</sub> P) <sub>2</sub> [Fe(dto) <sub>2</sub> I]	4.2–77	0.30(2)	3.33(2)	98

<sup>a</sup> Prepared from ferrous salt.

the absorption spectra of K<sub>2</sub>[M(dto)<sub>2</sub>] (M=Ni, Pd, Pt) in solution and in the crystalline state [105].

Deskin compared the nickel complexes of dithio-oxalate, dithiomalonate and trithiocarbonate by UV–VIS spectroscopy and evaluated the formation constants log *K<sub>f</sub>* in aqueous solution (total ionic strength 0.1), which gave a series: dto > dithiomalonate > trithiocarbonatē. The log *K<sub>f</sub>* of [Ni(dto)<sub>2</sub>]<sup>2-</sup> (9.43) is also

found to be markedly higher than that of  $[\text{Ni}(\text{ox})_2]^{2-}$  (6.51) [119], indicating that oxalate is a rather weaker ligand for Ni(II) than is dithio-oxalate. Deskin's value of the formation constant of  $[\text{Ni}(\text{dto})_2]^{2-}$  has been doubted by Pearson and Sweigart, who found a much higher value ( $16.1 \pm 0.2$ ) determined using three different methods in aqueous solution [86]. The formation of mixed ligand complexes containing the dto ligand has been followed spectrophotometrically, as well as the kinetics and mechanisms of the formation of  $[\text{Ni}(\text{dto})_2]^{2-}$ . Ligand exchange reactions were also studied spectrophotometrically involving dto as nucleophile, as leaving group and as *trans* group in Ni(II), Pd(II) and Cu(II) complexes [17,86,110,112].

In 1964, Latham et al. came up with a detailed interpretation of the electronic spectra of the planar  $(\text{Bu}_4\text{N})_n[\text{M}(\text{dto})_2]$  ( $n = 2$ : M = Ni, Pd, Pt;  $n = 1$ : M = Au) on the basis of a molecular orbital (MO) theory treatment [89]. Figure 20 shows the relative energies of the most important levels of  $[\text{Ni}(\text{dto})_2]^{2-}$  with an interchange of both  $4a_g(d_{x^2-y^2})$  and  $3b_{3g}(d_{yz})$  metal orbitals proposed by Coucouvanis and co-workers [28]. The bands in the absorption spectrum (taken in acetonitrile) were assigned as follows:  $44\,400\text{ cm}^{-1}$  ( $\epsilon = 16\,700$ )  $\text{L}(\sigma) \rightarrow \text{M}$ ;  $38\,600\text{ cm}^{-1}$  (18 800)  $\text{L}(\pi) \rightarrow \text{L}(\pi^*)$ ;  $33\,400\text{ cm}^{-1}$  (24 000)  $\text{L}(\pi) \rightarrow \text{M}$ ;  $19\,900\text{ cm}^{-1}$  (3700)  $\text{M} \rightarrow \text{L}(\pi^*)$  and a shoulder at  $17\,700\text{ cm}^{-1}$  (1300)  $\text{M} \rightarrow \text{L}(\pi^*)$  and/or  $d_{x^2-y^2} \rightarrow d_{xy}$  (see also assignments made in [74] for pyridinium salts of  $[\text{Ni}(\text{dto})_2]^{2-}$ ). The spectra of  $[\text{Pd}(\text{dto})_2]^{2-}$ ,  $[\text{Pt}(\text{dto})_2]^{2-}$  and  $[\text{Au}(\text{dto})_2]^-$  are similar. The assignments were used to place dto in the spectrochemical series and to compare its position with those of other bidentate (geminal or vicinal) sulphur-donor ligands. The position of dto is consistently highest in the Ni(II), Pd(II) and Pt(II) complexes containing dithio ligands, and maleonitriledithiolate (mnt), another well-investigated 1,2-dithiolate ligand, is consistently lowest.

Carlin and Canziani estimated the octahedral splitting parameter  $D_q$  from the transition  ${}^4A_{2g} \rightarrow {}^4T_{2g}$  in  $[\text{Cr}(\text{dto})_3]^{3-}$  [91]. They found a value of  $1700\text{ cm}^{-1}$  (oxalate  $1754\text{ cm}^{-1}$ ) and a Racah parameter  $B = 406\text{ cm}^{-1}$  (oxalate  $522\text{ cm}^{-1}$ ), supporting the theory that dithio-oxalate is actually a weaker field ligand than is oxalate. The sulphur ligand atoms, on the other hand, cause a reduction in the  $B$  parameter, i.e. in the electron–electron repulsion (compared with the free  $\text{Cr}^{3+}$ ) because of higher covalency. The absorption spectra, including circular dichroism studies, of  $[\text{Co}(\text{dto})_3]^{3-}$ ,  $[\text{Cr}(\text{dto})_3]^{3-}$ ,  $[\text{Co}(\text{en})_2(\text{dto})]^+$ ,  $[\text{Co}(\text{en})_3]^{3+}$  and  $\text{dto}^{2-}$ , are compared by Hidaka and Douglas to figure out the origin of the individual bands [93]. The absorption bands of  $[\text{Co}(\text{en})_3]^{3+}$  are shifted towards lower frequencies and have increased intensities when en is replaced by dto to give  $[\text{Co}(\text{en})_2(\text{dto})]^+$  and  $[\text{Co}(\text{dto})_3]^{3-}$ . The bands assigned to  $d \rightarrow d$  transitions gain intensity through Co–S bonding. Kanamori et al. could show by a resonance Raman study [103] that in  $[\text{Co}(\text{dto})_3]^{3-}$  a charge-transfer (CT) transition situated at about  $20\,000\text{ cm}^{-1}$  overlaps with two  $d \rightarrow d$  transitions producing two shoulders at  $17\,000$  and  $21\,800\text{ cm}^{-1}$  [93].

The position of dto in the spectrochemical series is mentioned also by Jørgensen [120] and by Lockyer and Martin [121].

The reaction of  $[\text{M}(\text{dto})_2]^{2-}$  with  $\text{R}_x\text{SnCl}_{4-x}$  [28,108] introduces a bathochromic shift of the  $\text{M} \rightarrow \text{L}$  CT bands (assigned by Gray and co-workers), as can be seen from Fig. 20. Possible interchange of the  $4a_g$  ( $d_{x^2-y^2}$ ) and  $3b_{3g}$  ( $d_{yz}$ ) orbitals [28,30,107,108] has no fundamental influence on this statement.

The question whether the reaction of  $[\text{M}(\text{dto})_3]^{3-}$  ( $\text{M} = \text{Cr}, \text{Co}$ ) with  $[\text{M}'(\text{PPh}_3)_2]^+$  ( $\text{M}' = \text{Cu}, \text{Ag}$ ) (yielding  $\{\text{M}[(\text{dto})\text{M}'(\text{PPh}_3)_2]_3\}$ ) causes an increase of the ligand field strength cannot be answered directly. However, the  $10\text{Dq}$  transitions ( $17\,000\text{ cm}^{-1}$  for  $\text{Co(III)}$  and  $17\,700\text{ cm}^{-1}$  for  $\text{Cr(III)}$ ) in the spectra of the “parent” complexes are not observed in the spectra of the “adducts”. This is presumably because of a hypsochromic shift as the result of increasing ligand field strength ( $\text{S}, \text{S}$ -chelation  $\rightarrow$   $\text{O}, \text{O}$ -chelation) which moves the  $d \rightarrow d$  bands below the intense charge-transfer absorption [27].

As shown first by Dwyer and Sargeson,  $[\text{M}(\text{dto})_3]^{3-}$  ( $\text{M} = \text{Cr}, \text{Co}$  and  $\text{Rh}$ , but not  $\text{Fe}$ ) are stable and sufficiently inert to be separated enantioselectively via  $\text{Ca}[\text{Co}(\text{en})_2(\text{NO}_2)_2][\text{M}(\text{dto})_3]$  [92]. Using also  $(-)\text{-cis}[\text{Co}(\text{en})_2(\text{NO}_2)_2]\text{Cl}$ , Hidaka and Douglas [93] prepared  $(+)\text{-KBa}[\text{M}(\text{dto})_3]$  for circular dichroism studies and found an absolute configuration which was checked and verified later by X-ray structure analysis on  $(+)\text{-KCa}[\text{Co}(\text{dto})_3] \cdot 4\text{H}_2\text{O}$  by Butler and Snow [58] as the  $\Lambda$ -configuration (according to the IUPAC Information Bulletin 1968). This is the same configuration as found for the oxalate complex  $(-)[\text{Co}(\text{ox})_3]^{3-}$ . Both the oxalate and the dithio-oxalate compound have a dominant positive circular dichroism in the region of the  $^1A_{1g} \rightarrow ^1T_{1g}$  transition. The correlation between two observed circular dichroism bands and the stereochemical chirality was investigated by McCaffery et al., including  $(-)[\text{Co}(\text{dto})_3]^{3-}$  and  $(+)\text{-}[\text{Rh}(\text{dto})_3]^{3-}$ . For the latter two complexes a right-handed helical stereochemistry was found [122].

#### (e) IR, Raman and resonance Raman spectra

The classical IR treatment of dto complexes ( $\text{Ni}, \text{Pd}, \text{Pt}, \text{Co}$ ) was made by Fujita and Nakamoto in 1964 [106] and accepted by most subsequent authors. They interpreted the IR spectra of  $\text{K}_2[\text{Pt}(\text{dto})_2]$  on the basis of a normal coordinate analysis. Because the spectra are little sensitive to the nature of the metal, the IR spectra of the other dto complexes are very similar. A band near  $1600\text{ cm}^{-1}$  has been assigned to the C–O stretching vibrations, a band at about  $1080\text{ cm}^{-1}$  to a combination of the C–C and C–S stretching vibrations, a band around  $940\text{ cm}^{-1}$  to a combination of C–O and C–S stretching vibrations, and a band at  $570\text{ cm}^{-1}$  to the C–S stretching vibration. Absorptions at  $436, 422$  and  $322\text{ cm}^{-1}$  were attributed to Pt–S vibrations.

Later on Czernuszewicz et al. measured the IR and Raman spectra of both the “red” and the “black” form of  $\text{K}_2[\text{Ni}(\text{dto})_2]$  and made complete vibrational assignments on the basis of  $^{58}\text{Ni}$ ,  $^{62}\text{Ni}$  isotope data and a thorough normal coordinate calculation for “red”  $\text{K}_2[\text{Ni}(\text{dto})_2]$  [31]. The calculated frequencies were in excellent agreement with the observed values with an average error of less than 1%. The

assignments above  $500\text{ cm}^{-1}$  are close to those reported for  $[\text{Pt}(\text{dto})_2]^{2-}$  ( $1602$ ,  $1585\text{ cm}^{-1}$ :  $\nu_{\text{C-O}}$ ;  $1084\text{ cm}^{-1}$ :  $\nu_{\text{C-C}}$  coupled with  $\nu_{\text{C-S}}$ ;  $933\text{ cm}^{-1}$ :  $\nu_{\text{C-S}}$  coupled with  $\delta_{\text{O=C-S}}$ ;  $615\text{ cm}^{-1}$ :  $\nu_{\text{C-S}}$  coupled with  $\nu_{\text{C-C}}$ ,  $\nu_{\text{C-O}}$ ). In contrast to Coucouvanis et al. [28], who found only one band ( $360\text{ cm}^{-1}$ ) shifted by  $^{58}\text{Ni}$ – $^{62}\text{Ni}$  isotope substitution, Czernuszewicz et al. reported that three far-IR bands showed significant isotopic shifts at  $362.0$ ,  $352.6$  and  $435.0\text{ cm}^{-1}$ . The former two are dominated by Ni–S stretching coordinates and the last contains also a significant Ni–S stretching contribution (32%).

The IR and resonance Raman spectra of the “red” and the “black” form of  $\text{K}_2[\text{Ni}(\text{dto})_2]$  differ mainly in the position of three band sets: in the “black” compound the  $\nu_{\text{C-O}}$  vibrations occur at lower frequencies ( $1500$  and  $1488\text{ cm}^{-1}$ ), whereas the band due to the coupled  $\nu_{\text{C-C}} + \nu_{\text{C-S}}$  is found at higher frequency ( $1144\text{ cm}^{-1}$ ). These observed frequency shifts are completely consistent with the bond length changes reported for the X-ray structures (see Table 6). Increasing C–O bond distances cause decreasing frequencies of those modes which involve C–O stretching, and the opposite change is found in the C–S stretching region. Drastic changes of the vibrational frequencies in the third spectral region where the Ni–S stretching vibrations are expected show that force constants and thus vibrational frequencies are obviously rather more sensitive measures of bond lengths than are X-ray data, because Ni–S bond lengths in both X-ray structures are found to be nearly identical [28,60].

The IR and Raman spectra of the non-coordinated dithio-oxalate ion ( $\text{K}_2\text{dto}$ , as well as  $\text{Cs}_2i\text{-dto}$ ,  $\text{K}_2i\text{-dto}$ ,  $\text{K}_2\text{trto} \cdot \text{KCl}$  and  $\text{K}_2\text{mto} \cdot \text{H}_2\text{O}$ ) in the solid state and in solution were published by Mattes et al. in 1977. The fundamentals were also assigned on the basis of a normal coordinate analysis and Raman polarization data [43]. Taking into account the differences between a coordinated and a non-coordinated dto ligand, these results support the assignments for  $[\text{M}(\text{dto})_2]^{2-}$  very well.

As already mentioned in Sect. C, vibrational spectroscopy is especially useful to detect the mode of coordination of the dto ligand and ligand “flipping” from *S,S*- to *O,O*-chelation, which can be followed sometimes in solution [27,29].

Also, when going from the “parent” complex  $\text{K}_2[\text{Ni}(\text{dto})_2]$  to the trinuclear heterobimetallic complex  $(\text{BzPh}_3\text{P})_2[\text{Ni}(\text{dtoSnCl}_4)_2]$ , containing two dto bridges *S,S*-chelated to Ni(II) and *O,O*-chelated to Sn(IV), the IR and Raman spectra are changed dramatically:  $\bar{\nu}_{\text{C-S}}$  drops by more than  $100\text{ cm}^{-1}$  while the coupled  $\nu_{\text{C-C}} + \nu_{\text{C-S}}$  shifts upward by  $70\text{ cm}^{-1}$  and  $\bar{\nu}_{\text{Ni-S}}$  jumps  $17\text{ cm}^{-1}$ . The ground-state MO involves overlap of the Ni(4s) and S( $\sigma$ ) orbitals as well as the metal  $d_{xz}$  ( $d_{yz}$ ) with a  $\pi$  orbital located at the S atom. As electron density is moved toward the oxygen parts of the dto ligands the  $\text{S} \rightarrow \text{Ni}(\sigma)$  bonding is expected to decrease, but apparently  $d_{xz} \rightarrow \pi$  back donation increases, thus stabilizing the Ni–S bond length. This interpretation agrees very well with decreasing C–C and C–S bond lengths [28] (see also Table 6).

In the binuclear complexes  $(\text{BzPh}_3\text{P})_2[(\text{dto})\text{Ni}(\text{dtoSnX}_4)]$ , containing only one dto bridge, the perturbation for this bridging dto is even greater than that observed for dto in the trinuclear compounds. It should also be mentioned that the high-

energy  $\nu_{C-O}$  ( $1640\text{ cm}^{-1}$ ) in the terminal dto is appreciably different from that of the same vibration in the spectrum of  $[\text{Ni}(\text{dto})_2]^{2-}$  [28].

There is IR evidence that dto is thermally and photolytically degraded, producing COS gas [46,51,88,104] (see above).

Resonance Raman spectroscopy is very useful and selective in cases of more than one chromophore in the molecule. Nakamoto and co-workers investigated the trinuclear heterometal complex  $(\text{BzPh}_3\text{P})_2[\text{Ni}(\text{dtoSnI}_4)_2]$ , which has two different chromophoric groups ( $\text{NiS}_4$  and  $\text{SnO}_2\text{I}_4$ ) [30]. Four electronic transitions are observed at  $15.9$ ,  $18.5$ ,  $27.9$  and  $30.0 \times 10^3\text{ cm}^{-1}$ . Using  $611.4$ ,  $514.5$ ,  $457.5$  and  $488.0\text{ nm}$  for the excitation energies, the authors showed that two electronic bands at  $15.9$  and  $18.5 \times 10^3\text{ cm}^{-1}$  are due to the  $\text{Ni} \rightarrow \text{dto}$  charge transfer transitions as assigned earlier by Coucouvanis et al. [28]. The third band at  $27.9 \times 10^3\text{ cm}^{-1}$  is due to the charge transfer within the  $\text{SnO}_2\text{I}_4$  unit, and the last one at  $33.0 \times 10^3\text{ cm}^{-1}$  must be due to a  $\pi \rightarrow \pi^*$  transition of the  $\text{Ni}(\text{dto})_2$  moiety, since the parent complex exhibits a similar absorption.

Nakamoto and his group also used excitation profile studies of two totally symmetric vibrations ( $\nu_1$  and  $\nu_4(\text{Ni-S stretch})$ ) of both forms of  $[\text{Ni}(\text{dto})_2]^{2-}$  ("red" and "black") to reveal the presence of three transitions lying underneath the visible absorption envelope [31], as claimed by Latham et al. [89] on the basis of MO calculations. The unusual nature of the excitation profiles of the "red"  $[\text{Ni}(\text{dto})_2]^{2-}$  gives a useful restriction of the order of the highest occupied molecular orbitals (HOMO) and supports the order published in ref. 89.

A resonance Raman study on the visible absorption spectrum of tris(dithio-oxalato)cobaltate(III) was carried out by Kanamori et al. [103]. They found that the Raman bands due to the totally symmetric coupled  $\nu_{C-C} + \nu_{C-S}$  ( $113\text{ cm}^{-1}$ ) and  $\nu_{\text{Co-S}}$  ( $349\text{ cm}^{-1}$ ) exhibit resonance enhancements with excitation lines in the visible region. The resonance Raman excitation profiles of these two modes reveal the presence of a charge-transfer transition in the visible absorption spectrum (see also the previous chapter).

#### (f) ESCA spectra

The only two reports concerning X-ray photoelectron spectra on dithio-oxalate complexes are by Folkesson and Jonson in 1984 [87] and Bellitto et al. in 1989 [77(a)]. Findings for the potassium and  $\text{Ph}_4\text{P}^+$  salts of the Fe, Co, Ni, Pd, Pt, and Cu complexes (the Co complexes are described as Co(II) bis-chelates; however, they are probably Co(III) tris-chelates) are presented in the first paper.

The effective charges have been calculated from the measured binding energies (see Table 10) using the following relations for the ligand atoms

$$E_b(\text{S } 2p) = 3.38q_s + 163.8$$

$$E_b(\text{C } 1s) = 11.5q_c + 284.8$$

$$E_b(\text{O } 1s) = 6.04q_o + 535.6$$



TABLE 10

Binding energies (eV) of dithio-oxalate and tetrathio-oxalate complexes

Compound	S 2p	O 1s	C 1s	P 2p	M 2p <sub>1/2</sub>	M 2p <sub>3/2</sub>	M 3d <sub>5/2</sub>	M 4f <sub>7/2</sub>	Ref.
K <sub>3</sub> [Fe(dto) <sub>3</sub> ]	163.3	531.0	287.9			708.6			87 <sup>a</sup>
K <sub>2</sub> [Co(dto) <sub>2</sub> ] <sup>b</sup>	162.2	530.8	287.6			778.8			87 <sup>a</sup>
K <sub>2</sub> [Ni(dto) <sub>2</sub> ]	162.1	530.4	286.3			852.8			87 <sup>a</sup>
K <sub>2</sub> [Pd(dto) <sub>2</sub> ]	162.3	530.5	286.5				336.9		87 <sup>a</sup>
	162.4						337.4		77(a) <sup>c</sup>
K <sub>2</sub> [Pt(dto) <sub>2</sub> ]	162.4	530.6	286.3					72.1	87 <sup>a</sup>
K <sub>2</sub> [Cu(dto) <sub>2</sub> ] <sup>d</sup>	162.6	530.7	286.6			931.3			87 <sup>a</sup>
(Et <sub>4</sub> N) <sub>2</sub> [Pt(dto) <sub>2</sub> ]	162.8						75.6	72.2	77(a) <sup>c</sup>
(Ph <sub>4</sub> P) <sub>3</sub> [Fe(dto) <sub>3</sub> ]	163.1	531.4	288.0	133.5		708.3			87 <sup>f</sup>
(Ph <sub>4</sub> P) <sub>2</sub> [Co(dto) <sub>2</sub> ] <sup>e</sup>	162.8	531.4	288.2	133.3		779.3			87 <sup>f</sup>
(Ph <sub>4</sub> P) <sub>2</sub> [Ni(dto) <sub>2</sub> ]	162.9	531.3	287.1	133.5		853.6			87 <sup>f</sup>
(Ph <sub>4</sub> P) <sub>2</sub> [Pd(dto) <sub>2</sub> ]	163.1	531.2	287.2	133.6			337.4		87 <sup>f</sup>
(Ph <sub>4</sub> P) <sub>2</sub> [Pt(dto) <sub>2</sub> ]	163.4	531.3	287.1	133.5				72.9	87 <sup>f</sup>
(Ph <sub>4</sub> P) <sub>2</sub> [Cu(dto) <sub>2</sub> ] <sup>d</sup>	163.1	531.3	287.2	133.6		931.9			87 <sup>f</sup>
(TTF) <sub>2</sub> [Pt(dto) <sub>2</sub> ]	161.8					342.2			87 <sup>f</sup>
	164.0 <sup>a</sup>						337.4		77(a) <sup>c</sup>

TABLE 10 (continued)

Compound	S 2p	O 1s	C 1s	P 2p	M 2p <sub>1/2</sub>	M 2p <sub>3/2</sub>	M 3d <sub>5/2</sub>	M 4f <sub>7/2</sub>	Ref.
(TTF) <sub>2</sub> [Pt(dto) <sub>2</sub> ]	162.6 164.7 <sup>h</sup>						75.5	72.3	77(a) <sup>e</sup>
(TTF) <sub>3</sub> [Pt(dto) <sub>2</sub> ]	162.5 163.8 <sup>i</sup> 164.8 <sup>h</sup>						75.4	72.1	77(a) <sup>e</sup>
TTF[Cu(dto) <sub>2</sub> ] <sup>d</sup>	162.1 164.0 <sup>h</sup>				955.8	935.6			77(a) <sup>e</sup>
{[(Ph <sub>3</sub> P) <sub>2</sub> Cu] <sub>2</sub> (tto)}	162.6			131.7		932.4			177
{Cu[(tto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> }	163.1			131.9		932.4			177
{Pd[(tto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub> }	162.7			131.8		932.6	337.6		177

<sup>a</sup> Referred to E<sub>b</sub>(K 2p<sub>3/2</sub>) = 292.2 eV.<sup>b</sup> More likely K<sub>3</sub>[Co(dto)<sub>3</sub>].<sup>c</sup> Referred to E<sub>b</sub>(K 2p<sub>3/2</sub>) = 292.9 eV.<sup>d</sup> Obviously reduction to Cu(I) upon irradiation during measurements.<sup>e</sup> Referred to E<sub>b</sub>(C 1s) = 285.0 eV.<sup>f</sup> Referred to E<sub>b</sub>(C 1s) = 285.5 eV.<sup>g</sup> More likely (Ph<sub>4</sub>P)<sub>3</sub>[Co(dto)<sub>3</sub>].<sup>h</sup> TTF<sup>+</sup>.<sup>i</sup> TTF<sup>0</sup>.

Table 11 summarizes all estimated effective charges. As shown, the charge at the oxygen atoms is nearly the same in all dto complexes. A similar situation occurs for the dto carbons (exceptions are found, however, for the tris-chelates  $[M(dto)_3]^{3-}$ ,  $M = Fe, Co$ ). The largest variation in charge is estimated for the sulphur ligator atoms. Taking into account that no charge transfer takes place between the cation and the complex anion, the charge at the central metal ion ( $q_M$ ) could be calculated from the charges on the ligand atoms, giving the following series:  $q_{Fe}, q_{Co} < q_{Ni}$  and  $q_{Ni} > q_{Pd} > q_{Pt}$ . The first-row trend implies a general increase of covalence in the  $M-S$  bond as the d orbitals are gradually filled. The trend in the nickel triad is expected from the increased softness of the metal ions: the greater the tendency towards a  $S \rightarrow M(\sigma)$  bonding, the smaller  $q_M$  will be. This effect is also reflected by the decreasing negative charges on the sulphur donor atoms.

The authors could show that the copper(II) in  $[Cu(dto)_2]^{2-}$  is actually reduced during X-ray bombardment.

In the second paper the ESCA data are used to confirm the oxidation state of the metal ions in the donor-acceptor systems  $(TTF)_m[M(dto)_2]$ ,  $(TTF)_2[Pt(dto)_2]$ ,  $(TTF)_3[Pt(dto)_2]$  and  $(TTF)_2[Pd(dto)_2]$ . They contain the metal ions in the oxidation state +2. The S 2p peak analysis obtained by deconvolution gave three values (162.5, 163.8 and 164.8 eV) in the case of  $(TTF)_3[Pt(dto)_2]$  attributed to the S 2p orbitals of the anion sulphur, of the  $(TTF^+)_2$  cation and the neutral TTF molecule found by crystal structure analysis.  $(TTF)_{1.0}[Cu(dto)_2]$  does not show shake-up satellites (expected for a paramagnetic Cu(II) compound). It seems that the initial Cu(III) complex is reduced at its surface to a Cu(I) compound during X-ray exposure, accompanied by elimination of gaseous COS.

#### (g) Redox behaviour

Although the  $[M(dto)_2]^{2-}$  complexes can be formally classified as 1,2-dithiolene complexes, they do not exhibit the extensively studied electron transfer behaviour

TABLE 11

Estimated effective charges on the atoms in various dithio-oxalate complexes [87]<sup>a</sup>

Compound	$q_s$	$q_c$	$q_o$	$q_M$
$(Ph_4P)_3[Fe(dto)_3]$	-0.29	+0.28	-0.70	+0.8(2)
$(Ph_4P)_2[Co(dto)_2]^b$	-0.29	+0.30	-0.70	+0.8(2)
$(Ph_4P)_2[Ni(dto)_2]$	-0.26	+0.20	-0.71	+1.1(2)
$(Ph_4P)_2[Pd(dto)_2]$	-0.21	+0.21	-0.73	+0.9(2)
$(Ph_4P)_2[Pt(dto)_2]$	-0.12	+0.20	-0.71	+0.5(2)
$(Ph_4P)_2[Cu(dto)_2]^c$	-0.21	+0.21	-0.71	+0.8(2)

<sup>a</sup>The electronic charge is used as a unit.


<sup>b</sup>More likely  $(Ph_4P)_3[Co(dto)_3]$ .

<sup>c</sup>Obviously reduction to Cu(I) upon irradiation during measurements.

typical of the latter [123,124]. Copper bis-dithio-oxalate is the only binary dithio-oxalate complex which undergoes a reversible one-electron transition



The electrochemical oxidation occurs for example in  $\text{CH}_2\text{Cl}_2$  (at scan rates between 0.02 and  $20 \text{ V s}^{-1}$ ), at a potential  $E_{1/2} = +0.128(3) \text{ V}$  vs. the SCE. The electrochemical reduction process of  $[\text{Cu}(\text{dto})_2]^{-}$  is also reversible over the same scan rate scale:  $E_{1/2} = +0.135(2) \text{ V}$  [51]. The "oxidation product"  $[\text{Cu}(\text{dto})_2]^{-}$  was described originally as  $[\text{Cu}(\text{dto})(\text{SCO})_2]^{-}$  because addition of  $\text{PPh}_3$  splits off COS gas ([88], see Sect. D (iii)(b)). The reason is the redox-substitution reaction



Thus, even in the presence of a low concentration of  $\text{PPh}_3$  the cathodic peak disappears and the anodic peak shifts to more negative values.

The oxidation of  $[\text{Cu}(\text{dto})_2]^{2-}$  can also be effected chemically in DMF or  $\text{CH}_2\text{Cl}_2$  by either Cu(II) or Fe(III) ions [88]. Kinetic studies were carried out using major changes in the absorption spectra of  $(\text{Bu}_4\text{N})_2[\text{Cu}(\text{dto})_2]$  ( $2.54 \times 10^{-4} \text{ M}$  solution) after adding  $\text{Bu}_4\text{N}(\text{FeCl}_4)$  ( $4.36 \times 10^{-3} \text{ M} > [\text{Bu}_4\text{N}(\text{FeCl}_4)]_0 \geq 0$ ) solutions. The reaction proceeds in a 1 : 1 molar ratio. Kinetic measurements were also carried out for the  $(\text{Ph}_4\text{P})_2[\text{Cu}(\text{dto})_2] - \text{Bu}_4\text{N}(\text{FeCl}_4)$  and  $(\text{Bu}_4\text{N})_2[\text{Cu}(\text{dto})_2] - (\text{Bu}_4\text{N})_2(\text{CuCl}_4)$  systems. The higher (by more than 10 times) second-order rate constant in the case of Cu(II) oxidation suggests that the metal ion of the oxidant markedly influences the rate-determining step [51].

A compound  $\text{TTF}[\text{Ni}(\text{dto})_2]$  is mentioned in [77(b)]. The details given, however, do not give sufficient confirmation for a Ni(III) species.

The only reversible one-electron reduction is reported for  $\text{Bu}_4\text{N}[\text{TcO}(\text{dto})_2]$  (voltammetry at Hg pool or rotating Pt electrode; SCE; acetonitrile; 0.1 M tetrabutylammonium perchlorate (TBAP);  $-0.75 \text{ V}$ ). A second irreversible step follows at  $-1.56 \text{ V}$ . The analogous Re compound exhibits two irreversible reduction steps ( $-0.94$  and  $-1.35 \text{ V}$ ). None of the reduction products is accessible by conventional chemical techniques [95].

Unlike the strongly irreversible reductions of the parent bis(dithio-oxalato)metalates  $[\text{M}(\text{dto})_2]^{2-}$  ( $\text{M} = \text{Ni}$ :  $-2.03$ ;  $\text{Pd}$ :  $-2.05$ ;  $\text{Pt}$ :  $< -2.05 \text{ V}$ ), the 1 : 1 "adducts"  $[(\text{dto})\text{M}(\text{dtoSnCl}_4)]^{2-}$  undergo a single reversible one-electron reduction (cyclovoltammetric data taken in  $\text{CH}_2\text{Cl}_2$ ; 0.1 M LiCl; Ag/AgCl;  $\text{M} = \text{Ni}$ :  $-0.53$ ;  $\text{Pd}$ :  $-0.54$ ;  $\text{Pt}$ :  $-0.15 \text{ V}$ ) whereas the 2 : 1 compounds  $[\text{M}(\text{dtoSnCl}_4)_2]^{2-}$  undergo reversible reduction in two steps ( $\text{M} = \text{Ni}$ :  $-0.15$ ,  $-0.51$ ;  $\text{Pd}$ :  $-0.23$ ,  $-0.52$ ;  $\text{Pt}$ :  $-0.15$ ,  $-0.54 \text{ V}$ ) [107] (see also ref. 108). Using other Sn(IV) halides, only the compound  $[(\text{dto})\text{Ni}(\text{dtoSnF}_4)]^{2-}$  shows a reversible reduction (cyclovoltammetry in  $\text{CH}_2\text{Cl}_2$ ; TBAP; Ag/AgI:  $-0.99 \text{ V}$ ). Under similar conditions the bromides and iodides  $[\text{M}(\text{dtoSnX}_4)_2]^{2-}$  gave irreversible reductions ( $\text{Ni}/\text{SnBr}_4$ :  $-0.55$ ;  $\text{Ni}/\text{SnI}_4$ :  $-0.42$ ;  $\text{Pd}/\text{SnBr}_4$ :  $-0.63$ ;  $\text{Pd}/\text{SnI}_4$ :  $-0.37 \text{ V}$ ) [28].

It has been suggested that the (reversible) reductions are ligand rather than transition metal based, where a ligand antibonding orbital ( $b_{1u}$ ) takes the added electrons whose energy is lowered relatively to that in the parent bis(dithio-oxalato)metalates by coordination of  $SnX_4$  to the “diketone ends” of the dto ligands (see Fig. 20) [28,108].

Cyclovoltammetric measurements on mixed ligand compounds  $[L_2M(dto)]$  ( $M = Ni, Pd, Pt$ ;  $L_2 = 1,2$ -bis(diphenylphosphino)ethane, 1,2-bis(diethylphosphino)ethane or two  $PMe_3$ ) also revealed chemically reversible reductions for all Pt compounds. Most of the Ni and Pd complexes containing the bisphosphine ligands show some degree of chemical reversibility. The ease of reduction follows the order  $Ni > Pd > Pt$ , as a result of the increasing stability of  $d^8$  square-planar structures by increasing crystal field strength on descending the metal triad. Controlled potential coulometry investigations for  $[(Ph_2PCH_2CH_2PPh_2)Pt(dto)]$  showed 0.95(1) electron transferred in the reduction [46(b),109].

$[Ni(dto)_2]^{2-}$  is reduced irreversibly at the dropping mercury electrode in two steps (aqueous solution, supporting electrolyte KCl), whereas  $[Ni(CN)_4]^{2-}$  exhibits under similar conditions only one irreversible reduction. All three original steps disappeared when a mixture of both complexes was investigated by polarography. A new well-defined step was described as proof for the formation of the mixed ligand complex  $[(dto)Ni(CN)_2]^{2-}$  [111].


In contrast to, for example, platinum bis-oxalate  $[Pt(ox)_2]^{2-}$ , which can be partially oxidized to form a highly one-dimensional conducting system [125,126], attempts to synthesize similar dto systems failed. However, oxidation of “red”  $K_2[Ni(dto)_2]$  by iodine in acetone or acetonitrile (see above) yielded a blue “adduct” (formulated as  $K_2[Ni(dto)_2I]$ ) which obviously exists in equilibrium with  $K_2[Ni(dto)_2]$ . The blue compound could not be isolated in the solid state. The “oxidation” using  $K_2Cr_2O_7$  resulted in a blend of the parent “red” form and a polymorphic “black” form, both, however, containing nickel(II) (see above).

#### (h) Photo- and thermochemical behaviour

Most information concerning the behaviour of dithio-oxalate complexes towards light and heat is qualitative.  $K[Au(dto)_2]$  was found to be unstable to heat and light [89]. Also, tris-chelates  $[M(dto)_3]^{3-}$  ( $M = Cr, Fe, Co$ ) are reported to decompose very fast in the presence of light [92]. As already mentioned (see Sect. D (iii)(f)), the copper ions in  $[Cu(dto)_2]^{2-}$  and  $[Cu(dto)_2]^-$  are actually reduced to Cu(I) during X-ray bombardment in ESCA experiments [77(a),87]. It seems, however, that the Cu(III) complex  $[Cu(dto)_2]^-$  is appreciably more light-sensitive than the Cu(II) complex. Thus the cleavage of the C–C bond in one of the coordinated dto ligands of this compound is light-induced. Recording the IR spectra, the first rapid scan in the narrow IR region 2020–2060  $cm^{-1}$  showed no evidence of absorption at 2055  $cm^{-1}$  (COS). However, subsequent repetitive scans over the same region produced a band appearing at 2055  $cm^{-1}$ , which increased in intensity as a function of

exposure time to the IR light source. It appears, therefore, that light, including the visible light component of the IR light source, promotes an intramolecular two-electron  $\text{dto} \rightarrow \text{Cu(III)}$  transfer in  $[\text{Cu(dto)}_2]^-$  with cleavage of the C–C bond and generation of COS [51,88].

The photochemistry of  $[\text{M(dto)}\text{L}_2]$  ( $\text{L} = \text{phosphines}$ ) was investigated in detail [46]. All compounds ( $\text{M} = \text{Ni, Pd, Pt}$ ) are slightly light-sensitive in the solid state and thermally stable only below  $50^\circ\text{C}$ . The thermal and photochemical sensitivity decreases in the sequence  $\text{Ni} > \text{Pd} > \text{Pt}$ . After the discovery of light-induced C–C bond splitting in dto complexes, attempts were made to generate COS in this way within the coordination sphere of coordinatively unsaturated zero-valent metals. However, the electronic spectra of the nickel and palladium species show only a decrease in the initial absorbing complex concentration on UV photolysis. The platinum complexes show isosbestic behaviour which varies with the solvent used (in  $\text{CH}_3\text{CN}$ , isosbestic behaviour is observed even with extended photolysis times; in  $\text{CH}_2\text{Cl}_2$  there is initially an isosbestic point which disappears on continued irradiation). Secondary photolysis or thermal decomposition according to the equation



occurs in halogenated solvents (e.g.  $\text{CH}_2\text{Cl}_2$ ) when a solution of  $[\text{Ni(dto)}\text{L}_2]$  ( $\text{L}_2 = \text{R}_2\text{PCH}_2\text{CH}_2\text{PR}_2$ , with  $\text{R} = \text{Et}$  or  $\text{Ph}$ ) is irradiated; IR absorption at  $2030\text{ cm}^{-1}$  attributed to free COS starts growing immediately. This peak is soon followed by new absorptions attributable to nickel carbonyls resulting from the carbonylating ability of COS. Two of the photolysis products in  $\text{CH}_3\text{CN}$  were identified as  $[\text{Ni}(\text{CO})_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)]$  and  $[\text{Ni}(\eta^2\text{-COS})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)]$ . The same metal species are formed in the photochemical and thermal reactions. Therefore one can suggest that the photochemistry of these metal bis(phosphine)dithio-oxalates proceeds through a coordinatively unsaturated zero-valent species which reacts with available COS.

Some bis- and tris-dithio-oxalato complexes are patented as photosensitive components in photoimaging materials [127,128].

(i) *Ligand exchange reactions and mixed ligand complexes*

The formation of  $[\text{Ni(dto)(CN)}_2]^{2-}$  in water according to the equation



was detected polarographically in 1959 [111]. This reaction was also confirmed spectrophotometrically [110]. However, using identical conditions, it was impossible to extend the reaction to  $\text{Pd(II)}$  and  $\text{Pt(II)}$  complexes.

Also in solution, a large number of mixed ligand complexes of  $\text{Ni(II)}$ ,  $\text{Pd(II)}$  and  $\text{Cu(II)}$  containing the dto ligand have been generated and investigated by Pearson


et al. [17,86,112]. They studied the kinetics of ligand exchange reactions of bis(dithiolato)metalate(II) complexes with dithiolates or cyanide as the nucleophiles mainly in aqueous solution at 25°C using a stopped-flow technique. The reactions in which dto is involved either as the nucleophile, as the leaving group or as the *trans* group are summarized in Table 12 (abbreviations of the ligands are listed in a footnote to the table). Most experiments have been carried out with Ni(II). The following conclusions were drawn.

The stability constant of  $[\text{Ni(dto)}_2]^{2-}$  (16.1) fits better in a series of constants of 1,1-dithiolato complexes (cpd: 20.3, ned: 14.4, cdc: 13.7, *i*-mnt: 12.2) than of 1,2-dithiolato (dithiolene) compounds (tfd = 1,2-trifluoromethylethylene-1,2-dithiolate: > 36, and mnt: < 25). This is somewhat surprising because dto forms five-membered chelate rings like dithiolenes which are considerably more stable than the analogous four-membered rings formed by the 1,1-dithiolates.

Keeping the *trans* group and the nucleophile constant, the ease of displacement of the leaving group decreases in the series *i*-mnt, ned > cds > dto > cpd > mnt, following (with the exception of ned) the inverted series of stability constants.

The general order for the dithiolato nucleophiles, keeping the leaving group and the *trans* group constant, is mnt > ned > cpd > cdc > dto, which only very roughly parallels the stability series.

It was proposed that substitution mechanisms both with bidentate and  $\text{CN}^-$  nucleophiles follow an associative path. This means that the first step should in all cases be the rapid formation of a five-coordinate adduct. Considering the reactions



the stability of the cyanide adducts decreases as L is changed: *i*-mnt ( $k(\text{M}^{-1}) > 1000$ ) > cdc (850) > ned (160) > dto (< 50). The adduct constants agree with the assumption that the larger the  $\pi$  density in the  $4p_z$ , the smaller is the tendency for a nucleophile to form a bond along the *z* direction.

The effect of the *trans* group, keeping the nucleophile and the leaving group constant, on the kinetics of nickel(II) dithiolate substitution reactions decreases in the order *i*-mnt > cdc > dto > cpd > mnt and ned > cpd, agreeing nicely with the stability series of the five-coordinate adducts. This leads to the suggestions that the rate of ligand exchange reactions increases as the adduct stability increases and that the major part of the *trans* influence operates via the stability of the five-coordinate adduct which must form before substitution can occur.

The exchange behaviour of palladium(II) complexes largely parallels the behaviour of nickel(II) complexes. The fact that only one reaction is observed when  $[\text{Pd(dto)}_2]^{2-}$  reacts with cyanide is probably due to the apparent instability of  $[\text{Pd(dto)(CN)}_2]^{2-}$ . The reaction sequence appears to be as shown in the equations




TABLE 12

Second-order rate constants for ligand exchange reactions of Ni(II), Pd(II) and Cu(II) complexes with bidentate (or tridentate) ligands including dithio-oxalate as ligand and/or nucleophile and first-order rate constants with cyanide, determined in aqueous solution at 25°C<sup>a</sup>

Reactant <sup>b</sup>	Nucleophile <sup>b</sup>	Product	$k_{\text{obsd}}$ (10 <sup>-3</sup> M <sup>-1</sup> s <sup>-1</sup> )	$k_{\text{obsd}}$ (s <sup>-1</sup> ) <sup>c</sup>	Ref.
[Ni(dto) <sub>2</sub> ] <sup>2-</sup>	mnt <sup>2-</sup>	[Ni(dto)(mnt)] <sup>2-</sup>	0.33		86
[Ni(dto)(mnt)] <sup>2-</sup>	mnt <sup>2-</sup>	[Ni(mnt) <sub>2</sub> ] <sup>2-</sup>	0.019		86
[Ni(dto)(cpd)] <sup>2-</sup>	mnt <sup>2-</sup>	[Ni(cpd)(mnt)] <sup>2-</sup>	0.11		86
[Ni(dto)(cdc)] <sup>2-</sup>	mnt <sup>2-</sup>	[Ni(dto)(mnt)] <sup>2-</sup>	23		86
[Ni(dto)(i-mnt)] <sup>2-</sup>	mnt <sup>2-</sup>	[Ni(dto)(mnt)] <sup>2-</sup>	53		86
[Ni(cdc) <sub>2</sub> ] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto)(cdc)] <sup>2-</sup>	6.0		86
[Ni(dto)(cdc)] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto) <sub>2</sub> ] <sup>2-</sup>	0.55		86
[Ni(cdc)(mnt)] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto)(mnt)] <sup>2-</sup>	0.0057		86
[Ni(cdc)(CN) <sub>2</sub> ] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto)(CN) <sub>2</sub> ] <sup>2-</sup>	0.25		86
[Ni(i-mnt) <sub>2</sub> ] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto)(i-mnt)] <sup>2-</sup>	28		86
[Ni(dto)(i-mnt)] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto) <sub>2</sub> ] <sup>2-</sup>	1.6		86
[Ni(i-mnt)(mnt)] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto)(mnt)] <sup>2-</sup>	0.025		86
[Ni(i-mnt)(CN) <sub>2</sub> ] <sup>2-</sup>	dto <sup>2-</sup>	[Ni(dto)(CN) <sub>2</sub> ] <sup>2-</sup>	0.80		86




$[\text{Ni}(i\text{-mnt})(\text{cpd})]^{2-}$	dto <sup>2-</sup>	$[\text{Ni}(\text{dto})(\text{cpd})]^{2-}$	0.52	86
$[\text{Ni}(\text{cpd})_2]^{2-}$	dto <sup>2-</sup>	$[\text{Ni}(\text{dto})(\text{cpd})]^{2-}$	0.00087	86
$[\text{Ni}(\text{dto})(i\text{-mnt})]^{2-}$	cdc <sup>2-</sup>	$[\text{Ni}(\text{dto})(\text{cdc})]^{2-}$	≈10	86
$[\text{Ni}(\text{dto})(\text{mnt})]^{2-}$	cpd <sup>2-</sup>	$[\text{Ni}(\text{mnt})(\text{cpd})]^{2-}$	0.0049	86
$[\text{Ni}(\text{dto})_2]^{2-}$	en	$[\text{Ni}(\text{en})_3]^{2+}$	0.15	86
$[\text{Ni}(\text{dto})_2]^{2-}$	dien	$[\text{Ni}(\text{dien})_2]^{2+}$	0.10	86
$[\text{Pd}(i\text{-mnt})_2]^{2-}$	dto <sup>2-</sup>	$[\text{Pd}(\text{dto})(i\text{-mnt})]^{2-}$	0.0125	112
$[\text{Pd}(\text{ned})_2]^{2-}$	dto <sup>2-</sup>	$[\text{Pd}(\text{dto})(\text{ned})]^{2-}$	0.197	112
$[\text{Pd}(\text{dto})(\text{ned})]^{2-}$	dto <sup>2-</sup>	$[\text{Pd}(\text{dto})_2]^{2-}$	0.0038	112
$[\text{Cu}(\text{dto})(\text{mnt})]^{2-}$	mnt <sup>2-</sup>	$[\text{Cu}(\text{mnt})_2]^{2-}$	≈140	86
$[\text{Ni}(\text{dto})_2]^{2-}$	CN <sup>-</sup>	$[\text{Ni}(\text{dto})(\text{CN})_2]^{2-}$	$40 \times 10^6 [\text{CN}^-]^2$	86
$[\text{Ni}(\text{dto})(\text{CN})_2]^{2-}$	CN <sup>-</sup>	$[\text{Ni}(\text{CN})_4]^{2-}$	$12 \times 10^6 [\text{CN}^-]^2$	86
$[\text{Ni}(\text{dto})(\text{mnt})]^{2-}$	CN <sup>-</sup>	$[\text{Ni}(\text{mnt})(\text{CN})_2]^{2-}$	$3.1 \times 10^4 [\text{CN}^-]^2$	86
$[\text{Pd}(\text{dto})_2]^{2-}$	CN <sup>-</sup>	$[\text{Pd}(\text{CN})_4]^{2-}$	$2.7 \times 10^4 [\text{CN}^-]^2$	112

<sup>a</sup> Ionic strength 0.2 M (NaClO<sub>4</sub>) for nickel complexes and 0.1 M for palladium complexes.



<sup>b</sup> Abbreviations: mnt = maleonitriledithiolate, cpd = 1-cyano-1-phenylethylene-2,2-dithiolate, cdc = *N*-cyanodithiocarbamate, *i*-mnt = 1,1-dicyanoethylene-2,2-dithiolate, ned = 1-nitroethylene-2,2-dithiolate, en = ethylenediamine, dien = diethylenetriamine.

<sup>c</sup>  $k_{\text{obsd}}$  is the pseudo-first-order rate constant obtained with cyanide in excess.



Comparing the rate of the first reaction with the rate of the corresponding reaction of  $[\text{Ni}(\text{dto})_2]^{2-}$ , a ratio  $K_{\text{Ni}}/K_{\text{Pd}} = 1470$  has been found. This rate increase on going from palladium to nickel is unusually large, because ratios of the order of 10 are expected [129].

The species  $[\text{Cu}(\text{dto})_2]^{2-}$  reacts with  $\text{mnt}^{2-}$  according to the equations



The reaction is very fast with an approximate second-order rate constant  $k \approx 1.4 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ , indicating that copper(II) dithio-oxalate reacts, as expected, more than 7000 times faster than nickel(II) dithio-oxalate with  $\text{mnt}^{2-}$  (see Table 12).

In addition to UV–VIS spectroscopy, EPR is suitable for detecting copper(II) mixed ligand systems, as mentioned before [118]. In solution,  $[\text{Cu}(\text{dto})_2]^{2-}$  reacts with Cu(II) benzoylthio- or selenoureas forming mixed ligand species which contain a five-membered (dto) and a six-membered (urea ligand) chelate ring [116].

Some mixed ligand Co(III) complexes containing dto and a nitrogen donor ligand have been reported [36,93,104]. Obviously the  $\text{N}_4\text{S}_2$  donor atom set in  $[\text{Co}(\text{en})_2(\text{dto})]\text{X}$  and  $[\text{Co}(\text{phen})_2(\text{dto})]\text{X}$  is particularly stable and forms easily. A special case of a mixed ligand complex is found in  $[\text{In}_2(\text{dto})_5]^{4-}$ , in which two different terminal “cis-dtos” and a planar *trans* bridging dto were detected by X-ray structure analysis [41]. There is also a huge number of dto complexes containing additional halogenides, methyl, carbonyl, phosphines, oxygen, sulphur, nitrogen or nitrosyl as ligands:  $\{\text{XSn}[(\text{dto})\text{Cu}(\text{PPh}_3)_2]_3\}$  ( $\text{X} = \text{Cl}, \text{Br}, \text{Me}$ ) [29],  $\{\text{X}_2\text{Sn}[(\text{dto})\text{Cu}(\text{PPh}_3)_2]_2\}$  ( $\text{X} = \text{Cl}$  and/or  $\text{Me}$ ) [29],  $\{\text{X}_2\text{Cl}_2\text{Sn}[(\text{dto})\text{Cu}(\text{PPh}_3)_2]\}^-$  ( $\text{X} = \text{Cl}$  and/or  $\text{Me}$ ) [29],  $\{\text{Sn}_3\text{Cl}_4[(\text{dto})\text{Cu}(\text{PPh}_3)_2]_2\}$  [29],  $[\text{Cu}(\text{dto}\text{SnCl}_4)_2\text{Cl}]^{3-}$  [88],  $\{[\text{Mn}(\text{CO})_5]_2(\text{dto})\}$  [25],  $\{[(\text{Co})_3\text{Re}(\text{dto})]_2(\text{dto})\}$  [26],  $[\text{Mo}_2\text{O}_2\text{S}_2(\text{dto})_2]^{2-}$  [21,94],  $[(\text{S}_2)\text{MoO}(\mu\text{-S})_2\text{MoO}(\text{dto})]^{2-}$  [94],  $[\text{MA}(\text{dto})_2]^n$  ( $\text{M} = \text{Mo}, \text{A} = \text{O}, n = 2$  [39];  $\text{M} = \text{Tc}, \text{Re}, \text{A} = \text{O}, n = 1$  [26,55,95];  $\text{M} = \text{Tc}, \text{A} = \text{N}, n = 2$  [55,56];  $\text{M} = \text{Fe}, \text{A} = \text{Cl}, \text{Br}, \text{I}, \text{NO}, n = 2$  [96–99]),  $[(\text{PMe}_3)_2\text{M}(\text{dto})]$  ( $\text{M} = \text{Ni}, \text{Pd}, \text{Pt}$ ) [46(b)],  $[(\text{PPh}_3)_2\text{Cu}(\text{dto})]^-$  [29,51],  $[(\text{R}_2\text{PCH}_2\text{CH}_2\text{PR}_2)\text{M}(\text{dto})]$  ( $\text{M} = \text{Ni}, \text{Pd}, \text{Pt}, \text{R} = \text{Et}, \text{Ph}$ ) [46(b),109],  $[(\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_3)\text{Ni}(\text{dto})]$  [46(b)],  $[\text{Pd}_3(\mu_3\text{-S})_2(\text{dto})(\text{PMe}_3)_4]$  [46(b)]. Mixed ligand systems are also formed when only one of two or three dto ligands present reacts with coordinatively unsaturated metal complex species. Thereby ligand scrambling reactions (*S,S* to *O,O* coordination) can occur depending on the relative softness of the metal species:  $[(\text{dto})\text{M}(\text{S}_2\text{C}_2\text{O}_2\text{SnX}_4)]^{2-}$  ( $\text{M} = \text{Ni}; \text{X} = \text{F}, \text{Cl}$  or  $\text{M} = \text{Pd}, \text{Pt}; \text{X} = \text{Cl}$ ) [28,107,108] and  $\{(\text{dto})_2\text{Sn}[(\text{O}_2\text{C}_2\text{S}_2)\text{Cu}(\text{PR}_3)_2]\}^-$  ( $\text{R} = \text{Ph}, \text{Tol}$ ) [29,33].

(j) *Analytical use of dto complexes*

In their classical paper in 1909, Jones and Tasker already pointed to the analytical usefulness of dithio-oxalate for determining Ni, Co and Fe ions [2]. In the 1920s, techniques were developed to determine nickel colorimetrically in neutral or acidic aqueous solution (also in tissues, excreta and foodstuffs) with dithio-oxalate [130,131]. The complex composition depending on the relative concentrations of  $\text{Ni}^{2+}$  and  $\text{dto}^{2-}$  in aqueous solution (Job's method) and the validity of Beer's law were checked and formation constants were evaluated (necessary for unambiguous analytical use) [86,119]. Welcher devotes a separate chapter in his reference book [132] to dithio-oxalate. Besides a procedure to synthesize potassium dithio-oxalate, he describes reactions of this reagent with metal ions, especially nickel, cobalt and iron. The colour of nickel dithio-oxalate is approximately three times more intense than that of the permanganate ion, thus producing a sensitivity for nickel of up to  $1:4 \times 10^7$ . The colour reaction with cobalt is as intense as that of nickel and may therefore interfere. The sensitivity of the iron reaction, however, is only about  $1:10^5$ .

In 1982, two papers were published dealing with ion-pair extraction of  $[\text{M}(\text{dto})_2]^{2-}$  complexes from aqueous solution into chloroform by use of bulky organic cations Q, forming 1:2 ion pairs  $\text{Q}_2[\text{M}(\text{dto})_2]$  [133,134]. The extraction constants  $K_{\text{ex}} = [\text{Q}_2[\text{Ni}(\text{dto})_2]]$  (in  $\text{CHCl}_3$ )/ $[\text{Q}^+]^2[\text{Ni}(\text{dto})_2]^{2-}$  (in  $\text{H}_2\text{O}$ ) were determined for dodecyltrimethylammonium ( $\log K_{\text{ex}} = 8.97$ ), tetrabutylammonium (9.38), *N*-dodecylpyridinium (11.37), hexadecyltrimethylammonium (12.88), benzyldimethyldodecylammonium (14.75), *N*-hexadecylpyridinium (14.82), benzyldimethyltetradecylammonium (16.46), and tetrahexylammonium (18.05) at ionic strength 0.3 and 25°C. A linear correlation was obtained between  $\log K_{\text{ex}}$  and the number of carbon atoms constituting the  $\text{Q}^+$  ions. However, the data for cations with an aromatic group and those without follow two independent lines. Empirical parameters for estimating extraction constants have been deduced: the values  $\Delta \log K_{\text{ex}}$  (the contribution of the functional group to  $\log K_{\text{ex}}$ ) were 0.47 for the methylene group, 0.82 for the methyl group, 3.18 for the phenyl group, 0.95 for the ammonium ion, 4.52 for the pyridinium ion, and  $-9.79$  for the  $[\text{Ni}(\text{dto})_2]^{2-}$  anion.

The simultaneous spectrophotometric determination of palladium(II) and platinum(II) can be carried out after ion-pair extraction of  $[\text{Pd}, \text{Pt}(\text{dto})_2]^{2-}$  from water into chloroform using benzyldimethyltetradecylammonium perchlorate as a quaternary ammonium ion buffer (the concentration of the benzyldimethyltetradecylammonium ion  $\text{Q}^+$  in the aqueous phase is controlled by the combined use of the  $\text{Q}^+\text{ClO}_4^-$  ion pair in the chloroform phase and the perchlorate ion in the aqueous phase). The extraction constants are 12.15 for  $\text{Q}_2\text{dto}$ , 16.69 for  $\text{Q}_2[\text{Pd}(\text{dto})_2]$ , and 16.83 for  $\text{Q}_2[\text{Pt}(\text{dto})_2]$  at ionic strength 0.3 ( $\text{Na}_2\text{SO}_4$ ) and 25°C. Possible interfering metal ions can be masked by use of ethylenediaminetetraacetic acid [134].

(iv) *Trithio-oxalate*  $[\text{SOC-CS}_2]^{2-}$ , *trto*

Trithio-oxalate has the inverted O/S topology of monothio-oxalate. Therefore, in contrast to *mto*, the soft parts of the ligand should govern the coordination modes.

Potassium trithio-oxalate seems to be the most unstable mixed-chalcogen potassium thio-oxalate. Although the synthesis of potassium trithio-oxalate, its vibrational spectra and the X-ray structure of  $K_2trto$  KCl had already been published by Mattes and co-workers by the middle seventies [9,15,43], the coordination chemistry of  $trto$  was started only in 1988 [19,135], when the first bi-, tri- and tetranuclear complexes were isolated with trithio-oxalate as bridging ligand (see Table 13).

The reaction of trithio-oxalate (potassium,  $Ph_4As^+$  or  $BzPh_3P^+$  salt) with coordinatively unsaturated species such as  $(Ph_3P)_2M^+$  ( $M = Cu, Ag, Rh$ ) results in the formation of binuclear homometal compounds of the type  $\{[(Ph_3P)_2M]_2(trto)\}$ . Tri- or tetranuclear heterometal complexes of the general formula  $\{[(Ph_3P)_2Cu(trto)]_nM\}$  can be obtained by reaction of trithio-oxalate with the corresponding metal salt followed by extraction of the resulting complex solution with  $(Ph_3P)_3CuCl$  in dichloromethane. The formal composition does not differ from that of compounds containing monothio-, 1,2-dithio- or tetrathio-oxalate as bridging ligands linking identical or different metal ions. The IR spectra, however, are more complicated. A band at  $1530\text{ cm}^{-1}$  (potassium salt [43]) or  $1520\text{ cm}^{-1}$  ( $Ph_4As^+$  salt [135]) has been assigned to the C–O stretching vibration. This absorption was found, e.g., in  $\{[(Ph_3P)_2Ag]_2(trto)\}$  at a higher frequency ( $1563\text{ cm}^{-1}$ ), unexpected for a bridging ligand. Surprisingly, the homologous copper complex shows a bathochromically shifted C–O stretching mode ( $1475\text{ cm}^{-1}$ ).

Until now only the X-ray structure of the binuclear silver complex is known (see Fig. 21) [19(d),(e)], exhibiting an unexpected result: whereas one silver ion is

TABLE 13

Trithio-oxalate complexes [19(b),135]

Compound	M.p. (°C)	Colour (solid state)	IR spectra $\bar{\nu}_{C-O}$ ( $\text{cm}^{-1}$ )	Electronic spectra $\bar{\nu}_{\max}$ ( $10^3\text{ cm}^{-1}$ ) ( $\epsilon_0$ )
$\{[(Ph_3P)_2M]_2(trto)\}$				
$M = Cu$	206–8	Violet	1475s	13.5 (8700) 22.0 (2100)
Ag	167	Olive-green	1563s	21.0 (3300) 27.0 (2000)
Rh	162–4	Brown	1610br	
$\{M'[(trto)Cu(PPh_3)_2]_2\}$				
$M' = Cu$	125–7	Violet	1583s 1520s	13.0 (16200) 25.5 (300)
Ni	165	Green	1570s 1520s	13.2 (13500) 29.3 (23400)
Pd	160–2	Blue-violet	1580s 1528s	12.9 (12600) 23.8 (7800)
$\{Fe[(trto)Cu(PPh_3)_2]_3\}$	164–6	Violet	1600br	13.0 (22400) 27.0 (40700)




Fig. 21. Molecular structure of  $[(\text{Ph}_3\text{P})_2\text{Ag}(\text{trto})\text{Ag}(\text{PPh}_3)_2]$  (phenyl rings omitted) [19(e)].

bound “normally”, i.e. side-on, to two sulphur atoms of trto, the other is bound end-on to two sulphur atoms. The latter coordination mode has been detected for the first time for thio-oxalate ligands. The bridging trithio-oxalate ligand is far from being planar. The dihedral angle between the COS and the  $\text{CS}_2$  group is  $50(1)^\circ$ . The C–O bond length ( $1.25(2) \text{ \AA}$ ) is markedly longer than those of the non-coordinated C–O entities in monothio-oxalate complexes ( $1.09(1)–1.23 \text{ \AA}$ ) [21,36–38]. There are numerous complexes where sulphur ligand atoms of unsaturated dithio ligands are bound to more than one metal ion in oligonuclear species. Examples are  $\{[(\text{Ph}_3\text{P})_2\text{Ag}]_2(\text{mnt})\}$  [136,137],  $\text{Ni}(i\text{-mnt})_2[\text{Ag}(\text{PPh}_3)_2]_2$ ,  $\{\text{Ni}(\text{mnt})_2[\text{Ag}(\text{PPh}_3)_2]_2\}$  and  $[\text{Ni}(\text{mnt})_2[\text{Ag}(\text{PPh}_2\text{Et})_2]_2]$  [138,139] with Ag–S (mean  $2.808 \text{ \AA}$ ) and C–S distances (mean  $1.77 \text{ \AA}$ ) somewhat longer than those found for  $\{[(\text{Ph}_3\text{P})_2\text{Ag}]_2(\text{trto})\}$  (Ag–S, with S bound to two Ag:  $2.680(2)$  and  $2.725(2) \text{ \AA}$  and C–S  $1.68(1) \text{ \AA}$ ) (see also Table 15). On the other hand, the Ag–P bond lengths are almost identical (mean values  $2.474$  and  $2.477(2) \text{ \AA}$ , respectively).

The anion  $\text{MoOS}_3^{2-}$ , tetrahedrally configured but formally comparable with trto, forms a compound  $\{[(\text{Ph}_3\text{P})\text{Au}](\text{MoOS}_3)[\text{Au}(\text{PPh}_3)_2]\}$  with an  $(\text{MoOS}_3)$  bridge using also exclusively the three sulphur atoms, i.e. a similar arrangement to that found in  $\{[(\text{Ph}_3\text{P})_2\text{Ag}]_2(\text{trto})\}$  with non-coordinated oxygen [140].

One argument that the C=O of the trto ligand remains uncoordinated follows from the shape of the highest occupied molecular orbitals (HOMOs). MO calculations (ab initio self-consistent field (SCF)) for trithio- and tetrathio-oxalate [19(e)] show, in agreement with the extended Hückel calculations for the  $\text{C}_2\text{S}_4$  unit [141], four in-plane  $\sigma$  donor orbitals (see Fig. 22). In trto a group of three donor orbitals, mainly localized at the sulphur atoms, is significantly separated from one low lying orbital localized at the oxygen. This indicates that the donor ability via the sulphur atoms is dominant.




Fig. 22. In-plane  $\sigma$  donor orbitals of *i*-dto, trto and tto [19(e)].

(v) Tetrathio-oxalate  $[\text{S}_2\text{C}-\text{CS}_2]^{2-}$ , tto

The discovery of the first solely sulphur-compound-based superconductor bis(ethylenedithio)tetrathiafulvalene (BEDT-TTF or in short “ET”) and its radical cation salts has prompted several laboratories to start synthetic programmes for new extended multisulphur  $\pi$  donors and to investigate their inorganic complexes [142].

Obviously the  $\text{C}_2\text{S}_4$  subunits in TTF, BEDT-TTF and related compounds are fundamentally important for that class of “organic metals” and lead directly to the simple tetrathio-oxalate ligands. A rising interest in developing new transition metal tetrathio-oxalates and in understanding their structural and electronic relationship

is also motivated by their possible use in syntheses of extended low-dimensional metal tetrathiolene arrays with metallic or semiconducting properties [141].

Tetrathio-oxalate is unique among all thio-oxalates with respect to its highly symmetrical topology and, in connection with this, to the chemical consequences of its “all sulphur” constitution. All four sulphurs can serve as effective donor sites and the  $C_2S_4$  unit can formally or really accept or release electrons, and does so especially when switched in tune with the electronic system of a transition metal in an appropriate symmetry. It is also unique with respect to the manifold chemical individuals which contain the  $C_2S_4$  entity in the same topology as in the free ligand, (i.e.  $S_2C_2S_2$ ) regardless of its overall charge. Moreover, there is also a huge number of compounds which have the same net content ( $C_2S_4$ ) but different connectivity.

Examples of “non-tetrathio-oxalate  $C_2S_4$ ” are:

(1) Metal promoted head-to-tail dimerization of two coordinated  $CS_2$  molecules, see Fig. 23 [143–146], where subsequent splitting into coordinated “ $C_2S_3$ ” and S has been observed [147]. Dimerization of  $[Pt(\eta^2-CS_2)(dppe)]$  (dppe = 1,2-bis(diphenylphosphino)ethane) in the presence of excess methyl iodide also forms a dimethylated “ $C_2S_4$ ” unit [148]. Even pseudo-trimerization exists with dual head-to-tail linking of three  $CS_2$  molecules with central  $sp^3$  carbon [149], in contrast to the




Fig. 23. Selection of head-to-tail coupled “ $(CS_2)_2$ ” ligands.

formal trimerization of  $\text{CS}_2$  giving a C–C–C connection [150]. Head-to-tail linking at metal centres was also found for  $\text{CSSe}$  [151] and  $\text{CSe}_2$  by Werner and co-workers [151,152]. Tributylphosphine reacts with  $\text{CSe}_2$  yielding  $n\text{-Bu}_3\text{P}=\text{C}_3\text{Se}_4$  with a head-to-tail coupled  $\text{C}_2\text{Se}_4$  moiety [153].

(2) Linking of two metal ions by two non-coupled  $\text{CS}_2$  bridges [154,155]. Trimeric or polymeric species are postulated [156].

(a) *Metal-promoted head-to-head dimerizations of  $\text{CS}_2$*

The first authentic and structurally proven example of coordinated  $\text{C}_2\text{S}_4$  was realized by Maj et al. [157] (see Fig. 24). The condensation proceeded through activation of  $\text{CS}_2$  coordinated at electron rich nickel pentamethylcyclopentadienyl-dicarbonyl  $[\text{Cp}^*\text{Ni}(\text{CO})_2]_2$ , which leads to a side-on bridging  $\text{C}_2\text{S}_4$  linking with  $\text{Cp}^*\text{Ni}$  fragments. After reaction of  $\text{CS}_2$  with oligonuclear cluster carbonyls like  $\text{Fe}_2(\text{CO})_9$ , Broadhurst et al. identified, among many other products, a tetranuclear iron complex in low yield, in which the central  $\mu\text{-C}_2\text{S}_4$  unit links, end-on-like, two  $\text{Fe}_2$  fragments [158] (Fig. 25). A titanium(II)-activated carbon–carbon bond forma-




Fig. 24. Molecular structure of  $[(\eta^5\text{-Me}_5\text{C}_5)\text{Ni}(\text{C}_2\text{S}_4)\text{Ni}(\eta^5\text{-Me}_5\text{C}_5)]$  [157].




Fig. 25. Molecular structure of  $[(\text{CO})_6\text{Fe}_2(\text{C}_2\text{S}_4)\text{Fe}_2(\text{CO})_6]$  [158].



tion by reductive head-to-head coupling of  $\text{CS}_2$  is reported by Harris et al. [159]. Starting from  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO})_2]$  in  $\text{CS}_2$  solution, dark purple  $\{[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}]_2(\mu\text{-C}_2\text{S}_4)\}$  is formed at room temperature after about one day (see Fig. 26). The  $\text{C}_2\text{S}_4$  core is described by the authors as “tetrathio-oxalate (or ethylene tetrathiolate)”. C–C bond formation to a tetrathiolate moiety also occurs by reduction of cationic  $\eta^2$ -coordinated methylthioformate with borohydride [160,161]. The reaction yields the coordinated dimethyl ester of tetrathio-oxalate. The remarkably short C–C bond length of 1.390(15) Å shows, however, that the ligand coordinates as a 1,2-dimercaptoethylene derivative rather than as tetrathio-oxalate (see the following chapter). The formation of  $[(\text{triphos})\text{Rh}(\mu\text{-C}_2\text{S}_4)\text{Rh}(\text{triphos})](\text{BPh}_4)_2 \cdot \text{CH}_2\text{Cl}_2$  (triphos =  $\text{MeC}(\text{CH}_2\text{PPh}_2)_3$ ) by addition of  $\text{CS}_2$  to  $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$  and triphos in THF–benzene in the presence of  $\text{NaBPh}_4$ , is comparable with the previous reaction [162,163]. The  $\text{C}_2\text{S}_4$  unit is planar (see Fig. 27), but the  $\text{Rh}(\mu\text{-C}_2\text{S}_4)\text{Rh}$  core is not completely planar, each Rh atom being about 0.3 Å out of the  $\text{C}_2\text{S}_4$  plane. According




Fig. 26. Molecular structure of  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{C}_2\text{S}_4)\text{Ti}(\eta^5\text{-C}_5\text{H}_5)_2]$  [159].




Fig. 27. Structure of the cation in  $[(\text{triphos})\text{Rh}(\text{C}_2\text{S}_4)\text{Rh}(\text{triphos})](\text{BPh}_4)_2 \cdot \text{CH}_2\text{Cl}_2$  [162].

to chemical behaviour, these authors also claim the analogous ethenetetrathiolate-bridged binuclear Ir compound  $[(\text{triphos})\text{Ir}(\mu\text{-C}_2\text{S}_4)\text{Ir}(\text{triphos})]^{2+}$  [163].

Metal-promoted head-to-head coupling is also reported for  $\text{CSe}_2$ , from the same laboratory, yielding  $[(\text{triphos})\text{Rh}(\mu\text{-CSe}_2)_2\text{Rh}(\text{triphos})]^{2+}$  [163,164]. However, Jensen and Huge-Jensen in 1973 had already reported two compounds  $[(\text{Ph}_3\text{P})_2\text{Ni}(\text{CSe}_2)_2]$  and  $[(\text{Ph}_3\text{P})_2\text{RhCl}(\text{CSe}_2)_2]$  with possibly coupled  $\text{CSe}_2$  moieties [165]. The description of coordinated  $\text{C}_2\text{S}_4$  units is often not clear. Some so-called thio-oxalate compounds should rather be interpreted as ethenetetrathiolate (ett) derivatives. The C–C bond length is a suitable indicator as to whether there is oxalate or ethenetetrathiolate behaviour; see Tables 14 and 15.

(b) *Dimethyl tetrathio-oxalate,  $\text{ttoMe}_2$ , and dimethyl ethenetetrathiolate,  $\text{ettMe}_2$  complexes*

The first authentic tetrathio-oxalic acid derivative, namely the dimethyl ester, was isolated in 1976 by Kissel et al. by photolytic decarbonylation of the heterocycle 4,5-bis(methylthio)-1,3-dithiole-2-one [166] (see Fig. 28). Surprisingly, all attempts to prepare higher esters have so far failed [4]. Dimethyl tetrathio-oxalate reacts with electron rich metal centres forming neutral chelates, for example  $[\text{Ni}(\text{ettMe}_2)_2]$  or  $[\text{Mo}(\text{ettMe}_2)_3]$ , or mixed ligand chelates such as  $[\text{Pd}(\text{PPh}_3)_2(\text{ettMe}_2)]$  [167,168] (see Fig. 29). However, in all attempts to coordinate the  $\text{ttoMe}_2$  ligand, a thermal or photochemical redox reaction takes place, where the destiny of the ligand is to be transformed into an ethenetetrathiolate derivative ( $\text{ettMe}_2$ ) indicated by short C–C bonds with typical dithiolene behaviour (intense absorptions in low energy visible and near IR region, reversible one-electron transitions).

TABLE 14  
Coordination modes of  $\text{C}_2\text{S}_4$  species

Compound	Bond distance (Å)		Mode <sup>a</sup>	Ref.
	C–C	C–S		
1 $(\text{Ph}_4\text{P})_2(\text{C}_2\text{S}_4) \cdot 6\text{H}_2\text{O}$	1.46(2)	1.713(9) 1.691(10)	tto	10
2 $\{[(\text{CO})_6\text{Fe}_2]_2(\text{C}_2\text{S}_4)\}$	1.332	1.775	ett/end-on	158
3 $\{[\eta^5\text{-Cp}]\text{Ni}\}_2(\text{C}_2\text{S}_4)\}$	1.36(1)	1.718(2)	ett/side-on	157
4 $\{[(\text{triphos})\text{Rh}]_2(\text{C}_2\text{S}_4)\}^{2+}$	1.36(3)	1.76(2) 1.73(2)	ett/side-on	163
5 $\{[(\text{Ph}_3\text{P})_2\text{Cu}]_2(\text{C}_2\text{S}_4)\}$	1.53(8)	1.668(6) 1.689(6)	tto/side-on	188
6 $\{[(\text{C}_3\text{OS}_4)\text{Cu}]_2(\text{C}_2\text{S}_4)\}^{2-}$	1.467(10)	1.673(5) 1.678(5)	tto(?)/side-on	189(b)
7 $\{[(\eta^5\text{-Cp})\text{Ti}]_2(\text{C}_2\text{S}_4)\}$	1.41	1.736	ett/side-on	159

<sup>a</sup> tto = tetrathio-oxalate-like; ett = ethenetetrathiolate-like.

TABLE 15

Selected bond distances (Å) and angles (°) of *i*-dto, *trto* and *tto* complexes

Compound	Bond distances				Angles			Coord. mode	Ref.
					Intraligand				
	M-S	M-O	C-S	C-O	C-C	SMS SMO		Torsion	
1 [Ph <sub>3</sub> P] <sub>2</sub> Cu( <i>i</i> -dto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ] <sup>a</sup>	2.323(4)	2.05(1)	1.65(2)	1.28(2)	1.52(2)	85.2(3)	18.5(2)	Side-on	19(e)
	2.307(5)	2.105(9)		1.27(2)		84.3(3)			
2 [(Ph <sub>3</sub> P) <sub>2</sub> Ag(trto)Ag(PPh <sub>3</sub> ) <sub>2</sub> ]	2.680(2)		1.683(10)	1.25(2)	1.52(2)	81.82(9)	50(1)	Side-on/ end-on	19(e)
	2.545(3)		1.696(8)			66.87(8)			
	2.725(2) <sup>b</sup>		1.688(10)						
	2.645(3) <sup>b</sup>								
3 [(Ph <sub>3</sub> P) <sub>2</sub> Cu(tto)Cu(PPh <sub>3</sub> ) <sub>2</sub> ]	2.308(2)		1.668(6)		1.531(8)	89.18(7)	Planar	Side-on	188
	2.275(2)		1.689(6)						
4 [(C <sub>3</sub> OS <sub>4</sub> )Cu(tto)Cu(C <sub>3</sub> OS <sub>4</sub> )] <sup>2-</sup>	2.258(2)		1.673(5)		1.467(10)	89.0(1)	Planar	Side-on	189
	2.245(2)		1.678(5)						

<sup>a</sup>The analogous Ag complex  $[(\text{Ph}_3\text{P})_2\text{Ag}(i\text{-dto})\text{Ag}(\text{PPh}_3)_2]$  is isostructural.<sup>b</sup>End-on coordination.




Fig. 28. Photolytic synthesis of tetrathio-oxalate dimethyl ester [166].




Fig. 29. Molecular structure of  $[(PPh_3)_2Pd(ettMe_2)]$  (phenyl rings omitted) [168].

The synthesis of  $ettMe_2$  complexes from the components is more versatile, especially for mixed ligand chelates, than the in situ preparation starting with 4,5-bis(methylthio)-1,3-dithiole-2-one [169].

The monoanionic species with, formally, Ni(III) or Pd(III) can also be reached by hydrolytic cleavage of 4,5-bis(methylthio)-1,3-dithiole-2-one and subsequent reaction of the generated 1,2-bis(methylthio)ethene-1,2-dithiolate with an appropriate M(II) starting material (for example  $NiCl_2 \cdot 6H_2O$ ) [170]. The conversion of  $ttoMe_2$  into  $ettMe_2$  during coordination to metal centres is comparable with the formation of Diels–Alder-like [4 + 2]-cycloaddition products, where the  $ttoMe_2$  behaves as a dithiabutadiene [171].

(c) *Authentic tetrathio-oxalate — synthesis and complex behaviour*

There have been several claims for tetrathio-oxalate. The first goes back to 1927 [172]. 4,5-Dimethylthio-1,3-dithiole-2-thione ( $Me_2dmit$ ), which is formed by chemical reduction of  $CS_2$  and subsequent methylation, was wrongly seen as the dimethyl ester of tetrathio-oxalic acid because of similar elemental analyses. There is even a patent in 1973 which claims the preparation of tetrathio-oxalate [173]. The reduction was repeated using different reductants and media (e.g. electrochemically in DMF [174]). In all these cases, the dimethylthioether  $Me_2dmit$  was formed instead

of  $\text{ttoMe}_2$ . In 1981 Lodmell et al. proved by HPLC that  $\text{tto}$  is indeed formed during the electroreduction of  $\text{CS}_2$  [175]. The preparation is surprisingly simple: the important factors are the solvent, the  $\text{CS}_2$  concentration, the supporting electrolyte and the cathode material. Use acetonitrile instead of the commonly used DMF,  $\text{Et}_4\text{NBr}$  or  $\text{KI}$  instead of  $\text{Bu}_4\text{NX}$ , and a mercury pool electrode, and it can be prepared. After applying a potential slightly more negative than the reduction potential of  $\text{CS}_2$  ( $-1.15$  V vs. SCE) the corresponding  $\text{C}_2\text{S}_4^{2-}$  salt is formed and precipitates, thus possibly preventing undesired follow-up reactions. The  $\text{Et}_4\text{N}^+$  salt can be purified by reprecipitation from concentrated aqueous solution by acetone. The orange crystals can be stored for a long time if moisture is excluded [5,10,176]. For organic follow-up reactions see, e.g., [5]. The X-ray structure of  $(\text{Ph}_4\text{P})_2\text{C}_2\text{S}_4 \cdot 6\text{H}_2\text{O}$  (crystallized from water) shows that the dianion of  $\text{tto}$  is far from being planar [10]. The dihedral angle between the two  $\text{CS}_2$  units is  $79.5^\circ (\pm 1.0^\circ)$ . This is analogous to potassium 1,2-dithio-oxalate ( $76.5^\circ$ ). The C–C bond length ( $1.46$  Å) is noticeably shorter than in the other thio-oxalates and oxalate ( $1.55$ – $1.58$  Å) (see Table 1). Tetrathio-oxalate can act as a multisulphur bridging ligand with electron delocalization capacity. So far, five types of tetrathio-oxalate complexes are known (see Table 16).

**Polymeric complexes.** Deeply coloured precipitates are formed from aqueous solutions of  $(\text{Et}_4\text{N})_2\text{C}_2\text{S}_4$  after adding heavy metal ions. With  $\text{M}^{2+}$  ions, the materials obtained are insoluble in common solvents, non-melting up to  $300^\circ\text{C}$  and obviously polymeric. Their metal-to-ligand ratio roughly corresponds to a 1:1 composition [177–181]. Obtained from aqueous solutions [182], or from methanol or methanol–DMF solutions under anaerobic conditions [183,184], polymeric (or oligomeric) tetrathio-oxalates  $[\text{M}(\text{tto})]_n$  ( $\text{M} = \text{Cu}, \text{Ni}, \text{Pd}, \text{Pt}$ ) and  $\{\text{A}[\text{M}(\text{tto})]\}_n$  ( $\text{M} = \text{Cu(I)}$ ;  $\text{A} = \text{Na}^+, \text{Et}_4\text{N}^+$ ) display typical semiconducting behaviour with pressed-pellet conduc-

TABLE 16

Tetrathio-oxalate complexes

Compound	M.p. ( $^\circ\text{C}$ )	Colour	Electronic spectra $\bar{\nu}_{\text{max}}$ ( $10^{-3} \text{ cm}^{-1}$ ) ( $\epsilon_0$ )	Ref.
$[(\text{Ph}_3\text{P})_2\text{Cu}]_2(\text{tto})$	212–215	Blue	15.9 (11700)	177,178,188
$[(\text{Ph}_3\text{P})_2\text{Ag}]_2(\text{tto})$	115–118	Red–brown	20.8 (9000)	177,178
$[(\text{Ph}_3\text{P})_2\text{Au}]_2(\text{tto})$	157–160	Light brown		177,178
$[(\text{Ph}_3\text{P})_2\text{Rh}]_2(\text{tto})$	142–143	Dark brown		44
$\{\text{Cu}[(\text{tto})\text{Cu}(\text{PPh}_3)_2]_2\}$	184–187	Blue	15.5 (5500)	177,178,191
$\{\text{Ni}[(\text{tto})\text{Cu}(\text{PPh}_3)_2]_2\}$	162–165	Violet	18.7 (18600)	177,178,191
$\{\text{Zn}[(\text{tto})\text{Cu}(\text{PPh}_3)_2]_2\}$	181–184	Blue		177,178,191
$\{\text{Pd}[(\text{tto})\text{Cu}(\text{PPh}_3)_2]_2\}$	203–207	Black	25.1 (16800)	177,178,191
$\{\text{VO}[(\text{tto})\text{Cu}(\text{PPh}_3)_2]_2\}$	145–147	Blue–black		44
$\{\text{Ni}[(\text{tto})\text{Ag}(\text{PPh}_3)_2]_2\}$	125–128	Dark brown		190
$\{\text{Fe}[(\text{tto})\text{Cu}(\text{PPh}_3)_2]_3\}$	173–176	Blue	16.5 (15000)	177,178,191

tivities in the  $10^{-1}$ – $10^2 \Omega^{-1} \text{ cm}^{-1}$  range. Recently, Jolly and Reynolds published the preparation of the first semiconducting and optically polarizing poly(nickel tetrathio-oxalate)/poly(vinyl alcohol) composites made by the reaction of tetrathio-oxalate with nickel ions in a poly(vinyl alcohol) matrix [185]. Structural data on the  $[\text{M}(\text{tto})]_n$  powders have been obtained by the large angle X-ray scattering technique (LAXS) [184]. Polymeric ethenetetrathiolates have been prepared and found to be non-stoichiometric [186].

**Mononuclear complexes.** In the presence of a large excess of precipitating cation and with very slow addition of metal salt solution, mononuclear anionic complexes can be isolated [187].

**Binuclear complexes.** By treating aqueous solutions of tto with dichloromethane, solutions of  $(\text{Ph}_3\text{P})_3\text{MCl}$  ( $\text{M} = \text{Cu}, \text{Ag}, \text{Rh}$ ) deeply coloured tto-bridged binuclear metal(I) neutral chelates are formed in the organic phase, from which they can be isolated after concentration and subsequent addition of 2-propanol [177–180]. The X-ray structure of  $\{[(\text{Ph}_3\text{P})_2\text{Cu}]_2(\text{C}_2\text{S}_4)\}$  revealed the first authentic coordination compound containing a real tetrathio-oxalate ( $\text{C}-\text{C}$  bond length  $1.531(8) \text{ \AA}$ ) [188] (see Fig. 30).

Coordinatively halfside-saturated  $(\text{Ph}_3\text{P})_2\text{M}^+$  moieties act as “lids”, which prevent the formation of polymeric compounds. Starting from tetrathiapentalenedione (TPD), by incomplete basic hydrolysis in the presence of  $\text{Cu(II)}$  the compound  $(\text{Ph}_4\text{As})_2[(\text{C}_3\text{OS}_4)\text{Cu}(\mu\text{-C}_2\text{S}_4)\text{Cu}(\text{C}_3\text{OS}_4)]$  can be isolated (see Fig. 31). The  $\text{C}-\text{C}$  bond length of the bridging  $\text{C}_2\text{S}_4$  unit ( $1.467(10) \text{ \AA}$ ) [189] does not alone allow a clear decision as to whether the bridging ligand is tto- or ett-like: it is very close to




Fig. 30. Molecular structure of  $[(\text{Ph}_3\text{P})_2\text{Cu}(\text{C}_2\text{S}_4)\text{Cu}(\text{PPh}_3)_2]$  (phenyl rings omitted) [188].




Fig. 31. Structure of the anion in  $(\text{Ph}_4\text{As})_2[(\text{C}_3\text{OS}_4)\text{Cu}(\text{C}_2\text{S}_4)\text{Cu}(\text{C}_3\text{OS}_4)]$  [189(b)].

that in  $(\text{Ph}_4\text{P})_2(\text{tto})$  (1.46(2) Å) but markedly shorter than in  $\{[(\text{Ph}_3\text{P})_2\text{Cu}]_2(\text{tto})\}$  (1.53(8) Å) (see Table 15).

**Tri- and tetranuclear complexes.** Treating the mononuclear complexes with a  $\text{CH}_2\text{Cl}_2$  solution of  $(\text{Ph}_3\text{P})_3\text{CuCl}$  [178] or  $(\text{Ph}_3\text{P})_3\text{AgCl}$  [190], intensely coloured trinuclear complexes  $\{\text{M}^{\text{II}}[(\text{tto})\text{M}^{\text{I}}(\text{PPh}_3)_2]_2\}$  are formed. The reactive extraction of the wet insoluble material obtained by the reaction between tto and metal ions with solutions of  $(\text{Ph}_3\text{P})_3\text{MCl}$  (in, e.g.,  $\text{CH}_2\text{Cl}_2$ ) gives complexes identical to those from the previous procedure.  $(\text{Ph}_3\text{P})_2\text{M}^+$  cations probably act not only as “chain-stoppers” but also as “chain-breakers”.

Starting with trivalent metal ions (e.g. Fe(III)), tetranuclear complexes  $\{\text{M}^{\text{III}}[(\text{tto})\text{M}^{\text{I}}(\text{PPh}_3)_2]_3\}$  can also be isolated [178]. The tetrathiolate units  $\text{C}_2\text{S}_4$  (also  $\text{C}_2\text{Se}_4$ ) were thoroughly analyzed by several groups from a theoretical MO calculation point of view also including isolobal analogies for the extended system [141,163,164]. Tetrathio-oxalate complexes are intensely coloured with extinction coefficients of the order of  $10^4 \text{ mol}^{-1} \text{ cm}^{-1}$  in the visible region, independently of whether they are mono- or oligonuclear or which central metal ion they contain. Many are light-sensitive. Obviously the unit  $\text{L}_2\text{M}^{\text{I}}(\text{tto})$  is responsible for the transition. For instance,  $[(\text{PPh}_3)_2\text{Cu}^{\text{I}}(\text{C}_2\text{S}_4)\text{Cu}^{\text{I}}(\text{PPh}_3)_2]$  is deep blue with an absorption band at about  $1.5 \times 10^4 \text{ cm}^{-1}$ , which is typical of all tto systems with a terminating  $\text{L}_2\text{Cu}^+$  fragment linked to tto. The results of an extended Hückel calculation for the entity are shown in Fig. 32 [191]. The tto is linked to the  $\text{CuL}_2$  fragment via the  $a_1$  and  $b_2$  orbitals. Interaction of the occupied fragment orbital  $b_1(d_{xz})$  with the tto lowest unoccupied molecular orbital (LUMO) yields the bonding  $b_1$  MO with mainly metal character. The remaining four d orbitals of the metal interact only weakly with the occupied tto orbitals and are, therefore, only slightly destabilized. The typical absorption mentioned above is thus caused by a transition from the d block to the  $b_1$  LUMO and is of metal-to-tto charge-transfer type.

*(d) Related ligand systems containing the  $\text{C}_2\text{S}_4$  core*

Apart from tetrathio-oxalate and its dimethyl ester, several other sulphur ligands have a  $\text{C}_2\text{S}_4$  building block. The coordination chemistry of these ligand systems would require at least one review to itself. There are close relations as well




Fig. 32. Qualitative orbital interaction diagram of  $d^{10}$ - $ML_2$  fragments with  $C_2S_4^{2-}$  ( $C_{2v}$  symmetry) [191].

as significant differences in structure and topology between these “non-oxalate” systems and tetrathio-oxalate. Therefore some characteristic compounds should be mentioned, without any claim of completeness.

Ethenetetrathiolate (ett), the reduced tetrathio-oxalate, has a coordination chemistry very close to that of tetrathio-oxalate and its esters. Obviously, sharp differentiation between tto and ett is difficult. Most of the published complexes containing a bridging head-to-head coupled  $C_2S_4$  ligand are in fact ett complexes with a formal  $C_2S_4^{4-}$  moiety and an unambiguous C–C double bond (see Table 14) [157–164, 167–170, 186, 189, 192]. Lappert and co-workers [193, 194] and Abel et al. [195] reported a series of compounds with ethenetetrathiolate derivatives as ligands.

The dianions 1,4-dithiin-2,3-dithiolate (DDT) [196, 197] and 5,6-dihydro-1,4-dithiin-2,3-dithiolate (DDDT) [198–210], which are cyclic analogues of  $Me_2ett$ , and a follow-up product of tetrathio-oxalate during the reductive dimerization of excess  $CS_2$ , 1,3-dithiole-2-thione-4,5-dithiolate (dmit) [211], also contain a  $C_2S_4$  unit with a C–C double bond. All these ligands show typical dithiolene behaviour and an extensive coordination chemistry, and have attracted the attention of numerous laboratories by virtue of their solid state properties.

Tetrathiofulvalene (TTF), as part of the classic organic superconductor TTF–TCNQ, stimulated this development of chalcogen rich (S, Se, Te) systems, but its



ligand behaviour is far from that of tetrathio-oxalate (see, e.g., [212] and references therein).

Finally, the coordination chemistry of the thio-oxalates is related to that of thiosquarates [213–229] and tetrachalcogeno metalates [230–232], with a more or less rigid planar or tetrahedral arrangement, respectively, preventing the flexibility found for thio-oxalates in their complexes.

#### ACKNOWLEDGEMENTS

We thank Mrs. Hannelore Morgner, Mrs. Ingeborg Krüger and Dr. Rhett Kempe for their help in the preparation of the manuscript.

#### REFERENCES

- 1 H. Kolbe and E. Drechsel, *Justus Liebigs Ann. Chem.*, 146 (1868) 140.
- 2 H.O. Jones and H.S. Tasker, *J. Chem. Soc.*, 95 (1909) 1904.
- 3 E.G. Cox, W. Wardlaw and K.C. Webster, *J. Chem. Soc.*, (1935) 1475.
- 4 W. Thiel and R. Mayer, *Sulfur Rep.*, 8 (1988) 1.
- 5 P. Jeroschewski and P. Hansen, *Sulfur Rep.*, 7 (1986) 1.
- 6 D.J. Hodgson and J.A. Ibers, *Acta Crystallogr. Sect. B*, 25 (1969) 469; see also H. Küppers, *Acta Crystallogr. Sect. B*, 29 (1973) 318.
- 7 R. Mattes and W. Meschede, *Chem. Ber.*, 109 (1976) 1832.
- 8 R. Mattes, W. Meschede and W. Stork, *Chem. Ber.*, 108 (1975) 1.
- 9 W. Meschede and R. Mattes, *Chem. Ber.*, 109 (1976) 2510.
- 10 H. Lund, E. Hoyer and R. Grønbaek-Hazell, *Acta Chem. Scand. Ser. B*, 36 (1982) 207.
- 11 U. Niemer and R. Mattes, *Chem. Ber.*, 111 (1978) 2118.
- 12 R. Mattes, W. Meschede and U. Niemer, *Chem. Ber.*, 110 (1977) 2584.
- 13 C. Matz and R. Mattes, *Z. Naturforsch. Teil B*, 33 (1978) 461.
- 14 (a) R.G. Delaplane and J.A. Ibers, *Acta Crystallogr. Sect. B*, 25 (1969) 2423.  
(b) T.M. Sabine, G.W. Cox and B.M. Craven, *Acta Crystallogr. Sect. B*, 25 (1969) 2437.
- 15 W. Stork and R. Mattes, *Angew. Chem.*, 87 (1975) 452.
- 16 M. Leitheiser and D. Coucouvanis, *J. Inorg. Nucl. Chem.*, 39 (1977) 811.
- 17 R.G. Pearson and P. Ellgen, *Inorg. Chem.*, 6 (1967) 1379.
- 18 P. Jeroschewski, *Z. Chem.*, 21 (1981) 412.
- 19 (a) W. Dietzsch, A. Mewes and E. Hoyer, *Z. Chem.*, 28 (1988) 341.  
(b) W. Dietzsch, L. Golič, A. Mewes and E. Hoyer, *Proc. Conf. Coord. Chem.*, 12th, Smolenice, Bratislava, 1989, p. 85  
(c) R.-M. Olk, W. Dietzsch, L. Golič, R. Kirmse, J. Reinhold, J. Sieler, P. Strauch and E. Hoyer, *Proc. Int. Conf. Coord. Chem.*, 26th, Porto, 1988, Poster B127.  
(d) P. Strauch, W. Dietzsch, K. Trübenbach, L. Golič, A. Franke and E. Hoyer, *Proc. Int. Conf. Coord. Chem.*, 28th, 1990, Gera, P 6–30.  
(e) P. Strauch, W. Dietzsch, L. Golič, J. Sieler, A. Franke, I. Münzberg, K. Trübenbach, R. Kirmse, J. Reinhold and E. Hoyer, manuscript in preparation.  
(f) I. Münzberg, Diploma work, University of Leipzig, 1990.  
(g) P. Strauch, I. Münzberg, W. Dietzsch and R. Kirmse, *Z. Anorg. Allg. Chem.*, 613 (1992) 119.
- 20 A.E. Martell and R.J. Motekaitis, *J. Am. Chem. Soc.*, 110 (1988) 8059.

- 21 K. Mennemann and R. Mattes, *J. Chem. Res.*, Miniprint, (1979) 1343.
- 22 T. Shibahara, S. Ooi and H. Kuroya, *Bull. Chem. Soc. Jpn.*, 55 (1982) 3742.
- 23 G.J. Kruger, E.L.J. Breet and R. van Eldik, *Inorg. Chim. Acta*, 19 (1976) 151.
- 24 K. Zhou, J. Huang and J. Lu, *Jiegou Huaxue (Struct. Chem.)*, 2 (1983) 269.
- 25 H. Weber and R. Mattes, *Chem. Ber.*, 112 (1979) 95.
- 26 R. Mattes and H. Weber, *Z. Anorg. Allg. Chem.*, 474 (1981) 216.
- 27 D. Coucouvanis and D. Piltingsrud, *J. Am. Chem. Soc.*, 95 (1973) 5556.
- 28 D. Coucouvanis, N.C. Baenziger and S.M. Johnson, *J. Am. Chem. Soc.*, 95 (1973) 3875.
- 29 M. Leitheiser and D. Coucouvanis, *Inorg. Chem.*, 16 (1977) 1611.
- 30 R. Czernuszewicz, D.P. Strommen and K. Nakamoto, *Inorg. Chim. Acta*, 34 (1979) L211.
- 31 R.S. Czernuszewicz, K. Nakamoto and D.P. Strommen, *J. Am. Chem. Soc.*, 104 (1982) 1515.
- 32 F.J. Hollander and D. Coucouvanis, *Inorg. Chem.*, 13 (1974) 2381.
- 33 F.J. Hollander, M. Leitheiser and D. Coucouvanis, *Inorg. Chem.*, 16 (1977) 1615.
- 34 C.J. Weschler, J.C. Sullivan and E. Deutsch, *J. Am. Chem. Soc.*, 95 (1973) 2720.
- 35 C.J. Weschler, J.C. Sullivan and E. Deutsch, *Inorg. Chem.*, 13 (1974) 2360.
- 36 J.D. Lydon, K.J. Mulligan, R.C. Elder and E. Deutsch, *Inorg. Chem.*, 19 (1980) 2083.
- 37 G.J. Gainsford, W.G. Jackson and A.M. Sargeson, *Aust. J. Chem.*, 33 (1980) 707.
- 38 K. Mennemann and R. Mattes, *Angew. Chem.*, 89 (1977) 269.
- 39 K. Mennemann and R. Mattes, *J. Chem. Res.*, Miniprint, (1979) 1372.
- 40 Z. Chen, J.Z. Zhang, J.M. Shen and S.L. Ding, *Proc. Int. Conf. Coord. Chem.*, 25th, Nanjing, 1987, p. 338.
- 41 W. Dietzsch, N. Bulc, L. Golič and J. Šiftar, *Z. Chem.*, 21 (1981) 269.
- 42 N.B. Kalinichenko, L.E. Kordonskij, V.K. Belyaeva, I.N. Marov, E. Hoyer and W. Dietzsch, *Zh. Neorg. Khim.*, 35 (1990) 1237.
- 43 R. Mattes, W. Stork and J. Kahlenberg, *Spectrochim. Acta Part A*, 33 (1977) 643.
- 44 A. Müller, Diploma work, University of Leipzig, 1985.
- 45 L. Golič, N. Bulc and W. Dietzsch, *Polyhedron*, 2 (1983) 1201.
- 46 (a) W.C. Trogler, in A.B.P. Lever (Ed.), *Excited States and Reactive Intermediates — Photochemistry, Photophysics and Electrochemistry*, ACS Symp. Ser. 307, Washington, DC, 1986, pp. 177–196.  
(b) R.L. Cowan, D.B. Pourreau, A.L. Rheingold, S.J. Geib and W.C. Trogler, *Inorg. Chem.*, 26 (1987) 259.
- 47 D. Coucouvanis, in A. Müller and E. Diemann (Eds.), *Transition Met. Chem.*, Proc. Workshop, 1980, Verlag Chemie, Weinheim, 1981, pp. 59–87.
- 48 G. Kiel, M. Dräger and U. Reuter, *Chem. Ber.*, 107 (1974) 1483.
- 49 M.A. Pellinghelli, A. Tiripicchio and M. Tiripicchio Camellini, *Cryst. Struct. Commun.*, 3 (1974) 159.
- 50 L. Golič, N. Bulc and W. Dietzsch, *Inorg. Chem.*, 21 (1982) 3560.
- 51 T. Imamura, M. Ryan, G. Gordon and D. Coucouvanis, *J. Am. Chem. Soc.*, 106 (1984) 984.
- 52 M.G. Kanatzidis, N.C. Baenziger and D. Coucouvanis, *Inorg. Chem.*, 24 (1985) 2680.
- 53 A. Gleizes and M. Verdager, *J. Am. Chem. Soc.*, 106 (1984) 3727.
- 54 L. Golič, N. Bulc and W. Dietzsch, *Acta Crystallogr. Sect. C*, 42 (1986) 811.
- 55 S.F. Colmanet and M.F. Mackay, *Inorg. Chim. Acta*, 147 (1988) 173.
- 56 U. Abram, R. Münze, R. Kirmse, K. Köhler, W. Dietzsch and L. Golič, *Inorg. Chim. Acta*, 169 (1990) 49.
- 57 L. Qian, P. Singh, H.K. Ro and W.E. Hatfield, *Inorg. Chem.*, 29 (1990) 761.
- 58 K.R. Butler and M.R. Snow, *Inorg. Nucl. Chem. Lett.*, 8 (1972) 541; *Acta Crystallogr. Sect. B*, 31 (1975) 354.

- 59 F. Maury and A. Gleizes, *Inorg. Chim. Acta*, 41 (1980) 185.
- 60 A. Gleizes, F. Clery, M.F. Bruniquel and P. Cassoux, *Inorg. Chim. Acta*, 37 (1979) 19.
- 61 A. Gleizes, F. Maury and J. Galy, *Nouv. J. Chim.*, 8 (1984) 521.
- 62 A. Gleizes, F. Maury, P. Cassoux and J. Galy, *Z. Kristallogr.*, 155 (1981) 293.
- 63 J.C. Trombe, A. Gleizes and J. Galy, *C. R. Acad. Sci.*, 294–II (1982) 1369.
- 64 J.C. Trombe, A. Gleizes and J. Galy, *Inorg. Chim. Acta*, 87 (1984) 129.
- 65 C. Frasse, J.C. Trombe, A. Gleizes and J. Galy, *C. R. Acad. Sci.*, 300–II (1985) 403.
- 66 J.C. Trombe, C. Frasse and A. Gleizes, *C. R. Acad. Sci.*, 301–II (1985) 483.
- 67 J.C. Trombe, A. Gleizes and J. Galy, *C. R. Acad. Sci.*, 302–II (1986) 21.
- 68 J.C. Trombe, A. Gleizes, F. Dahan and J. Galy, *C. R. Acad. Sci.*, 303–II (1986) 907.
- 69 J.C. Trombe, A. Gleizes and J. Galy, *Nouv. J. Chim.*, 9 (1985) 55; see also GIT Fachz. Lab., 31 (1987) 89.
- 70 A. Gleizes and M. Verdaguer, *J. Am. Chem. Soc.*, 103 (1981) 7373.
- 71 P.D.W. Boyd, J. Hope, C.L. Raston and A.H. White, *Aust. J. Chem.*, 43 (1990) 601.
- 72 F. Maury, A. Gleizes and J. Galy, *C. R. Acad. Sci., Ser. C*, 291 (1980) 49.
- 73 R. Enjalbert, A. Gleizes, J.C. Trombe, M.L. Gutiérrez-Campo and P. Román, *J. Mol. Struct.*, 131 (1985) 1.
- 74 (a) P. Román, C. Bao, J.M. Gutiérrez-Zorilla and A. Vegas, *J. Crystallogr. Spectrosc. Res.*, 18 (1988) 207.  
(b) P. Román, J.M. Gutiérrez-Zorilla, A. Luque, J.I. Beitia, C. Guzmán-Miralles and M. Martínez-Ripoll, *Acta Crystallogr. Sect. C*, 47 (1991) 48.
- 75 J.C. Trombe, A. Gleizes and J. Galy, *C. R. Acad. Sci.*, 300-II (1985) 5.
- 76 J.M. Gutiérrez-Zorilla López, P. Román and A. Gleizes, *C. R. Acad. Sci.*, 305-II (1987) 1353.
- 77 (a) C. Bellitto, M. Bonamico, V. Fares, P. Imperatori and S. Patrizio, *J. Chem. Soc. Dalton Trans.*, (1989) 719.  
(b) P. Delhaes and M. Drillon (Eds.), *Organic and Inorganic Low-Dimensional Crystalline Materials*, NATO Adv. Study Inst. Ser. Ser. B, Vol. 168, Plenum, New York, 1987, p. 337.
- 78 M. Verdaguer, A. Gleizes, J.P. Renard and J. Seiden, *Phys. Rev. B*, 29 (1984) 5144.
- 79 J.D. Forrester, A. Zalkin and D.H. Templeton, *Inorg. Chem.*, 3 (1964) 1507.
- 80 D. Coucouvanis, F.J. Hollander and M.L. Caffery, *Inorg. Chem.*, 15 (1976) 1853.
- 81 R. Eisenberg, *Prog. Inorg. Chem.*, 12 (1970) 295.
- 82 C. Mahadevan, *J. Crystallogr. Spectrosc. Res.*, 16 (1986) 347.
- 83 O. Kahn, *Proc. Indian Acad. Sci. Chem. Sci.*, 98 (1987) 33.
- 84 C.S. Robinson and H.O. Jones, *J. Chem. Soc.*, 101 (1912) 62.
- 85 L.K. White and R.L. Belford, *J. Am. Chem. Soc.*, 98 (1976) 4428.
- 86 R.G. Pearson and D.A. Sweigart, *Inorg. Chem.*, 9 (1970) 1167.
- 87 B. Folkesson and B. Jonson, *J. Electron Spectrosc. Relat. Phenom.*, 34 (1984) 319.
- 88 D. Coucouvanis, *J. Am. Chem. Soc.*, 93 (1971) 1786.
- 89 A.R. Latham, V.C. Hascall and H.B. Gray, *Inorg. Chem.*, 4 (1965) 788.
- 90 R. Kirmse, W. Dietzsch, J. Stach, R.-M. Olk and E. Hoyer, *Z. Anorg. Allg. Chem.*, 548 (1987) 133.
- 91 R.L. Carlin and F. Canziani, *J. Chem. Phys.*, 40 (1964) 371.
- 92 F.P. Dwyer and A.M. Sargeson, *J. Am. Chem. Soc.*, 81 (1959) 2335.
- 93 J. Hidaka and B.E. Douglas, *Inorg. Chem.*, 3 (1964) 1724.
- 94 (a) D. Coucouvanis, A. Toupadakis and A. Hadjikyriacou, *Inorg. Chem.*, 27 (1988) 3272.  
(b) D. Coucouvanis, A. Toupadakis, Sang-Man Koo and A. Hadjikyriacou, *Polyhedron*, 8 (1989) 1705.
- 95 A. Davison, C. Orvig, H.S. Trop, M. Sohn, B.V. DePamphilis and A.G. Jones, *Inorg. Chem.*, 19 (1980) 1988.

- 96 D. Coucouvanis, R.E. Coffman and D. Piltingsrud, *J. Am. Chem. Soc.*, 92 (1970) 5004.
- 97 T. Birchall and K.M. Tun, *Inorg. Chem.*, 15 (1976) 376.
- 98 D. Niarchos, A. Kostikas, A. Simopoulos, D. Coucouvanis, D. Piltingsrud and R.E. Coffman, *J. Chem. Phys.*, 69 (1978) 4411.
- 99 W.V. Sweeney and R.E. Coffman, *J. Phys. Chem.*, 76 (1972) 49.
- 100 R.G. Kooser, *J. Phys. Chem.*, 80 (1976) 1601.
- 101 W.V. Sweeney, D. Coucouvanis and R.E. Coffman, *J. Chem. Phys.*, 59 (1973) 369.
- 102 R.E. DeSimone, *J. Am. Chem. Soc.*, 95 (1973) 6238.
- 103 K. Kanamori, A. Ono, S. Ino, I. Kanesaka, K. Kawai and J. Hidaka, *Bull. Chem. Soc. Jpn.*, 62 (1989) 2720.
- 104 H. Hennig, W. Kumpf, K. Jurdeczka and R. Benedix, *J. Prakt. Chem.*, 319 (1977) 444.
- 105 S. Kida, K. Nakamoto and R. Tsuchida, *Nippon Kagaku Zasshi*, 72 (1951) 749.
- 106 J. Fujita and K. Nakamoto, *Bull. Chem. Soc. Jpn.*, 37 (1964) 528.
- 107 G.A. Bowmaker, P.D.W. Boyd and G.K. Campbell, *Inorg. Chem.*, 21 (1982) 3565.
- 108 D. Coucouvanis, *J. Am. Chem. Soc.*, 92 (1970) 707.
- 109 G.A. Bowmaker, P.D.W. Boyd and G.K. Campbell, *Inorg. Chem.*, 21 (1982) 2403.
- 110 S. Kida, *Bull. Chem. Soc. Jpn.*, 33 (1960) 1204.
- 111 N. Maki, S. Kida and R. Tsuchida, *Bull. Chem. Soc. Jpn.*, 32 (1959) 573.
- 112 R.G. Pearson and M.J. Hynes, *J. Coord. Chem.*, 1 (1971) 245.
- 113 S.A. Cotton and J.F. Gibson, *J. Chem. Soc. A*, (1971) 803.
- 114 (a) A.R. Butler, C. Glidewell and M.-H. Li, *Adv. Inorg. Chem.*, 32 (1988) 335.  
(b) J.A. McCleverty, N.M. Atherton, J. Locke, E.J. Wharton and C.J. Winscom, *J. Am. Chem. Soc.*, 89 (1967) 6082.  
(c) G.R. Davies, J.A.J. Jarvis, B.T. Kilbourn, R.H.B. Mais and P.G. Owston, *J. Chem. Soc. A*, (1970) 1275.
- 115 J. Stach, R. Kirmse, W. Dietzsch, R.-M. Olk, L. Golič and N. Bulc, manuscript in preparation.
- 116 J. Stach, R. Kirmse, A. Heinrich, W. Dietzsch, J. Hartung and L. Beyer, *Z. Chem.*, 23 (1983) 453.
- 117 S. Sakaguchi and A.W. Addison, *J. Am. Chem. Soc.*, 99 (1977) 5189.
- 118 W. Dietzsch, J. Reinhold, R. Kirmse, E. Hoyer, I.N. Marov and V.K. Belyaeva, *J. Inorg. Nucl. Chem.*, 39 (1977) 1377.
- 119 W.A. Deskin, *J. Am. Chem. Soc.*, 80 (1958) 5680.
- 120 C.K. Jørgensen, *J. Inorg. Nucl. Chem.*, 24 (1962) 1571.
- 121 T.N. Lockyer and R.L. Martin, *Prog. Inorg. Chem.*, 27 (1980) 223.
- 122 A.J. McCaffery, S.F. Mason and R.E. Ballard, *J. Chem. Soc.*, (1965) 2883.
- 123 J.A. McCleverty, *Prog. Inorg. Chem.*, 10 (1968) 49.
- 124 E. Hoyer, W. Dietzsch and W. Schroth, *Z. Chem.*, 11 (1971) 41.
- 125 F.N. LeCrone, M.J. Minot and J.H. Perlstein, *Inorg. Nucl. Chem. Lett.*, 8 (1972) 173.
- 126 T.W. Thomas, C. Hsu, M.M. Labes, P.S. Gomm, A.E. Underhill and D.M. Watkins, *J. Chem. Soc. Dalton Trans.*, (1972) 2050.
- 127 Agency of Industrial Sciences and Technology, *Jpn. Kokai Tokkyo Koho*, JP 57 154 239 [82 154 239]; *Chem. Abstr.*, 99 (1983) 203 629d.
- 128 Agency of Industrial Sciences and Technology, *Jpn. Kokai Tokkyo Koho*, JP 58 200 234 [83 200 234]; *Chem. Abstr.*, 100 (1984) 148 635q.
- 129 (a) F. Basolo, *Adv. Chem. Ser.*, 49 (1965) 81.  
(b) R.G. Pearson and M.J. Hynes, *K. Tek. Hoegsk. Handl.*, (248–296) (1972) 459.
- 130 L.T. Fairhall, *J. Ind. Hyg.*, 8 (1926) 528.
- 131 J.H. Yoe and F.H. Wirsing, *J. Am. Chem. Soc.*, 54 (1932) 1866.

- 132 F.J. Welcher, *Organic Analytical Reagents*, Vol. IV, Van Nostrand, New York, 1955, p. 142.
- 133 H. Matsunaga and T. Yotsuyanagi, *Nippon Kagaku Kaishi*, (1982) 785.
- 134 H. Matsunaga and T. Yotsuyanagi, *Bunseki Kagaku*, 31 (1982) 289.
- 135 A. Mewes, Diploma work, University of Leipzig, 1988.
- 136 F. Estevan, P. Lahuerta and I. Solana, *Polyhedron*, 4 (1985) 1965.
- 137 D.D. Heinrich, J.P. Fackler, Jr. and P. Lahuerta, *Inorg. Chim. Acta*, 116 (1986) 15.
- 138 D. Coucouvanis, N.C. Baenziger and S.M. Johnson, *Inorg. Chem.*, 13 (1974) 1191.
- 139 F.J. Hollander, Y.L. Ip and D. Coucouvanis, *Inorg. Chem.*, 15 (1976) 2230.
- 140 J.M. Charnock, S. Bristow, J.R. Nicholson, C.D. Garner and W. Clegg, *J. Chem. Soc. Dalton Trans.*, (1987) 303.
- 141 S. Alvarez, R. Vicente and R. Hoffmann, *J. Am. Chem. Soc.*, 107 (1985) 6253.
- 142 (a) J.M. Williams, H.H. Wang, T.J. Emge, U. Geiser, M.A. Beno, P.C.W. Leung, K.D. Carlson, R.J. Thorn, A.J. Schultz and M.-H. Whangbo, *Prog. Inorg. Chem.*, 35 (1987) 51.  
(b) P. Cassoux, L. Valade, H. Kobayashi, A. Kobayashi, R.A. Clark and A.E. Underhill, *Coord. Chem. Rev.*, 110 (1991) 115.
- 143 H. Werner, O. Kolb, R. Feser and U. Schubert, *J. Organomet. Chem.*, 191 (1980) 283.
- 144 M. Cowie and S.K. Dwight, *J. Organomet. Chem.*, 214 (1981) 233.
- 145 (a) J.A. Ibers, *Chem. Soc. Rev.*, 11 (1982) 57.  
(b) M.G. Mason, P.N. Swepston and J.A. Ibers, *Inorg. Chem.*, 22 (1983) 411.
- 146 (a) E. Carmona, *J. Organomet. Chem.*, 358 (1988) 283.  
(b) E. Carmona, J. Cámpora, M.A. Muñoz, M. Paneque and M.L. Poveda, *Pure Appl. Chem.*, 61 (1989) 1701.  
(c) E. Carmona, A. Galindo, A. Monge, M.A. Muñoz, M.L. Poveda and C. Ruiz, *Inorg. Chem.*, 29 (1990) 5074.
- 147 S.J.N. Burgmayer and J.L. Templeton, *Inorg. Chem.*, 24 (1985) 3939.
- 148 S. Myrvold, O.A. Nassif, G. Semelhago, A. Walker and D.H. Farrar, *Inorg. Chim. Acta*, 117 (1986) 17.
- 149 C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *J. Chem. Soc. Chem. Commun.*, (1983) 545.
- 150 (a) H.G. Raubenheimer, J.C. Viljoen, S. Lotz, A. Lombard and G. Kruger, *J. Chem. Soc. Chem. Commun.*, (1981) 749.  
(b) H.G. Raubenheimer, S. Lotz, G.J. Kruger, A. van A. Lombard and J.C. Viljoen, *J. Organomet. Chem.*, 336 (1987) 349.
- 151 O. Kolb and H. Werner, *J. Organomet. Chem.*, 268 (1984) 49.
- 152 M. Ebner and H. Werner, *Chem. Ber.*, 119 (1986) 482.
- 153 H.P. Fritz, G. Müller, G. Reber and M. Weis, *Angew. Chem.*, 97 (1985) 1057.
- 154 C. Bianchini, C.A. Ghilardi, A. Meli, S. Midollini and A. Orlandini, *J. Chem. Soc. Chem. Commun.*, (1983) 753.
- 155 D.H. Farrar, R.R. Gukathasan and S.A. Morris, *Inorg. Chem.*, 23 (1984) 3258.
- 156 M.C. Zonnevylle, R. Hoffmann, C. Mealli and C. Bianchini, *Gazz. Chim. Ital.*, 117 (1987) 117.
- 157 J.J. Maj, A.D. Rae and L.F. Dahl, *J. Am. Chem. Soc.*, 104 (1982) 4278.
- 158 P.V. Broadhurst, B.F.G. Johnson, J. Lewis and P.R. Raithby, *J. Chem. Soc. Chem. Commun.*, (1982) 140.
- 159 H.A. Harris, A.D. Rae and L.F. Dahl, *J. Am. Chem. Soc.*, 109 (1987) 4739.
- 160 D. Touchard, J.-L. Fillaut, P. Dixneuf, C. Mealli, M. Sabat and L. Toupet, *Organometallics*, 4 (1985) 1684.
- 161 D. Touchard, J.-L. Fillaut, D.V. Khasnis, P. Dixneuf, C. Mealli, D. Masi and L. Toupet, *Organometallics*, 7 (1988) 67.

- 162 C. Bianchini, C. Mealli, A. Meli and M. Sabat, *Inorg. Chem.*, 23 (1984) 4125.
- 163 C. Bianchini, C. Mealli, A. Meli, M. Sabat and P. Zanello, *J. Am. Chem. Soc.*, 109 (1987) 185.
- 164 C. Bianchini, C. Mealli, A. Meli and M. Sabat, *J. Chem. Soc. Chem. Commun.*, (1984) 1647.
- 165 K.A. Jensen and E. Høge-Jensen, *Acta Chem. Scand.*, 27 (1973) 3605.
- 166 (a) T. Kissel, R. Matusch and K. Hartke, *Z. Chem.*, 16 (1976) 318.  
(b) T. Kissel, Ph.D Thesis, Marburg University, 1976.
- 167 C. Keller, D. Walther, J. Reinhold and E. Hoyer, *Z. Chem.*, 28 (1988) 410.
- 168 C. Kraffert, D. Walther, K. Peters, O. Lindqvist, V. Langer, J. Sieler, J. Reinhold and E. Hoyer, *Z. Anorg. Allg. Chem.*, 588 (1990) 167.
- 169 E. Fanghänel and H. Poleschner, *J. Prakt. Chem.*, 323 (1981) 1.
- 170 C.T. Vance and R.D. Bereman, *Inorg. Chim. Acta*, 149 (1988) 229.
- 171 (a) K. Hartke, T. Kissel, J. Quante and G. Henssen, *Angew. Chem.*, 90 (1978) 1016.  
(b) K. Hartke, J. Quante and T. Kämpchen, *Justus Liebigs Ann. Chem.*, (1980) 1482.  
(c) K. Hartke, T. Kissel, J. Quante and R. Matusch, *Chem. Ber.*, 113 (1980) 1898.
- 172 B. Fetkenheuer, H. Fetkenheuer and H. Lecus, *Chem. Ber.*, 60 (1927) 2528.
- 173 H.D. Hartzler, U.S. Pat. 3,717,619; *Chem. Abstr.*, 78 (1973) 137 673b.
- 174 S. Wawzonek and S.M. Heilmann, *J. Org. Chem.*, 39 (1974) 511.
- 175 J.C. Lodmell, W.C. Anderson, M.F. Hurley and J.Q. Chambers, *Anal. Chim. Acta*, 129 (1981) 49.
- 176 P. Jeroschewski, *Z. Chem.*, 21 (1981) 412.
- 177 P. Strauch, Diploma work, Leipzig University, 1983.
- 178 P. Strauch, W. Dietzsch and E. Hoyer, *Z. Chem.*, 23 (1983) 448.
- 179 E. Hoyer, *Comments Inorg. Chem.*, 2 (1983) 261.
- 180 E. Hoyer, R.-M. Olk and P. Strauch, in J.J. Ziolkowski (Ed.), *Coordination Chemistry and Catalysis*, World Scientific, Singapore, 1988, pp. 337–368.
- 181 J.R. Reynolds, F.E. Karasz, C.P. Lillya and J.C.W. Chien, *J. Chem. Soc. Chem. Commun.*, (1985) 268.
- 182 P. Jeroschewski, D.Sc. Thesis, Rostock University, 1984.
- 183 J.R. Reynolds, J.C.W. Chien and C.P. Lillya, *Macromolecules*, 20 (1987) 1184.
- 184 (a) T. Vogt, C. Faulmann, R. Soules, P. Lecante, A. Mosset, P. Castan, P. Cassoux and J. Galy, *J. Am. Chem. Soc.*, 110 (1988) 1833.  
(b) C.A. Jolly, F. Wang, S. Krichene, J.R. Reynolds, P. Cassoux and C. Faulmann, *Synth. Met.*, 29 (1989) F189.  
(c) J.R. Reynolds, C.A. Jolly, S. Krichene, P. Cassoux and C. Faulmann, *Synth. Met.*, 31 (1989) 109.
- 185 C.A. Jolly and J.R. Reynolds, *Chem. Mater.*, 2 (1990) 479.
- 186 H. Polescher, W. John, F. Hoppe, E. Fanghänel and S. Roth, *J. Prakt. Chem.*, 325 (1983) 957.
- 187 P. Strauch, M. Frieese and E. Hoyer, *Ger. Pat. DD249 908*; *Chem. Abstr.*, 108 (1988) 230989g.
- 188 L.K. Hansen, J. Sieler, P. Strauch, W. Dietzsch and E. Hoyer, *Acta Chem. Scand. Ser. A*, 39 (1985) 571.
- 189 (a) J. Ribas, R. Vicente, X. Solans and M. Verdaguer, *Proc. Int. Conf. Coord. Chem.*, 24th, Athens, 1986, p. 436.  
(b) R. Vicente, J. Ribas, S. Alvarez, A. Segui, X. Solans and M. Verdaguer, *Inorg. Chem.*, 26 (1987) 4004.
- 190 P. Strauch, M. Frieese, E. Hoyer, *Ger. Pat. DD236 089*; *Chem. Abstr.*, 106 (1987) 20655c.

- 191 J. Reinhold, G. Stich, P. Strauch, R. Benedix, J. Sieler and E. Hoyer, *Z. Chem.*, 27 (1987) 29.
- 192 S. Alvarez, M. Julve and M. Verdaguer, *Inorg. Chem.*, 29 (1990) 4500.
- 193 (a) B. Cetinkaya, P.B. Hitchcock, M.F. Lappert and P.L. Pye, *J. Chem. Soc. Chem. Commun.*, (1975) 683.  
(b) B. Cetinkaya, P.B. Hitchcock, M.F. Lappert, P.L. Pye and D.B. Shaw, *J. Chem. Soc. Dalton Trans.*, (1979) 434.
- 194 M.F. Lappert, D.B. Shaw and G.M. McLaughlin, *J. Chem. Soc. Dalton Trans.*, (1979) 427.
- 195 E.W. Abel, K. Kite and B.L. Williams, *J. Chem. Soc. Dalton Trans.*, (1983) 1017.
- 196 T. Nakamura, T. Nogami and Y. Shirota, *Bull. Chem. Soc. Jpn.*, 60 (1987) 3447.
- 197 H. Kim, A. Kobayashi, Y. Sasaki, R. Kato, H. Kobayashi, T. Nakamura, T. Nogami and Y. Shirota, *Bull. Chem. Soc. Jpn.*, 61 (1988) 2559.
- 198 C.T. Vance, R.D. Bereman, J. Bordner, W.E. Hatfield and J.H. Helms, *Inorg. Chem.*, 24 (1985) 2905.
- 199 H.J. Keller, R. Niebl, D. Nöthe, D. Schweitzer and A. Weber, *Physica B*, 143 (1986) 301.
- 200 R. Kato, H. Kobayashi, A. Kobayashi and Y. Sasaki, *Bull. Chem. Soc. Jpn.*, 59 (1986) 627.
- 201 H. Kim, A. Kobayashi, Y. Sasaki, R. Kato and H. Kobayashi, *Bull. Chem. Soc. Jpn.*, 61 (1988) 579.
- 202 A.J. Schultz, H.H. Wang, L.C. Soderholm, T.L. Sifter, J.M. Williams, K. Bechgaard and M.-H. Wangbo, *Inorg. Chem.*, 26 (1987) 3757.
- 203 U. Geiser, A.J. Schultz, H.H. Wang, M.A. Beno and J.M. Williams, *Acta Crystallogr. Sect. C*, 44 (1988) 259.
- 204 C.T. Vance and R.D. Bereman, *Inorg. Chim. Acta*, 149 (1988) 229.
- 205 J.H. Welch, R.D. Bereman and P. Singh, *Inorg. Chem.*, 27 (1988) 3680.
- 206 J.H. Welch, R.D. Bereman, P. Singh and C. Moreland, *Inorg. Chim. Acta*, 158 (1989) 17.
- 207 J.H. Welch, R.D. Bereman, P. Singh, D. Haase, W.E. Hatfield and M.L. Kirk, *Inorg. Chim. Acta*, 162 (1989) 89.
- 208 J.H. Welch, R.D. Bereman and P. Singh, *Inorg. Chim. Acta*, 163 (1989) 93.
- 209 C.T. Vance, J.H. Welch and R.D. Bereman, *Inorg. Chim. Acta*, 164 (1989) 191.
- 210 (a) S.S. Nagapetyan, V.E. Shklover, L.V. Vetoshkina, A.I. Kotov, L.Yu. Ukhin, Yu.T. Struchkov and E.B. Yagubskii, *Mater. Sci.*, 14 (1988) 5.  
(b) E.B. Yagubskii, A.I. Kotov, L.I. Buravov, A.G. Khomenko, V.E. Shklover, S.S. Naha-petyan, Yu.T. Struchkov, L.V. Vetoshkina and L.Yu. Ukhin, *Synth. Met.*, 35 (1990) 271.
- 211 R.-M. Olk, B. Olk, W. Dietzsch, R. Kirmse and E. Hoyer, *Coord. Chem. Rev.*, 117 (1992) 99.
- 212 A.R. Siedle, T.J. Kistenmacher, R.M. Metzger, Chiun-Shiung Kuo, R.P. van Duyne and T. Cape, *Inorg. Chem.*, 19 (1980) 2048.
- 213 D. Coucouvanis, F.J. Hollander, R. West and D. Eggerding, *J. Am. Chem. Soc.*, 96 (1974) 3006.
- 214 (a) F.J. Hollander and D. Coucouvanis, *J. Am. Chem. Soc.*, 96 (1974) 5646.  
(b) F.J. Hollander and D. Coucouvanis, *J. Am. Chem. Soc.*, 99 (1977) 6268.
- 215 D. Coucouvanis, D.G. Holah and F.J. Hollander, *Inorg. Chem.*, 14 (1975) 2657.
- 216 D. Coucouvanis, D. Swenson, N.C. Baenziger, D.G. Holah, A. Kostikas, A. Simopoulos and V. Petrouleas, *J. Am. Chem. Soc.*, 98 (1976) 5721.
- 217 A. Avdeef and J.P. Fackler, Jr., *Inorg. Chem.*, 17 (1978) 2182.
- 218 D. Altmeyen and R. Mattes, *Acta Crystallogr. Sect. B*, 36 (1980) 1942.
- 219 D. Coucouvanis, D. Swenson, N.C. Baenziger, C. Murphy, D.G. Holah, N. Sfarnas, A. Simopoulos and A. Kostikas, *J. Am. Chem. Soc.*, 103 (1981) 3350.

- 220 P. Arrizabalaga, G. Bernardinelli, P. Castan, M. Geoffroy and R. Soules, *C. R. Acad. Sci.*, 304-II (1987) 559.
- 221 J.-F. Petit, J.C. Trombe, A. Gleizes and J. Galy, *C. R. Acad. Sci.*, 304-II (1987) 1117.
- 222 J.-J. Bonnet, P. Cassoux, P. Castan, J.-P. Laurent and R. Soules, *Mol. Cryst. Liq. Cryst.*, 142 (1987) 113.
- 223 P. Arrizabalaga, G. Bernardinelli, M. Geoffroy and P. Castan, *Inorg. Chim. Acta*, 154 (1988) 35.
- 224 R. Krause and R. Mattes, *Z. Naturforsch. Teil B*, 45 (1990) 490.
- 225 R. Krause and R. Mattes, *Z. Naturforsch. Teil B*, 45 (1990) 1457.
- 226 F. Götzfried, W. Beck, A. Lerf and A. Sebal, *Angew. Chem.*, 91 (1979) 499.
- 227 P.G. Jones, G.M. Sheldrick, A. Fügner, F. Götzfried and W. Beck, *Chem. Ber.*, 114 (1981) 1413.
- 228 F. Götzfried, R. Grenz, G. Urban and W. Beck, *Chem. Ber.*, 118 (1985) 4179.
- 229 R. Grenz, F. Götzfried, U. Nagel and W. Beck, *Chem. Ber.*, 119 (1986) 1217.
- 230 D. Coucouvanis, *Acc. Chem. Res.*, 14 (1981) 201.
- 231 A. Müller, E. Diemann, R. Jostes and H. Bögge, *Angew. Chem.*, 93 (1981) 957.
- 232 A. Müller, H. Bögge, U. Schimanski, M. Penk, K. Nieradzick, M. Dartmann, E. Krickemeyer, J. Schimanski, C. Römer, M. Römer, H. Dornfeld, U. Wienböcker, W. Hellmann and M. Zimmermann, *Monatsh. Chem.*, 120 (1989) 367.