

# $\eta^2$ -Alkyne copper(I) and silver(I) compounds; from polymeric $[M^I R]_n$ to monomeric $[M^I R]$ units ( $M = Cu, Ag$ )

Heinrich Lang \*, Katrin Köhler, Sabine Blau

*Ruprecht-Karls-Universität Heidelberg, Anorganisch-Chemisches Institut, Im Neuenheimer Feld 270,  
D-69120 Heidelberg, Germany*

Received 5 May 1994

## Contents

Abstract	114
1. Introduction	114
2. Main-group element substituted $\eta^2$ -alkyne copper(I) and silver(I) compounds	115
2.1. Type A molecules, $[(\eta^2\text{-RC}\equiv\text{CR})\text{Cu}^I\text{X}]_2$	115
2.1.1. Synthesis	116
2.1.2. Reactivity	116
2.1.3. Spectroscopy and bonding	118
2.2. Type B molecules, $[(\eta^2\text{-RC}\equiv\text{CR})(\text{Cu}^I\text{X})_2]_2$	119
2.3. Type C molecules, $\{[\text{E}(\eta^2\text{-C}\equiv\text{CR})_2] (\text{M}^I\text{X})_2\}_n$	126
2.4. Type D and E molecules	128
2.4.1. Type D molecules, $[(\text{TBC})(\text{M}^I\text{OSO}_2\text{CF}_3)_n] (n = 1, 3); [(\text{TBC})_2\text{AgOSO}_2\text{CF}_3]$	129
2.4.2. Type E molecules, $(\eta^2\text{-RC}\equiv\text{CR})\text{Cu}^I\text{X}$	131
2.4.2.1. Synthesis	131
2.4.2.2. Spectroscopy and bonding	134
3. Organometallic substituted $\eta^2$ -alkyne copper(I) and silver(I) compounds	135
3.1. Type G molecules, $\text{M}_2\text{M}'_4 (\text{C}\equiv\text{CR})_8 (\text{PPh}_3)_2 (M = \text{Ir, Rh, Pt}; M' = \text{Cu, Ag})$	136
3.2. Type H molecules, $\{[\text{L}_n\text{M}(\text{C}\equiv\text{CR})_2]\text{M}'\text{X}\}_2$	142
3.3. Type I molecules, $\{\text{L}_n\text{M}(\text{C}\equiv\text{CR})_2\}\text{M}'\text{X} (M' = \text{Cu, Ag})$	145
3.3.1. Synthesis	146
3.3.2. Spectroscopy and bonding	149
3.3.3. Reactivity	153
3.3.3.1. $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti} (\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuCl}$	154
3.3.3.2. $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti} (\text{C}\equiv\text{CSiMe}_3)_2\}\text{M}' (\text{OSO}_2\text{CF}_3) (M' = \text{Cu, Ag})$	158
3.4. Type A, J and K molecules	158
3.4.1. Type J and K molecules, $[(\eta^2\text{-RC}\equiv\text{CML}_n)_2]\text{Cu}^I\text{X} (\text{J}), (\eta^2\text{-RC}\equiv\text{CML}_n)\text{Cu}^I\text{X} (\text{K})$	159
3.4.2. Organometallic type A molecules, $[(\eta^2\text{-RC}\equiv\text{CML}_n)\text{Cu}^I\text{X}]_2$	162
Acknowledgements	163
References	164

\* Corresponding author.

## Abstract

This paper deals with the breakdown of polymeric organo-copper(I) and -silver(I) compounds of general type  $[M^I R]_n$  ( $M = Cu, Ag$ ;  $R$  = singly bonded organic or inorganic ligand) to aggregates with lower polynuclearity and to mononuclear  $M^I X$  species. For the breakdown of the polynuclear structures of organo-copper(I) and -silver(I) compounds, main-group element substituted and organometallic substituted alkynes, diynes, and triynes are used. The synthesis, reactivity and structural features of  $[M^I R]_x$  ( $x = 1-4$ ) compounds are discussed.

**Keywords:** Copper(I) compounds; silver(I) compounds; Alkynes; Diynes; Triynes

---

## 1. Introduction

Organo-copper(I) and organo-silver(I) compounds of general type  $[M^I R]_n$  ( $M = Cu, Ag$ ;  $R$  = singly bonded organic ligand) possess wide potential as reagents for the stereoselective and regiospecific introduction of organic building blocks in organic and organometallic synthesis [1]. In most of these reactions the organo-copper(I) or organo-silver(I) reagents have been prepared in situ and used in excess. Therefore, the yield, based on organo-group 11 metal reagents, is sometimes low and depends on the reaction conditions used.

At present, it is still not clear whether  $[M^I R]_n$  compounds have any relation to the reactive species in copper(I)- or silver(I)-mediated organic reactions. One of the reasons is that organo-group 11 metal compounds do not exist as single discrete aggregates in solution: the  $[M^I R]_n$  compounds synthesized and characterized in the solid state are in equilibrium in solution with species with lower association.






One approach to this problem is to use mononuclear and therefore well defined organo-copper(I) and organo-silver(I) species.

So far, it has been found that nitrogen-, phosphorus-, oxygen- or sulphur-containing molecules are suitable for breaking down polynuclear structures of organo-copper(I) and organo-silver(I) compounds to aggregates with a lower polynuclearity [2]. However, only a few examples of mononuclear organo-copper(I) and organo-silver(I) species  $[M^I R]$  ( $M = Cu, Ag$ ;  $R$  = singly bonded organic ligand, e.g. 2,4,6- $Ph_3C_6H_2$  or 2,4,6- $tBu_3C_6H_2$ ) are known [3]. Recently, it was shown that main-group element and organometallic substituted alkynes and 1,4-diynes are also very capable of breaking down polynuclear copper(I) and silver(I) compounds to mononuclear compounds.

This paper focuses on the synthesis and reactivity of  $(\eta^2\text{-alkyne})\text{-copper(I)}$  and  $\text{-silver(I)}$  species and is divided into two parts: (1) the synthesis and reactivity of main-group element substituted alkyne, diyne and triyne copper(I) and silver(I) compounds (Section 2) and (2) the synthesis and reactivity of organometallic substituted alkyne and 1,4-diyne copper(I) and silver(I) compounds (Section 3).

## 2. Main-group element substituted $\eta^2$ -alkyne copper(I) and silver(I) compounds

Depending on the nature of the main-group substituted alkynes and of the copper(I) compounds  $[\text{Cu}^I\text{X}]_n$  ( $\text{X}$  = singly bonded organic or inorganic ligand) used, different coordination modes are observed. This section describes the synthesis of structural type A–E molecules.




In compounds of structural types A–E, alkyne units are  $\eta^2$ -coordinated to  $\text{Cu}^I\text{X}$  fragments.

So far, it has been found that alkynes  $\text{RC}\equiv\text{CR}$ , cyclic monoalkynes and cyclic triynes are able to form monomeric (type D and E), dimeric (type A) and tetrameric (type B) copper(I) alkyne complexes, while main-group bridged 1,4-diynes or cyclic diynes form polymers of type C.

### 2.1. Type A molecules, $[(\eta^2\text{-RC}\equiv\text{CR})\text{Cu}^I\text{X}]_2$

Structural type A molecules are dimers, which contain two copper atoms and two bridging X units in a centrosymmetric  $\text{Cu}_2\text{X}_2$  ring [4–17]. Each copper atom possesses a trigonal planar geometry.




The  $C_2$  unit of the alkyne ligand  $RC\equiv CR$  is symmetrically  $\eta^2$ -coordinated to one copper atom; the  $Cu_2X_2$  ring is coplanar with the  $CuC_2$  fragments.

### 2.1.1. Synthesis

There are two methods for the synthesis of type A molecules: (a) direct synthesis from alkynes and (b) nucleophilic substitution of  $[(\eta^2\text{-alkyne})Cu^IX]_2$  ( $X = Cl, Br, I$ ) compounds.

(a) *Direct synthesis route.* The dimeric compounds **2** are best synthesized by treatment of  $RC\equiv CR$  (**1a**,  $R = SiMe_3$ ; **1b**,  $R = Et$ ; **1c**,  $R = Ph$ ) or tmtch (3,3,6,6-tetramethyl-1-thia-4-cycloheptyne,  $C_{10}H_{16}S$ ; **1d**) with  $[Cu^IX]_n$  ( $X = Cl, Br, I, OSO_2CF_3, \dots$ ) (Table 1) [4–17].



(b) *Nucleophilic substitution route.* Compounds of type A are also formed by the reaction of  $[(\eta^2\text{-tmtch})Cu^IX]_2$  (**2j**) and  $NaX$  ( $X = OPh, O^tBu$ ) in thf; the dimeric complexes **2m** and **2n** are formed in  $\approx 50\%$  yield (Table 1) [10].

### 2.1.2. Reactivity

An overview of the diverse reactivity of type A compounds (**2a**, **2b** and **2j–2l**) is given in Scheme 1.

The nucleophilic substitution reaction of  $[(\eta^2\text{-Me}_3SiC\equiv CSiMe_3)CuX]_2$  (**2a**,  $X = Cl$ ; **2b**,  $X = Br$ ) and  $[(\eta^2\text{-tmtch})CuX]_2$  (**2j**,  $X = Cl$ ; **2k**,  $X = Br$ ; **2l**,  $X = I$ ) with  $[AgO_2CR]_n$  (route (a)),  $Na[acac]$  ( $acac = \text{acetylacetonate}$ ) (route (b)),  $Li[(Me_3SiN)_2CPh]$  (route (c)) or  $LiC_5H_5$  (route (d)) leads to the formation of type B and E molecules (Scheme 1) [13,14,18].

With the Lewis bases  $SMe_2$  (route (e)) and  $PPh_3$  (route (f)), the monomeric adduct  $[(\eta^2\text{-tmtch})Cu(I)(SMe_2)]$  (a type E molecule) [9,11] or the dimeric complexes  $\{(\eta^2\text{-tmtch})[Cu(\mu\text{-X})_2(PPh_3)_2]\}$  (**3a**,  $X = Cl$ ; **3b**,  $X = Br$ ; **3c**,  $X = I$ ) are formed (Scheme 1) [12].

In a similar manner, **2a** and **2b** react with the organometallic chelate  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(C\equiv CSiMe_3)_2$  to yield the monomeric bimetallic compounds  $\{(\eta^5\text{-}$

Table 1


Compounds  $[(\eta^2\text{-RC}\equiv\text{CR})\text{CuX}]_2$  (2)


Compound	R <sup>a</sup>	X	Ref.
2a	SiMe <sub>3</sub>	Cl	[5–7,13,14]
2b	SiMe <sub>3</sub>	Br	[5,7,13,14]
2c	SiMe <sub>3</sub>	OSO <sub>2</sub> CF <sub>3</sub>	[13,14]
2d	Et	O <sub>2</sub> CCF <sub>3</sub>	[15]
2e	Ph	O <sub>2</sub> CCF <sub>3</sub>	[15]
2f	Ph	O <sub>2</sub> CPh	[16]
2g	Ph	O <sub>2</sub> C(2-ClC <sub>6</sub> H <sub>4</sub> )	[17]
2h	Ph	O <sub>2</sub> C(2-BrC <sub>6</sub> H <sub>4</sub> )	[17]
2i	SiMe <sub>3</sub>	O <sub>2</sub> CCH <sub>3</sub>	[13,14]
2j	tmtch	Cl	[9–11]
2k	tmtch	Br	[9]
2l	tmtch	I	[9]
2m	tmtch	OPh	[10]
2n	tmtch	O <sup>t</sup> Bu	[10]
2o	tmtch	SPh	[8,10]
2p	tmtch	C≡CPh	[10]

<sup>a</sup> tmtch = 3,3,6,6-tetramethyl-1-thia-4-cycloheptyne.

$\text{C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuX}$  ( $\text{X}=\text{Cl}, \text{Br}$ ) as shown by route (g) (Scheme 1) [13,19]. However, with  $\text{LiC}\equiv\text{CSiMe}_3$  dimeric **2b** yields the copper(I) acetylide  $[\text{CuC}\equiv\text{CSiMe}_3]_n$  (**4**) (route (h), Scheme 1) [13]. The formation of a compound of structural type A in which the  $\text{Me}_3\text{SiC}\equiv\text{C}$  ligands are  $\sigma, \pi$  bridging two  $(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CSiMe}_3)\text{Cu}$  building blocks was not observed [13].

When isoelectronic  $\text{CN}^-$  is used instead of  $\text{Me}_3\text{SiC}\equiv\text{C}^-$ , the products formed are  $[\text{CuCN}]_n$  and  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  (**1a**). Whereas the substitution of  $\text{Cl}^-$  in  $[(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CSiMe}_3)\text{CuCl}]_2$  (**2a**) by  $\text{CN}^-$  was not successful, the direct reaction of the





Scheme 1. Reaction of 2a, 2b, 2j, 2k and 2l with different substrates.

highly strained cycloalkyne tmtch with  $[\text{CuCN}]_n$  affords the tetranuclear complex  $[(\eta^2\text{-tmtch})\text{CuCN}]_4$  (**5**) in high yields [11].

Complex **5** contains a slightly puckered  $[\text{CuCN}]_4$  ring. Each cyanide ion is  $\sigma$  and dative bonded to two different copper atoms, thus forming an almost linear  $\text{Cu}-\text{CN}\rightarrow\text{Cu}$  unit [11]. This coordination mode is typical for copper(I) cyanide [20–22]. A similar compound containing  $\text{CuCl}$  building blocks instead of  $\text{CuCN}$  units can be isolated by the reaction of  $\text{MeC}\equiv\text{CMe}$  with  $[\text{CuCl}]_n$  [4].

For a detailed discussion of type B, E and I molecules (Scheme 1), see Sections 2.2, 2.4 and 3.3.

### 2.1.3. Spectroscopy and bonding

In the binuclear compounds  $[(\eta^2\text{-RC}\equiv\text{CR})\text{CuX}]_2$  ( $\text{R} = \text{SiMe}_3$ , Et, Ph) and  $[(\eta^2\text{-tmtch})\text{CuX}]_2$  (type A molecules), the alkyne ligands are  $\eta^2$ -coordinated to  $\text{Cu}^{\text{I}}\text{X}$  moieties. This is evidenced by the IR and  $^{13}\text{C}$  NMR data (Table 2). In the IR spectra the  $\nu(\text{C}\equiv\text{C})$  absorption decreases from 2200/2170 or 2110  $\text{cm}^{-1}$  in the free alkynes tmtch and  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  to 2015/1950  $\text{cm}^{-1}$  in **2j–2o** and **3a–3c** or 1940–1960  $\text{cm}^{-1}$  in **2a–2c** (Table 2) [5–14]. The same behaviour is found for all other compounds of type A (Tables 1 and 2).

It is found that the copper–alkyne bond is stronger in the  $[(\eta^2\text{-tmtch})\text{CuX}]_2$  compounds **2j–2o** than in the analogous  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  substituted compounds **2a–2c**, as given by the  $\Delta\nu$  value of  $\sim 190\text{ cm}^{-1}$  (**2j–2o**) or  $\sim 150\text{ cm}^{-1}$  (**2a–2c**) (Table 2). In addition, the  $\eta^2$ -coordination of the alkyne ligands in compounds **2**

Table 2  
Selected IR and  $^{13}\text{C}\{^1\text{H}\}$  NMR data for compounds **2** and **3**

Compound	IR $\nu(\text{C}\equiv\text{C})$ ( $\text{cm}^{-1}$ )	$^{13}\text{C}\{^1\text{H}\}$ NMR $\delta(\text{C}\equiv\text{C})$ (ppm)	Ref.
<b>2a</b>	1949	115.2	[5–7,13,14]
<b>2b</b>	1953	116.4	[5,13,14]
<b>2c</b>	1953	–	[13]
<b>2d</b>	2070	88.2	[15]
<b>2e</b>	–	95.7	[15]
<b>2j</b>	2007; 1981	109.1	[9–12]
<b>2k</b>	2005; 1981	110.5	[12]
<b>2l</b>	2015; 1980	112.2	[12]
<b>2m</b>	1955	107.8	[10]
<b>2o</b>	1970; 1950	–	[8,10]
<b>3a</b>	2001; 1980	109.5	[12]
<b>3b</b>	2000; 1979	110.5	[12]
<b>3c</b>	2011; 1980	112.5	[12]

and **3** to a  $\text{Cu}^{\text{I}}\text{X}$  building block leads to changes in the  $^{13}\text{C}$  NMR feature of the  $\text{RC}\equiv\text{CR}$  ligands (Table 2), the carbon signals of the  $\text{RC}\equiv\text{CR}$  units in type **A** molecules being shifted slightly to lower field compared with the corresponding starting materials (Table 2) [8–19].

The lengthening of the  $\text{C}\equiv\text{C}$  triple bond and the bending of the  $\text{R}-\text{C}\equiv\text{C}-\text{R}$  units from the linear arrangement in the free ligands, due to  $\eta^2$ -coordination of the alkyne ligand to a  $\text{Cu}^{\text{I}}\text{X}$  building block, is corroborated by X-ray structure analysis (Table 3).

A typical example of a type **A** molecule (complex **2b**) is shown in Fig. 1 [7,13].

Fig. 1 shows that compounds of type **A** are dimeric. Each  $\text{C}_2$  unit of the acetylene group is symmetrically  $\pi$  bonded to one copper atom. The  $\text{CuC}_2$  building block is coplanar with the four-membered  $\text{Cu}_2\text{Br}_2$  ring and each copper atom possesses a trigonal planar environment. As a result of the  $\eta^2$ -coordination of the  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  ligand to  $\text{Cu}^{\text{I}}\text{Br}$ , the  $\text{Si}-\text{C}\equiv\text{C}-\text{Si}$  unit is distorted compared with the non-coordinated  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  ligand (Table 3). At the same time, the  $\text{C}\equiv\text{C}$  triple bond of the alkyne is lengthened from 1.182 Å in  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  to 1.24 Å in **2b** (Table 3). The copper–copper distance of 3.05 Å is significantly larger than the sum of the van der Waals radii (1.54 Å) [23], implying that there is no copper–copper bond.

## 2.2. Type **B** molecules, $[(\eta^2\text{-RC}\equiv\text{CR})(\text{Cu}^{\text{I}}\text{X})_2]_2$

As shown in Section 2.1, acetylenes can be used to break down the oligomeric structure of copper(I) carboxylates to a dimeric structure (type **A** molecules) (Table 1) [14–17]. However, depending on the nature of the R and R' groups on the alkynes  $\text{RC}\equiv\text{CR}'$  and on the stoichiometry of the copper(I) carboxylate and alkyne used,

Table 3  
X-ray data for compounds 2 and 3

Compound	Ref.	Cu...Cu (Å)	Cu-X (Å)	Cu-    (Å)	C≡C (Å)	X-Cu-X (°)	Cu-X-Cu (°)	C≡C-R (°)
<b>2a</b>	[5]	—	2.279(1) 2.281(1)	2.017(3) 2.020(3)	1.227(5)	96.64(3)	93.36(3)	166.3(3) 164.2(3)
<b>2b</b>	[7,13]	3.05	2.407(1)	2.043(7) 2.045(7)	1.24(1)	89.9	—	165.0(6) 162.9(6)
<b>2f</b>	[16]	2.782(3)	1.99(1); 1.93(1) 1.99(1); 1.95(1)	1.93(2); 1.94(1) 1.95(1); 1.93(1)	1.28(2) 1.21(2)	100.4(4) 99.6(4)	—	157(1); 158(2) 158(1); 160(1)
<b>2j</b>	[10]	3.021(1)	2.267(2) 2.283(2)	1.937(5) 1.951(5)	1.220(6)	96.8(1)	83.2(1)	147.9(5) 147.3(5)
<b>2m</b>	[10]	2.940(2)	1.945(2)	1.929(3)	1.247(5)	81.8(2)	98.2(2)	146.9(3)
<b>2o</b>	[8]	2.904(1)	2.293(1); 2.284(1) 2.295(1); 2.283(1)	1.970(3); 1.962(3) 1.962(3); 1.969(3)	1.230(5) 1.223(4)	91.89(4) 91.87(4)	78.53(3) 78.95(4)	147.8(4); 147.1(4) 145.9(4); 147.7(4)
<b>2p</b>	[10]	2.386(1)	2.014(3); 2.022(3)	1.940(4) 1.951(4)	1.220(5)	107.5(1)	72.5(1)	147.2(4) 148.1(4)
<b>3a<sup>a</sup></b>	[12]	3.134(2)	2.244(2); 2.242(3) 2.414(3); 2.416(2)	1.942(7) 1.946(6)	1.22(1)	100.1(1)	84.5(1)	148.1(7) 147.3(6)
<b>3b<sup>b</sup></b>	[12]	3.186(2)	2.370(2); 2.384(2) 2.610(2); 2.586(2)	1.95(1) 1.97(1)	1.22(2)	103.4(1) 91.7(1)	79.4(1) 79.6(1)	149(1) 148(1)
<b>3c<sup>c</sup></b>	[12]	3.157(1)	2.544(1); 2.541(1) 2.704(1); 2.714(1)	1.989(8) 1.982(8)	1.23(1)	108.6(1) 99.3(1)	73.9(1) 73.8(1)	146.1(8) 149.3(8)

<sup>a</sup> Cu-P 2.255(3)/2.254(2) Å; P-Cu-P 125.9(1)°.

<sup>b</sup> Cu-P 2.264(3)/2.264(3) Å; P-Cu-P 126.8(1)°.

<sup>c</sup> Cu-P 2.268(2)/2.261(2) Å; P-Cu-P 125.7(1)°.





Fig. 1. Molecular geometry and atom labelling scheme for **2b**. Selected bond distances and angles are given in Table 3 [7,13].



other coordination modes can be realized, i.e. structural type **B** molecules [10,13–17,24,25].

The reaction of the alkynes  $\text{RC}\equiv\text{CR}'$  ( $\text{R}, \text{R}' = \text{Me}, \text{Et}, \text{CO}_2\text{Me}, \text{CO}_2\text{Et}, \text{tmthc}$ ) with  $[\text{CuO}_2\text{CR}'']_n$  yields the tetranuclear complexes  $[(\eta^2\text{-RC}\equiv\text{CR}')(\text{CuO}_2\text{CR}'')]_2$  (**6**) (Table 4) together with dimeric  $[(\eta^2\text{-RC}\equiv\text{CR}')\text{CuO}_2\text{CR}'']_2$  (**2d–2h**) (Section 2.1; Table 1).


The formation of the dimeric compounds  $[(\eta^2\text{-RC}\equiv\text{CR}')\text{CuO}_2\text{CR}'']_2$  (**2d–2h**) ( $\text{R}=\text{R}' = \text{Et}, \text{Ph}$ ) (Section 2.1; Table 1) and the tetranuclear compounds  $[(\eta^2\text{-RC}\equiv\text{CR}')(\text{CuO}_2\text{CR}'')]_2$  (**6**) (Table 4) clearly depends on the donor strength of the acetylenes used; weak donor ligands, such as dimethylacetylene dicarboxylate and diethylacetylene dicarboxylate, lead to the formation of coordination compounds in which the initial tetrameric copper(I) carboxylate moiety is retained.

With electron-donating substituents, such as Ph or Et, on the alkyne, the dimeric compounds described in Section 2.1 (type **A** molecules) are formed (Table 1).

Table 4  
Compounds **6** and **7**

Compound	R	R'	R''	Ref.
<b>6a</b>	Et	Et	CF <sub>3</sub>	[15,24]
<b>6b</b>	CO <sub>2</sub> Me	CO <sub>2</sub> Me	CF <sub>3</sub>	[15]
<b>6c</b>	CO <sub>2</sub> Me	Me	CF <sub>3</sub>	[15]
<b>6d</b>	CO <sub>2</sub> Me	CO <sub>2</sub> Me	Ph	[17]
<b>6e</b>	CO <sub>2</sub> Me	CO <sub>2</sub> Me	2-ClC <sub>6</sub> H <sub>4</sub>	[17]
<b>6f</b>	CO <sub>2</sub> Me	CO <sub>2</sub> Me	2-BrC <sub>6</sub> H <sub>4</sub>	[17]
<b>6g</b>	CO <sub>2</sub> Et	CO <sub>2</sub> Et	Ph	[17]
<b>6h</b>	CO <sub>2</sub> Et	CO <sub>2</sub> Et	2-ClC <sub>6</sub> H <sub>4</sub>	[17]
<b>6i</b>	CO <sub>2</sub> Et	CO <sub>2</sub> Et	2-BrC <sub>6</sub> H <sub>4</sub>	[17]
<b>6j</b>		tmrch <sup>a</sup>	Me	[10]
<b>6k</b>		tmrch <sup>a</sup>	Ph	[10]
<b>7</b>	SiMe <sub>3</sub>	SiMe <sub>3</sub>	Me	[13,25]

<sup>a</sup> tmrch = 3,3,6,6-tetramethyl-1-thia-4-cycloheptyne.




When the tetrameric starting compound  $[\text{CuO}_2\text{CPh}]_4$  is reacted with isocyanates, azo compounds and quinoline as nitrogen donor ligands, the  $[\text{Cu}(\mu\text{-O}_2\text{CPh})_2]$  building block is also formed [26].

As expected, the reaction of  $[\text{CuO}_2\text{CCF}_3]_n$  with  $\text{EtC}\equiv\text{CEt}$  yields both types of compound (type **A** and **B** molecules) [15,24]. The product obtained can be controlled by the stoichiometry of the starting materials used: with an excess of 3-hexyne in the reaction mixture the only product isolated is the dinuclear compound  $[(\eta^2\text{-EtC}\equiv\text{CEt})\text{CuO}_2\text{CCF}_3]_2$  (**2d**), whereas with no excess of 3-hexyne the tetranuclear compound  $[(\eta^2\text{-EtC}\equiv\text{CEt})(\text{CuO}_2\text{CCF}_3)_2]_2$  (**6a**) is formed [15,24].

In one special case it could be shown that tetrameric compounds of type **B** are formed from dimeric structural type **A** complexes. Treatment of ethereal solutions of  $[\text{CuO}_2\text{CCH}_3]_n$  with  $\text{Me}_3\text{SiC}\equiv\text{SiMe}_3$  (**1a**) yields together with dimeric  $[(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CSiMe}_3)\text{CuO}_2\text{CCH}_3]_2$  (**2i**) the tetrameric coordination compound  $[(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CSiMe}_3)(\text{CuO}_2\text{CCH}_3)_2]_2$  (**7**) [13,25].

It has been confirmed independently that compound **2i** dimerizes in solution, by  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  elimination, to form **7**. This dimerization is evidenced by the IR spectroscopic data: the  $\nu(\text{C}\equiv\text{C})$  vibration of **2i** at  $1952\text{ cm}^{-1}$  is shifted to  $1731\text{ cm}^{-1}$  in **7** [13,25]. These data are consistent with the  $\eta^2$ -coordination of the  $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$  unit to one copper atom in **2i** and to two copper atoms in **7**. Similar IR



spectroscopic data for the  $\text{C}\equiv\text{C}$  stretching vibration are found for other alkyne-bridged dicopper(I) complexes containing macrocycles such as tropocoronand-6,6 [27,28].

The solid-state structures of selected examples of compounds **6** and **7** have been determined by X-ray diffraction studies (Fig. 2, Table 5).

Fig. 2 shows two structural isomers (**B**<sub>1</sub> and **B**<sub>2</sub>) of type **B** molecules. The formation of these two products is surprising, since the reaction conditions for the synthesis of type **B** molecules used are identical. The common feature of compounds **6** and **7** is that each of the two acetylenes, and also each of the four carboxylate units, act as a four-electron donor, bridging two adjacent copper atoms. The two structural isomers of type **B** compounds (**B**<sub>1</sub> and **B**<sub>2</sub>) differ in the arrangement of the four copper atoms with respect to each other. In the tetrameric compounds **6** the four copper atoms form a distorted tetrahedral core, whereas in **7** a square-planar arrangement is found. In **6** two short (2.76–3.00 Å) and four long (3.00–3.17 Å)  $\text{Cu}\cdots\text{Cu}$  distances are found [10]; in **7** the  $\text{Cu}\cdots\text{Cu}$  distances are 3.02 Å [13,25].

The observed bond weakening by IR spectroscopy (see above) of the  $\text{C}\equiv\text{C}$  triple bonds in compounds **6** and **7**, relative to the starting materials  $\text{RC}\equiv\text{CR}'$ , was confirmed by X-ray diffraction studies (Fig. 2, Table 5). The  $\text{C}\equiv\text{C}$  distances of 1.26–1.30 Å found in compounds **6** and **7** are significantly longer than in the free acetylenes  $\text{RC}\equiv\text{CR}'$ , but they are shorter than in the alkyne-bridged dinuclear copper(I) tropocoronand-6,6 compound  $[\text{Cu}_2(\eta^2\text{-RC}\equiv\text{CR})(\text{TC-6,6})]$  ( $\text{R}=\text{CO}_2\text{Me}$ ,  $\text{CO}_2\text{Et}$ ; TC-6,6 = tropocoronand-6,6) [27,28]. The longer  $\text{C}\equiv\text{C}$  distances found in tetrameric **6** and **7** (Table 5) compared with dimeric **2** (Table 3) are consistent with





Fig. 2. Molecular geometry and atom labelling scheme for **6j** (top) and **7** (bottom) [10,13,25]. Selected bond distances (Å) and angles (°) are given in Table 5.

bridging alkyne units serving as 'four-electron donor' ligands,  $\pi$ -bonded to two copper(I) centres. As a result of the  $\eta^2$ -coordination of the alkyne ligands to the copper atoms in **6** and **7**, the angles  $R-C\equiv C-R$  are considerably reduced ( $143$ – $158^\circ$ ) (Table 5). In comparison with the dinuclear  $Cu_2(\eta^2-R-C\equiv C-R)(TC-6,6)$  compounds ( $R-C\equiv C-R$  angles  $139.0^\circ$ ,  $138.2^\circ$ ), both structural features imply that  $\pi$  back-bonding in compounds **6** and **7** is less important.

Table 5  
X-ray data for compounds **6** and **7**

Compound	Ref.	Cu...Cu (Å)	Cu—O (Å)	Cu←    (Å)	C≡C (Å)	O—Cu—O (°)	C≡C—R (°)
<b>6a</b>	[24]	2.820(3); 3.136(3)	—	1.97(2); 1.99(2)	1.26(2)	—	152(2)
		3.130(3); 3.079(3)		1.97(3); 2.00(2)	1.26(2)		155(2)
		3.166(3); 2.799(3)		1.98(2); 2.00(2)			148(2)
<b>6b</b>	[17]			1.92(2); 1.98(2)			156(2)
		2.960(2); 2.998(2)	1.927(3); 1.899(4)	1.976(4); 1.968(4)	1.280(5)	98.3(2)	147.1(4)
		2.980(2); 3.001(2)	1.914(3); 1.922(3)	1.951(3); 1.950(4)	1.295(4)	96.6(2)	146.4(4)
		3.129(3); 2.941(2)	1.915(3); 1.910(3)	1.948(3); 1.950(3)		96.2(2)	148.5(3)
			1.915(3); 1.907(3)	1.963(3); 1.952(3)		96.6(2)	151.6(3)
<b>6j</b>	[10]	3.103(3); 3.151(3)	1.932(4)	1.974(5)	1.259(7)	—	142.7(5)
		2.774(3); 2.762(3)					
		3.131(3); 3.157(3)					
<b>7</b>	[13,25]	3.02(1); 3.02(1)	1.941(3); 1.99(1)	1.982(4); 1.985(4)	1.278(6)	106.2(6)	158.6(4)
			1.99(1); 1.945(3)	1.995(4); 2.007(5)		111.8(5)	157.0(4)



Scheme 2. Possible routes for the formation of type B compounds: (a) weak donors (type **B<sub>1</sub>** molecules); (b) strong donors (type **A** and **B<sub>2</sub>** molecules).

The formation of compounds **2d–2i**, **6** and **7** from copper carboxylates and alkynes is best explained by the two possible reaction pathways given in Scheme 2. The reaction of copper(I) carboxylates with acetylenes proceeds most likely via initial attack of the acetylene at one of the copper atoms (Scheme 2).


With weak donor ligands, such as  $\text{RO}_2\text{CC}\equiv\text{CCO}_2\text{R}$  ( $\text{R} = \text{Me}, \text{Et}$ ) the formation of tetrameric **6** proceeds via route (a) (Scheme 2). The donor strength of these acetylenes is not large enough to break copper–oxygen bonds in the tetrameric copper(I) carboxylate starting compounds, thus generating a three-coordinated copper(I) centre. The  $\pi$ -bonded acetylene moiety then interacts with the opposite copper atom in the  $\text{Cu}_4$  core in the starting compound; a perpendicularly oriented acetylene bridging unit is formed. Subsequent complexation of a second acetylene ligand then yields complex **6** (Scheme 2).

The formation of the dimeric complex **2i** and the tetrameric compound **7** is explained by pathway (b) by using a strong donor acetylene ligand (Scheme 2): the donor strength of  $\text{RC}\equiv\text{CR}$  ( $\text{R} = \text{Ph}, \text{Et}, \text{SiMe}_3$ ) is large enough to break copper–oxygen bonds in tetrameric copper(I) carboxylates. Thus, bond breaking of  $\text{Cu}(1,3)-\text{O}$  and simultaneous inverse bond rebuilding yields the intermediate **F** (Scheme 2); **F** then reacts further with alkynes to form dimeric **2i** or tetrameric **7** [13,25].


### 2.3. Type C molecules, $\{[E(\eta^2\text{-C}\equiv\text{CR})_2](\text{M}^{\text{I}}\text{X})_2\}_n$

In general, alkynes can be used to break down the polymeric structures of  $\text{Cu}^{\text{I}}\text{X}$  compounds,  $[\text{Cu}^{\text{I}}\text{X}]_n$  ( $\text{X} = \text{singly bonded organic or inorganic ligand}$ ) (Sections 2.1

and 2.2) [4–25]. Similarly, diynes can be used. However, oligomers are obtained in which the diyne units are linked together by  $\text{Cu}^{\text{I}}\text{X}$  building blocks, thus forming type C molecules [13,29–33].



Polymers containing  $\text{M}^{\text{I}}\text{OSO}_2\text{CF}_3$  ( $\text{M} = \text{Cu}, \text{Ag}$ ) entities can be obtained, as shown by Gleiter et al. [29], when cyclic diynes, such as 1,6-cyclodecadiyne (**8a**), 1,7-cyclododecadiyne (**8b**) and 1,8-cyclotetradecadiyne (**8c**) are reacted with  $[\text{M}^{\text{I}}\text{OSO}_2\text{CF}_3]_n$  in thf [29]. Compounds **9** (**9a**,  $\text{M} = \text{Cu}$ ,  $n = 3$ ; **9b**,  $\text{M} = \text{Ag}$ ,  $n = 3$ ; **9c**,  $\text{M} = \text{Cu}$ ,  $n = 4$ ; **9d**,  $\text{M} = \text{Ag}$ ,  $n = 4$ ; **9e**,  $\text{M} = \text{Ag}$ ,  $n = 5$ ) are formed in high yields.



The spectroscopic data (IR,  $^{13}\text{C}$  NMR) for polymeric **9** indicate a weak interaction between the  $\text{C}\equiv\text{C}$  triple bonds and the copper(I) or silver(I) ions [29].

The solid-state structures of compounds **9** were elucidated by X-ray crystallographic studies, carried out on **9b–e** [29]. The molecular structure of **9b** is shown in Fig. 3 [29]. In the crystal structure of **9b** the double strands of the 1,6-cyclodecadiyne rings are in a chair conformation and linked by  $\text{Ag}(\text{OSO}_2\text{CF}_3)_2\text{Ag}$  units (Fig. 3, right) to form a two-dimensional network (Fig. 3, left).

In compounds **9c–e** the twelve-membered rings of the cyclododecadiynes are coupled by copper(I) or silver(I) ions. Since each trifluoromethanesulphonate ligand in these compounds uses two oxygen atoms to coordinate two different metal centres, a three-dimensional network is formed [29].

In contrast to the above-described reaction of cyclic diynes with  $\text{M}^{\text{I}}\text{OSO}_2\text{CF}_3$ , the reaction of the  $\text{Me}_2\text{Si}$ -bridged 1,4-diyne  $\text{Me}_3\text{SiC}\equiv\text{C}-\text{Me}_2\text{Si}-\text{C}\equiv\text{CSiMe}_3$  (**10**) with  $[\text{Cu}^{\text{I}}\text{X}]_n$  ( $\text{X} = \text{Cl}, \text{Br}$ ) in  $\text{CH}_2\text{Cl}_2$  yields the polymer  $\{\text{Me}_2\text{Si}(\text{C}\equiv\text{CSiMe}_3)_2(\text{CuX})_2\}_n$  (**11a**,  $\text{X} = \text{Cl}$ ; **11b**,  $\text{X} = \text{Br}$ ) of structural type C [13,30].

In polymeric **11** each  $\text{Me}_3\text{SiC}\equiv\text{C}$  building block is  $\eta^2$ -coordinated to a  $\text{Cu}^{\text{I}}\text{X}$  unit and the  $\text{Cu}^{\text{I}}\text{X}$  moieties dimerize to form  $\text{Cu}_2\text{X}_2$  rings. As a result of the formation of  $\text{Cu}_2\text{X}_2$  units, two  $\text{Me}_2\text{Si}(\text{C}\equiv\text{CSiMe}_3)_2$  building blocks are linked together, thus yielding a polymeric structure.

The  $\eta^2$ -coordination of the  $\text{Me}_3\text{SiC}\equiv\text{C}$  units to  $\text{Cu}^{\text{I}}\text{X}$  is indicated by the IR





Fig. 3. Molecular geometry and atom labelling scheme for **9b** (left, polymer; right,  $\text{Ag}_2(\text{OSO}_2\text{CF}_3)_2$  bridge) [29]. Selected bond distances (Å) and angles (°) are as follows: C1–C2/C6–C7 1.211, Ag–Cl/Ag–C2/Ag–C6/Ag–C7 2.410; C1–C2–C3 170.7, D1–Ag–D2 131.2 (D=mid-point of the  $\text{C}_2$  units).



spectroscopic data. In the IR spectra of **11a** and **11b** the  $\nu(\text{C}\equiv\text{C})$  vibration decreases to  $1960 \text{ cm}^{-1}$  compared with  $2104 \text{ cm}^{-1}$  in **10** [13,30].

A similar structure to that of compounds **11a** and **11b** is found in  $\{\text{Hg}(\text{C}\equiv\text{CPh})_2(\text{CuCl})_2\}_n$  (**11c**) in which the  $\text{Me}_2\text{Si}$  bridge head of the 1,4-diyne in compounds **11a** and **b** is replaced with a mercury atom [31].


Another polymer, containing  $\text{Cu}_2\text{Cl}_2$  units, can be prepared by the reaction of  $[\text{Cu}^{\text{I}}\text{Cl}]_n$  with  $\text{HC}\equiv\text{C}(\text{CH}_2)_4\text{C}\equiv\text{CH}$  (**12**) [32]. The resulting  $\{\text{C}_8\text{H}_{10}(\text{CuCl})_2\}_n$  polymer (**13**) was characterized by IR spectroscopy and a crystal structure determination. In **13**, layers of fused four- and eight-membered  $\text{Cu}^{\text{I}}\text{Cl}$  rings are cross-linked by **12** to form a three-dimensional network [32].

The polymeric structures of **11a** and **11b** can be depolymerized using the organic chelate acetylacetonate (acac); monomeric **14** is formed in quantitative yields [13,30].



For a detailed discussion of monomeric  $(\eta^2\text{-alkyne})\text{copper(I)}$   $\beta$ -diketonates see Section 2.4.

#### 2.4. Type D and E molecules

The breakdown of polymeric copper(I) and silver(I) compounds into dimeric or tetrameric compounds by alkynes is described in Sections 2.1 and 2.2 [4–25]. In




For a detailed discussion of structural type I molecules, see Section 3.



X = Singly bonded ligand; L = dative bonded ligand.

2.4.1. Type **D** molecules,  $[(TBC)(M^I\text{OSO}_2\text{CF}_3)_n]$  ( $n=1,3$ );  $[(TBC)_2\text{AgOSO}_2\text{CF}_3]$


The reaction of TBC (1,2:5,6:9,10-tribenzocyclododeca-1,5,9-triene-3,7,11-triyne) (**15**) with  $[\text{CuOSO}_2\text{CF}_3]_n$  yields  $[(TBC)\text{-CuOSO}_2\text{CF}_3]$  (**16**) and  $[(TBC)\text{-(CuOSO}_2\text{CF}_3)_3]$  (**17**) [41,43].



In **16**, the copper(I) ion is chelated by the three  $\text{C}_2$  building blocks of TBC and semi-coordinated by an oxygen of the triflate anion ( $\text{Cu-O}$  2.549 Å) [41].

In **17**, each  $\text{C}\equiv\text{C}$  triple bond of the macrocyclic trialkyne ligand TBC is  $\eta^2$ -coordinated to a copper(I) triflate building block. Complex **17** can best be described as a cofacial bimacrocyclic: the TBC ligand is  $\pi$ -bonded to a cyclic trimer of  $\text{CuOSO}_2\text{CF}_3$  moieties in which the three copper atoms are bridged by an  $\text{O-S-O}$  building block, thus forming a 12-membered macrocyclic ring with  $\text{Cu-O-S-O}$  as a repeat unit [43].

In contrast, when  $[\text{AgOSO}_2\text{CF}_3]_n$  instead of  $[\text{CuOSO}_2\text{CF}_3]_n$  is reacted with TBC, the silver(I) trifluoromethanesulphonate sandwich complex  $[(TBC)_2\text{AgOSO}_2\text{CF}_3]$  (**18**) is formed [42].




Compound **18** is the first example of a 12-membered macrocyclic sandwich complex [42]. A single X-ray diffraction study of **18** (Table 7) shows that the asymmetric unit contains one  $(TBC)_2\text{Ag}^+$  eclipsed sandwich together with one  $(TBC)_2\text{Ag}^+$  staggered sandwich. The  $\text{Ag-C}$  distances in the eclipsed conformation range from 2.5 to 2.9 Å, whereas in the staggered conformation they are 2.7–2.8 Å. The much smaller range of the  $\text{Ag-C}$  interatomic distances in the latter indicate that in the

staggered situation the six  $C_2$  units are each donating 1.33 electrons, or that there is an electrostatic interaction [42]. In contrast, in the eclipsed conformation four of the six alkynes behave as a two-electron donor while two  $C_2$  building blocks are non-interacting. Similar metal to ligand interactions are found in ruthenium and rhodium bis(arene) complexes [50–52].

#### 2.4.2. Type E molecules, $(\eta^2-RC\equiv CR)Cu^IX$

**2.4.2.1. Synthesis.** As shown in Section 2.3, polymeric  $\{Me_2Si(C\equiv CSiMe_3)_2(CuX)_2\}_n$  (**11a**,  $X=Cl$ ; **11b**,  $X=Br$ ) reacts with  $Na[acac]$  to form the monomeric compound  $[Me_2Si(C\equiv CSiMe_3)_2][Cu(acac)_2]$  (**14**) [13,30].

Using similar reaction conditions, the dimeric complexes  $[(\eta^2-RC\equiv CR)CuX]_2$  ( $X=Cl, Br$ ) (type A molecules) can be cleaved by  $\beta$ -diketonates to afford the monomeric  $(\eta^2-RC\equiv CR)Cu(\beta\text{-diketonate})$  compounds **19** in high yields [13,14,53–55].




- 19a:**  $R = SiMe_3, R' = R'' = Me$   
**19b:**  $R = Ph, R' = R'' = CF_3$   
**19c:**  $R = SiMe_3, R' = R'' = CF_3$   
**19d:**  $R = Me, R' = R'' = CF_3$   
**19e:**  $R = SiMe_3, R' = tBu, R'' = (CF_2)_2CF_3$   
**19f:**  $R = Me, R' = tBu, R'' = (CF_2)_2CF_3$

The spectroscopic data for compounds **19a–19f** are consistent with chelation of the  $\beta$ -diketonate ligand through both oxygens to the copper atom (Section 2.4.2.2).


Recently, it was found that compounds **19** deposit copper films by chemical vapour deposition (CVD) through a thermally induced disproportionation reaction that generates  $Cu^0$ ,  $Cu^{II}(\beta\text{-diketonate})_2$  and  $RC\equiv CR$  [13,14,53,55].

Further, Olbrich et al. [18] have shown that dimeric  $[(\eta^2\text{-tmtch})CuCl]_2$  (**2j**) (tmtch = 3,3,6,6-tetramethyl-1-thia-4-cycloheptyne) reacts with lithium  $N,N'$ -bis-(trimethylsilyl)benzaminidate to yield the monomeric compound **20** [18].



Compound **20** is monomeric in the solid state and the copper atom possesses a trigonal planar environment, where the nitrogen atoms of the  $\text{PhC}(\text{NSiMe}_3)_2$  ligand and the  $\text{C}_2$  unit of the tmthc group are considered to occupy the three coordination sites (see Table 7) [18].


However, using the dimetallated macrocyclic tropocoronand-6,6 instead of  $\text{Li}[\text{PhC}(\text{NSiMe}_3)_2]$ , the alkyne-bridged dicopper(I) complexes **21a–21d** can be isolated in moderate yields [27,28].




There are three routes for the synthesis of complexes **21**: two methods employing labile copper(I) acetonitrile or carbon monoxide intermediates, and a third route in which the formation of complex **21** was achieved by generating the dicopper(I) tropocoronand unit in the presence of an excess of  $\text{RC}\equiv\text{CR}$  [27,28].

The structures in the solid state of compounds **21c** and **21d** have been determined by X-ray diffraction (see Table 7) [27,28]. It was found that the  $\text{C}\equiv\text{C}$  triple bond lengths of the  $\text{RC}\equiv\text{CR}$  groups are 1.314 Å (**21c**) and 1.320 Å (**21d**), substantially longer than the values found in other copper(I) alkyne complexes (see above). These larger  $\text{C}\equiv\text{C}$  distances are consistent with the bridging alkyne unit  $\text{RC}\equiv\text{CR}$  serving as a four-electron donor. Similar  $\text{C}\equiv\text{C}$  bond lengths were found in complexes of the type  $(\eta^2\text{-RC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$  [56–60].

Complexes of type **E** are also formed by the addition of bidentate neutral chelating ligands such as 1,10-phenanthroline, 2,2'-bipyridine or di-2-pyridylamine to  $[\text{Cu}^I\text{X}]_n$  ( $\text{X} = \text{BF}_4, \text{ClO}_4, \text{PF}_6, \text{SbF}_6$ ) compounds in the presence of an excess of  $\text{RC}\equiv\text{CR}'$  (Table 6) [44–47].



Reger and Huff [47] showed that compounds  $\{(\eta^2\text{-RC}\equiv\text{CR}')[\text{Cu}(\text{bpy})]_2\}\text{X}_2$  (**23**) can be isolated by repeated precipitation from solutions which initially contain  $[(\eta^2\text{-RC}\equiv\text{CR}')\text{Cu}(\text{bpy})]\text{X}$  (**22q–22s**) (bpy = 2,2'-bipyridine) and no additional  $\text{RC}\equiv\text{CR}'$  (Table 6).

**23**


Whereas in compounds **22** the  $\text{C}_2$  unit of the alkyne is  $\eta^2$ -coordinated to one copper atom in complexes **23** the  $\text{C}_2$  building block is  $\pi$ -bonded to two (bpy)Cu fragments. The bonding situation in compounds **23** is similar to that found in the alkyne-bridged dicopper(I) complexes of tropocoronand-6,6 [27,28].

Table 6  
Compounds **22** and **23**


Compound	$\text{RC}\equiv\text{CR}'$		X	NN <sup>a</sup>	Ref.
	R	R'			
<b>22a</b>	H	H	$\text{BF}_4$	NHpy <sub>2</sub>	[44,45]
<b>22b</b>	H	H	$\text{ClO}_4$	phen	[46]
<b>22c</b>	H	Ph	$\text{ClO}_4$	phen	[46]
<b>22d</b>	H	$\text{CO}_2\text{Et}$	$\text{ClO}_4$	phen	[46]
<b>22e</b>	H	OEt	$\text{ClO}_4$	phen	[46]
<b>22f</b>	H	n-C <sub>7</sub> H <sub>15</sub>	$\text{ClO}_4$	phen	[46]
<b>22g</b>	Ph	Ph	$\text{ClO}_4$	phen	[46]
<b>22h</b>	$\text{CO}_2\text{Me}$	$\text{CO}_2\text{Me}$	$\text{ClO}_4$	phen	[46]
<b>22i</b>	H	Ph	$\text{ClO}_4$	5-NO <sub>2</sub> -phen	[46]
<b>22j</b>	H	Ph	$\text{ClO}_4$	5-Cl-phen	[46]
<b>22k</b>	H	Ph	$\text{ClO}_4$	5-Me-phen	[46]
<b>22l</b>	H	Ph	$\text{ClO}_4$	5,6-Me <sub>2</sub> phen	[46]
<b>22m</b>	Et	Et	$\text{PF}_6$	bpy	[47]
<b>22n</b>	H	<sup>n</sup> Pr	$\text{PF}_6$	bpy	[47]
<b>22o</b>	Me	$\text{CO}_2\text{Me}$	$\text{PF}_6$	bpy	[47]
<b>22p</b>	Ph	Ph	$\text{PF}_6$	bpy	[47]
<b>22q</b>	$\text{CO}_2\text{Et}$	$\text{CO}_2\text{Et}$	$\text{PF}_6$	bpy	[47]
<b>22r</b>	H	Ph	$\text{SbF}_6$	bpy	[47]
<b>22s</b>	$\text{CO}_2\text{Et}$	$\text{CO}_2\text{Et}$	$\text{SbF}_6$	Me <sub>2</sub> bpy	[47]
<b>23a</b>	$\text{CO}_2\text{Et}$	$\text{CO}_2\text{Et}$	$\text{PF}_6$	bpy	[47]
<b>23b</b>	$\text{CO}_2\text{Et}$	$\text{CO}_2\text{Et}$	$\text{SbF}_6$	Me <sub>2</sub> bpy	[47]
<b>23c</b>	H	Ph	$\text{SbF}_6$	bpy	[47]

<sup>a</sup> NHpy<sub>2</sub> = 2-pyridylamine; phen = 1,10-phenanthroline; bpy = 2,2'-bipyridine.


In compounds **19–23** a bidentate chelating ligand is used for the stabilization of monomeric  $\eta^2$ -alkyne copper(I) moieties. In addition, the tris-chelating oxygen ligand  $\{(\eta^5\text{-C}_5\text{H}_5)\text{Co}[\text{P}(\text{O})\text{R}'_2]_3\}^-$  ( $\text{R}' = \text{OMe}, \text{O}^i\text{Pr}$ ) reacts with  $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{PF}_6$  and  $\text{RC}\equiv\text{CR}$  ( $\text{R} = \text{Ph}, \text{CO}_2\text{Me}$ ) to yield the  $\eta^2$ -alkyne copper(I) complexes of composition  $(\eta^2\text{-RC}\equiv\text{CR})\text{Cu}\{(\eta^5\text{-C}_5\text{H}_5)\text{Co}[\text{P}(\text{O})\text{R}'_2]_3\}$  (**24a**,  $\text{R} = \text{Ph}, \text{R}' = \text{O}^i\text{Pr}$ ; **24b**,  $\text{R} = \text{CO}_2\text{Me}, \text{R}' = \text{OMe}$ ), as shown by Lenders and Kläui [48]. The  $\eta^2$ -bonded  $\text{RC}\equiv\text{CR}$  ligand in **24** can be easily exchanged by carbon monoxide and alkenes [48].



Another type of a monomeric  $\eta^2$ -alkyne copper(I) compound can be obtained by the nucleophilic cleavage of dimeric type **A** molecules. Thus, the reaction of  $[(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CSiMe}_3)\text{CuCl}]_2$  (**2a**) and  $[(\eta^2\text{-tmtch})\text{CuCl}]_2$  (**2j**) with  $\text{Li}[\text{C}_5\text{R}_5]$  ( $\text{R} = \text{H}, \text{CH}_3$ ) leads to the formation of monomeric **25a–25c** [6,18,49].



**25a:**  $\text{R} = \text{H}$   
**25b:**  $\text{R} = \text{Me}$




**25c**

The species **25a–c** represent the first “mixed sandwich” compounds of copper(I), as members of a long isoelectronic series in which  $(\eta^5\text{-C}_5\text{R}_5)\text{M}$  building blocks interact with 2–5-electron donor ligands [6].

With the Lewis base  $\text{SMe}_2$ , the breakdown of the dimeric structure of  $[(\eta^2\text{-tmtch})\text{CuI}]_2$  (**21**) could be induced, forming the monomeric base-stabilized compound  $[(\eta^2\text{-tmtch})\text{Cu(I)SMe}_2]$  (**26**) [11].

Compound **26** is the first example of a type **E** molecule in which the  $\eta^2$ -alkyne copper(I) moiety is not stabilized by a bi- or tridentate chelating ligand.



**2.4.2.2. Spectroscopy and bonding.** As already pointed out in Section 2.1.3, the most remarkable feature in the IR spectra of the  $\eta^2$ -alkyne copper(I) complexes **16–26** is that these compounds all exhibit a significant  $\nu(\text{C}\equiv\text{C})$  absorption band shifted to lower wavenumbers than the corresponding  $\nu(\text{C}\equiv\text{C})$  vibration in the free acetylenes  $\text{RC}\equiv\text{CR}'$  [4–21,24–49,53–55]. This phenomenon is typical of  $\text{C}\equiv\text{C}$  triple bonds  $\eta^2$ -coordinated to transition metal centres [34,58–60].

In compounds **21** and **23**, the  $\text{C}_2$  unit of the alkyne ligands is  $\eta^2$ -coordinated to two copper(I) centres, thus acting as a four-electron donor. Hence the  $\nu(\text{C}\equiv\text{C})$  vibration is shifted to lower wavenumbers ( $1630\text{--}1750\text{ cm}^{-1}$ ) as compared with compounds **16–20**, **22** and **24–26** ( $1700\text{--}2000\text{ cm}^{-1}$ ) in which the  $\text{C}_2$  unit acts as a two-electron donor. This is consistent with the lengthening of the  $\text{C}\equiv\text{C}$  triple bond of the alkyne ligands from  $1.20\text{--}1.25\text{ \AA}$  in compounds **16–20**, **22** and **24–26** to  $\sim 1.32\text{ \AA}$  in compounds **21** and **23** (Fig. 4, Table 7).

Some selected examples of compounds **16–26** are shown in Fig. 4.

Fig. 4 shows that compounds of type **E** are monomeric. The copper atoms possess essentially a trigonal planar environment. As a result of the  $\eta^2$ -coordination of the alkyne ligands to copper(I), the  $\text{R}-\text{C}\equiv\text{C}-\text{R}'$  units are distorted compared with the free situation in which they are almost linear (Table 7) [11,18,27,28,41,43–46].

### 3. Organometallic substituted $\eta^2$ -alkyne copper(I) and silver(I) compounds

As shown in Section 2, alkynes ( $\text{RC}\equiv\text{CR}'$ ), 1,4-diynes ( $\text{RC}\equiv\text{C}-\text{E}-\text{C}\equiv\text{CR}$ ;  $\text{E}$  = main-group element building blocks such as  $\text{SiMe}_2$ ) and cyclic triynes (TBC) can be used for the breakdown of polymeric  $[\text{Cu}^{\text{I}}\text{X}]_n$  compounds to discrete  $[\text{Cu}^{\text{I}}\text{X}]_x$  ( $x = 1\text{--}4$ ) aggregates. However, the application of organometallic substituted acetylides  $\text{L}_n\text{M}-\text{C}\equiv\text{CR}$  or organometallic bridged 1,4-diynes  $\text{RC}\equiv\text{C}-\text{ML}_n-\text{C}\equiv\text{CR}$  ( $\text{R}$  = singly bonded organic ligand;  $\text{ML}_n$  = organometallic building block) afford other coordination modes, i.e. type **G–K** molecules.

As shown schematically in Scheme 3, cluster **G** contains as substructures type **H**, **J** and **K** molecules (routes (a)). Structural type **H** molecules can formally be broken down in to type **I** or **J**-molecules (route (b)). The reversion/re-building of this breakdown is the formation of a type **A** molecule from two structural type **K** compounds (route (c)).




Fig. 4. Selected X-ray structures of type E molecules. The interatomic bond distances and angles are given in Table 7.

This Section describes the synthesis and reactivity of all compounds of these structural types.


### 3.1. Type G molecules, $M_2M'_4(C\equiv CR)_8(PPh_3)_2$ ( $M=Ir, Rh, Pt$ ; $M'=Cu, Ag$ )

The reaction of Vaska's complex  $trans-(PPh_3)_2(CO)IrCl$  (**27a**) or  $(PPh_3)_3RhCl$  (**27b**) with  $[M^1C\equiv CR]_n$  yields  $\{[(PPh_3)_2(RC\equiv C)_2]M[(\eta^2-C\equiv CR)_2]M^1[PPh_3]\}$  ( $M=Ir, M^1=Cu$ ;  $M=Rh, M^1=Ag$ ) (type I molecules; see Section 3.3) as byproducts, together with the hexanuclear clusters  $M_2M'_4(C\equiv CR)_8(PPh_3)_2$  (**28**) ( $M=Ir, Rh$ ;  $M^1=Cu, Ag$ ) of type G [61–66].

In clusters of type G the six metal atoms (four Cu (Ag) and two Ir or Rh atoms) define an octahedron in which each iridium or rhodium atom is bonded apically to a phosphine ligand  $PR_3$ . Four  $RC\equiv C$  ligands are  $\sigma$ -bonded to each iridium or rhodium atom, and two  $RC\equiv C$  ligands (one from each Ir, Rh) form a  $\pi$  linkage with one of the four equatorial copper or silver atoms on the octahedron meridian.

Table 7  
X-ray data for compounds 16, 17, 19–22, 25 and 26

Compound	Ref.	Cu–X (Å)	Cu←    (Å)	C≡C (Å)	X–Cu–X (°)	C≡C–R (°)
16	[41]	2.549(5)	2.060(4)	1.22(1)	–	177.8(6)
17	[43]	1.955(5)–2.006(5)	1.991(8)–2.001(7)	–	94.4(3)–101.5(3)	163.0(8)–166.0(8)
19b	[54]	1.969(6)	1.959(8)	1.25(1)	93.8(2)	158.4(8)
		1.954(6)	1.956(8)			160.4(8)
20	[18]	2.004(2)	1.928(2)	1.248(3)	–	146.0(2)
		2.019(2)	1.942(2)			144.3(2)
21c	[27,28]	1.912(5); 1.901(5)	1.946(7); 1.938(7)	1.314(9)	84.5(2)	139.0(7)
		1.910(6); 1.926(6)	1.935(7); 1.952(7)		84.1(2)	138.2(4)
21d	[27]	1.912(3); 1.907(3)	1.937(4); 1.955(4)	1.320(6)	83.7(2)	138.7(4)
		1.907(3); 1.910(3)	1.961(4); 1.939(4)		84.0(2)	136.9(4)
22a	[44,45]	1.968(3)	1.971(4)	1.19(1)	96.8(2)	–
22b	[46]	1.979(4); 1.978(4)	1.930(5); 1.961(5)	1.190(7)	84.9(2)	–
22c	[46]	1.997(8); 2.000(7)	1.92(1); 1.99(1)	1.22(1)	85.3(3)	163(1)
22d	[46]	1.978(6); 1.977(6)	1.925(9); 1.934(7)	1.19(1)	84.5(2)	154.7(8)
25c	[18]	2.247(3); 2.168(2)	1.954(3); 1.937(2)	1.236(4)	–	148.1(2)
		2.251(3); 2.180(2)				146.4(2)
		2.169(3)				
26	[11]	2.508(1) (Cu–I) 2.278(2) (Cu–S)	1.975(6); 1.971(6)	1.241(8)	107.1(1)	148.9(6) 147.0(6)




Complexes **28**, i.e. clusters of type **G**, can be prepared in good yields by the nucleophilic substitution of the transition metal halides *trans*-(PPh<sub>2</sub>R)<sub>2</sub>(CO)MCl (**27a**, M = Ir, R = Ph; **27c**, M = Ir, R = Me; **27d**, M = Rh, R = Me; **27e**, M = Rh, R = Ph), (PPh<sub>3</sub>)<sub>3</sub>RhCl (**27b**) or (tht)<sub>2</sub>PtCl<sub>2</sub> (tht = tetrahydrothiophene) (**27f**) with [M<sup>I</sup>C≡CR]<sub>n</sub> (M<sup>I</sup> = Cu, Ag) (Table 8) [61–66].

Conspicuous in this method of synthesis is that the yields of clusters **28** decrease on changing from iridium to rhodium and from copper to silver. The clusters **28g–28n**, **28s** and **28t** (Table 8), containing rhodium, are apparently less stable than the corresponding iridium clusters. The lower stability of the rhodium clusters gives rise to acetylide couplings of two RC≡C ligands, generating symmetrically substituted 1,3-diynes [61–66]. It is possible to replace the PPh<sub>3</sub> ligand with PPh<sub>2</sub>Me in order to increase the solubility of clusters **28** (Table 8).

The hexanuclear clusters Ir<sub>2</sub>Cu<sub>4</sub>(PPh<sub>3</sub>)<sub>2</sub>(C≡CR)<sub>8</sub> (**28a**, R = Ph; **28b**, R = 4-MeC<sub>6</sub>H<sub>4</sub>) react with Fe<sub>2</sub>(CO)<sub>9</sub> in benzene to yield the octanuclear clusters Ir<sub>2</sub>Cu<sub>4</sub>Fe<sub>2</sub>(C≡CR)<sub>8</sub>(PPh<sub>3</sub>)<sub>2</sub>(CO)<sub>8</sub> (**29a**, R = Ph; **29b**, R = 4-MeC<sub>6</sub>H<sub>4</sub>) [64,65]. In **29** the Ir<sub>2</sub>Cu<sub>4</sub> skeleton is retained. A possible coordination mode of **29** is shown.

By the reaction of **27b** with [M<sup>I</sup>C≡CC<sub>6</sub>F<sub>5</sub>]<sub>n</sub> (M<sup>I</sup> = Ag) two further products, {[PPh<sub>3</sub>]<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>C≡C)<sub>2</sub>}Rh(η<sup>2</sup>-C≡CC<sub>6</sub>F<sub>5</sub>)<sub>2</sub>}Ag(PPh<sub>3</sub>) **39e** (type **I** molecule; see Section 3.3) and the trinuclear compound RhAg<sub>2</sub>(C≡CC<sub>6</sub>F<sub>5</sub>)<sub>5</sub>(PPh<sub>3</sub>)<sub>3</sub> (**30**), are formed [61,63–65].

The synthesis and reactivity of compound **39e** is described in Section 3.3.



Scheme 3. Substructural features of cluster G.

The structures of the heteronuclear clusters  $M_2M'_4(C\equiv CR)_8(PPh_3)_2$  ( $M=Ir, Rh$ ;  $M'=Cu, Ag$ ) (**28**) and  $RhAg_2(C\equiv CC_6F_5)_5(PPh_3)_3$  (**30**) were established by X-ray diffraction studies (Fig. 5) [61,63–65].

The four copper and two iridium atoms in **28a** form a distorted octahedron with the four copper atoms taking equatorial positions. Each iridium atom is surrounded by an apically bonded  $PPh_3$  group and four coplanar  $\sigma$ -bonded  $RC\equiv C$  ligands. It is also found that each acetylenic fragment simultaneously participates in an asymmetric  $\pi$  interaction with a copper atom on the octahedral meridian. Thus, each







Table 8  
Clusters 28

Compound	ML	M <sup>I</sup>	R	Ref.
28a	Ir(PPh <sub>3</sub> )	Cu	Ph	[61,62,64]
28b	Ir(PPh <sub>3</sub> )	Cu	4-MeC <sub>6</sub> H <sub>4</sub>	[64]
28c	Ir(PPh <sub>3</sub> )	Cu	4-FC <sub>6</sub> H <sub>4</sub>	[64]
28d	Ir(PPh <sub>3</sub> )	Cu	C <sub>6</sub> F <sub>5</sub>	[64]
28e	Ir(PPh <sub>2</sub> Me)	Cu	Ph	[64]
28f	Ir(PPh <sub>2</sub> Me)	Cu	C <sub>6</sub> F <sub>5</sub>	[64]
28g	Rh(PPh <sub>3</sub> )	Cu	Ph	[64]
28h	Rh(PPh <sub>3</sub> )	Cu	4-MeC <sub>6</sub> H <sub>4</sub>	[64]
28i	Rh(PPh <sub>3</sub> )	Cu	4-FC <sub>6</sub> H <sub>4</sub>	[64]
28j	Rh(PPh <sub>3</sub> )	Cu	C <sub>6</sub> F <sub>5</sub>	[64]
28k	Rh(PPh <sub>2</sub> Me)	Cu	C <sub>6</sub> H <sub>5</sub>	[64]
28l	Rh(PPh <sub>2</sub> Me)	Cu	4-MeC <sub>6</sub> H <sub>4</sub>	[64]
28m	Rh(PPh <sub>2</sub> Me)	Cu	4-FC <sub>6</sub> H <sub>4</sub>	[64]
28n	Rh(PPh <sub>2</sub> Me)	Cu	C <sub>6</sub> F <sub>5</sub>	[64]
28o	Pt	Cu	Ph	[66]
28p	Pt	Cu	<sup>t</sup> Bu	[66]
28q	Ir(PPh <sub>3</sub> )	Ag	Ph	[65]
28r	Ir(PPh <sub>3</sub> )	Ag	C <sub>6</sub> F <sub>5</sub>	[65]
28s	Rh(PPh <sub>3</sub> )	Ag	Ph	[63,65]
28t	Rh(PPh <sub>3</sub> )	Ag	C <sub>6</sub> F <sub>5</sub>	[63,65]
28u	Pt	Ag	Ph	[66]
28v	Pt	Ag	<sup>t</sup> Bu	[66]
28w	Pt	Au	<sup>t</sup> Bu	[66]

copper atom is  $\pi$ -bonded to two acetylenic ligands, one below and one above the equatorial Cu<sub>4</sub> plane.

As a result of the  $\eta^2$ -coordination of the alkynyl units to a copper atom, the C $\equiv$ C triple bonds of the RC $\equiv$ C building blocks are lengthened from approximately 1.20 Å in L<sub>n</sub>MC $\equiv$ CR compounds to 1.23 Å in **28a**. This fits to the IR spectroscopically



made observation that all clusters **28** show absorption bands due to  $\nu(\text{C}\equiv\text{C})$  vibrations in the  $2000\text{ cm}^{-1}$  region [61–66].

The formal oxidation states of the metal atoms in the heteronuclear clusters **28** are  $\text{Ir}^{\text{IV}}/\text{Rh}^{\text{IV}}$  combined with  $\text{Cu}^0/\text{Ag}^0$ , hence these clusters are electron rich, containing six electrons more than required to fill the metal bonding and non-bonding orbitals. These six electrons are assumed to be located in metal–metal antibonding orbitals, thus giving an overall bond order of  $3/4$  [61–67].




Fig. 5. Molecular geometry and atom labelling scheme for **28a** (left) and **30** (right) [61,63–65]. Selected bond distances (Å) and angles (°) are as follows. **28a**: Ir–Cu (av. (average)) 2.87(5), Cu–Cu (av. 2.74(7), Ir–P (av.) 2.26(1), Ir–C (av.) 2.04(4), Cu–C<sub>α</sub> (av.) 1.99(4), Cu–C<sub>β</sub> (av.) 2.19(4), C≡C (av.) 1.23(5); C<sub>α</sub>–Ir–C<sub>α</sub> (av.) 89.6(1), Ir–C<sub>α</sub>–C<sub>β</sub> (av.) 170(4), C≡C–Ph (av.) 165(3). **30**: Rh–Ag1 3.1021(8), Rh–Ag2 3.0871(8), Ag1–Ag2 5.087(1), Rh–C<sub>α</sub> (av.) 2.02, C=C (av.) 1.21(8), Ag1–C≡C(A) 2.642, Ag1–C≡C(B) 2.572, Ag1–C≡C(C) 2.534, Ag2–C≡C(A) 2.822, Ag2–C≡C(D) 2.751, Ag2–C≡C(E) 2.396; P3–Rh–C1A 173.8, C<sub>α</sub>–Rh–C<sub>α</sub> (av.) 89.6, C≡C(A)–Ag1–C≡C(B) 93.75, C≡C(B)–Ag1–C≡C(C) 89.79, C≡C(C)–Ag1–C≡C(A) 83.30, C≡C(A)–Ag2–C≡C(D) 81.77, C≡C(D)–Ag2–C≡C(E) 89.40, C≡C(E)–Ag2–C≡C(A) 94.20.


According to Fig. 5, the trinuclear complex  $\text{RhAg}_2(\text{C}\equiv\text{CC}_6\text{F}_5)_5(\text{PPh}_3)_3$  (**30**) prepared from  $(\text{Ph}_3\text{P})_3\text{RhCl}$  (**27b**) and  $[\text{AgC}\equiv\text{CC}_6\text{F}_5]_n$  contains a central rhodium atom, which is octahedrally coordinated by one  $\text{PPh}_3$  ligand (apical position) and five  $\sigma$ -bonded alkynyl moieties [61,63–65]. Three alkynyls are  $\eta^2$ -coordinated to an  $\text{Ag}[\text{PPh}_3]$  building block, one being involved with both silver atoms.

The rhodium–silver distances of 3.1021(8) and 3.0871(8) Å in **30** are considerably larger than those expected for rhodium–silver single bonds (2.79 Å) [61,67]. In the absence of a metal–metal bond, the metal atoms appear to be present in their most common oxidation states, i.e.  $\text{Rh}^{\text{III}}$  and  $\text{Ag}^{\text{I}}$ . Thus, the rhodium atom shows its expected regular octahedral coordination geometry and the silver atoms are tetrahedrally coordinated. Consequently, it seems that the compound  $\text{RhAg}_2(\text{C}\equiv\text{CC}_6\text{F}_5)_5(\text{PPh}_3)_3$  (**30**) is zwitterionic, consisting of a  $[(\text{Ph}_3\text{P})\text{Rh}^{\text{III}}-(\text{C}\equiv\text{CC}_6\text{F}_5)_5]^{2-}$  unit and two  $[(\text{Ph}_3\text{P})\text{Ag}^{\text{I}}]^+$  moieties. These components are kept together by  $\eta^2$ -coordination of the alkynyl units to  $[\text{Ag}(\text{PPh}_3)]^+$  building blocks.

### 3.2. Type **H** molecules, $\{[L_n\text{M}(\text{C}\equiv\text{CR})_2]\text{M}^{\text{I}}\text{X}\}_2$


Another common feature of complexes isolated from reactions involving transition metal halides  $L_n\text{MCl}$  and copper(I) or silver(I) acetylides  $[\text{M}^{\text{I}}\text{C}\equiv\text{CR}]_n$  ( $\text{M} = \text{Cu}$ ,  $\text{Ag}$ ) (Section 3.1) are molecules of structural type **H**.

Type **H** molecules can be considered as substructures of clusters **28** (type **G** molecules, Section 3.1) in which the two bis(alkynyl) complex fragments  $L_nM(C\equiv CR)_2$  are bridged by two  $M'X$  ( $M = Cu, Ag$ ) building blocks.



H

A procedure for the synthesis of a type **H** molecule containing ruthenium and copper as metal atoms is given by the reaction of  $(\eta^5\text{-C}_5\text{H}_5)(\text{PPh}_3)_2\text{RuCl}$  (**27g**) with  $[\text{CuC}\equiv\text{CC}_6\text{F}_5]_n$  [68].



31

Another route to structural type **H** molecules is the reaction of the square-planar  $d^8$  platinum ( $\text{Pt}^{\text{II}}$ ) compounds  $A_2[\text{Pt}_2(\mu\text{-X}')_2(\text{C}_6\text{X}_5)_4]$  (**32**) ( $A = \text{NBu}_4, \text{PMePh}_3$ ;  $X' = \text{Cl, I}$ ;  $X = \text{F, Cl}$ ) with  $[\text{AgC}\equiv\text{CR}]_n$  ( $R = \text{Ph, 'Bu}$ ) [69,70].

Compounds **31a–31g** can also be synthesized by treatment of  $A_2[\text{cis}(\text{C}_6\text{X}_5)_2\text{Pt}(\text{C}\equiv\text{CR})_2]$  (**33**) ( $A = \text{NBu}_4, \text{PMePh}_3$ ;  $R = \text{Ph, 'Bu}$ ;  $X = \text{F, Cl}$ ) with  $[\text{Cu}^{\text{I}}\text{Cl}]_n$ ,  $[\text{Ag}^{\text{I}}\text{Cl}]_n$  or  $[\text{Ag}^{\text{I}}\text{ClO}_4]_n$  [70].

All compounds synthesized by these methods are listed in Table 9.

The most remarkable feature in the IR spectra of compounds **31a–31g** is that they




Table 9  
Compounds **31a–31g** [70]

Compound	M	X	R	A
<b>31a</b>	Ag	F	Ph	NBu <sub>4</sub>
<b>31b</b>	Ag	F	<sup>t</sup> Bu	NBu <sub>4</sub>
<b>31c</b>	Ag	Cl	Ph	NBu <sub>4</sub>
<b>31d</b>	Ag	Cl	<sup>t</sup> Bu	NBu <sub>4</sub>
<b>31e</b>	Ag	F	Ph	PMePh <sub>3</sub>
<b>31f</b>	Cu	F	Ph	PMePh <sub>3</sub>
<b>31g</b>	Cu	F	<sup>t</sup> Bu	NBu <sub>4</sub>


all exhibit a  $\nu(C\equiv C)$  absorption band in the 2035–2000  $cm^{-1}$  region [70]. This is typical for  $C\equiv C$  triple bonds  $\eta^2$ -coordinated to a copper(I) or silver(I) fragment (see above).

The X-ray crystal structure analysis which was carried out on representative compound **31a** shows that two square-planar platinum building blocks,  $cis-(C_6F_5)_2Pt(C\equiv CPh)_2$ , are linked together by two silver atoms. The  $C_2$  units of the phenylethynyl ligands are  $\eta^2$ -coordinated to the silver atoms in such a way that these two  $PhC\equiv C$  ligands are asymmetrically  $\pi$ -bonded to each silver atom.

As a consequence of the  $\eta^2$ -coordination of the  $PhC\equiv C$  group to silver, the alkynyl ligands are not linear ( $Pt-C\equiv C$ , 176.3°, 176.6°;  $C\equiv C-Ph$ , 173.4°, 172.8°) [70]. The  $C_2$  units, associated with each silver atom, are oriented perpendicular to one another. The  $Pt\cdots Ag$  distances of 3.10 and 3.15 Å are consistent with the absence of bonding character. Furthermore, the  $Ag\cdots Ag$  distance of 2.939 Å is at the high end of the range observed for silver–silver systems (2.740–3.085 Å) [71].

Molecules of structural type **H** can be considered as dimeric species of type **I** molecules (for a detailed discussion of structural type **I** molecules, see Section 3.3).


Thus, the anionic tetranuclear platinum–silver compounds **31a** and **31b** react with  $\text{PR}'_3$  ( $\text{R}' = \text{Ph}$ , Et) to give the anionic platinum–silver compounds **34** in which the alkynyl groups act as  $\sigma, \pi$  bridges between platinum and silver atoms [69].




The structures of compounds **34** were established by a single-crystal diffraction study of **34a** [69]. The structure of complex **34a** reveals that the two metal atoms (platinum and silver) are asymmetrically bridged by two phenylethynyl groups, each of which forms a  $\sigma$  bond to platinum and a  $\pi$  bond to silver.

### 3.3. Type I molecules, $\{\text{L}_n\text{M}(\text{C}\equiv\text{CR})_2\}\text{M}^{\text{I}}\text{X}$ ( $n^{\text{I}} = \text{Cu}, \text{Ag}$ )

In general, organometallic bis( $\sigma$ -alkynyl) compounds of type **L** can be used for the synthesis of binuclear complexes (type **I** molecules) containing  $\sigma, \pi$ -alkynyl ligands as bridging groups between two metal atoms.




Compounds of type  $\text{RC}\equiv\text{C}-\text{L}_n\text{M}-\text{C}\equiv\text{CR}$  ( $\text{L}_n\text{M} = (\eta^5\text{-C}_5\text{H}_4\text{R}')_2\text{Ti}$ , etc.) (**L**) can be considered as organometallic  $\pi$  tweezers [34,72,73]. In compounds of type **I**, the  $\text{RC}\equiv\text{C}-\text{L}_n\text{M}-\text{C}\equiv\text{CR}$  fragments act as host molecules with the  $\text{M}^{\text{I}}\text{X}$  ( $\text{M} = \text{Cu}, \text{Ag}$ ) building blocks as guests. The aim is to use simultaneously the steric stabilization of the organic groups **R** and the chelating effect of type **L** compounds for the preparation of stable type **I** molecules.



### 3.3.1. Synthesis

There are three general routes for the synthesis of type I molecules: (a) direct synthesis from complexes of type L, (b) nucleophilic substitution of  $\text{L}_n\text{M}(\text{PPh}_3)\text{Cl}$  and  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}'\text{X}$  ( $\text{X}=\text{Cl}$ ,  $\text{OSO}_2\text{CF}_3$ ) compounds and (c) breakdown of dimeric complexes  $(\text{NBu}_4)_2[\text{Pt}_2\text{Ag}_2(\text{C}\equiv\text{CR})_4(\text{C}_6\text{F}_5)_4]$  under the influence of Lewis bases.

(a) *Direct synthesis route.* The bis(alkynyl) titanocenes  $\text{Me}_3\text{SiC}\equiv\text{C}-\text{L}_2\text{Ti}-\text{C}\equiv\text{CSiMe}_3$  ( $\text{L}=\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$ ) (**35**) [74] have been used as organometallic chelating ligands (organometallic  $\pi$  tweezers) for the synthesis of  $\{\text{L}_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{M}'\text{X}$  compounds (**36**,  $\text{M}'=\text{Cu}$ ; **37**,  $\text{M}'=\text{Ag}$ ;  $\text{X}=\text{singly bonded ligand}$ ) (Table 10) [13,25,30,34–40]. They are best synthesized using the appropriate  $[\text{M}'\text{X}]_n$  starting materials at  $25^\circ\text{C}$  in thf or  $\text{Et}_2\text{O}$  (Table 10).




In a similar manner, ansa-[bis(alkynyl)] titanocenes react with  $[\text{CuCl}]_n$  to yield the binuclear complexes **38a–38c** [75].

In general, the chelate effect of type L compounds,  $\text{RC}\equiv\text{C}-\text{L}_n\text{M}-\text{C}\equiv\text{CR}$  [76], is suitable for stabilizing low-valent  $\text{M}(\text{CO})$ ,  $\text{M}[\text{PR}_3]$  ( $\text{M}=\text{Ni}$  [77–79],  $\text{Co}$  [79,80],


Table 10

Compounds  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{M}^1\text{X}$  (36, 37)

Compound	M <sup>1</sup> X	Ref.	Compound	M <sup>1</sup> X	Ref.
36a	CuCl	[82,85]	37a	AgCl	[36]
36b	CuBr	[40,85]	37b	AgBr	[36]
36c	CuI	[40,85]	37c	AgI	[36]
36d	CuCN	[40,85]	37d	AgCN	[36]
36e	CuSCN	[40,85]	37e	AgSCN	[36]
36f	CuOC(O)Me	[13,14]	37f	AgNO <sub>2</sub>	[36]
36g	CuOC(O)Ph	[13,14]	37g	AgNO <sub>3</sub>	[36]
36h	CuC≡CPh	[37,40]	37h	AgClO <sub>4</sub>	[36]
36i	CuC≡CSiMe <sub>3</sub>	[37,40]	37i	AgS <sub>2</sub> C(NEt <sub>2</sub> )	[36]
36j	CuC≡C <sup>t</sup> Bu	[37,40]	37j	AgO <sub>2</sub> CMe	[13,14]
36k	CuPh	[38]	37k	AgO <sub>2</sub> CPh	[13,14]
36l	CuTol <sup>a</sup>	[38]	37l	AgMes <sup>b</sup>	[38]
36m	CuMes <sup>b</sup>	[38]	37m	AgOTf <sup>d</sup>	[38]
36n	CuCH <sub>2</sub> SiMe <sub>3</sub>	[86]	37n	AgBF <sub>4</sub>	[13]
36o	CuSR <sup>c</sup>	[86]			
36p	CuOTf <sup>d</sup>	[38,40]			
36q	CuBF <sub>4</sub>	[38,40]			
36r	CuPF <sub>6</sub>	[38,40]			
36s	Cu(acac)	[13]			

<sup>a</sup> Tol = 4-MeC<sub>6</sub>H<sub>4</sub>.<sup>b</sup> Mes = 2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub>.<sup>c</sup> R = 2-Me<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>.<sup>d</sup> OTf = trifluoromethanesulphonate, OSO<sub>2</sub>CF<sub>3</sub>.

38a: R = H

38b: R = SiMe<sub>3</sub>


38c

Pt [81]) and  $\text{M}(\text{C}_6\text{F}_5)_2$  ( $\text{M} = \text{Pd}, \text{Pt}$ ) [69] moieties. In addition, the organometallic 1,4-diynes  $\text{RC}\equiv\text{C}-\text{L}_n\text{M}-\text{C}\equiv\text{CR}$  can be used for the stabilization of monomeric  $\text{MCl}_2$  ( $\text{M} = \text{Fe}, \text{Co}, \text{Ni}$ ) [82,83] and  $\text{Mg}(\text{Cl})(\text{thf})$  [84] units.


(b) *Nucleophilic substitution route.* Compounds of type I are also formed by the reaction of  $[(\text{Ph}_3\text{P})_2\text{ML}_n]\text{Cl}$  (**27a**,  $\text{ML}_n = \text{Ir}(\text{CO})$  [64,65]; **27b**,  $\text{ML}_n = \text{Rh}(\text{PPh}_3)$  [64,65]; **27h**,  $\text{ML}_n = \text{Re}(\text{CO})_3$  [87,88]; **27g**,  $\text{ML}_n = \text{Ru}(\eta^5\text{-C}_5\text{H}_5)$  [87,88]) with  $[\text{M}^1\text{C}\equiv\text{CR}]_n$  ( $\text{M} = \text{Cu}, \text{Ag}$ ) (Table 11).

Table 11  
Compounds 39

Compound	[ML' <sub>n</sub> ]	R	M <sup>I</sup>	Ref.
39a	Re(CO) <sub>3</sub> (PPh <sub>3</sub> )	C <sub>6</sub> F <sub>5</sub>	Cu	[87,88]
39b	Ru(η <sup>5</sup> -C <sub>5</sub> H <sub>5</sub> )	<i>p</i> -C <sub>6</sub> H <sub>4</sub> Me	Cu	[87,88]
39c	Ru(η <sup>5</sup> -C <sub>5</sub> H <sub>5</sub> )	<i>p</i> -C <sub>6</sub> H <sub>4</sub> F	Cu	[87,88]
39d	Ir(PPh <sub>3</sub> ) <sub>2</sub> (C≡CR) <sub>2</sub>	C <sub>6</sub> F <sub>5</sub>	Cu	[64,65]
39e	Rh(PPh <sub>3</sub> ) <sub>2</sub> (C≡CR) <sub>2</sub>	C <sub>6</sub> F <sub>5</sub>	Ag	[64,65]
39f	Ir(PPh <sub>3</sub> ) <sub>2</sub> (C≡CR) <sub>2</sub>	C <sub>6</sub> F <sub>5</sub>	Ag	[64,65]



(c) *Breakdown of dimeric complexes with Lewis bases.* Another route to type I molecules is given by the reaction of the tetranuclear platinum–silver complex (NBu<sub>4</sub>)<sub>2</sub>[Pt<sub>2</sub>Ag<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>(C≡CR)<sub>4</sub>] (**31a**, R = Ph; **31b**, R = <sup>t</sup>Bu) with PR'<sub>3</sub> (R' = Ph, Et) [69]. The binuclear complexes **34** are formed in good yields (see also Section 3.2).



In general, the organometallic 1,4-diynes RC≡C–L<sub>n</sub>M–C≡CR (L<sub>n</sub>M = see above) can be used as organometallic chelating ligands for the stabilization of M<sup>I</sup>X (M = Cu, Ag) units (Tables 10 and 11).

The binuclear compounds {L<sub>n</sub>M(C≡CR)<sub>2</sub>}MX (**34** and **36–39**) are the first exam-

ples of stable molecules containing monomeric bis( $\eta^2$ -alkyne) $M^IX$  ( $M = Cu, Ag$ ) species.

### 3.3.2. Spectroscopy and bonding

In binuclear compounds of type  $\{L_nM(C\equiv CR)_2\}M^IX$  (**34** and **36–39**), the organo-metallic bis(alkynyl) building blocks  $L_nM(C\equiv CR)_2$  [ $L_nM = (\eta^5-C_5H_4SiMe_3)_2Ti$ ,  $Re(CO)_3(PPh_3)_3$ ,  $(\eta^5-C_5H_5)Ru$ ,  $Ir(PPh_3)_2(C\equiv CC_6F_5)_2$ ,  $Rh(PPh_3)_2(C\equiv CC_6F_5)_2$ ] act, through their alkynyl groups, as organometallic chelating ligands to the  $Cu^IX$  and  $Ag^IX$  entities. Spectroscopic data (IR,  $^1H$ ,  $^{13}C$  NMR) of compounds **34** and **36–39** also indicate that both alkynyl ligands are  $\eta^2$ -coordinated to  $M^IX$  moieties (Table 12).

In the IR spectra, the  $\nu(C\equiv C)$  absorption decreases from  $2012\text{ cm}^{-1}$  in the parent compound  $(\eta^5-C_5H_4SiMe_3)_2Ti(C\equiv CSiMe_3)_2$  (**35**) to  $1850\text{--}1960\text{ cm}^{-1}$  in compounds **36** and **37** (Table 12). This is in agreement with the increasing participation of the backbonding component in alkyne to copper (silver) bonding.

As Table 12 shows, changing the monoanionic ligand  $X$  in  $\{(\eta^5-C_5H_4SiMe_3)_2Ti(C\equiv CSiMe_3)_2\}M^IX$  ( $M = Cu$ , **36**;  $M = Ag$ , **37**) from a weaker (e.g.  $X = OSO_2CF_3$ ) to a stronger  $\sigma$  donor (e.g.  $X = 2,4,6-Me_3C_6H_2$ ) increases the back-donation component of the  $\eta^2$ -alkyne–metal interaction in the total bonding. The reduction of the  $Ti-C\equiv C$  and  $C\equiv C-Si$  angles from  $177/176.5^\circ$  in the parent compound  $(\eta^5-C_5H_4SiMe_3)_2Ti(C\equiv CSiMe_3)_2$  (**35**) to  $171.8/167^\circ$  in **37m**,  $166.6/162.9^\circ$  in **37i**,  $166.8/157.4^\circ$  in **36p** and  $163.7/155.5^\circ$  in **36m** also supports this conclusion, i.e. bending of the substituents of the  $C_2$  unit increases with increasing  $\sigma$  donation of the ligand  $X$  in  $\{(\eta^5-C_5H_4SiMe_3)_2Ti(C\equiv CSiMe_3)_2\}M^IX$ . Further, it is found that

Table 12  
Selected IR and  $^{13}C\{^1H\}$  NMR data for compounds **35–37**

Compound	IR $\nu(C\equiv C)$ ( $\text{cm}^{-1}$ )	$^{13}C\{^1H\}$ NMR $\delta C\equiv C$ (ppm)	Ref.
<b>35</b>	2012	$C_\beta$ 135.4; $C_\alpha$ 172.5	[74]
<b>36d</b>	1901	$C_\beta$ 132.8; $C_\alpha$ 173.6	[13,85]
<b>36f</b>	1901	$C_\beta$ 132.4; $C_\alpha$ 170.9	[13,14]
<b>36g</b>	1903	—	[13,14]
<b>36m</b>	1856	$C_\beta$ 125.0; $C_\alpha$ 201.8	[38]
<b>36n</b>	1930	$C_\beta$ 125.4; $C_\alpha$ 202.9	[86]
<b>36o</b>	1897	$C_\beta$ 120.5; $C_\alpha$ 180.4	[86]
<b>36p</b>	1923	$C_\beta$ 140.2; $C_\alpha$ 162.2	[86]
<b>36r</b>	1934	$C_\beta$ 136.2; $C_\alpha$ 163.2	[38,40]
<b>37j</b>	1944	$C_\beta$ 137.5; $C_\alpha$ 155.8	[13,14]
<b>37k</b>	1951	$C_\beta$ 137.4 (d, $J_{AgC} = 15\text{ Hz}$ ) $C_\alpha$ 156.3 (d, $J_{AgC} = 15\text{ Hz}$ )	[13,14]
<b>37l</b>	1902	$C_\beta$ 125.8 (d, $J_{AgC} = 7\text{ Hz}$ ) $C_\alpha$ 184.5 (d, $J_{AgC} = 3\text{ Hz}$ )	[38]
<b>37m</b>	1956	$C_\beta$ 144.8; $C_\alpha$ 151.8	[38]
<b>37n</b>	1948	$C_\beta$ 127.2; $C_\alpha$ 149.3	[13]

the  $\eta^2$ -alkyne–copper bonding interaction is stronger than the corresponding  $\eta^2$ -alkyne–silver bonding interaction (Table 12) [38]: complexes **37** readily exchange silver(I) for copper(I) building blocks, thus forming compounds **36** [38].

The  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra of compounds **36** and **37** remain essentially unchanged in the temperature range 200–317 K and the data are consistent with monomeric species in solution (as confirmed by molecular weight determinations in solution by cryoscopy).

The silver compound  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{AgMes}$  (Mes = 2,4,6- $\text{Me}_3\text{C}_6\text{H}_2$ ) (**37l**) provides direct information concerning the Ag–C bonding of the  $\eta^1$ -bonded  $\text{C}_{ipso}$  atom of the mesityl unit. The  $\text{C}_{ipso}$  mesityl resonance is observed at 168.3 ppm as two doublets due to coupling with  $^{107}\text{Ag}$  and  $^{109}\text{Ag}$ :  $^1J(^{107}\text{Ag}, ^{13}\text{C}) = 142\text{ Hz}$  and  $^1J(^{109}\text{Ag}, ^{13}\text{C}) = 164\text{ Hz}$  [38]. The fact that these  $J(\text{Ag}, \text{C})$  couplings are observed indicates that the Ag– $\text{C}_{ipso}$  bond is stable and does not dissociate on the NMR time-scale. These data are the first  $^{107,109}\text{Ag}$  coupling constant values for a two-electron, two-centre ( $2e\text{--}2c$ ) silver–carbon bond;  $^3J(\text{Ag}, \text{C})$  of 7 Hz is also observed for the hydrogen-bearing carbon atoms of the mesityl ligands. Examples of  $^1J(\text{Ag}, \text{C})$  values have been reported for two-electron, three-centre bonded aryl silver compounds (118.3, 136.9 Hz in  $[\text{Ag}_2\text{Li}_2(2\text{-NMe}_2\text{CH}_2\text{C}_6\text{H}_4)_4]$  and 115.0, 132.0 Hz in  $[\text{Ag}_2\text{Li}_2(\text{C}_6\text{H}_5)_4]$  for  $^{107}\text{Ag}$  and  $^{109}\text{Ag}$ , respectively [89,90]). The larger silver–carbon coupling constant found in **37l** reflects the anticipated higher degree of s-orbital participation in the silver–carbon bond in this complex than in the electron-deficient Ag–C interactions in  $[\text{Ag}_2\text{Li}_2(2\text{-NMe}_2\text{CH}_2\text{C}_6\text{H}_4)_4]$  and  $[\text{Ag}_2\text{Li}_2(\text{C}_6\text{H}_5)_4]$  [89,90]. Interestingly, Ag– $^{13}\text{C}$  couplings are also seen for the two carbon signals of the  $\eta^2$ -bonded alkynyl ligands. The signal at 184.5 ppm ( $\text{Ti}\text{--}\text{C}\equiv\text{C}$ ) occurs as a doublet resulting from a  $^1J(\text{Ag}, ^{13}\text{C})$  splitting of 3 Hz, while the doublet at 125.8 ppm ( $\text{C}\equiv\text{C}\text{--}\text{Si}$ ) has a  $^1J(\text{Ag}, ^{13}\text{C})$  splitting of 7 Hz [38,86].

The structure in the solid state of selected examples from compounds **34–39** have been determined by X-ray diffraction (Table 13).

Fig. 6 shows that compounds of type **I** are monomeric. The metal atoms M (M = Cu, Ag) of the  $\text{M}'\text{X}$  building blocks (where X is a unidentate ligand) show essentially a trigonal planar environment comprising  $\eta^2$ -coordination of both alkynyl groups from the  $\text{L}_n\text{M}(\text{C}\equiv\text{CR})_2$  fragment and  $\eta^1$ -bonding of the ligand X. In complexes **36s** and **37f** (Table 13), where X is a bidentate ligand, the copper or silver atoms have a tetrahedral environment in which the ligand X, i.e. X = acetylacetonate (acac) (**36s**) or  $\text{NO}_2$  (**37f**), is almost perpendicular to the titanium–alkynyl–copper (**36s**) or the titanium–alkynyl–silver (**37f**) plane (interplanar angle  $\sim 87^\circ$ ).

As a result of the  $\eta^2$ -coordination of the alkynyl units in **36–38** to the  $\text{M}'\text{X}$  moiety, the angle  $\text{C}_\alpha\text{--Ti--C}_\alpha'$  between the  $\sigma$ -bonded alkynyl ligands is considerably smaller than that in  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  (**35**), where it is  $102.8^\circ$  (Table 13). The  $\text{Ti}\text{--}\text{C}\equiv\text{C}\text{--}\text{SiMe}_3$  units are distorted compared with the starting compounds, in which they are linear (Table 13).

The  $\eta^2$ -coordination of the alkynyl ligands to the  $\text{M}'\text{X}$  moiety in **36** and **37** is accompanied by a bond lengthening of the  $\text{C}\equiv\text{C}$  triple bonds of the  $\text{Me}_3\text{SiC}\equiv\text{C}$  building blocks from 1.203 and 1.214 Å in **35** to 1.23–1.27 Å in **36** and **37** (Table 13). It is found that the  $\text{C}\equiv\text{C}$  triple bond is not significantly lengthened upon changing

Table 13  
X-ray data for compounds  $\{L_nM(C\equiv CR)_2\}M'X$ , 35, 36, 37 and 39

Compound	Ref.	M...M' (Å)	M—C $\equiv$ C (Å)	M'—C $\equiv$ C (Å)	C $\equiv$ C (Å)	C—M—C (°)	M—C $\equiv$ C (°)	C $\equiv$ C—R (°)
<b>35</b>	[74]	—	2.124(5), 2.103(5)	—	1.214(6)	102.8(2)	175.8(4)	174.8(4)
<b>36a</b>	[82,85]	290.9(3)	209(1), 210.4(9)	C $_4$ : 205.5(9), 218(1) C $_5$ : 204(1), 221(1)	1.203(9) 1.23(2)	89.6141	178.2(5) 167.8(9)	178.3(5) 161(1)
<b>36f</b>	[13,14]	2.930(3)	2.10(1), 2.09(1)	C $_4$ : 2.08(1); 2.07(1) C $_5$ : 2.15(1); 2.12(1)	1.24(2) 1.25(2)	90.3(4)	168.6(9) 163.6(9)	163(1) 164.1
<b>36m</b>	[38]	2.941(2)	2.079(2)	C $_5$ : 2.067(2); C $_6$ : 2.083(2)	1.23(2)	—	166(1)	165.1
<b>36p</b>	[38,40]	2.998(1)	2.105(4), 2.110(4)	C $_4$ : 2.066(4); 2.066(4) C $_5$ : 2.143(5); 2.158(5)	1.250(3) 1.233(6)	89.30(9) 87.4(2)	163.6(2) 166.7(3)	155.5(2) 156.18(4)
<b>36s</b>	[13]	2.920(1)	2.081(4), 2.091(4)	C $_4$ : 2.079(4); 2.088(3) C $_5$ : 2.171(4); 2.145(3)	1.238(5) 1.235(5)	90.7(1)	166.9(3) 164.7(3)	158.1(4) 163.8(3)
<b>37f</b>	[36]	3.162(5)	2.12(2), 2.10(2)	C $_4$ : 2.29(2); 2.33(2) C $_5$ : 2.41(2); 2.43(2)	1.227(5) 1.26(2)	93.8(1)	165.8(3) 171(2)	164.9(3) 170(2)
<b>37l</b>	[38]	3.104(7)	2.090(8)	C $_5$ : 2.270(9) C $_6$ : 2.305(9)	1.27(3) 1.24(1)	—	170(2) 166.6(6)	171(2) 162.9(7)
<b>37m</b>	[38]	3.156(2)	2.141(6), 2.159(7)	C $_4$ : 2.291(6); 2.311(6) C $_5$ : 2.417(7); 2.478(7)	1.22(1) 1.225(8)	93.8(2)	170.8(5) 172.7(5)	163.7(3) 170.2(5)
<b>37n</b>	[38]	3.096(2)	2.114(7), 2.128(7)	C $_4$ : 2.294(7); 2.293(7) C $_5$ : 2.472(8); 2.447(7)	1.23(1) 1.222(9)	95.6(3)	170.7(6) 171.0(6)	168.8(6) 166.8(7)
<b>39a</b>	[87,88]	3.078(3)	2.39(2), 2.26(2)	C $_4$ : 1.98(2); 2.12(2) C $_5$ : 2.33(2); 2.34(2)	—	—	—	—




Fig. 6. Selected X-ray structures of type I molecules. The interatomic bond distances and angles are given in Table 13.

the  $\sigma$ -donating properties of the ligand X (Table 13). However, spectroscopic studies (IR and  $^{13}\text{C}$  NMR) show a clear correlation: the  $^{13}\text{C}$  chemical shift of the alkyne units is almost linear with the  $\text{C}\equiv\text{C}$  stretching frequency [38,86].

As a result of the steric constraints, and also the overlap of the orbitals participating in the bonding of the group 11 metal atoms with the alkynyl groups of the  $\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  entity, the two alkynyl ligands are in-plane bonded [38,86]. A similar bonding arrangement is present in the structure of  $[\text{cyclo}(\text{o-C}_6\text{H}_4\text{C}\equiv\text{C})_3]\text{Cu}(\text{OSO}_2\text{CF}_3)$  (**16**) ( $\text{C}\equiv\text{C}$  1.222 Å,  $\text{Cu}-\text{C}$  (mean) 2.060 Å,  $\text{Cu}$  0.18 Å above the  $[(\text{C}\equiv\text{C})_3]$  plane) (Section 2.4.1) [41–43].

In compounds **36** and **37** the central Group 11 metal has a formal 16-electron count and consequently the  $\text{Ti}-\text{Cu}$  and  $\text{Ti}-\text{Ag}$  distances ( $\text{Ti}-\text{Cu}$ , 2.92–2.99 Å;  $\text{Ti}-\text{Ag}$ , 3.10–3.26 Å) are considered to represent non-bonding situations [38,86].


The titanium–copper distances in **36** (see above) are shorter than the similar titanium–silver distances in **37**, owing to the smaller ionic radius of copper(I) (0.96 Å) than that of silver(I) (1.26 Å). Similar observations have been made for the copper to alkyne and silver to alkyne distances in the isostructural compounds  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}(\text{Mes})$  (**36m**)/ $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Ag}(\text{Mes})$  (**37l**) ( $\text{Mes}=2,4,6\text{-Me}_3\text{C}_6\text{H}_2$ ) and  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}(\text{OSO}_2\text{CF}_3)$  (**36p**)/ $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Ag}(\text{OSO}_2\text{CF}_3)$  (**37m**) (Table 13). It was found that the copper to alkyne interatomic distances were approximately 0.25 Å shorter than the corresponding bond lengths in the silver complexes, reflecting the smaller ionic radius of  $\text{Cu}^{\text{I}}$  over that of  $\text{Ag}^{\text{I}}$  (see above). Similarly, the copper–carbon distance ( $\text{C}_{\text{ipso}}-\text{Mes}$ ) in **36m** of 1.947 Å is 0.15 Å shorter than the corresponding silver–carbon ( $\text{C}_{\text{ipso}}-\text{Mes}$ ) distance of 2.099 Å in **37l** (Table 13).

The  $\text{Cu}-\text{C}_{\text{ipso}}$  and  $\text{Ag}-\text{C}_{\text{ipso}}$  mesityl bond lengths are in the range expected for two-electron, two-centre (2e–2c) bonding of carbon to copper. Reported (2e–2c)  $\text{Cu}-\text{C}$  distances include that for  $\text{sp}^3$ -hybridized carbon atoms, 2.04(1) Å in  $(\text{Ph}_3\text{P})_3\text{CuMe}$  [3b],  $\text{sp}^2$ -hybridized carbon atoms, 2.020(4) Å in  $\text{MeC}(\text{CH}_2\text{PPh}_2)_3\text{CuPh}$  [3a], 1.916(3) Å in  $\text{Cu}(2,4,6\text{-}^i\text{Bu}_3\text{C}_6\text{H}_2)(\text{Me}_2\text{S})$  [3c] and 1.890(6) Å in  $\text{Cu}(2,4,6\text{-Ph}_3\text{C}_6\text{H}_2)$  [3f], and  $\text{sp}$ -hybridized carbon atoms, 1.898(3) Å in  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}-\text{C}\equiv\text{CSiMe}_3$  (**36i**) [86]. A similar observation is valid for the analogous binding of carbon to silver (the  $\text{Ag}-\text{C}$  bond length for a  $\text{sp}^2$ -hybridized carbon is 1.902(5) Å in  $\text{Ag}(2,4,6\text{-Ph}_3\text{C}_6\text{H}_2)$  [3f]).

These results show that the striking property of  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  (**35**) to bind mononuclear metal fragments gives access to a novel type of mononuclear organo-copper and -silver species (type **I** molecules) with an organometallic chelating bis(alkynyl) ligand  $\text{RC}\equiv\text{C}-\text{L}_2\text{Ti}-\text{C}\equiv\text{CR}$ . It is important that in the  $\text{L}_2\text{Ti}(\text{C}\equiv\text{CR})_2-\text{CuX}$  interaction (see above) copper represents a metal with a low Lewis acidity (electrophilicity). Metals with a higher electrophilicity (e.g.  $\text{R}_2\text{Pt}$ ) tend to coordinate preferentially to the  $\text{C}_\alpha$  atom of the  $\text{Ti}(\text{C}\equiv\text{CR})_2$  fragment, giving rise to a bonding, better described as asymmetric bridging of  $\text{C}\equiv\text{CR}$  between the titanium and the platinum centres (a type **M** molecule) [91].


### 3.3.3. Reactivity

The binuclear compounds  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{M}^{\text{I}}\text{X}$  (**36a**,  $\text{M}=\text{Cu}$ ,  $\text{X}=\text{Cl}$ ; **36p**,  $\text{M}=\text{Cu}$ ,  $\text{X}=\text{OSO}_2\text{CF}_3$ ; **37m**,  $\text{M}=\text{Ag}$ ,  $\text{X}=\text{OSO}_2\text{CF}_3$ ) contain, as func-



tional ligands, a chloride or trifluoromethanesulphonate (OTf) group at the copper or silver atom. Substitution of these groups should lead to a versatile reaction chemistry.

**3.3.3.1.  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuCl}$ .** To study intermolecular vs. intra-molecular alkyne to copper coordination, the mononuclear bis( $\eta^2$ -alkyne)- $\eta^1$ -acetylide copper(I) compounds  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuC}\equiv\text{CR}$  (**36h**, R = Ph; **36i**, R = SiMe<sub>3</sub>; **37j**, R = <sup>t</sup>Bu; **36t**, R = 4-C<sub>6</sub>H<sub>4</sub>C≡CSiMe<sub>2</sub>H) were prepared: addition of  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  (**35**) to a solution or suspension of oligomeric or polymeric alkynyl copper(I) compounds,  $[\text{CuC}\equiv\text{CR}]_n$ , in a 1 : 1 molar ratio yields the quantitative formation of the monomeric compounds **36h–36j** and **36t** [37,40]. An alternative preparative route is the metathesis reaction of  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuCl}$  (**36a**) with  $\text{LiC}\equiv\text{CR}$  (R = Ph, <sup>t</sup>Bu, SiMe<sub>3</sub>, 4-C<sub>6</sub>H<sub>4</sub>C≡CSiMe<sub>2</sub>H).




The presence of both  $\eta^2$ -bonded alkyne ligands and a  $\eta^1$ -bonded acetylide in compounds **36h–36j** and **36t** is shown by two different C≡C stretching frequencies

in their IR spectra: through the  $\eta^2$ -coordination of the trimethylsilylethynyl moieties in **35** to a copper atom in **36h–36j** and **36t** the  $\nu(\text{C}\equiv\text{C})$  vibration is shifted from  $2012\text{ cm}^{-1}$  in **35** to  $1905\text{ cm}^{-1}$  in **36h**,  $1902\text{ cm}^{-1}$  in **36i**,  $1896\text{ cm}^{-1}$  in **36j** and  $1899\text{ cm}^{-1}$  in **36t** [37,40]. The  $\sigma$ -bonded acetylide ligand  $\text{C}\equiv\text{CR}$  absorbs in the infrared region at  $2095\text{ cm}^{-1}$  in **36h**,  $2095\text{ cm}^{-1}$  in **36i**,  $2035\text{ cm}^{-1}$  in **36j** and  $2152/2090\text{ cm}^{-1}$  in **36t** for  $\nu(\text{C}\equiv\text{C})$ , and these data clearly indicate that the copper acetylide ligand is not  $\pi$  coordinated.

Surprisingly, compound **35** reacts selectively with  $1/4[\text{CuO}^t\text{Bu}]_4$  in diethyl ether to yield quantitatively the dimeric complex  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)-(\text{C}\equiv\text{CCu})_2\}$  (**40**) [37,40]. In this case, the binuclear compound  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuO}^t\text{Bu}$  (**36u**) could not be isolated; elimination of  $^t\text{BuOSiMe}_3$  (detected by GC–MS) and formation of **40** are instantaneous and quantitative.

Other nucleophiles show a similar reactivity towards compounds **36h–36j**, although the reaction is much slower and less selective owing to competitive cleavage of the  $\text{Ti}-\text{C}\equiv\text{C}$   $\sigma$  bond [37,40].



However, when  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuC}\equiv\text{CSiMe}_3$  (**36i**) is heated homolytic cleavage of the copper–carbon  $\sigma$  bond of the  $\text{CuC}\equiv\text{CSiMe}_3$  fragment is observed [13]. The formation of  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  (**35**), copper(0) together with  $\text{Me}_3\text{Si}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{SiMe}_3$  (**41**) takes place. This observation represents the well known Glaser reaction in which two acetylenes are oxidative coupled to symmetrical 1,3-diynes in presence of copper(I).

Compound **40** has a decomposition point of  $157^\circ\text{C}$ ; this high thermal stability is remarkable, since bimetallic acetylide species of copper(I), e.g.  $\text{CuC}\equiv\text{CCu}$ , are usually very reactive or even explosive [92].

Cryoscopic molecular weight determination of **40** in benzene indicates that **40** is dimeric in solution. Variable-temperature  $^1\text{H}$  NMR experiments confirm that **40** maintains this aggregation state in solution; in the temperature range 207–353 K the  $^1\text{H}$  NMR spectra remain essentially identical [37,40].

To establish the solid-state structure of **40**, an X-ray crystallographic study was carried out [37,40].

The molecular structure of **40** (Fig. 7) consists of a dimer of the bimetallic acetylide  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)(\text{C}\equiv\text{CCu})]$ . The  $\text{C}\equiv\text{C}$  bond length of the alkynyl ligands within this building block is lengthened from 1.208 Å (av.) in **35** to 1.234 Å ( $\text{TiC}\equiv\text{CSi}$ ) and 1.243 Å ( $\text{TiC}\equiv\text{CCu}$ ) in **40**. Each copper atom exhibits a distorted trigonal planar geometry with two  $\eta^2$ - and one  $\eta^1$ -bonded alkynyl ligands. The  $\text{Ti}-\text{C}\equiv\text{C}-\text{Cu}$  unit is non-linear ( $\text{Ti1}-\text{C22}-\text{C23}=163.4^\circ$ ;  $\text{Cu1}-\text{C23}-\text{C22}=165.0^\circ$ ), since this acetylide unit is  $\eta^2$ -coordinated to a second copper atom.

Formally, the structural arrangement of the  $\text{CuC}_2$  building blocks in **40** corresponds to that realized in polymeric  $[\text{Cu}^1\text{C}\equiv\text{CR}]_n$  in which the alkynyl units  $\text{C}\equiv\text{CR}$  are both  $\sigma$ - and  $\pi$ -bonded to copper atoms, thus forming an infinite zig-zag chain [1]. Similar structural observations were made for tetrameric  $[(\text{PMe}_3)\text{Cu}(\text{C}\equiv\text{CPh})]_4$  [93].

Further examples of transmetalation  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuCl}$




Fig. 7. Molecular geometry and atom labelling scheme for **40** [37,40]. Selected bond distances (Å) and angles ( $^\circ$ ) are as follows:  $\text{Ti1}-\text{Cu1}$  2.911(1),  $\text{Cu1}-\text{Cu2}$  2.998(1),  $\text{Cu1}-\text{C22}$  2.107(2),  $\text{Cu2}-\text{C23}$  2.162(2),  $\text{Cu1}-\text{C23}$  1.920(3),  $\text{Cu2}-\text{C17}$  2.031(3),  $\text{Cu2}-\text{C18}$  2.087(3),  $\text{C17}-\text{C18}$  1.234(3),  $\text{C22}-\text{C23}$  1.243(4),  $\text{Ti1}-\text{C17}$  2.098(2),  $\text{Ti1}-\text{C22}$  2.080(3);  $\text{Ti1}-\text{C17}-\text{C18}$  164.5(2),  $\text{C17}-\text{C18}-\text{Si3}$  164.9(3),  $\text{Ti1}-\text{C22}-\text{C23}$  163.4(2),  $\text{C22}-\text{C23}-\text{Cu1}$  165.0(2),  $\text{C17}-\text{Ti1}-\text{C22}$  90.5(1),  $\text{Cu1}-\text{C23}-\text{Cu2}$  39.7(1).


(**36a**) with  $[\text{AgX}]_n$  [ $\text{X}=\text{SCN}$ , route (a);  $\text{X}=\text{CN}$ , route (b);  $\text{X}=\text{O}_2\text{CMe}$ ,  $\text{O}_2\text{CPh}$ , route (c), or  $\text{Na}[\text{acac}]$  ( $\text{acac}=\text{acetylacetonate}$ ), route (e)] are given in Scheme 4.

$\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}[\text{OC}(\text{O})\text{Ph}]$  (**36g**) eliminates  $\text{CO}_2$  if heated in high-boiling solvents and yields  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{CuPh}$  (**36k**) (Scheme 4), in which a monomeric  $\text{CuPh}$  fragment is stabilized by the chelating effect of the organometallic ligand  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  [13].

Another procedure for the synthesis of **36k** was discussed earlier (addition of  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  (**35**) to polymeric  $[\text{Cu}^{\text{I}}\text{Ph}]_n$ ) (Section 3.3.1, Table 10) [38,40,86].

These results as a whole show that the striking property of the organometallic chelate  $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2$  (**35**) to bind mononuclear metal fragments gives access to a novel type of mononuclear organo-copper(I) species with a chelating bisalkynyl ligand. Studies are currently being undertaken in which this property of a fixed bidentate alkynyl ligand is exploited for the isolation of mononuclear aryl copper fragments from heterocopper and cuprate reagents and, as an example, the quantitative isolation of  $35\cdot\text{Cu}\{2,6\text{-(Me}_2\text{NCH}_2)_2\text{C}_6\text{H}_3\}$  from  $\text{Cu}_4\{2,6\text{-(Me}_2\text{NCH}_2)_2\text{C}_6\text{H}_3\}_2\text{Br}_2$  has been achieved [38,86].


Finally, spectroscopic and chemical properties of these compounds, with an extended ligand to ligand electronic communication (alkyne-to-aryl), are being investigated by variation of the mesomeric and inductive properties of the aryl para substituents [94].



Scheme 4. Synthesis of compounds **36d**, **36e**, **36f**, **36g**, **36k** and **36s**.

3.3.3.2.  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{M}(\text{OSO}_2\text{CF}_3)$  ( $\text{M} = \text{Cu}$ ,  $\text{Ag}$ ). The complexes  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{M}^{\text{I}}\text{X}$  ( $\text{M} = \text{Cu}$ , **36**;  $\text{M} = \text{Ag}$ , **37**) contain a monomeric  $\text{M}^{\text{I}}\text{X}$  moiety in which the copper or silver atom possesses a trigonal planar environment. There are several complexes known in which copper or silver atoms show a tetrahedral coordination sphere [1,95]. Hence,  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}(\text{OSO}_2\text{CF}_3)$  (**36p**) was reacted with  $\text{N}\equiv\text{CR}$  ( $\text{R} = \text{Me}$ ;  $\text{R} = \text{Ph}$ ) in an attempt to produce a copper complex with a copper atom in a tetrahedral environment. The compounds  $[\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Cu}(\text{N}\equiv\text{CR})](\text{OSO}_2\text{CF}_3)$  were formed in high yields [40].

However,  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Ag}(\text{OSO}_2\text{CF}_3)$  (**37m**) reacts even at low temperature with thf,  $\text{N}\equiv\text{CR}$  ( $\text{R} = \text{Me}$ ,  $\text{Ph}$ ),  $\text{PR}_2\text{R}'$  ( $\text{R} = \text{R}' = \text{Ph}$ ;  $\text{R} = \text{C}\equiv\text{CPh}$ ,  $\text{R}' = \text{CH}_2\text{Ph}$ ) or  $\text{P}(\text{OMe})_3$  to afford compounds **42**, **43** and **44** in quantitative yields [13].



As a result of the coordination of the Lewis bases thf,  $\text{N}\equiv\text{CR}$  and  $\text{PR}_2\text{R}'$  to the silver(I) centre in the adducts **42**, **43** and **44**, the  $\nu(\text{C}\equiv\text{C})$  vibration in the IR spectra decreases from  $1956\text{ cm}^{-1}$  in  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Ag}(\text{OSO}_2\text{CF}_3)$  (**37m**) to  $1950\text{--}1870\text{ cm}^{-1}$  in **42–44**. This effect is in agreement with the increasing participation of the back-bonding components in the silver metal to alkyne bonding.

In order to establish the solid-state structure of compounds **42–44**, an X-ray diffraction study was carried out on a single crystal of representative  $\{(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2\}\text{Ag}[(\text{thf})(\text{BF}_4)]$  (**42b**) [13].

Fig. 8 shows that **42b** is monomeric, and that the silver atom possesses a pseudo-tetrahedral environment built up from two alkyne ligands, one  $\text{BF}_4$  group and one thf molecule [13].

#### 3.4. Type A, J and K molecules

Formally, type J and K molecules can be considered as sub-structures of type H molecules, which themselves are part of structural type G clusters (see above). Dimerization of K generates molecules of type A (see also Section 2.1).





Fig. 8. Molecular geometry and atom labelling scheme for **42b** [13]. Selected bond distances (Å) and angles (°) are as follows: Ti1–Ag1 3.096(2), Ti1–C17 2.114(7), Ti1–C22 2.128(7), Ag1–C17 2.293(6), Ag1–C18 2.447(7), Ag1–C22 2.294(7), Ag1–C23 2.472(8), Ag1–O1 2.389(5), Ag1–F2 2.47(2), C17–C18 1.23(1), C22–C23 1.222(9); Ti1–C17–C18 170.7(6), Ti1–C22–C23 171.0(6), C17–C18–Si3 168.8(6), C22–C23–Si4 166.8(7), Ti1–Ag1–O1 135.8(4), Ti1–Ag1–F2 138.7(3), C17–Ti1–C22 95.6(3).




### 3.4.1. Type **J** and **K** molecules, $[\eta^2\text{-RC}\equiv\text{CML}_n)_2]\text{Cu}^I\text{X}$ (**J**), $(\eta^2\text{-RC}\equiv\text{CML}_n)\text{Cu}^I\text{X}$ (**K**)

There are two general routes for the synthesis of type **K** molecules: (a) metathesis reaction of  $\text{L}_n\text{MCl}$  compounds and (b) direct synthesis from  $\text{L}_n\text{M}-\text{C}\equiv\text{CR}$  complexes.

(a) *Metathesis reaction route.* The reaction of transition metal halides  $\text{L}_n\text{MCl}$  [**27g**,  $\text{L}_n\text{M}=(\eta^5\text{-C}_5\text{H}_5)(\text{PPh}_3)_2\text{Ru}$ ; **27h**,  $\text{L}_n\text{M}=(\text{CO})_3(\text{PPh}_3)_2\text{Re}$ ] with copper(I) acetylides

$[\text{CuC}\equiv\text{CR}]_n$  ( $\text{R} = \text{Ph}$ , 4-MeC<sub>6</sub>H<sub>4</sub>, Me) affords the monomeric compounds **45** [68,96–99].



27

45

**45a:**  $\text{L}_n\text{M} = (\eta^5\text{-C}_5\text{H}_5)(\text{PPh}_3)_2\text{Ru}$ ;  $\text{R} = \text{Ph}$

**45b:**  $\text{L}_n\text{M} = (\eta^5\text{-C}_5\text{H}_5)(\text{PPh}_3)_2\text{Ru}$ ;  $\text{R} = 4\text{-MeC}_6\text{H}_4$


**45c:**  $\text{L}_n\text{M} = (\eta^5\text{-C}_5\text{H}_5)(\text{PPh}_3)_2\text{Ru}$ ;  $\text{R} = \text{Me}$

**45d:**  $\text{L}_n\text{M} = (\text{CO})_3(\text{PPh}_3)_2\text{Re}$ ;  $\text{R} = \text{Ph}$

(b) *Direct synthesis route.* Another possibility for the synthesis of structural type **K** molecules is given by the reaction of  $[\text{Mn}]-\text{C}\equiv\text{CR}$  ( $[\text{Mn}] = \text{fac-Mn}(\text{CO})_3(\text{dppe})$ ; **46a**,  $\text{R} = \text{Ph}$ ; **46b**,  $\text{R} = \text{tBu}$ ; **46c**,  $\text{R} = \text{CH}_2\text{OMe}$ ) with  $[\text{Cu}^+\text{Cl}]_n$  or  $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$  ( $\text{tht} = \text{tetrahydrothiophene}$ ) to yield compounds **47** (Scheme 5, route (a)) (Table 14) [97–99].

As shown in Scheme 5, complexes **47** are suitable for the synthesis of ionic compounds  $\{[\text{Mn}](\eta^2\text{-C}\equiv\text{CR})_2\text{M}^+\text{Y}^-\}$  (**48**) (type **J** molecules; route (b)) and  $\{[\text{Mn}](\eta^2\text{-C}\equiv\text{CR})\text{M}^+(\text{PR}_3)_2\text{Y}^-\}$  (**49**) (type **K** molecules; route (c)) (Table 14) [97,98].

Compound **48a** is formed by the reaction of **47a** with **46** and  $\text{TiPF}_6$  in  $\text{CH}_2\text{Cl}_2$  (route (b)). Compounds **48c** and **48d** can be synthesized on addition of solid  $[\text{AuCl}(\text{tht})]$  and  $\text{TiPF}_6$  to the starting material **46**, whereas for the synthesis of **48b** the addition of  $\text{TiPF}_6$  is not necessary [97,98]. On addition of Lewis bases, e.g.  $\text{PR}_3$ , to solutions of **48**, one organometallic  $[\text{Mn}]-\text{C}\equiv\text{CR}$  building block



Scheme 5. Synthesis of compounds **47**, **48** and **49**;  $[\text{Mn}] = \text{fac-Mn}(\text{CO})_3(\text{dppe})$ ;  $\text{R}' = o\text{-tolyl}$  [97–99].

is exchanged by  $\text{PR}'_3$ , yielding the monomeric ionic compounds  $[\{[\text{Mn}](\eta^2\text{-C}\equiv\text{CR})\}\text{M}'(\text{PR}'_3)]\text{Y}$  (**49a–49d**) (type **K** molecules) (route (c)). A further possibility for preparing compounds **49a–49d** is given by route (d) [97–99].

All complexes prepared at present are summarized in Table 14 [96–99].

Compared with the starting compounds **46**, the  $\nu(\text{C}\equiv\text{C})$  frequencies of complexes **47–49** are significantly lower owing to the  $\eta^2$ -coordination of the  $\text{RC}\equiv\text{C}$  ligands to  $\text{M}'\text{X}$  ( $\text{M}=\text{Cu}$ ,  $\text{Ag}$ ,  $\text{Au}$ ) building blocks [96–99]. This is the normal observation made for alkynes, changing from the uncoordinated to the  $\eta^2$ -coordinated state (see above).

The monomeric structures of compounds **45**, **47** and **49** are evidenced both by osmometric measurements and X-ray diffraction studies [68,96–99].

The X-ray diffraction studies additionally confirm the IR spectroscopically observed bond lengthening of the  $\text{C}_2$  building blocks (see above). The  $\text{C}\equiv\text{C}$  bond is lengthened from 1.20 Å in organometallic substituted alkynes to 1.23 Å in **47a**, 1.24 Å in **49b** and 1.25 Å in **45a** [68,96–99]. In addition, the  $\eta^2$ -coordination leads to a deformation of the  $\text{M}-\text{C}\equiv\text{C}-\text{R}$  fragment (**47a**,  $171.1^\circ$ ,  $165.6^\circ$ ; **49b**,  $171.0^\circ$ ,  $164.0^\circ$ ; **45a**,  $173.0^\circ$ ,  $165.0^\circ$ ) [68,96–99].


On replacement of the organic group  $\text{R}$  in transition metal substituted alkynes  $[\text{M}]-\text{C}\equiv\text{CR}$  with organometallic building blocks, bimetallic complexes of the type  $[\text{M}]-\text{C}\equiv\text{C}-[\text{M}]$  are obtained. One example is  $[\text{Re}]-\text{C}\equiv\text{C}-[\text{Re}]$  ( $[\text{Re}]=\text{(CO)}_5\text{Re}$ ) (**50**), synthesized by Beck and co-workers [100]. Treatment of **50** with  $[\text{M}'\text{L}_4]\text{Y}$  ( $\text{L}=\text{NCMe}$ ;  $\text{M}=\text{Cu}$ ,  $\text{Y}=\text{PF}_6$ ;  $\text{M}=\text{Ag}$ ,  $\text{Y}=\text{BF}_4$ ) yields the cationic compounds **51a** ( $\text{M}=\text{Cu}$ ,  $\text{Y}=\text{PF}_6$ ) and **51b** ( $\text{M}=\text{Ag}$ ,  $\text{Y}=\text{BF}_4$ ) (Scheme 6, route (a)) [101].

On addition of  $[\text{Re}]-\text{C}\equiv\text{C}-[\text{Re}]$  (**50**) to **51**, the cationic complexes **52** (**52a**,  $\text{M}=\text{Cu}$ ,  $\text{Y}=\text{PF}_6$ ; **52b**,  $\text{M}=\text{Ag}$ ,  $\text{Y}=\text{BF}_4$ ) are formed (Scheme 6, route (b)), whereas with  $[\text{M}'\text{L}_4]\text{Y}$  the metalla-carbon tetrahedranes **53** (**53a**,  $\text{M}=\text{Cu}$ ,  $\text{Y}=\text{PF}_6$ ; **53b**,  $\text{M}=\text{Ag}$ ,  $\text{Y}=\text{BF}_4$ ) can be obtained (Scheme 6, route (c)) [101].

Table 14

Compounds **47**, **48** and **49** [97–99]

Compound	M	X	R	Y
<b>47a</b>	Cu	Cl	Ph	–
<b>47b</b>	Cu	Cl	<sup>t</sup> Bu	–
<b>47c</b>	Cu	Cl	$\text{CH}_2\text{OMe}$	–
<b>47d</b>	Au	$\text{C}_6\text{F}_5$	<sup>t</sup> Bu	–
<b>47e</b>	Au	$\text{C}_6\text{F}_5$	$\text{CH}_2\text{OMe}$	–
<b>48a</b>	Cu	–	<sup>t</sup> Bu	$\text{PF}_6$
<b>48b</b>	Ag	–	<sup>t</sup> Bu	$\text{BF}_4$
<b>48c</b>	Au	–	Ph	$\text{PF}_6$
<b>48d</b>	Au	–	<sup>t</sup> Bu	$\text{PF}_6$
<b>49a</b>	Cu	–	<sup>t</sup> Bu	$\text{PF}_6$
<b>49b</b>	Cu	–	$\text{CH}_2\text{OMe}$	$\text{PF}_6$
<b>49c</b>	Ag	–	<sup>t</sup> Bu	$\text{BF}_4$
<b>49d</b>	Au	–	<sup>t</sup> Bu	$\text{PF}_6$




Scheme 6. Synthesis of compounds **51–53**;  $[Re]=Re(CO)_5$ ;  $L=N\equiv CMe$ ;  $M=Cu, Ag$ ;  $Y=PF_6, BF_4$  [100,101].

The  $\eta^2$ -coordination of the  $C_2$  unit in **50** to a copper(I) or silver(I) fragment leads to  $C\equiv C$  stretching vibrations in the  $1890\text{--}1950\text{ cm}^{-1}$  region, a phenomenon typical of this coordination mode (see above).

### 3.4.2. Organometallic type A molecules, $[(\eta^2-RC\equiv CML_n)Cu^IX]_2$

In structural type A compounds, the  $C_2$  unit of the organometallic group  $L_nM-C\equiv CR$  is  $\eta^2$ -coordinated to a  $Cu^IX$  fragment and through the formation of copper–halide bridges ( $Cu_2X_2$ ) a dimer is formed (for the synthesis of  $[(\eta^2-RC\equiv CR)Cu^IX]_2$  compounds, see Section 2.1).



A


Type A molecules can be obtained by the reaction of  $L_nMCl$  (**27**) with  $[CuC\equiv CR]_n$  or the reaction of  $L_nM-C\equiv CR$  (**46**) with  $Cu^IX$  (Table 15) [68,96,102–106].

Compounds **54** are dimeric, as indicated by osmometric measurements [68,96,102–106].

In order to establish the solid-state structure of compounds **54a–54j**, an X-ray diffraction study was carried out on a single crystal of **54b** (Fig. 9) [102–104]. Fig. 9 shows that complex **54b** is dimeric and consists of two asymmetric units related by

Table 15  
Compounds **54**

Compound	$L_nM$	X	R	Ref.
<b>54a</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{PPh}_3)_2\text{Ru}$	Cl	$\text{C}_6\text{H}_5$	[87,96]
<b>54b</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Cl	$\text{C}_6\text{H}_5$	[102–104]
<b>54c</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Cl	4-MeC <sub>6</sub> H <sub>4</sub>	[102]
<b>54d</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Cl	4-FC <sub>6</sub> H <sub>4</sub>	[102]
<b>54e</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Cl	$\text{C}_6\text{F}_5$	[102]
<b>54f</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Cl	Me	[102]
<b>54g</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Cl	SiMe <sub>3</sub>	[105,106]
<b>54h</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}$	Br	$\text{C}_6\text{H}_5$	[102]
<b>54i</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{Mo}$	Cl	SiMe <sub>3</sub>	[105,106]
<b>54j</b>	$(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{W}$	Cl	SiMe <sub>3</sub>	[105,106]



a centre of inversion. Each copper atom possesses a trigonal planar environment and is symmetrically  $\eta^2$ -coordinated by the  $\text{C}_2$  unit of the phenylethynyl group. The  $\text{Cu}_2\text{Cl}_2$  ring is coplanar with the  $\text{CuC}_2$  moiety. The  $\text{Cu}\cdots\text{Cu}$  distance of 3.075(4) Å indicates that there is no bonding character between these two metal atoms.

Through the  $\eta^2$ -coordination of the  $\text{PhC}\equiv\text{C}$  ligand to a  $\text{Cu}^I\text{Cl}$  building block, the bond length of the  $\text{C}\equiv\text{C}$  unit is lengthened from approximately 1.20 Å in the starting material to 1.23 Å in **54b**. In addition, it is found that the  $\text{Fe}-\text{C}\equiv\text{C}-\text{Ph}$  building block is distorted from linearity (Fig. 9). The same kind of structure is present in a series of analogous compounds  $[(\eta^2\text{-RC}\equiv\text{CR})\text{Cu}^IX]_2$  ( $\text{R}$ =singly bonded organic ligand;  $\text{X}=\text{Cl}, \text{Br}$ , etc.) which contain an organic group  $\text{R}$  instead of an organometallic moiety  $\text{ML}_n$  (Section 2.1).

## Acknowledgements

The creative contributions of Dr. Mathias Herres, Dipl.-Chem. Markus Weinmann, Dipl.-Chem. Gerd Rheinwald, Maurits D. Janssen and Susanne Ramspeck are gratefully acknowledged. The financial support from the Deutsche Forschungsgemeinschaft, the Volkswagenstiftung, the Fonds der Chemischen Industrie and the Hermann-Schlosser Stiftung Degussa AG, Frankfurt, was essential




Fig. 9. Molecular geometry and atom labelling scheme for **54b** [102–104]. Important selected interatomic distances (Å) and angles (°) are as follows: Cu–Cl 2.283(3), Cu–Cl' 2.291(3), Fe–C8 1.91(1), Cu–C8 2.02(1), Cu–C9 2.00(1), C8–C9 1.23(1), Fe–C8–C9 164.3(9), C8–C9–C91 161.1(1), Cu–Cl–Cu' 84.3(1), Cl–Cu–Cl' 95.7(1).

throughout. For many fruitful discussions we are grateful to Professor Dr. G. Huttner, Professor Dr. G. van Koten, Dr. D.M. Grove and Dr. N. Compton.

## References

- [1] B.H. Lipshutz and S. Sengupta, *Org. React.*, 41 (1992) 135; G.H. Posner, *An Introduction to Synthesis Using Organocopper Reagents*, Wiley-Interscience, New York, 1980; B.H. Lipshutz, *Synthesis*, (1987) 325; J.F. Normant, *Synthesis*, (1972) 63; W. Carruthers, in G. Wilkinson, F.G.A. Stone and E.W. Abel (eds.), *Comprehensive Organometallic Chemistry*, Vol. 7, Pergamon Press, Oxford, 1982, Chapter 49; C. Glaser, *Liebigs Ann. Chem.*, 154 (1870) 137; T. Sandmeyer, *Chem. Ber.*, 17 (1884) 1633; F. Ullman and J. Bielecki, *Chem. Ber.*, 43 (1901) 2174; W.R.H. Hurtley, *J. Chem. Soc.*, (1929) 1870; H. Gilman and J.M. Straley, *Recl. Trav. Chim. Pays-Bas*, 55 (1936) 821; M.S. Kharash and P.O. Tawney, *J. Am. Chem. Soc.*, 63 (1941) 2308; H.O. House, W.L. Respass and G.M. Whitesides, *J. Org. Chem.*, 31 (1966) 3138; G.M. Whitesides, W.F. Fischer, Jr., J. San Filippo, Jr., R.W. Bashe and H.O. House, *J. Am. Chem. Soc.*, 91 (1969) 4871; L.I. Zakharkin and L.P. Sorokina, *Zh. Org. Khim.*, 6 (1970) 2430; *J. Org. Chem. (USSR)*, 6 (1970) 2482; J.F. Normant and M. Bourgain, *Tetrahedron Lett.*, 11 (1970) 2659; H.O. House and M.J. Umen, *J. Am. Chem. Soc.*, 94 (1972) 5495; G.M. Whitesides and P.E. Kendall, *J. Org. Chem.*, 37 (1972) 3718; G.H. Posner, C.E. Whitten and J.J. Sterling, *J. Am. Chem. Soc.*, 95 (1973) 7788; H.O. House and M.J. Umen, *J. Org. Chem.*, 36 (1973) 3893; C.R. Johnson and G.A. Dutra, *J. Am. Chem. Soc.*, 95 (1973) 7783; A. Bruggink and A. McKillop, *Tetrahedron*, 31 (1975) 2607; H.O. House, *Acc.*

- Chem. Res., 9 (1976) 59; J. Berlan, J.P. Battioni and K. Koosha, *Tetrahedron Lett.*, 17 (1976) 3355; T. Olsson, M.T. Rahman and C. Ullenius, *Tetrahedron Lett.*, 18 (1977) 75; M. Nilsson, M.T. Rahman and C. Ullenius, *Acta Chem. Scand., Sect. B*, 31 (1977) 514; J. Berlan and K.J. Koosha, *J. Organomet. Chem.*, 152 (1978) 359; J. Berlan, J.P. Battioni and K.J. Koosha, *Bull. Soc. Chim. Fr.*, (1979) 183; C.P. Casey and M.C. Cesa, *J. Am. Chem. Soc.*, 101 (1979) 4236; R.A.J. Smith and D.J. Hannah, *Tetrahedron*, 35 (1979) 1183; S.R. Krauss and S.G. Smith, *J. Am. Chem. Soc.*, 103 (1981) 141; F. Leyendecker and D. Laucher, *Tetrahedron Lett.*, 24 (1983) 3517; E.J. Corey and N.W. Boaz, *Tetrahedron Lett.*, 25 (1984) 3063; B.H. Lipshutz, R.S. Wilhelm and J.A. Kozlowski, *Tetrahedron*, 40 (1984) 5005; F. Leyendecker and D. Laucher, *Nouv. J. Chim.*, 9 (1985) 13; E.J. Corey and N.W. Boaz, *Tetrahedron Lett.*, 26 (1985) 6015; J. Berlan and Y. Besace, *Tetrahedron*, 42 (1986) 4767; E.J. Corey, R. Naef and F.J. Hannon, *J. Am. Chem. Soc.*, 108 (1986) 7114; Y. Yamamoto, *Angew. Chem.*, 98 (1986) 956; *Angew. Chem., Int. Ed. Engl.*, 25 (1986) 947; G.M. Villacorta, C. Pulla and S.J. Lippart, *J. Am. Chem. Soc.*, 110 (1988) 3175; C. Ullenius and B. Christenson, *Pure Appl. Chem.*, 60 (1988) 57; S.H. Bertz and R.A.J. Smith, *J. Am. Chem. Soc.*, 111 (1989) 8276; A.E. Dorigo and K. Morokuma, *J. Am. Chem. Soc.*, 111 (1989) 6524; A.E. Dorigo and K. Morokuma, *J. Am. Chem. Soc.*, 111 (1989) 4635; D.M. Knotter, A.L. Spek and G. van Koten, *J. Chem. Soc., Chem. Commun.*, (1989) 1738; D.M. Knotter, G. van Koten, L.H. van Maaren, D.M. Grove and A.L. Spek, *Angew. Chem.*, 101 (1989) 351; *Angew. Chem., Int. Ed. Engl.*, 28 (1989) 341; B.H. Lipshutz, *Synlett*, (1990) 119; B.H. Lipshutz, E.L. Ellsworth and S.H. Dimock, *J. Am. Chem. Soc.*, 112 (1990) 5869.
- [2] P.P. Power, *Prog. Inorg. Chem.*, 39 (1991) 76; A.J. Jukes, *Adv. Organomet. Chem.*, 12 (1974) 215; G. van Koten and J.G. Noltes, in F.G.A. Stone and E.W. Abel (eds.), *Comprehensive Organometallic Chemistry*, Vol. II/14, Pergamon Press, Oxford, 1982, p. 709; A.M. Sladkov and I.R. Gol'ding, *Russ. Chem. Rev.*, 48 (1979) 868; A. Camus, N. Marsich, G. Nardin and L. Randaccio, *Inorg. Chim. Acta*, 23 (1977) 131; G. Bähr and P. Barba, in Houben-Weyl, *Methoden der Organischen Chemie*, Vol. XIII/1, Georg Thieme, Stuttgart, 1970, p. 735; D.W. Stephan, *Coord. Chem. Rev.*, 95 (1989) 41; E. Wehman, G. van Koten, J.T.B.H. Jastrzebski, M.A. Rotteveel and C.H. Stam, *Organometallics*, 7 (1988) 1477; E. Wehman, G. van Koten, D.M. Knotter, H. Spelten, D. Heijdenrijk, N.S. Mak and C.H. Stam, *J. Organomet. Chem.*, 325 (1987) 293; B. Lenders, D.M. Grove, W.J.J. Smeets, P. van der Sluis, A.L. Spek and G. van Koten, *Organometallics*, 10 (1991) 786; D.M. Knotter, W.J.J. Smeets, A.L. Spek and G. van Koten, *J. Am. Chem. Soc.*, 112 (1990) 5895; S. Gambarotta, C. Floriani, A. Chiesi-Villa and C. Guastini, *J. Chem. Soc., Chem. Commun.*, (1983) 1156.
- [3] S. Gambarotta, S. Strologo, C. Floriani, A. Chiesi-Villa and C. Guastini, *Organometallics*, 3 (1984) 1444; P.S. Coan, K. Folting, J.C. Huffman and K.G. Caulton, *Organometallics*, 8 (1989) 2724; P.W. Corfield and H.M.M. Shearer, *Acta Crystallogr.*, 20 (1966) 502; R. Uson, A. Laguna, A. Uson, P.G. Jones and K. Meyer-Base, *J. Chem. Soc., Dalton Trans.*, (1988) 341; X. He, M.M. Olmstead and P.P. Power, *J. Am. Chem. Soc.*, 114 (1992) 9668; R. Lingnau and J. Strähle, *Angew. Chem.*, 100 (1988) 409; *Angew. Chem., Int. Ed. Engl.*, 27 (1988) 436; T.A. Wark and D.W. Stephan, *Inorg. Chem.*, 26 (1987) 363; D.W. Stephan, *J. Chem. Soc., Chem. Commun.*, (1991) 129; T.T. Nadasdi and D.W. Stephan, *Organometallics*, 11 (1992) 116; P.S. Braterman and V.A. Wilson, *J. Organomet. Chem.*, 31 (1971) 131; H.J. Küppers, K. Wieghardt, Y.-H. Tsay, C. Krüger, B. Nuber and J. Weiss, *Angew. Chem.*, 99 (1987) 583; *Angew. Chem., Int. Ed. Engl.*, 26 (1987) 575; H. Maelger, F. Olbrich, J. Kopf, D. Abeln and E. Weiss, *Z. Naturforsch., Teil B*, 47 (1992) 1276; A. Haaland, K. Rypdal, H.P. Verne, W. Scherer and W.R. Thiel, *Angew. Chem.*, 106 (1994) 2515; *Angew. Chem., Int. Ed. Engl.*, 33 (1994) 2443.
- [4] F.L. Carter and E.W. Hughes, *Acta Crystallogr.*, 10 (1974) 801.
- [5] G.G. Aleksandrov, I.R. Gol'ding, S.R. Sterlin, A.M. Sladkov, Yu.T. Struchkov, I.A. Garbuzova and V.T. Aleksanyan, *Izv. Akad. Nauk SSR, Ser. Khim.*, 12 (1980) 2679 (*Engl. Transl.*, p. 1857).
- [6] D.W. Macomber and M.D. Rausch, *J. Am. Chem. Soc.*, 105 (1983) 5325, and references cited therein.
- [7] G. Maier, M. Hoppe, H.P. Reisenauer and C. Krüger, *Angew. Chem.*, 94 (1982) 445; *Angew. Chem., Int. Ed. Engl.*, 21 (1982) 437; *Angew. Chem., Suppl.*, (1982) 1061.
- [8] F. Olbrich, J. Kopf, E. Weiss, A. Krebs and S. Müller, *Acta Crystallogr., Sect. C*, 46 (1990) 1650.
- [9] F. Olbrich, G. Schmidt, U. Behrens and E. Weiss, *J. Organomet. Chem.*, 418 (1991) 41.

- [10] F. Olbrich, U. Behrens, G. Gröger and E. Weiss, *J. Organomet. Chem.*, 448 (1993) C10; F. Olbrich, U. Behrens and E. Weiss, *J. Organomet. Chem.*, 472 (1994) 365.
- [11] F. Olbrich, J. Kopf and E. Weiss, *J. Organomet. Chem.*, 456 (1993) 293.
- [12] F. Olbrich, U. Behrens, G. Schmidt and E. Weiss, *J. Organomet. Chem.*, 463 (1993) 249.
- [13] K. Köhler, Diplom-Thesis, University of Heidelberg, 1994.
- [14] H. Lang, K. Köhler and L. Zsolnai, *Chem. Ber.*, submitted for publication.
- [15] D.L. Reger and M.F. Huff, *Organometallics*, 9 (1990) 2807.
- [16] M. Pasquali, P. Leoni, C. Floriani and A. Gaetani-Manfredotti, *Inorg. Chem.*, 21 (1982) 4324.
- [17] H.L. Aalton, G. van Koten, E. Riethorst and C.H. Stam, *Inorg. Chem.*, 28 (1989) 4140.
- [18] F. Olbrich, G. Schmidt, E. Weiss and U. Behrens, *J. Organomet. Chem.*, 456 (1993) 299.
- [19] H. Lang and K. Köhler, *J. Organomet. Chem.*, submitted for publication.
- [20] J.C. Dyason, P.C. Healy, L.M. Engelhardt, C. Pakawatchai, V.A. Patrick and A.H. White, *J. Chem. Soc., Dalton Trans.*, (1985) 839.
- [21] D.T. Cromer, A.C. Larson and R.B. Roof, *Acta Crystallogr.*, 20 (1966) 279.
- [22] Holleman-Wiberg, *Lehrbuch der Anorganischen Chemie*, Walter de Gruyter, Berlin, 1976.
- [23] N.N. Greenwood and A. Earnshaw, *Chemistry of the Elements*, Pergamon Press, Oxford, 1984, p. 1368.
- [24] D.L. Reger, M.F. Huff, Th.A. Wolfe and R.D. Adams, *Organometallics*, 8 (1989) 848.
- [25] H. Lang, K. Köhler and L. Zsolnai, *Chem. Ber.*, in press.
- [26] M. Pasquali, C. Floriani, G. Venturi, A. Gaetani-Manfredotti and A. Chiesi-Villa, *J. Am. Chem. Soc.*, 104 (1982) 4092; A. Toth, C. Floriani, A. Chiesi-Villa and C. Guastini, *Inorg. Chem.*, 26 (1987) 236.
- [27] E.g. G.M. Villacorta, D. Gibson, I.D. Williams, E. Whang and S.J. Lippard, *Organometallics*, 6 (1987) 2426, and references cited therein.
- [28] G.M. Villacorta, D. Gibson, I.D. Williams and S.J. Lippard, *J. Am. Chem. Soc.*, 107 (1985) 6732.
- [29] R. Gleiter, M. Karcher, D. Kratz, M.L. Ziegler and B. Nuber, *Chem. Ber.*, 123 (1990) 1461.
- [30] H. Lang, M. Herres, K. Köhler, S. Blau, S. Weinmann, M. Weinmann, G. Rheinwald and W. Imhof, *J. Organomet. Chem.*, in press.
- [31] I.A. Garbuzova, I.R. Gol'ding, N.N. Travkin, A.M. Sladkov and V.T. Aleksanyan, *Koord. Khim.*, 5 (1979) 345.
- [32] M. Hakansson, K. Wettström and S. Jagner, *J. Organomet. Chem.*, 421 (1991) 347.
- [33] A.C. Ukwueze and S.R. Lander, *Spectrochim. Acta, Part A*, 40 (1984) 433; A.C. Ukwueze and S.R. Lander, *Synth. React. Inorg. Met.-Org. Chem.*, 14 (1984) 1013.
- [34] H. Lang and M. Herres, *Trends in Organometallic Chemistry*, Council of Scientific Research Integration, India, 1995, and references cited therein.
- [35] H. Lang, *Angew. Chem.*, 106 (1994) 569; *Angew. Chem., Int. Ed. Engl.*, 33 (1994) 547.
- [36] H. Lang, M. Herres and L. Zsolnai, *Organometallics*, 12 (1993) 5008.
- [37] M. Herres, M.D. Janssen, L. Zsolnai, D.M. Grove, H. Lang and G. van Koten, unpublished.
- [38] M.D. Janssen, M. Herres, A.L. Spek, D.M. Grove, H. Lang and G. van Koten, *J. Am. Chem. Soc.*, to be submitted.
- [39] H. Lang, M. Herres, L. Zsolnai and W. Imhof, *J. Organomet. Chem.*, 409 (1991) C7.
- [40] M. Herres, Diplom-Thesis, University of Heidelberg, 1990; M. Herres, Ph.D. Thesis, University of Heidelberg, 1993.
- [41] J.D. Ferrara, C. Tessler-Youngs and W.J. Youngs, *Organometallics*, 6 (1987) 676.
- [42] J.D. Ferrara, A. Djebli, C. Tessler-Youngs and W.J. Youngs, *J. Am. Chem. Soc.*, 110 (1988) 647.
- [43] J.D. Ferrara, C. Tessler-Youngs and W.J. Youngs, *Inorg. Chem.*, 27 (1988) 2201.
- [44] J.S. Thompson and J.F. Whitney, *J. Am. Chem. Soc.*, 105 (1983) 5488.
- [45] J.S. Thompson and J.F. Whitney, *Inorg. Chem.*, 23 (1984) 2813.
- [46] M. Munakata, S. Kitagawa, I. Kawada, M. Maekawa and H. Shimono, *J. Chem. Soc., Dalton Trans.*, (1992) 2225.
- [47] D.L. Reger and M.F. Huff, *Organometallics*, 11 (1992) 69.
- [48] B. Lenders and W. Kläui, *Chem. Ber.*, 123 (1990) 2233.
- [49] G.A. Carriedo, J.A.K. Howard and F.G.A. Stone, *J. Chem. Soc., Dalton Trans.*, (1984) 1555.
- [50] G. Huttner and K. Knoll, *Angew. Chem.*, 99 (1987) 765; *Angew. Chem., Int. Ed. Engl.*, 26 (1987) 743.

- [51] G. Huttner, D. Lange and E.O. Fischer, *Angew. Chem.*, 83 (1971) 579; *Angew. Chem., Int. Ed. Engl.*, 10 (1971) 556.
- [52] H. Wadepohl, *Angew. Chem.*, 104 (1992) 253; *Angew. Chem., Int. Ed. Engl.*, 31 (1992) 247.
- [53] A. Jain, K.M. Chi, T.T. Kodas, M.J. Hampden-Smith, J.D. Farr and M.F. Paffett, *Chem. Mater.*, 3 (1991) 995.
- [54] K.M. Chi, H.K. Shin, M.J. Hampden-Smith, T.T. Kodas and E.N. Duesler, *Inorg. Chem.*, 30 (1991) 4293.
- [55] K.M. Chi, T.S. Corbitt, M.J. Hampden-Smith, T.T. Kodas and E.N. Duesler, *J. Organomet. Chem.*, 449 (1993) 181.
- [56] D.M. Hoffman, R. Hoffmann and C.R. Fisel, *J. Am. Chem. Soc.*, 104 (1982) 3858.
- [57] F. Baert, A. Guelzim and P. Coppens, *Acta Crystallogr., Sect. B*, 40 (1984) 590.
- [58] H. Lang and L. Zsolnai, *Chem. Ber.*, 124 (1991) 259, and references cited therein; H. Lang and L. Zsolnai, *Z. Naturforsch., Teil B*, 45 (1990) 1529; H. Lang, H. Keller, W. Imhof and S. Martin, *Chem. Ber.*, 123 (1990) 417.
- [59] R.S. Dickson and P.J. Fraser, *Adv. Organomet. Chem.*, 12 (1974) 323.
- [60] R. Nast, *Coord. Chem. Rev.*, 47 (1982) 89.
- [61] O.M. Abu Salah and M.I. Bruce, *J. Chem. Soc., Chem. Commun.*, (1972) 858.
- [62] M.R. Churchill and S.A. Bezman, *Inorg. Chem.*, 13 (1974) 1418.
- [63] M.R. Churchill and B.G. DeBoer, *Inorg. Chem.*, 14 (1975) 2630.
- [64] O.M. Abu Salah and M.I. Bruce, *Aust. J. Chem.*, 29 (1976) 531; O.M. Abu Salah and M.I. Bruce, *Aust. J. Chem.*, 30 (1977) 2639.
- [65] O.M. Abu Salah and M.I. Bruce, *J. Chem. Soc., Chem. Commun.*, (1974) 688.
- [66] P. Espinet, J. Fornies, F. Martinez, M. Tomas, E. Lalinde, M. Moreno, A. Ruiz and A.J. Welch, *J. Chem. Soc., Dalton Trans.*, (1990) 791.
- [67] International Tables for X-Ray Crystallography, Vol. 3, Kynoch Press, Birmingham, 1962, p. 278 (based on  $d(\text{Rh}-\text{Rh})_{\text{min}} = 2.690 \text{ \AA}$  and  $d(\text{Ag}-\text{Ag})_{\text{min}} = 2.889 \text{ \AA}$  in the pure metals).
- [68] O.M. Abu Salah and M.I. Bruce, *J. Chem. Soc., Dalton Trans.*, (1975) 2311.
- [69] J. Fornies, E. Lalinde, F. Martinez, M.T. Moreno and A.J. Welch, *J. Organomet. Chem.*, 455 (1993) 271.
- [70] P. Espinet, J. Fornies, F. Martinez, M. Sotes, E. Lalinde, M.T. Moreno, A. Ruiz and A.J. Welch, *J. Organomet. Chem.*, 403 (1991) 253.
- [71] R. Uson, J. Fornies, B. Menjon, F.A. Cotton, L.R. Falvello and M. Tomas, *Inorg. Chem.*, 24 (1985) 4651, and references cited therein.
- [72] C.W. Chen and H.W. Whitlock, Jr., *J. Am. Chem. Soc.*, 100 (1978) 4921.
- [73] F. Vögtle, Th. Papkalla, H. Koch and M. Nieger, *Chem. Ber.*, 123 (1990) 1097; E. Weber and F. Vögtle, *Chem. Unserer Zeit*, 23 (1989) 210; F. Vögtle, H.G. Löhr, J. Franke and D. Worsch, *Angew. Chem.*, 97 (1985) 721; *Angew. Chem., Int. Ed. Engl.*, 24 (1985) 727; F. Vögtle and E. Weber, *Angew. Chem.*, 91 (1979) 813; *Angew. Chem., Int. Ed. Engl.*, 18 (1979) 753.
- [74] H. Lang and D. Seyferth, *Z. Naturforsch., Teil B*, 45 (1990) 212, and references cited therein.
- [75] H. Lang and S. Blau, in preparation; S. Blau, Ph.D. Thesis, University of Heidelberg, 1995.
- [76] G.L. Wood, C.B. Knobler and M.F. Hawthorne, *Inorg. Chem.*, 28 (1989) 382; A. Sebald, P. Fritz and B. Wrackmeyer, *Spectrochim. Acta, Part A*, 41 (1985) 1405; R. Jimenez, M.C. Barral, V. Moreno and A. Sandoz, *J. Organomet. Chem.*, 174 (1979) 281; A.J. Jenkins, M.F. Lappert and R.C. Shrivastava, *J. Organomet. Chem.*, 23 (1970) 165; J.H. Teuben and H.J. Liefde Meijer, *J. Organomet. Chem.*, 17 (1969) 87; H. Köpf and M. Schmidt, *J. Organomet. Chem.*, 10 (1967) 383.
- [77] K. Yasufuku and H. Yamazaki, *Bull. Chem. Soc. Jpn.*, 45 (1972) 2664.
- [78] H. Lang and W. Imhof, *Chem. Ber.*, 125 (1992) 1307.
- [79] H. Lang, M. Herres and L. Zsolnai, *Bull. Chem. Soc. Jpn.*, 66 (1993) 429.
- [80] H. Lang and L. Zsolnai, *J. Organomet. Chem.*, 406 (1991) C5.
- [81] M. Ciriano, J.A.K. Howard, J.L. Spencer, F.G.A. Stone and H. Wadepohl, *J. Chem. Soc., Dalton Trans.*, (1979) 1749.
- [82] H. Lang, M. Herres, L. Zsolnai and W. Imhof, *J. Organomet. Chem.*, 409 (1991) C7.
- [83] M. Herres and H. Lang, *J. Organomet. Chem.*, 480 (1994) 235.
- [84] S.I. Troyanov, V. Varga and K. Mach, *Organometallics*, 12 (1993) 2820.

- [85] H. Lang, M. Herres, K. Köhler and L. Zsolnai, *J. Organomet. Chem.*, in press.
- [86] M.D. Janssen, M. Herres, A. Spek, D.M. Grove, H. Lang and G. van Koten, in preparation.
- [87] O.M. Abu Salah, M.I. Bruce and A.D. Redhouse, *J. Chem. Soc., Chem. Commun.*, (1974) 855.
- [88] O.M. Abu Salah and M.I. Bruce, *J. Chem. Soc., Dalton Trans.*, (1975) 2311.
- [89] A.J. Leusink, G. van Koten, J.W. Marsman and J.G. Noltes, *J. Organomet. Chem.*, 55 (1973) 419.
- [90] J. Blenkins, H.K. Hofstee, J. Boersma and G.J.M. van der Kerk, *J. Organomet. Chem.*, 168 (1979) 251.
- [91] J.R. Berenguer, L.R. Falvello, J. Fornies, E. Lalinde and M. Thomas, *Organometallics*, 12 (1993) 6.
- [92] G. Bähr and P. Barba, in *Houben-Weyl, Methoden der Organischen Chemie*, Vol. XIII/1, Georg Thieme, Stuttgart, 1970, pp. 738–741, and references cited therein.
- [93] P.W.R. Corfield and H.M.M. Shearer, *Acta Crystallogr.*, 21 (1966) 957; E.A. Jeffery and T. Mole, *J. Organomet. Chem.*, 11 (1968) 393.
- [94] M.D. Janssen, M. Herres, K. Köhler, H. Lang and G. van Koten, in preparation.
- [95] G. van Koten, *J. Organomet. Chem.*, 400 (1990) 283.
- [96] M.I. Bruce, O.M. Abu Salah, R.E. Davis and N.V. Raghavan, *J. Organomet. Chem.*, 64 (1974) C48.
- [97] G.A. Carriedo, D. Miguel, V. Riera and X. Solans, *J. Chem. Soc., Dalton Trans.*, (1987) 2867.
- [98] G.A. Carriedo, D. Miguel, V. Riera, X. Solans, M. Font-Altaba and M. Coll, *J. Organomet. Chem.*, 299 (1986) C43.
- [99] X. Solans, J. Solans, C. Miravittles, D. Miguel, V. Riera and J.M. Rubio-Gonzalez, *Acta Crystallogr., Sect. C*, 42 (1986) 975.
- [100] W. Beck, B. Niemer, J. Breitmaier and J. Heidrich, *J. Organomet. Chem.*, 371 (1989) 79; J. Heidrich, M. Steinmann, M. Appel, W. Beck, J.R. Philipps and W.C. Troglor, *Organometallics*, 9 (1990) 1296.
- [101] S. Mihan and W. Beck, unpublished results.
- [102] O.M. Abu Salah and M.I. Bruce, *J. Chem. Soc., Dalton Trans.*, (1974) 2302.
- [103] M.I. Bruce, R. Clark, J. Howard and P. Woodward, *J. Organomet. Chem.*, 42 (1972) C107.
- [104] R. Clark, J. Howard and P. Woodward, *J. Chem. Soc., Dalton Trans.*, (1974) 2027.
- [105] H. Lang, S. Blau, G. Rheinwald and L. Zsolnai, *J. Organomet. Chem.*, in press.
- [106] G. Rheinwald, Diplom-Thesis, University of Heidelberg, 1990.