


Scheme



Scheme: Synthesis of nine compounds in two dimensions using variable mixing. Six three-component Grignard reagents are reacted with an electrophile (E) to give six libraries. Screening of libraries 1,x and x,2 will show whether compound E-R_{1,2} is significantly more active than the background.


active library members would normally require individual re-synthesis or so-called deletion synthesis [5]. To avoid the unnecessary synthesis of scores of inactive compounds, a method has recently been reported that allows easy identification of active compounds from libraries generated using multicomponent Grignard reagents [6]. In essence the method works by variable mixing of the Grignard reagents to give the desired number of library 'dimensions' (see Scheme).

To prepare n^2 compounds, n libraries with n compounds would be prepared and each product (Grignard reagent) would be assigned a coordinate x, y , where x is the library number and y the number of the member. If the synthesis is now repeated with the meaning of x and y reversed, (so that y depicts a library number, etc), n new libraries will result containing the same n^2 compounds. By screening the $2n$ libraries, library members with extraordinary activities will be revealed directly through the display of activity in any of their coordinate libraries (see Scheme).

As target compounds to test this methodology coupled to biological screening, 3-substituted tropane analogues were found suitable as (a) 3-phenyl tropanes (**iii**) are dopamine transport inhibitors and (b) phenyl tropanes are made by a 1,4-conjugate addition of Grignard reagents to methyl ecgonidine (**iv**). In this fashion by varying reagent combinations, a 25-compound library composed of 2×5 sublibraries of 5 compounds were prepared in solution.

Compounds were screened against the monoamine transporters hDAT, hSERT and hNET in a competitive binding assay. One of the most potent compounds obtained from this library was (**v**) which displayed a K_i binding of 19 nM to hDAT.

This work is of interest as the methodology allows for the rapid preparation and screening of homologous compounds and further work in this area is merited.



- 1 Fakhfakh, M.A. *et al.* (2001) Expedient preparation of 2-substituted quinolines. *Tetrahedron Lett.* 42, 3847–3850
- 2 Smith, P.W. *et al.* (1994) Synthesis and biological evaluation of a library containing potentially 1600 amides/esters. A strategy for rapid compound generation and screening. *Bioorg. Med. Chem. Lett.* 4, 2821–2824
- 3 Bülow, A. *et al.* (2004) Two- and three-dimensional combinatorial chemistry from multicomponent Grignard reagents. *Comb. Chem.* 6, 509–519

Paul Edwards

paul.edwards@graftinity.com

Erratum

In the 15th March 2005 issue of *Drug Discovery Today* (Vol. 10, No. 6, p.446), in the article entitled *The first [Foscarnet]–[TSAO-T] conjugates*, there were some errors in the accompanying figure. The correct version is shown opposite.

The editorial team apologize for any confusion this might have caused.

PII: S1359-6446(05)03479-3




NEUROSCIENCE

Towards a cure for Fragile X

Fragile X syndrome (FXS) is the most common inheritable cause of mental retardation. The loss of a single gene, *FMR1*, is sufficient to cause FXS, which is associated with neurological deficits ranging from cognitive impairment to autistic behaviour. It has been postulated that many of these FXS symptoms might be attributed to overactivation of the metabotropic glutamate receptors (mGluR). A fruit fly model for FXS that is based on the loss of *dfmr1*, the *Drosophila* homolog of the *FMR1* gene, displays neuronal and behavioural phenotypes that are parallel to symptoms observed in Fragile X patients. McBride *et al.* now report that enhanced mGluR activity is a conserved feature of the fly model for Fragile X and is responsible for some of the neuronal and behavioural phenotypes [1].

To suppress mGluR activity, the mGluR antagonist 2-methyl-6-(phenylethynyl)pyridine (MPEP) was added to fly food during larvae development and after eclosion. Memory deficit is one of the most prominent aspects of FXS and the analysis of memory phenotypes in *dfmr1* mutant flies was a focus of this study. In particular, courtship conditioning assessment was employed for the analysis of learning and memory phenotypes. A deficit in recall memory was evident in *dfmr1* mutant flies. This memory deficit was rescued by MPEP treatment, thereby implicating mGluR signaling as the underlying cause of the impaired cognitive function in the fruit fly Fragile X model.



CANCER BIOLOGY

Phosphorylation of BRCA2 regulates homologous recombination

Mutations in *BRCA2* cause a predisposition to early-onset breast cancers. Cells lacking *BRCA2* are hypersensitive to DNA-damaging agents and it is thought that tumours are caused by defects in DNA repair pathways that require homologous recombination. *BRCA2* probably mediates homologous recombination by binding to *RAD51*, however it is not clear how this interaction is regulated. Esashi *et al.* show that phosphorylation of Ser3291 of *BRCA2* inhibits *RAD51* binding to the C-terminus of *BRCA2* [2].

RAD51 initiates homologous recombination by forming filaments and mediating strand exchange. The authors made nine GST-*BRCA2* fusions encompassing the entire gene and found that fragments 2, 3 and 9 bound to *RAD51*. Fragments 2 and 3 contained BRC repeats, which bind *RAD51*, whereas fragment 9 contained a novel *RAD51*-binding site. It was shown that this fragment was phosphorylated by cyclin-dependent kinases.

The authors made three truncations of fragment 9 and found that only TR2 bound to *RAD51*. This fusion contained five potential S/T phosphorylation sites, so the authors made a series of Glu-substituted mutants to mimic phosphorylation. Only mutants containing Glu at position 3291 did not bind *RAD51*, showing that phosphorylation of Ser3291 inhibits binding to *RAD51*.

Phosphorylation of Ser3291 was examined *in vivo* using a phosphorylation-specific antibody. Phosphorylation increased as cells progressed from G2 to M, and decreased after cells were treated with ionising radiation. Overexpression of the TR2 fragment reduced the efficiency of recombinational DNA repair. These data show that phosphorylation of Ser3291 was functionally important to regulate the *RAD51*-*BRCA2* interaction.

It was proposed that when Ser3291 is dephosphorylated, it accepts *RAD51* monomers from the BRC repeats. *BRCA2* then facilitates the transfer of *RAD51* to single-stranded DNA. However, further research is required to understand how the interactions between the BRC repeats and *RAD51* are regulated.

- 2 Esashi, F. *et al.* (2005) CDK-dependent phosphorylation of *BRCA2* as a regulatory mechanism for recombinational repair. *Nature* 434, 598–604

Christian Noble

cnoble@nimr.mrc.ac.uk

issues, soon after its X-ray crystal structure bound with ZipA was solved. Instead of abandoning the 3-D data, Rush *et al.* chose to use the X-ray coordinates of the active compound as a prototype to find other molecules with similar shape [1]. To this end, they developed a new similarity algorithm, called rapid overlay of chemical structures (ROCS).

ROCS perceives similarity between two molecules based on their molecular shape only. It approximates their volumes with Gaussian functions instead of hard spheres, thereby resulting in mathematical equations that are analytic and differentiable, which allow for fast and robust global optimization of volume overlap by varying their relative orientation. A similarity function then measures the 'shape distance' between the pair of molecules at optimal overlap of volume. Molecules that ROCS deems near the prototype are similar in shape, but not necessarily built from the same atoms in the same way.

Wyeth's corporate 3-D database of available lead-like compounds was searched by ROCS with two molecular shape queries derived from the prototype. Hits with different scaffolds than the prototype were overlaid on the X-ray coordinates of the prototype bound with ZipA, and those hits with unfavorable intermolecular contacts were discarded, leaving 29 candidates. A final fluorescence polarization assay yielded three non-toxic active molecules with μM -range K_D . One of the three molecules was co-crystallized with ZipA and its binding mode pose was found to be very similar to the pose of the prototype. Two novel scaffolds without obvious IP concerns were extracted from the three molecules.

This study showed that the volume approach of ROCS can 'hop' via the 3-D molecular shape of a known active to scaffolds built from unexpected 2-D chemical structure. These new scaffolds can lead to drug candidates that are out of patented areas of chemistry space. Extensions to the ROCS algorithm might include the scoring of the receptor–ligand contacts based on the matching of classified atoms or functional groups, as in Masek's original MSC method. An additional application of ROCS could be to overlay the training set of molecules for CoMFA methodology, which is a well-known source of difficulty in CoMFA analyses.

- 1 Rush, *et al.* (2005) A shape-based 3-D scaffold-hopping method and its application to a bacterial protein–protein interaction. *J. Med. Chem.* 2005, 48, 1489–1495

Joseph Talafous

talafous@gmail.com

This study validates the relevant memory paradigm in the *Drosophila* model of FXS. Moreover, it raises the possibility that mGluR antagonists could similarly improve cognitive functions in Fragile X patients. In response to mGluR activation, *FMR1* mRNA is translated into its gene product FMRP, which binds a subset of mRNAs and acts as a regulator of translation. As many similarities are found between symptoms observed in Fragile X patients and phenotypes observed in the fly, and that deregulation of mGluR activity appears to be an evolutionarily conserved feature, this model allows for easy access to investigate the intricate relationship between mGluR, mGluR antagonists and the gene product of *dfmr1*.

- 1 McBride, S.M. *et al.* (2005) Pharmacological rescue of synaptic plasticity, courtship behavior, and mushroom body defects in a *Drosophila* model of Fragile X syndrome. *Neuron* 45, 753–764

Herman H. Cheung

herman@mgcheo.med.uottawa.ca



Same shape, different way

Bacterial resistance to available antibiotics is quickly becoming a public health threat. Wyeth is conducting a drug discovery program targeted at the ZipA–FtsZ protein–protein interaction that is required for proper cell division. Unfortunately, the most active compound identified from a fluorescence polarization HTS was found to have toxic side-effects and to have intellectual property (IP)