

200 nm and 100 nm are better enclosed than the 500 nm variants. The enclosure of spheres modified with chromatin binding peptides is indeed higher than the enclosure of the non-targeted spheres and the spheres modified with the mutated AT-hook. When polystyrene spheres were injected in the cytoplasm of HeLa cells, initially, the nanospheres spread homogeneously in the cytoplasm. Upon cell division, however, the nanospheres accumulated in a specific perinuclear region and enclosure in the nuclei of divided cells was never observed. Therefore, reaching the nucleoplasm seems to be very difficult and we question whether the chromatin binding peptides are able to target the nanospheres into the daughter nuclei of living cells. It thus seems that nuclear inclusion in the XEE assay does not represent the situation in living cells

#### Acknowledgements

N. Symens is a predoctoral fellow from the Institute for the Promotion of Innovation through Science and Technology in Flanders. K. Remaut is a postdoctoral fellow of the Research Foundation Flanders. The financial support of these institutes is acknowledged with gratitude.

doi:10.1016/j.drudis.2010.09.360

#### A8

##### mRNA delivery to cervical carcinoma and mesenchymal stem cells mediated by cationic carriers

Joanna Rejman\*, Geertrui Tavernier, Neda Beversad, Joseph Demeester, Stefaan C. De Smedt

Ghent University, Laboratory of General Biochemistry and Physical Pharmacy, Ghent Research Group on Nanomedicines, Harelbekestraat 72, 9000 Ghent, Belgium

\*Corresponding author.

E-mail: joanna.rejman@ugent.be (J. Rejman).

We complexed mRNA encoding luciferase (mLUC) to either one of the cationic lipids Lipofectamine (LF) or DOTAP/DOPE, or to linear poly(ethyleneimine), a cationic polymer (linPEI). After incubating the resulting lipo- or polyplexes with HeLa cells for different periods of time, we determined luciferase expression by a bioluminescence assay. Both extent and duration of luciferase expression were dependent on the type of complex used. With LF, mRNA expression lasted for about 9 days maximally, which is not significantly shorter than what can be achieved with pDNA polyplexes. When electroporation was used to transfer mLUC into the cells, luciferase expression lasted for 12 h only.

An important characteristic of mRNA-mediated transfection by means of all three complexes is that it could already be detected 30 min after adding the complexes to the cells. In order to estimate the number of positive cells, we transfected the cells with an mRNA encoding Green Fluorescent Protein (GFP) and compared the results with transfection by means of pDNA. With transfection by means of mRNA complexed to LF or DOTAP/DOPE a substantially larger fraction of cells (>80%) was transfected than with pDNA (40%). After establishing the characteristics of mRNA-mediated transfection by means of expression of reporter proteins, we tested the carriers for their ability to mediate expression of a functional protein in mesenchymal stem cells. For that purpose we complexed an mRNA encoding CXCR4, a receptor binding stromal derived factor 1, to the cationic lipids and the polymer. The resulting complexes were incubated with mesenchymal stem cells and CXCR4 expression was assayed. The fraction of CXCR4-positive cells was approximately 80% and 40% for mRNA-cationic lipoplexes and lin-PEI polyplexes respectively. The results of these experiments indicate that mRNA, under certain conditions, may be preferable to pDNA to achieve transfection, particularly in cases requiring transient protein expression.

doi:10.1016/j.drudis.2010.09.361

#### A9

##### Cellular uptake of long-circulating pH-sensitive liposomes: evaluation of the liposome and its encapsulated material penetration in cancer cells

Emilie Ducat\*, Julie Deprez, Olivier Peulen, Brigitte Evrard, Géraldine Piel  
Laboratory of Pharmaceutical Technology, CIRM, Department of Pharmacy, University of Liege, Belgium

\*Corresponding author.

E-mail: Emilie.Ducat@ulg.ac.be (E. Ducat).

Print 3G, a peptidic antagonist of oncoprotein involved in breast cancer, could reduce the angiogenic development of breast tumors, leading to tumor dormancy. The necessity of intravenous administration of Print 3G led to the development of long-circulating liposomes as drug carriers. Pegylated liposomes, too large to be collected by fenestrated organs, accumulate passively in solid tumors thanks to the EPR effect. The strategy was to combine the protective properties of PEG with the transfection properties of pH-sensitive lipids that could promote the uptake of liposomes by

cells and avoid lysosomal sequestration and degradation of entrapped materials such as peptides. In this study, we compare two formulations in terms of cellular uptake using confocal microscopy. The first one is composed of SPC:CHOL:mPEG-750-DSPE (47:47:6), used as 'standard' liposomes, and the second one composed of DOPE:CHEMS:CHOL:mPEG750-DSPE (43:21:30:6), used as pH-sensitive liposomes.

Firstly, we evaluated the penetration of an encapsulated model molecule, calcein, in Hs578t human breast cancer epithelial cells. When calcein was encapsulated in standard liposomes, its penetration was effective only in a few cells. On the contrary, the majority of cells were fluorescent when calcein-loaded pH-sensitive liposomes were applied on cells for three hours. Secondly, we studied the penetration of liposomes themselves in Hs578t cells using 25-[(nitrobenzoxadiazolyl)methylamino]nor-cholesterol (NBD-CHOL) as a fluorescent marker of the phospholipid membrane. The obtained results were comparable to those obtained with calcein: a higher penetration of liposome was observed for pH-sensitive liposomes. Finally, the cellular uptake of liposomes using both NBD-CHOL and rhodamine encapsulated in the inner cavity of vesicles was evaluated with Hs578t cells and compared with WI26 human diploid lung fibroblast cells. This experimental design allowed us to follow simultaneously the cell distribution of the encapsulated material and of the liposome itself. Confocal pictures obtained with pH-sensitive liposomes on both WI26 and Hs578t cells allowed us to visualize co-localized red and green of rhodamine and NBD-CHOL, with a higher degree of colocalization in an area close to the nucleus. In comparison with 'standard' liposomes, we observed a higher penetration of the encapsulated material and of the liposome itself in breast cancer cells. Moreover, we visualized a colocalization near the nucleus of liposomes components. From results obtained with fibroblastic cells, there was no difference in terms of cellular uptake between the two formulations. In perspective, we would like to compare these results, obtained with model molecules, with experiments performed with biotinylated Print 3G to assess its cellular distribution. Moreover, it would be interesting to correlate results obtained with confocal microscopy with a possible increase of the peptide efficacy against cancer cells when it is encapsulated in long-circulating pH-sensitive liposomes.

doi:10.1016/j.drudis.2010.09.362