

Prevalence of metabolic syndrome and associated risk factors among Turkish adults: Trabzon MetS study

Cihangir Erem · Arif Hacıhasanoglu · Orhan Deger ·
Murat Topbaş · İlgin Hosver · Halil Onder Ersoz ·
Gamze Can

Published online: 29 January 2009
© Humana Press 2009

Erratum to: Endocr 33(1):9–20 DOI 10.1007/s12020-008-9044-3

In Table 1 in the original publication (*Endocrine* 33(1): 9–20), the Parameter subheadings “Men” and “Women”

were mistakenly transposed. The data labeled as pertaining to Men apply to Women, and the data labeled as pertaining to Women apply to Men.

The online version of the original article can be found under doi: [10.1007/s12020-008-9044-3](https://doi.org/10.1007/s12020-008-9044-3).

C. Erem (✉) · A. Hacıhasanoglu · H. O. Ersoz
Division of Endocrinology and Metabolism, Department of
Internal Medicine, Karadeniz Technical University, Biyokimya
Anabilim Dalı, Endokrinoloji ve Metabolizma Hastalıkları Bilim
Dalı, Trabzon 61080, Turkey
e-mail: cihangirerem@hotmail.com; cihangirerem@yahoo.com

O. Deger · I. Hosver
Department of Biochemistry, Karadeniz Technical University,
Biyokimya Anabilim Dalı, Endokrinoloji ve Metabolizma,
Hastalıkları Bilim Dalı, Trabzon 61080, Turkey

M. Topbaş · G. Can
Department of Public Health, Karadeniz Technical University,
Halk Sağlığı Anabilim Dalı, Endokrinoloji ve Metabolizma,
Hastalıkları Bilim Dalı, Trabzon 61080, Turkey

C. Erem · O. Deger · M. Topbaş
The Trabzon Endocrinological Studies Group, Karadeniz
Technical University, İç Hastalıkları Anabilim Dalı,
Endokrinoloji ve Metabolizma Hastalıkları Bilim Dalı, Trabzon
61080, Turkey