

JCO 2000;18:2828), and may significantly influence systemic treatment. The objective of this study was to analyze the impact of ITL on axillary lymph node involvement, relapse, and mortality risk, in patients both with and without systemic and loco-regional treatment.

Retrospective analysis was conducted of 2,414 patients, who underwent R0 resection of the primary tumor and systematic axillary lymph node dissection (at least 5 lymph nodes resected) for UICC I-III stage breast cancer. Patients with unknown ITL, multifocal tumor spread, central ITL, or tumor location within 15° of the border between outer and inner quadrants were excluded from the study. Median observation time was 6.7 years.

The ITL was within or between the medial quadrants of the breast in 33.6% of the patients (n=810) and in the lateral hemisphere of the breast in 66.4% (n=1,604). Tumor size, histopathological grading, and estrogen receptor status were balanced between patients with lateral and medial ITL. Metastatic axillary lymph node involvement was significantly associated with a lateral tumor location ($P < .0001$). The mean number of axillary lymph node metastases was increased by 29% in cases with lateral ITL (2.2 vs. 1.7, $P = .003$). In a multivariate logistic regression analysis, allowing for ITL, estrogen receptor status, grading and tumor size, ITL was confirmed as significant risk factor ($P = .02$) for axillary lymph node involvement. ITL, however, did not correlate with either disease free survival (DFS) or overall survival (OS), by univariate (DFS: $P = .41$; OS: $P = .57$) or by multivariate analysis (DFS: $P = .16$; OS: $P = .98$).

In contrast to previous reports, we conclude that there is no sufficient evidence to support any independent prognostic significance of ITL in early breast cancer. However, medial tumor location may lead to underestimation of the axillary lymph node involvement, and, thus, inadequate systemic treatment.

427

POSTER

Potential of radiation-induced subcutaneous fibrosis by concomitant use of tamoxifen in adjuvant breast cancer treatment

D. Azria¹, W. Jeanneret Sozzi², A. Zouhair², P.A. Coucke², A. Kramar³, R.O. Mirimanoff², M. Ozsahin². ¹INSERM EMI 0227 CRLC Val d'Aurelle, Radiation Oncology, Montpellier, France; ²Centre Hospitalier Universitaire Vaudois (CHUV), Radiation Oncology, Lausanne, Switzerland; ³CRLC Val d'Aurelle, Biostatistics, Montpellier, France

In this study, we wanted to assess whether concomitant administration of tamoxifen (TMX) and adjuvant radiation therapy (RT) increases the risk of developing subcutaneous fibrosis after conservative or radical surgery in breast cancer patients. Therefore, we evaluated 149 women with breast cancer treated using adjuvant RT after conservative or radical surgery who took part among 399 patients with miscellaneous cancers included in the KFS 00539-9-1997/SKL 00778-2-1999 prospective study evaluating the predictive value of CD4 and CD8 T-lymphocyte apoptosis on the development of radiation-induced late effects. Median age was 57 years (range: 26-82). RT consisted of 50-Gy whole-breast or thoracic wall irradiation in 2-Gy fractions using either Co60 (n = 97) or 6-MV (n = 52) photons completed with a localized external electron boost up to 66 Gy. Adjuvant TMX concomitant with RT was prescribed at a dose of 20 mg/day for a period of five years in 91 patients (61%). All patients receiving TMX were hormonal receptor positive, and none of them received adjuvant chemotherapy. There were 20 premenopausal and 71 postmenopausal patients (median age: 60 years; range: 36-82). Acute and late toxicities were assessed according to CTC 2.0 and RTOG/EORTC grading systems, respectively. Breast volume and skin dose was estimated using physical RT parameters. In a median follow-up of 29 months (range: 23-73), 144 patients are alive with (n = 5) or without disease. Five patients died from breast cancer without any grade 3 side effects (1, 13, 33, 38, and 41 months). Acute toxicity was observed in all but 4 patients (3%). One hundred six (71%), 34 (23%), and 5 (3%) patients experienced grade 1, 2, and 3 acute side effects, respectively. No statistically significant difference was observed between the TMX and no TMX groups in terms of acute toxicity ($p = 0.58$). There was no significant correlation between the early and late toxicity ($R^2 = 0.05$). Thirty-five patients out of 91 (38%) in the TMX group and 15 out of 58 (26%) in the no TMX group experienced grade 2 or 3 late skin toxicity ($p = 0.11$). However, grade 2 or 3 subcutaneous fibrosis was significantly higher in patients treated with concomitant TMX (42 patients out of 91 (46%) in the TMX group vs. 10 out of 58 (17%) in the no TMX group; $p = 0.0002$). Breast volume and skin dose did not interfere with subcutaneous fibrosis. However, grade 3 telangiectasia incidence was more frequent but not statistically significant in patients with high breast volume irradiated with Co60 compared to 6-MV photons. We conclude that the concomitant use of TMX with RT increases significantly subcutaneous fibrosis. In patients treated with adjuvant hormonal treatment, TMX should be delayed until the completion of RT.

428

POSTER

The predictive value of specimen radiography to predict margins involvement in 188 breast intraclinical carcinomas.

C. Mazouni¹, R. Rouzier¹, C. Baileguier², M.C. Mathieu³, S. Delaloge⁴, H. Marsiglia⁵, J.-R. Garbay¹, J.-R. Garbay¹. ¹Institut Gustave Roussy, Surgery, Villejuif, France; ²Institut Gustave Roussy, Radiology, Villejuif, France; ³Institut Gustave Roussy, Pathology, Villejuif, France; ⁴Institut Gustave Roussy, Oncology, Villejuif, France; ⁵Institut Gustave Roussy, Radiotherapy, Villejuif, France

Background: This study was undertaken to evaluate the role of specimen radiography in predicting margins status for breast palpable lesions.

Material and methods: We retrospectively reviewed clinical, pathologic data and specimens radiography from 188 patients with DCIS referred in our centre between 1997 and 2000 for microcalcifications discovered at breast screening. The lesions were preoperatively localised by using a guide-wire. Specimen radiographic findings and clinico-pathological data were correlated with margin status.

Results: A total of 188 lesions revealed pure ductal carcinoma in situ (DCIS) in 125 (66%) and mixed carcinoma in 63 (34%). On specimen radiographs, the lesions were closed (<5mm) to one edge of lumpectomy in 74 (39%) cases. Histologic margins were positive in 86 cases (46%) and close (< 5 mm) in 51 (27%) cases. The factors associated with positive margins, in the univariate analysis, were a distance less than 5 mm from the tumour to the edge of the specimen radiograph ($p = 0.04$) and multifocality ($p = 0.05$). In the multivariate analysis (logistic regression), a radiologic margin <5mm was the only risk factor for close histologic margins. We therefore tested radiologic margin <5 mm as a potential tool to decrease the risk of close histological margins. Sensitivity, specificity, predictive positive and negative values are reported in Table 1.

Efficacy of specimen radiographs for detecting residual tumors

	Pathologic findings	
	Margins <5mm	Free margins >5mm
Incomplete excision	82 (43.6%)	6 (3.2%)
Complete excision	60 (32%)	40 (21.2%)
Se = 58% Sp = 87%	VPP = 93%	VPN = 40%

Conclusions: Specimen radiograph findings were found to be a predictive factors of margins involvement when tumor distance to the margin was less than 5 mm and may therefore lead the surgeon to perform an additional excision.

429

POSTER

Breast cancer: immediate breast reconstruction after radical surgery

D.D. Pak. P.A. Herzen Cancer Research Institute, General Oncology, Moscow, Russian Federation

Background: We reviewed the results of 11 years' experience with different techniques for immediate reconstruction in breast cancer patients.

Materials and methods: Between 1991 and 2002 a total of 611 breast cancer patients were operated. All patients were divided into 4 groups according to the type of breast reconstruction technique. Group I included 154 patients (stage IIa, IIb) who underwent quadrantectomy and regional lymph node dissection with immediate breast reconstruction using major pectoral (32) and latissimus dorsi (LD) (122) flaps. Group II - 386 women (stage IIa, IIb, IIIa) who underwent breast resection (70-90% of breast tissue and regional lymph nodes were removed). Breast was reconstructed using LD flaps. Group III - 46 patients (stage IIa, IIb, IIIa) who underwent modified radical mastectomy. Breast was reconstructed using LD (18) and pedicled TRAM (28) flaps. Group IV - 25 women (stage I, IIa, IIb, IIIa) who underwent skin/areola sparing mastectomy. In all cases breast was reconstructed using LD flap and silicone breast implant. Radiotherapy and drug therapy were administered depending on the stage of disease, receptor status, etc.

Results: In group I 5-year overall survival rate was 92.4% and 87.5% for stages IIa and IIb respectively. Local recurrence rate was 0% and 4.6%. Esthetic results: excellent - 21.6%, good - 49%, satisfactory - 26%, poor - 2%. In group II 5-year overall survival rates were as follows: IIa - 88.3%, IIb - 84.3%, IIIa - 72.4%. Esthetic results: excellent - 21.6%, good - 48%, satisfactory - 28.4%, poor - 2%. The rate of immediate postoperative complications was 7.3% (28 patients): total necrosis of the flap - 3, marginal necrosis - 2, wound suppuration - 2, bleeding - 3. In group III 5-year overall survival rates were as follows: IIa - 87.9%, IIb - 83.3%, IIIa - 73.6%. Local recurrence rate was 1.9%. Esthetic results: excellent - 21.7%, good - 49.6%, satisfactory - 28.4%, poor - 1%. The rate of immediate