

Methods and Materials: Sixteen head and neck cancer patients, irradiated between September 1999 and November 2000 using a conformal parotid-sparing technique, were included in this study. Before RT and seven months after RT a salivary gland scintigraphy was performed in all patients combined with a single photon emission computed tomography (SPECT). The salivary excretion function (SEF) was measured, after stimulation, in 8-12 transversal 5 mm SPECT slices of each parotid. Loss of salivary function in these areas (dSEF%) was calculated as a proportion of the excretion function before and after RT. Because the planning CT-scan and the SPECT analysis were performed in the same treatment position, doses to areas within the parotid gland could be matched with the dysfunction of that respective area. For each patient, dose-dysfunction plots were performed and curve fitting was done.

Results: At baseline level, all but one patient had a normal salivary excretion function at the level of both parotid glands with small variation between the functionality of the different areas within the same gland. Seven months after RT, the reduction in salivary excretion function reached statistical significance for both parotids and a huge variation in functionality of the different areas within the same gland could be seen. When plotting the dysfunction of the different areas within one gland at seven months after RT against the dose these areas received, a sigmoidal function could be fit in seven (7/15) patients to the plots of both parotids and in five (5/15) patients to the plot of one parotid gland. In three (3/15) patients no dose-response curve could be fit to the plots due to a total dysfunction of all partial volumes or an absence of difference in dose between the volumes. In nine (9/15) patients a large variety of functional responses could be obtained at low irradiation doses (10-15 Gy), ranging from an improvement of function to a total loss.

Conclusion: Salivary SPECT is a useful tool for the evaluation of functional loss after RT of different areas within the parotid gland. While these areas are acting as functional sub-units under normal conditions, their dysfunction after RT does not seem to be only the result of the absolute dose these volumes received but also of the mean dose to the entire parotid gland and the dose to the surrounding areas.

470

POSTER

Severe radiation-induced bowel complications are not uncommon in patients with uterine cervix cancer

E.-Y. Huang, C.-J. Wang, H.-C. Chen, L.-M. Sun, F.-M. Fang, S.-A. Yeh, H.-C. Hsu, C.-Y. Hsiung. *Kaohsiung Chang Gung Memorial Hospital, Radiation Oncology, Kaohsiung Hsien, Taiwan*

Background: To evaluate the factors of severe (Grade 3-4) bowel complications after external irradiation and high dose-rate intracavitary (HDR-IC) brachytherapy among patients with cervical cancer.

Materials and Methods: We reviewed 298 patients of stage IB-IVA cervical cancer managed by curative-intent radiotherapy from May 1993 through December 1997. External irradiation to whole pelvis (34.2-50 Gy/19-27 fractions) was delivered to all patients initially. Two hundred and three patients received additional bilateral parametrial boost (3.6-18 Gy/2-10 fractions) with 4-cm midline shielding. HDR-IC brachytherapy, 16-24 Gy/5 fractions to Point A, was given after external irradiation. External parametrial dose < 50 Gy, 50-54 Gy and > 54 Gy were categorized as low parametrial dose (LPMD), intermediate parametrial dose (IPMD) and high parametrial dose (HPMD) group, respectively. Cumulative rectal biologic effective dose (CRBED) < 85 Gy, 85-105 Gy and > 105 Gy were categorized as low cumulative rectal biologic effective dose (LCRBED), intermediate cumulative rectal biologic effective dose (ICRBED) and high cumulative rectal biologic effective dose (HCRBED) group, respectively. The actuarial rate of bowel complications was compared among groups. We used Cox regression for multivariate analysis of bowel complications.

Results: Grade 3-4 bowel complication rates were 16%. The rates were 7%, 13%, and 34% in the LPMD, IPMD, and HPMD group ($p=0.0001$), respectively. The rates were 5%, 11%, and 28% in the LCRBED, ICRBED, and HCRBED group ($p=0.0002$), respectively. In multivariate analysis of Grade 3-4 radiation-induced bowel complications, CRBED ($p=0.0010$) and external parametrial dose ($p=0.0007$) were independent factors.

Conclusions: Radiation-induced severe bowel complications are dependent on external parametrial dose and CRBED. We do not suggest external parametrial dose > 54 Gy and CRBED > 105 Gy for treatment of cervical cancer due to relatively high incidence of severe bowel complications.

471

POSTER

Radiation-induced changes in the cytoskeleton of human endothelial cells in relation to endothelial monolayer permeability.

D. Gabrys^{1,3}, K. Trott^{2,3}, G. Tozer³, C. Kanthou³. ¹Center of Oncology M.S.-Curie Memorial Institute, Radiotherapy Department, Gliwice, Poland; ²St. Bartholomew's and the Royal London School of Medicine and Dentistry, Department of Radiation Biology, London, United Kingdom; ³Gray Cancer Institute, Tumour Microcirculation Group, Northwood, United Kingdom

Treatment of solid tumours by radiotherapy damages the tumour cells as well as the microvasculature of the tumour and surrounding normal tissues. An increase in vascular permeability is a well known effect of radiation, which contributes toward changes in the interstitial space that lead to reduction of parenchymal cell function, necrosis and fibrosis. A response to radiation by normal tissue endothelium is thought to be a major limitation for its use in cancer treatment.

The aims of the present study were to investigate the early effects of radiation on the cytoskeleton of cultured endothelial cells and relate these to changes in endothelial monolayer permeability.

Human dermal microvascular endothelial cells (DMEC) were irradiated using a Pantak X-ray machine. Using immunofluorescence techniques, DMEC were found to respond to various doses of radiation (0.5 - 20 Gy) by a rapid (within minutes) increase in actin reorganization into stress fibers, accompanied by changes in the distribution of the adherens junction protein, VE-cadherin. Increased endothelial stress fiber formation, cell contraction and redistribution of junctional proteins can lead to intercellular gap formation and changes in monolayer permeability. Therefore, changes in permeability were quantified by monitoring the passage of fluorescently-labelled dextran through DMEC monolayers grown on microporous filters. Radiation was found to induce a significant increase in DMEC permeability suggesting that the changes in the actin cytoskeleton and distribution of cadherins were associated with increased monolayer permeability. Activation of the GTPase Rho and its associated Rho kinase have been recognised as key regulators of the actin cytoskeleton, intercellular junction integrity and permeability. Analysis of the mechanisms involved in irradiation-induced actin reorganisation, re-distribution of VE-cadherin and increased permeability revealed that these effects of radiation were dependent on activation of Rho kinase, since they were blocked by the highly specific Rho kinase inhibitor Y-27632. Simvastatin, a 3-hydroxy-3-methylglutaryl CoA reductase inhibitor used clinically for the treatment of hypercholesterolemia, is known to inactivate Rho by inhibiting its geranylgeranylation. Simvastatin was found to inhibit the radiation mediated changes in actin and junctional proteins as well as increased permeability suggesting the involvement of Rho.

These data provide an insight into possible mechanisms involved in radiation-induced changes in vascular permeability. Further investigations are needed in order to elucidate whether compounds such as Y-27632 and simvastatin may be useful in counteracting some of the side effects of radiation *in vivo*.

472

POSTER

Efficacy and morbidity of linear accelerator radiosurgery for cerebral arteriovenous malformations: a comparison with the natural history

M.A. Bollet¹, L. Picard², I. Buchheit¹, R. Anxionnat², A. Cordebar², N. Jay³, E. Desandes³, C. Marchal¹, P. Aleotti¹, P. Bey⁴. ¹Centre Alexis Vautrin, Radiotherapy, Vandoeuvre, France; ²University Hospital of Nancy, Neurosurgery, Nancy, France; ³Centre Alexis Vautrin, Statistics, Vandoeuvre, France; ⁴Faculty Of Medicine, Nancy, France

Background: To report the results of arc-therapy radiosurgery in terms of efficacy. To compare the adverse-effects rate with the one expected from the natural history.

Material and methods: At the University Hospital of Nancy, 217 patients have been treated for cerebral ArterioVenous Malformation (AVM) by linear accelerator radiosurgery since 1992. We report here the results of a retrospective study of the 118 first patients (55 men, 63 women) treated between 01/07/92 and 30/06/98. The mean follow-up was 46 months (5-105). The mean age was 35 years (13-65). AVMs had poor prognosis features at initial presentation: existence of previous therapeutic failures (85%), high Spetzler-Martin grade (67% of grade III or higher), large size (57% > 14cc) and a high rate of initial hemorrhage (54%). Patients had already been treated by previous embolizations with a mean number of 4 procedures (1-11) in 84% of patients (99/118); 79% by embolizations alone (93/118) and 5% by partial microsurgery and embolization (6/118).