

21 INVITED Legal and ethical requirements of human tissue research

C. Trouet. *Pharma.be, Brussels, Belgium*

The increasing possibilities for using tissues for research and the developments in genetics and biotechnology have made stored human biological materials more important than ever. Using stored human biological materials raises a lot of legal and ethical questions (commercialization, protection of privacy, implementation of informed consent, new findings, role of research ethics committees etc.). Research with human subjects and research with personal data is covered by detailed European regulation. The research use of human biological materials however has not been regulated in a detailed manner so far. After discussing the relevant regulation on the level of the European Union and the Council of Europe (EU Directive for the legal protection of biotechnological inventions; Council of Europe Convention on Human Rights and Biomedicine), we will briefly indicate the major issues in retrieving tissues for future research use and in using existing tissue banks for research purposes. A special focus will be on the implementation of the informed consent requirement in daily practice.

Wednesday 29 September 10:15–12:00

WORKSHOP 5 Tumour vaccines

22 INVITED Human dendritic cell subsets and their implications for clinical studies

C.G. Figdor, J. de Vries, J. Lesterhuis, G.A. Adema, C. Punt.
Departments of Tumor Immunology and Medical Oncology, Nijmegen Centre for Molecular Life Sciences, Radboud University Medical Centre, Nijmegen, The Netherlands

Dendritic cells (DC) are the professional antigen presenting cells of the immune system. They possess the unique capacity to take up and process antigen, migrate to the draining lymphnode, and present antigen to resting lymphocytes. This is the reason why several groups have embarked on the use of these cells for vaccination purposes. We have investigated the capacity of immature and mature monocyte-derived dendritic cells (DC) pulsed with melanoma-associated peptides (gp100 and tyrosinase) to induce a primary cytotoxic T lymphocyte response *in vivo*. Advanced melanoma patients were vaccinated with peptide- and keyhole limpet hemocyanin (KLH)-pulsed DC, either immature or matured DC Blood, Delayed type hypersensitivity (DTH) reactions against antigen-pulsed DC and tumor sites have been investigated for immunological responses and will be discussed. Results indicate that mature DC are superior to immature DC in the induction of immunological responses in melanoma patients. In other studies we have compared the capacity of immature and mature monocyte-derived DC to migrate to draining lymph nodes. To this end DC were radiolabeled with indium. Melanoma patients who were vaccinated with DC-vaccines received in addition to their first vaccine, radiolabeled DC to study their migratory capacity *in vivo*. In some cases lymph-nodes were resected and autoradiography was used to determine the number of DC that migrated into the T cell area of the lymph node. Results show that mature DC migrate much better when compared to immature DC. In this presentation I will not only talk about the current state of the art but also discuss issues that still have to be investigated and new opportunities that will broaden the use of this form of immunotherapy in the clinic.

23 INVITED Development of heat shock cancer vaccine strategies

G. Parmiani. *Istituto Nazionale Tumori, Unit of Immunotherapy of Human Tumors, Milan, Italy*

Heat Shock Proteins (HSPs) play an essential role in protein metabolism and exert stimulatory activities on innate and adaptive immunity. Vaccination with tumor-derived HSPs induces CD8⁺ T cell-mediated tumor regressions in animal models. I will show that HLA-A*0201-restricted CD8⁺ T cells recognizing antigens (Ags) expressed by human melanoma (e.g. Melan-A/MART-1) or colon carcinoma (CEA and EpCAM) were triggered to release IFN γ and to mediate cytotoxicity by HLA-A*0201 matched antigen presenting cells pulsed with HSP96 purified from tumor cells expressing the relevant Ag. Such activation occurred in class I

HLA-restricted fashion and appeared to be significantly higher than that achieved by direct peptide loading. Immunization with autologous tumor-derived HSP-peptide complex 96 (HSPPC-96) in either stage IV metastatic melanoma or liver metastasis-resected colon-rectal carcinoma (CRC) subjects, induced a significant increase in the recognition autologous and/or HLA-compatible tumor cells in approximately 50% of patients in both trials. Moreover, anti-Melan-A/MART-1₂₇₋₃₅, anti-CEA₅₇₁₋₅₇₉ and anti-EpCAM₂₆₃₋₂₇₁ T cell reaction was elicited in 3 out of 5 and in 2 out of 5 HLA-A*0201 melanoma and CRC patients tested, respectively. Analysis of NK activity also demonstrated an increase in vaccinated patients in both studies. The increments in Ag-specific T cell responses were associated with a favourable disease course after HSPPC-96 vaccination. A more recent phase II trial was conducted in metastatic melanoma patients with HSPPC-96 and GM-CSF as adjuvant. However, the addition of GM-CSF did not increase the frequency of patients that mounted a tumor-specific T cell response nor the clinical outcome as compared to the previous study of melanoma patients given the HSPPC-96 only. Altogether, our data provide evidence that HSPPC-96 derived from human tumors can activate the innate immunity and present antigenic peptides to patients' CD8⁺ T cells and activate them both *in vitro* and *in vivo*. Thus HSPPC-96 appears to represent an important tool for vaccination in cancer patients. Phase III prospective studies are ongoing in melanoma and renal cell carcinoma and will determine the efficacy of HSPPC-96-based vaccination in these diseases.

24 INVITED Experimental strategies to enhance the potency of dendritic cell-based cancer vaccines for clinical use

J.J. Mulé. *H. Lee Moffitt Cancer Centre, Translational Science and Technology Development Melanoma Research and Treatment, Tampa, USA*

The clinical application of immunotherapy for cancer is rapidly moving forward in multiple areas, which incorporates the adoptive transfer of antitumor-reactive T cells and the use of "therapeutic" vaccines. Both clinical and immunologic endpoints have shown new promise to the field. Novel dendritic cell-based vaccine strategies designed in the laboratory and proven in preclinical animal tumor models are now entering the clinic, with the intent to provide therapeutic efficacy. Improvements on this approach involve breaking tolerance to tumor "self" antigens by inhibiting regulatory cells, boosting T cell co-stimulation, and combinations of recombinant cytokines and other defined molecules with "immuno-enhancing" activities. This review presentation revisits our earlier reported dendritic cell-based vaccine trials in cancer patients and focuses on current and future approaches in the clinic.

Based on encouraging data from our murine studies (1), we initiated and completed two phase I clinical trials of autologous tumor lysate/KLH-pulsed dendritic cells (TP-DCs) in pediatric and adult patients with advanced solid tumors under an IND approved by the FDA in 1996 (2-4). The DCs were produced from adherent PBMC of leukapheresis collects by culture in GM-CSF and IL-4; they were characterized as immature by phenotypic marker profiling. Both immunologic and very modest patient clinical responses in melanoma were noted in these early studies. Autoimmune manifestations were not detected. Although immunologic assays (e.g., IFN-gamma ELISPOTs) revealed evidence of the induction of peripheral blood T cell reactivity to both KLH and tumor lysates post-immunization, particularly in our pediatric patients, durable and complete regressions of established tumors were not achieved through the administration of TP-DCs alone. Based on our recent animal studies, we have now embarked on a new series of clinical trials to evaluate potential improvements in our DC-based vaccine strategy that incorporate the intratumoral delivery of KLH-pulsed DCs (5); the systemic administration of IL-2 (6,7); the use of locally-produced secondary lymphoid tissue chemokine (SLC; 8-10); and the setting of lymphopenia-induced, homeostatic-driven T cell proliferation (11). In this regard, we are planning to conduct the following clinical trials in advanced cancer patients: 1) a phase I trial of escalating doses of Fludarabine/Cy followed by intranodal delivery of MHC class I/II peptide-pulsed DCs in patients with chemotherapy-naïve metastatic melanoma; 2) a randomized phase II trial of Fludarabine/Cy followed by intranodal delivery of peptide-pulsed DCs with or without autologous lymphocyte infusion; and 3) a phase I clinical study assessing autologous DCs gene-modified to secrete SLC as a vaccine to enhance T cell priming in patients with advanced melanoma or colorectal cancer.

Although vaccinations involving TP-DCs have been performed, little, if any, information is available on the effects of phagocytic uptake of tumor lysates on DC biology and function, and how this activity can be influenced to enhance the therapeutic potency of tumor vaccines. We have investigated gene expression pattern differences between unpulsed DCs (UP-DCs) and melanoma TP-DCs, using Affymetrix MG-U74Av2 oligonucleotide arrays, which contain ~12,000 genes and ESTs (expressed sequence tag) (12). Upon 24 hr tumor lysate pulsing, the levels of 87 transcripts increased at