

(n = 22) visceral metastases had a median survival of 8 months vs. 12.5 months, respectively (p = 0.014). The group of patients without visceral metastases could be further subdivided by adding the gene expression values. Addition of each single gene expression separated the survival curve for these patients as exemplified in the figure, where patients with low expression values had a median survival time of 30.5 months vs. 10 months for patients with high expression values (p = 0.0001). Addition of further gene expression profiles resulted in further separation of the survival curve. **Conclusions:** We have identified five genes with a stronger prognostic impact than the known clinical prognostic factors. Confirmation of results is ongoing.

801

ORAL

Endoglin and CD31 expression in relation to prognosis in conventional renal cell carcinoma

J. Sandlund¹, Y. Hedberg², A. Bergh², K. Grankvist³, B. Ljungberg⁴, T. Rasmussen¹. ¹Umeå University, Dept. of Radiation Sciences, Oncology, Umeå, Sweden; ²Umeå University, Dept. of Medical Biosciences, Pathology, Umeå, Sweden; ³Umeå University, Dept. of Medical Biosciences, Clinical Chemistry, Umeå, Sweden; ⁴Umeå University, Dept. of Surgical and Perioperative Sciences, Urology and Andrology, Umeå, Sweden

Background: Quantification of intratumoural microvessel density by endoglin (CD105) or CD31 (PECAM-1) staining has prognostic significance in selected neoplasms. Endoglin is a cell membrane glycoprotein expressed on tumour-associated vascular endothelium and it is a marker of angiogenesis. CD31 is a transmembrane glycoprotein belonging to the immunoglobulin superfamily of cell adhesion molecules presenting on the surface of various blood cells and endothelial cells. In this study, the prognostic information of endoglin and CD31 expression in human renal cell carcinoma (RCC) was evaluated.

Material and methods: Tumour samples from 162 patients with conventional RCC treated between 1982 and 1997 were analysed. The tumour samples were assessed using the tissue microarray technique and immunohistochemically stained for endoglin and CD31. The expression was related to gender, age, TNM stage, nuclear grade, tumour size, and survival data.

Results: The expression of endoglin was inversely associated with TNM stage (p = 0.019), and nuclear grade (p = 0.006). The expression of CD31 was inversely associated to TNM stage (p = 0.03) and to nuclear grade (p = 0.018). Furthermore, a correlation between the expression of endoglin and CD31 was seen (r = 0.439, p < 0.001). No correlation was found between endoglin or CD31 expression and gender, age, or tumour size. The material was subdivided in quartiles depending on the endoglin and CD31 expression. Patients with high CD31 expression (the highest quartile) had better prognosis compared to those with lower expression (p = 0.015). Endoglin showed a similar trend (p = 0.06). A multivariate analysis of prognostic factors showed that TNM stage and nuclear grade were independent predictors of prognosis. Endoglin or CD31 expression did not add further prognostic information.

Conclusion: The expression of endoglin and CD31 in conventional RCC is inversely related to stage and grade. Furthermore a correlation between the expression of endoglin and CD31 was observed. When comparing the endoglin and CD31 expression in conventional RCC, a higher sensitivity of one of the angiogenic factors over the other cannot be suggested. TNM stage and nuclear grade remains the strongest predictors of prognosis in RCC, but the results indicate that angiogenesis is related to prognosis.

802

ORAL

High detection rate of circulating tumor cells in blood of patients with prostate cancer using telomerase activity

K. Fizazi¹, L. Morat², L. Chauveinc², D. Prapotnich³, B. Escudier¹, X. Cathelineau³, F. Roze³, G. Vallancien³, L. Sabatier², J.C. Soria¹. ¹Institut Gustave Roussy, Medicine, Villejuif, France; ²Commissariat à l'Energie Atomique, Biology, Fontenay, France; ³Institut Mutualiste Montsouris, Urology, Paris, France

Objectives: To study whether a method using telomerase activity allows to isolate circulating tumor cells (CTC) in patients with prostate cancer.

Material and methods: Peripheral blood mononuclear cells (PBMC) were isolated from blood by using Ficoll/hypaque. Immunomagnetic beads coated with an epithelial-specific antibody (BerEP4) were used to harvest epithelial cells from PBMC. Telomerase activity was detected in harvested epithelial cells using the Telomerase-PCR-ELISA method.

Results: Blood samples from 107 patients with prostate cancer were studied. CTC were detected in 19/24 (75%) patients with advanced prostate cancer. In contrast, CTC were not detected in blood samples from 19

healthy male volunteers. CTC could be identified even in patients with a very low serum PSA (<0.1 ng/mL). CTC were detected in 55/70 (79%) when tested in patients with localized prostate cancer who were planned to be treated by radical prostatectomy (n = 30) or brachytherapy (n = 40). CTC could also be detected in 3/13 patients (23%) with an undetectable prostate specific antigen (PSA) at least 1 year after radical prostatectomy, which is consistent with the expected relapse rate in this setting.

Conclusion: CTC can be detected using telomerase activity in a large majority of patients with prostate cancer, including those with a localized stage. This method appears to be more sensitive than RT-PCR methods to detect CTC. Potential applications include tumor monitoring after definitive local therapy and accessibility to malignant material in the metastatic setting.

803

ORAL

Doppler ultrasonography with perfusion software and contrast agent injection as an early evaluation tool of metastatic renal cancers treated with the RAF kinase and VEGFR inhibitor: a prospective study

M. Lamuraglia¹, L. Chami¹, B. Escudier², B. Schwartz³, J. Leclère¹, N. Lassau¹. ¹Institut Gustave-Roussy, Imaging, Villejuif, France; ²Institut Gustave-Roussy, Medicine, Villejuif, France; ³Bayer, West Haven, USA

Objectives: The objective of this study was to evaluate Doppler Ultrasonography with perfusion software (Vascular Recognition Imaging, Amplio, Toshiba) and contrast agent injection (Sonovue® –Bracco) (DUSVRI) as a predictor of tumor response to new treatment BAY 43–9006 under investigation in phase III trials for the treatment of metastatic renal cancer.

Material and methods: Tumor vascularization in accessible targets was prospectively studied, (double-blind study with a placebo) with DUSVRI. The examinations were performed before administering BAY 43–9006 (day 1) and at 3 and 6 weeks. The percentage of contrast uptake was evaluated in each tumor by two radiologists. Results were compared to CT scan studies at 6 weeks.

Results: 30 patients were included and a total of 85 examinations were performed: 30 before randomization, 28 at 3 weeks and 27 at 6 weeks. The results showed a decrease in tumor vascularization in 10 patients out of 28 patients at 3 weeks and in 10 patients out of 27 patients at 6 weeks. The final results concerning 30 patients will be presented with a correlation with the CT-scan response at 6 weeks.

Conclusion: DUSVRI is a new cost-effective and simple non invasive imaging technique. Its effectiveness in predicting the efficacy of BAY will be presented.

804

ORAL

Cancer nanotechnology: drug encapsulated nanoparticle-aptamer bioconjugates for targeted delivery to prostate cancer cells

O.C. Farokhzad^{1,2}, J. Cheng², B. Teply², A. Khademhosseini², S. Jon³, E. Levy-Nissenbaum^{1,2}, R. Langer². ¹Brigham and Women's Hospital, Harvard Med School, Department of Anesthesiology, Boston, Massachusetts, USA; ²Massachusetts Institute of Technology, Division of Health Sciences and Technology, Cambridge, Massachusetts, USA; ³Gwangju Institute of Science & Technology, Department of Life Science, Gwangju, South Korea

Introduction: Nucleic acid ligands (aptamers) are potentially well suited for the therapeutic targeting of drug encapsulated controlled release polymer nanoparticles in a cell- or tissue-specific manner. We used Prostate Cancer (PCa) cells as a model to test this hypothesis.

Methods: We synthesized poly(lactic acid)-block-poly(ethylene glycol) controlled release copolymer with a terminal carboxylic acid functional group (PLA-PEG-COOH), and encapsulated rhodamine-labeled dextran (as a model drug) within PLA-PEG-COOH nanoparticles using the double emulsion method. We generated nanoparticle-aptamer bioconjugates using nuclease stabilized RNA aptamers that bind to the Prostate Specific Membrane Antigen (PSMA), a well known PCa tumor-marker which is over-expressed on prostate acinar epithelial cells. These bioconjugates were examined for targeted delivery and uptake by LNCaP (PSMA+) and PC3 (PSMA-) model PCa cells under a range of physiologic shear stress conditions using microfluidic channels.

Results: Nanoparticles had the following desirable characteristics: 1) negative surface charge ($-50 \text{ mV} \pm 3 \text{ mV}$, Mean \pm SD, N = 3), which may minimize non-specific interaction with the negatively charged nucleic acid aptamers, 2) carboxylic acid groups on the particle surface for potential modification and covalent conjugation to amine-modified aptamers, 3) presence of PEG on particle surface which enhances circulating half-life while contributing to decreased uptake in non-targeted cells. Nanoparticles were conjugated to PSMA aptamers to develop the first example of a