

1063

POSTER

Ki-67 and vascular endothelial growth factor (VEGF) in prediction of tumor growth and lymph node metastasis in head and neck squamous cell carcinomas

V. Boonkitticharoen¹, B. Kulapaditharom², J. Leopairut³, P. Kraiphibul¹, N. Larbcharoen³, W. Cheewaruangroj², C. Chintrakarn², M. Dhanachai¹, L. Pochanukul¹. ¹Ramathibodi Hospital, Mahidol University, Department of Radiology, Bangkok, Thailand; ²Ramathibodi Hospital, Mahidol University, Department of Otolaryngology, Bangkok, Thailand; ³Ramathibodi Hospital, Mahidol University, Department of Pathology, Bangkok, Thailand

Background: The TNM staging is an important guide for treatment but does not explain biology of tumor growth and progression during clinical course. Molecular markers reflecting growth and metastatic potentials of a cancer would be helpful in directing better therapeutic decision.

Materials and methods: One hundred surgical specimens of head and neck squamous cell carcinomas were stained by immunohistochemistry for expression of Ki-67 (K) and VEGF(V). Cutoff points for high and low marker expression were determined to allow grouping of the bifunctional markers into $K_{low}V_{low}$, $K_{high}V_{low}$, $K_{low}V_{high}$ and $K_{high}V_{high}$. Bifunctional marker profiles were analyzed for their associations with pathological T and N stages, tumor site and differentiation.

Results: Analysis of likelihood ratio (LR) to advanced T stage and lymph node metastasis when either or both markers were highly expressed revealed that $K_{high}V_{low}$ (LR=2.63, $p<0.001$) or $K_{low}V_{high}$ (LR=3.13, $p<0.001$) posed significant risk to T3/T4. Simultaneously enhanced expression of both markers not only facilitated tumor growth to T3/T4 (LR=2.86, $p<0.001$) but also promoted lymph node metastasis (LR=2.31, $p<0.001$). In two clinically extreme groups: patients with small tumors (T1/T2) and positive lymph nodes at presentation mostly expressed $K_{high}V_{high}$ ($p<0.001$) as opposed to patients with large tumors (T3/T4) without node involvement in whom tumors tended to have low expression of either or both markers ($p<0.05$). Differences in bifunctional marker profiles were observed among tumor sites ($p<0.05$). Distributions of tumor staging in these sites could be accounted for this finding. For tumor differentiation, moderately and poorly differentiated tumors inclined to express $K_{high}V_{high}$ while well differentiated tumors showed no inclination for any marker profiles ($p<0.05$).

Conclusion: Different patterns of Ki-67 and VEGF expression could help to explain biology of tumor growth and lymph node metastasis during pretreatment clinical course.

1064

POSTER

Overexpression of HSP27 confers chemoresistance which is associated with cell growth delay and inactivation of p38 kinase in Hep-2 laryngeal cancer cells.

D. Sun¹, K. Cho¹, M. Kim¹, S. Cho¹, J. Lee². ¹The Catholic University of Korea, Dept of Otolaryngology-HNS, Seoul, Korea; ²The Catholic University of Korea, Dept of Biochemistry, Seoul, Korea

Among the heat shock protein (HSP) families, Hsp70 and Hsp27 have been implicated in the tumorigenesis and chemoresistance, probably by regulation of cell proliferation and prevention of apoptosis. Hsp27 levels were often increased in large populations of tumors of head and neck, but the mechanisms of its chemoresistance are not clear yet. Here, we investigated the role of Hsp27 in resistance to cytotoxic stress using Hep-2 human laryngeal cancer cells overexpressing Hsp27. Hsp27 overexpression induced cellular resistance to heat shock at 45°C for 1 hour as well as against several cytotoxic agents, such as cisplatin, staurosporin and H_2O_2 . But there was no difference in the sensitivity against irradiation or serum starvation. Moreover, Hsp27 overexpressing Hep-2 cells showed delay in cell growth as compared with control cells. To examine if the decreased cell proliferation in Hsp27 overexpressing cells contributed to the chemoresistance, control Hep-2 cells were synchronized at late G1 phase by mimosine treatment. Synchronization of Hep-2 cells exhibited resistance to cisplatin, staurosporin and H_2O_2 , but not to irradiation and serum starvation, correlating the protection effect shown in Hsp27 overexpressing cells. Therefore, the protective effect of Hsp27 against several cytotoxic agents was based on the cell growth delay. Interestingly, the cell growth and the cytotoxicity of Hep-2 cells were suppressed by SB203580, a specific inhibitor of p38 MAP kinase, but not by PD098059, a specific inhibitor of ERK, or specific inhibitor of JNK. Furthermore, overexpression of Hsp27 lead to suppression of drug-induced p38 kinase activation and subsequent Hsp27 phosphorylation, while ERK and JNK activities were not changed by overexpression of Hsp27. Taken together, our results suggest that overexpression of Hsp27 in Hep-2 cells confers chemoresistance which is associated with the cell growth delay and p38 kinase inactivation. However, the substrate of p38 kinase which induces growth inhibition should be

identified to define the molecular mechanism by which Hsp27 induces growth inhibition in laryngeal cancer cells.

1065

POSTER

Clinical significance of Bcl-2 expression at the invasive front of oral squamous cell carcinomas

K. Kato¹, S. Kawashiri¹, A. Tanaka¹, E. Yamamoto¹. ¹Kanazawa University Graduate School of Medical Science, Department of Oral & Maxillofacial Surgery, Kanazawa, Japan

Background: Apoptosis is a genetically determined process playing an active role in tissue size regulation, morphogenesis and removing damaged cells that could be potentially dangerous for their host. Proteins of the Bcl-2 family play a key role in the control of apoptosis and carry out both proapoptotic and antiapoptotic functions. The previous study has reported that bcl-2 expression was the risk factor of oral squamous cell carcinoma. The present study evaluated the prognostic value of antiapoptotic protein expression at the invasive front of oral squamous cell carcinomas, considering the clinicopathological findings.

Materials and Methods: Forty-five specimens of oral squamous cell carcinoma were randomly selected. Bcl-2 expression was evaluated by immunohistochemistry in formalin fixed, paraffin embedded pretreated specimens at the invasive front of oral squamous cell carcinoma. Clinicopathological data were gathered, and patient survival was analyzed.

Results: Immunohistochemical staining showed that nineteen of the forty-five specimens (42.2%) examined were positive for Bcl-2. Positive cells tended to be in the outer layer of round tumor nests and cord-like microtumor nests at the invasive front of oral squamous cell carcinoma. None of the Bcl-2 expression correlated significantly with age, gender, primary sites, T category, and N category. However, there was a correlation between Bcl-2 expression and the degree of cell differentiation ($p<0.05$). Cases showing recurrence of oral squamous cell carcinoma demonstrated high rates of Bcl-2 expression, and there was a significant correlation between Bcl-2 expression and recurrence of oral squamous cell carcinoma ($p<0.05$). Furthermore, the five-year survival rate of Bcl-2 positive cases was significantly lower than that of Bcl-2 negative cases ($p<0.05$).

Conclusions: Cell differentiation is generally analogous to the normal cells and tissue. Cases showing a low degree of cell differentiation demonstrated the high rates of Bcl-2 expression. Cells deviating from normal might control the function of apoptosis. Therefore, Bcl-2 expression at the invasive front of oral squamous cell carcinomas is a significant indicator of prognosis.

1066

POSTER

Relevance of the immunoexpression of galectin-3 in tumor cell compartments and its impact on the metastasis potential of the well-differentiated thyroid carcinoma

E. Stabenow¹, A. Ab'Saber², E. Parra-Cuentas², L. de Matos³, E. Eher², M. Tavares¹, A. Ferraz¹. ¹Hospital das Clínicas – School of Medicine – University of Sao Paulo, Head and Neck Surgery, Sao Paulo, Brazil; ²Hospital das Clínicas – School of Medicine – University of Sao Paulo, Pathology, Sao Paulo, Brazil; ³School of Medicine – ABC's Foundation, Santo Andre, Brazil

Papillary and follicular carcinomas are primary epithelial malignant neoplasias of the thyroid gland, originating from follicular cells and are considered to be well-differentiated carcinomas (WDC). The diagnosis and therapy of choice are based on their classification, according to prognostic factors that are considered together with the results of biological-marker investigation. As applied to immunohistochemical investigations, galectin-3 has been amply studied and a wide range of functions that it carries out in the cellular compartments have been described. However, its role in WDC remains controversial. To investigate if galectin-3 can aid in the diagnosis and treatment choice of WDC, digital quantification of its immunoexpression in the different cell compartments was carried out in order to determine its difference in relation to following: local tissue free of tumors and the neoplastic tissue, tumors with local tissue invasion and metastasis and its relation to the cell proliferation index, apoptosis and angiogenesis. A retrospective study was carried out with 109 cases of WDC from 1995 to 1999 in "Hospital das Clínicas" of the University of São Paulo Medical School. A survey of clinical data and follow up of 73±32 months was carried out, and in addition, morphological evaluation and digital immunohistochemical analysis with galectin-3, Ki-67, caspase-3 and CD-34 antibodies were done. It was demonstrated that: 1) there was no galectin-3 expression in the non-neoplastic epithelial tissue and there was nucleolar, nuclear and cytoplasmatic expression of galectin-3 in the tumor cells, with positive correlation among them; 2) according to a multivariate analysis, there was greater nucleolar expression as an independent factor associated to the presence of lymphatic metastasis ($P=0.028$) and greater cytoplasmatic expression associated to the presence of lymphatic

invasion ($P=0.013$); greater cytoplasmatic expression was revealed with a univariate analysis in the female gender ($P=0.0025$); with the presence of extra-thyroid extension ($P=0.037$), perineural invasion ($P=0.012$) and multicentric tumors ($P=0.005$); 3) the greatest cytoplasmatic expression was associated to the greater Ki-67 expression and to the 3 compartments with the greater caspase-3 expression. In conclusion, it was shown the possible application of the galectin-3 as a tumor marker in well differentiated thyroid carcinoma and nucleolus expression, as an indicator of its metastasis potential.

1067

POSTER

Angiogenesis as an indicator of metastatic potential in papillary thyroid carcinoma

E. Stabenow¹, A. Ab'Saber², E. Parra-Cuentas², L. de Matos³, E. Eher², M. Tavares¹, V. Capelozzi², A. Ferraz¹. ¹Hospital das Clínicas – School of Medicine – University of Sao Paulo, Head and Neck Surgery, Sao Paulo, Brazil; ²Hospital das Clínicas – School of Medicine – University of Sao Paulo, Pathology, Sao Paulo, Brazil; ³School of Medicine – ABC's Foundation, Santo Andre, Brazil

Background: Angiogenesis is new blood vessel formation, a process that can lead to tumor development. Microvessel count has been correlated to metastasis in some neoplasias.

Purpose: To determine if measurement of microvessel density is useful in predicting metastasis to the cervical lymph node and prognosis in patients with papillary thyroid carcinoma.

Methods: A retrospective analysis was performed in 30 patients that had undergone total thyroidectomy. They were divided in 2 groups of 15 patients – with and without metastatic disease. Immunohistochemistry was used to detect expression of CD34 in archival paraffin-embedded papillary thyroid tumors, and microvessel density was calculated based on it. Association between microvessel density and the presence of metastasis, according to histological subtype, disease recurrence, and AMES prognostic index groups was determined through statistical analysis. **Results:** The median microvessel density for the patient group without metastasis (200.0 microvessels/mm²) was apparently, but not significantly, less than that observed among metastatic disease patients (254.4 microvessels/mm²) ($P=0.20$). When papillary carcinoma subtypes were analyzed, this difference became significant ($P=0.02$): the follicular variant exhibited a greater microvessel density than the other subtypes, independent of metastasis presence. There was an apparent, but not significant, tendency for a larger median microvessel density in the group of patients that presented recurrence (294.4 microvessels/mm² vs 249.6 microvessels/mm², $P=0.11$). There was no relationship between risk level and microvessel density: in the low- and high-risk groups, the median MVD was 304.0 microvessels/mm² and 229.6 microvessels/mm², respectively ($P=0.27$).

Conclusions: The results suggest that angiogenesis is more intense among metastatic tumors in the classic and the tall cell variants, indicating that microvessel count can be an indicator of the potential for metastasis in these subtypes of papillary thyroid carcinoma. Patients that developed recurrent disease had a tendency to exhibit higher angiogenesis; however, there was no association between microvessel density and the AMES prognostic index.

1068

POSTER

Human papillomavirus types in head and neck cancer: an analysis of patients with unresectable head and neck squamous cell carcinomas (HNSCC) treated with curative radiochemotherapy (RCT)

K. Lindel¹, E. de Villiers², A. Dietz³, F.X. Bosch⁴, B. Helmke⁵, K.J. Weber¹, J. Debus¹. ¹University of Heidelberg, Dept. of Radiation Oncology/Radiotherapy, Heidelberg, Germany; ²DKFZ, Division of Tumour Virus Characterisation, Heidelberg, Germany; ³University of Leipzig, Clinic of Rhino-otolaryngology and Head and Neck Surgery, Leipzig, Germany; ⁴University of Heidelberg, Clinic of Rhino-otolaryngology and Head and Neck Surgery, Heidelberg, Germany; ⁵University of Heidelberg, Institute of Pathology, Heidelberg, Germany

Introduction: Over the past 15 years, human papillomavirus (HPV), the necessary cause of cancer of the cervix, has also been etiologically linked with a subset of HNSCC. The true prevalence of HPV and its influence on clinical outcome remain uncertain. Highly sensitive detection techniques, such as nested polymerase chain reaction (PCR) identified HPV DNA in a substantial proportion (30%-50%) of nonmelanoma skin cancer. Cutaneous HPV types might play a role in head and neck cancer but clinical data are missing.

Purpose: This study was performed to evaluate retrospectively HPV prevalence, genotypes and influence on clinical outcome in a subset of

patients with unresectable HNSCC treated with RCT in a multicenter randomized German trial at the University Hospital of Heidelberg, Germany. **Methods:** Thirty-six paraffin embedded biopsies of patients treated with curative RCT (total RT-dose 69.9, 600 mg/m²/dy 5-FU; 70 mg/m² carboplatin days 1–5 and 29–33) were analysed with single-phase primers FAP 69/64 and CP 65/70 and a nested PCR of FAP and CP to detect the majority of cutaneous and mucosal HPV types. The experimental findings were correlated with overall survival (OS), progression-free survival (PFS), distant metastases-free survival (DMFS).

Results: Ten out of 36 tumors were positive for high risk cutaneous HPV-types (1 ct HPV 5, 3 HPV 8, 4 ct HPV 9, 1 HPV 20, 2 ct HPV 48, 2 ct HPV 50, 2 HPV 76, HPV 79). Seven biopsies showed multiple HPV-DNA. HPV-infection was independent of tumor site (oropharynx versus hypopharynx). Kaplan-Meier survival estimates showed no difference in OS, PFS and DMFS between patients with HPV positive compared to negative tumors.

Conclusion: The role of cutaneous HPV in oncogenetic processes is not yet clarified, but they might also play a role in HNSCC. In contrary to our previous data which demonstrated HPV 16 positive HNSCC have a better prognosis than HPV negative tumors, there seems to be no evidence of a positive influence on clinical outcome or preferred tumor localization for tumors positive for cutaneous HPV.

1069

POSTER

Loss of heterozygosity on chromosomes 2q21 and 19p13.2 in oxyphilic thyroid cancers

K. Stankov^{1,2}, A. Pastore², L. Toschi², J. McKay³, F. Lesueur⁴, J.L. Kraimps⁵, D. Bonneau⁶, M. Volante⁷, M. Papotti⁸, G. Romeo².

¹Clinical center Novi Sad, Biochemistry, Novi Sad, Serbia; ²University of Bologna Medical School, Medical Genetics, Bologna, Italy; ³IARC, Genome Analysis Group, Lyon, France; ⁴Centre National de Genotypage, Evry, France; ⁵Jean Bernard Hospital, Endocrine Surgery, Poitiers, France; ⁶Centre Hospitalier Universitaire d'Angers, Génétique Médicale, Angers, France; ⁷University of Turin, Biomedical Sciences and Oncology, Torino, Italy; ⁸San Luigi Hospital Orbassano, Torino, Italy

Hürthle thyroid tumors are characterized by frequent numerical chromosomal aberrations, including aneuploidy or polyploidy, losses and gains of some chromosomal regions and DNA fragmentation. In recent years great attention has been paid to the combined analysis of morphological and genetic features of oxyphilic tumors and to the elucidation of their pathogenesis. We analyzed for loss of heterozygosity (LOH) of the candidate regions for TCO (Thyroid Tumor with Cell Oxyphilia) and NMTC1 (Non-Medullary Thyroid Carcinoma 1), two loci already mapped on chromosomes 19p13.2 and 2q21 respectively. Matched normal and tumor DNA samples from 70 patients with sporadic oxyphilic thyroid tumors and 20 with sporadic follicular tumors were subjected to microsatellite analysis using 10 markers on 19p13.2 and 6 markers on 2q21. This approach led us to the observation of a more significant LOH in oxyphilic than in follicular tumors. Allelic loss in tumor samples was evenly distributed in both 19p13.2 and 2q21 regions, in accordance with the established linkage of TCO and NMTC1 for inherited tumors. In order to investigate the possible contribution of both susceptibility loci in oxyphilic tumors, the family which led to the original mapping of TCO locus was reanalyzed for the markers in the 2q21 region. This led to the exclusion of linkage with the NMTC1 locus and to the refutation of the digenic inheritance hypothesis at least in this family.

1070

POSTER

Cimetidine: a lethal weapon against malignant salivary gland tumor?

M. Fukuda¹, M. Ishikawa¹, N. Suka¹, Y. Horiuchi¹, K. Kusama², H. Sakashita¹. ¹Meikai University, Second Department of Oral and Maxillofacial Surger, Sakado, Saitama, Japan; ²Meikai University, Department of Oral Pathology, Sakado, Saitama, Japan

Background: It has been reported that cimetidine inhibits the growth of glandular tumor such as colorectal cancer, however, the effect of cimetidine against salivary gland tumor is still unknown. We demonstrated previously that human salivary gland tumor (HSG) cells spontaneously express neural cell adhesion molecule (NCAM) and that the proliferative activity of HSG cells could be controlled via homophilic binding mechanism. In the present study, we investigated whether cimetidine could prevent the growth of HSG cells.

Materials and methods: The HSG cell line, derived from a human submandibular salivary gland, was established by Shirasuna et al. and three human oral squamous cell carcinoma cell lines, HSC-2, 3 and Ca9-22 were respectively maintained in 25 cm² culture flasks at 37°C in a humidified 5% CO₂ incubator until required. Northern-blot, Western-blot analysis and MTT assay were performed. Morphological observation of