

POSTER SESSION

Translational research 1

401

Influence of TP53 gene status on treatment response in breast cancer cells

Poster

L. Fernández¹, S. Ghayad², P. Cohen², P. Hainaut¹, M. Olivier¹¹International Agency for Research on Cancer (CIRC/IARC), Lyon, France; ²INSERM U590, Lyon, France

Breast cancer is a heterogeneous disease for which reliable diagnostic and prognostic markers are still lacking. Currently, estrogen receptor (ER), progesterone receptor (PR) and HER2 statuses are used in the clinic to select patients for anti-hormone and anti-HER2 therapies respectively. However, a significant proportion of patients (up to 30% of patients treated with anti-hormone therapy) will not respond to treatment. It is thus essential to find markers that would predict treatment failures to improve patient management and survival. TP53 gene mutation status has been repeatedly found to be a factor of poor prognosis in breast cancer. It has also been related to poor response to chemotherapy. TP53 is a tumour suppressor gene that is inactivated by mutation in a large proportion of cancers. It encodes a nuclear protein involved in many cellular pathways controlling cell proliferation, cell survival and genomic integrity in response to various types of stress, including DNA-damage caused by most chemotherapeutic agents. While the role of p53 in the cytotoxic effects of chemotherapeutic agents is well characterized, its possible role in the anti-proliferative and apoptotic effects of anti-hormone treatments has not been specifically investigated. In a large series of breast cancers, we have previously shown that TP53 gene mutation status was a factor of poor prognosis independently of the classical clinical markers. In addition, we found an interaction between TP53 mutation status and progesterone receptor expression that suggested a possible influence of p53 status on the response to anti-hormone therapy (Olivier et al., Clin. Cancer Res., 2006). To investigate this hypothesis in a cellular model, we have selected two isogenic breast cancer cell-lines that only differ in their p53 status and we have analyzed possible cross-talks between p53 and ER pathways. We have observed that cells with inactivated p53 are more resistant to the cytotoxic effects of tamoxifen and resume proliferation under estrogen-deprived conditions. Moreover, estrogen-deprivation or ER silencing reduces basal and doxorubicin-induced p53 levels. These results show that cross-talks between p53 and ER pathways do exist and may play an important role in the responses to both anti-hormone and chemotherapy treatments.

402

Impact of TP53 and KRAS mutations on cisplatin-based adjuvant chemotherapy in non-small-cell lung cancer

Poster

X. Ma¹, E. Brambilla², J.C. Soria³, A. Dunant³, A.L. Vataire³, P. Hainaut¹, and the IALT-Bio study group¹Molecular Carcinogenesis and Biomarkers Group, International Agency for Research on Cancer (IARC), Lyon, France; ²Department of Pathology, Centre Hospitalier Universitaire Albert Michallon, France; ³Department of Medicine (J.-C.S.), Biostatistics and Epidemiology (A.D., A.-L.V.), Institute Gustave Roussy, France

Background: Cisplatin-based adjuvant chemotherapy improves survival among patients with completely resected non-small-cell lung cancer (NSCLC) (Ronald H. Blum, NEJM 2004), however there are few prognostic and predictive markers to select patients for adjuvant chemotherapy. Mutations of TP53 and KRAS are most frequent in NSCLC. Aim: The purpose of this study is to analyze the impact of TP53 and KRAS mutations in NSCLC patients treated with cisplatin-based adjuvant chemotherapy enrolled in the International Adjuvant Lung Cancer Trial (IALT). Methods: Genomic DNA was extracted from 783 paraffin-embedded sections as described previously. Mutations in exon5 to 8 of TP53 were determined by sequencing and confirmed by another separate sequencing. KRAS mutation was detected by sequencing and confirmed by restriction fragment length polymorphisms analysis. The prognostic and predictive values of TP53 and KRAS status on survival were studied using a Cox model. Results: TP53 Mutation was found in 46% (240/524) of patients. TP53 status had no prognostic or predictive significance in all cancer grouped together or in adenocarcinoma patients. But in non-adenocarcinoma patients, there was a borderline significant interaction between TP53 status and treatment effect on disease-free survival

(p=0.05). The effect of chemotherapy was different in TP53 mutated and wild type patients, with a trend for benefit in TP53 wild type patients and a trend for harm in TP53 mutated patients. The prevalence of KRAS mutation was 14% (98/716). The prognostic effect of KRAS on disease free survival was different among 3 histology groups (p=0.03), adenocarcinoma, squamous-cell carcinoma and non-adenocarcinoma/ non-squamous-cell carcinoma, with worst prognostic effect of KRAS mutation vs. wild type among non-adenocarcinoma/ non-squamous-cell carcinoma. Mutation of KRAS was not predictive of the effect of chemotherapy. Conclusion: Patients with non-adenocarcinoma and TP53 wild type appear to benefit from cisplatin-based adjuvant chemotherapy, whereas chemotherapy seems to be harmful in patients with mutated TP53. KRAS mutation is a bad-prognosis factor in non-adenocarcinoma/non-squamous-cell carcinoma patients. Thus mutation detection may help in assigning patients to different treatment protocol. Key words: TP53; KRAS; Cisplatin; Adjuvant Chemotherapy; Non-small-cell Lung Cancer (NSCLC).

403

Role of PI3K/Akt /m-TOR pathway in cell response to Cisplatin treatment in Non-Small-Cell-Lung-Cancer - correlation to p53 status

Poster

S. Arandelovic¹, M.P. Cros¹, P. Hainaut¹¹Molecular Carcinogenesis and Biomarkers Group, International Agency for Research on Cancer, Lyon, France

Combination chemotherapy regimens in NSCLC are usually based on cisplatin, although clinical outcome is poor mostly due to resistance to treatment. Functions that protect against genomic instability (p53 or DNA repair mechanisms) may be implicated in modulating the risk of acquiring resistance to cisplatin. Biochemical alterations such as activation of signal transduction pathways that promote cellular survival may also confer resistance. Clinical studies have shown that activation of PI3K / Akt / mTOR pathway occurs in the majority of pre-cancerous lesions in smokers, and may have predictive significance for non-small cell lung cancer patients. The aim of this study was to elucidate importance of PKB (protein kinase B)/ Akt signaling in NSCLC cells (A549, H1299 and H1975) in mediating resistance to cisplatin treatment, by blockade of signaling cascade either using kinase inhibitors or by down-regulating Akt by siRNA. Data obtained by Western blot using antibodies against Akt and phospho Ser-473 Akt (Cell Signaling Tech.) revealed that wortmannin (PI3K- inhibitor) alone (2 M) and in combination with various doses of cisplatin, inhibits Akt phosphorylation, as observed after 4 h of action. Phosphorylation status of Akt was slightly decreased after 16 h treatment with cisplatin and kinase inhibitors wortmannin, SH-5 (Akt-inhibitor) (10 M). Also, protein level of p53 as well as phosphorylation of p53-Ser-15 were significantly reduced. Dual staining by PI / BrdU (bromodeoxyuridine) and FACS analysis showed that both SH-5 and wortmannin affected cell cycle progression but did not significantly increase the proportion of cell death in comparison to cisplatin alone. Cell survival studies using SRB assay indicated that wortmannin and SH-5 did not sensitize cell to cisplatin treatment. On the contrary rapamycin (inhibitor of m-TOR) induced synergistic effect in combination with cisplatin on p53 wild-type / p53 null / p53 mutant - cell models, when used at sub-toxic concentration (10 nM). Immunocytochemical staining of p53 and confocal microscopy revealed that rapamycin when combined with cisplatin increases nuclear localization of p53, while Akt inhibitor SH-5 antagonized cisplatin action. In conclusion, m-TOR may be a promising target for sensitization of cells to cisplatin treatment independently of functional p53 status in NSCLC cells. Key words NSCLC, cisplatin, p53, PI3K-Akt-m-TOR pathway, rapamycin.

404

High resolution melting allows sensitive high-throughput assessment of methylation in tumour samples

Poster

T.K. Wojdacz¹, E. Algar², A. Dobrovic³, L.L. Hansen¹¹University of Aarhus, Institute of Human Genetics, Aarhus, Denmark; ²Royal Children's Hospital, Children's Cancer Centre Murdoch Children's Research Institute, Melbourne, Australia; ³Peter MacCallum Cancer Centre, Molecular Pathology Research Laboratory, Melbourne, Australia

Despite the overwhelming interest in methylation analysis, readily performed methodologies that are both sensitive and specific are not generally available. Methylated DNA and unmethylated DNA acquire different sequences after bisulphite treatment resulting in PCR products with markedly different melting temperatures. We thus aimed to develop the emergent methodology of high resolution melting analysis (HRM) as a sensitive and specific method for the detection of methylation. We used our new strategy of primer design, which corrects for PCR bias and allows us to use annealing temperature to ensure methylation independent amplification [1]. We then developed a panel of locus specific MS-HRM (Methylation Sensitive – High Resolution Melting) assays. Amplification and melting analysis were done in the same tube using a real time PCR