

Herceptest 3+ or FISH positive) on survival in this otherwise very good prognostic group. The group were 89% ER positive, with 72% smaller than 20mm. 80% were aged over 50 and 10% received chemotherapy and 91% received endocrine therapy (tamoxifen).

Results: The overall hazard ratio (HR) for HER2 positivity was 6.78 (95% CI 2.9-15.7,  $p < 0.001$ ) with 5yr breast cancer specific survival rates of 96% (HER2 negative) and 68% (HER2 positive). This reduction in survival in HER2 positive cases persisted when patients were split into subgroups by ER status, tumour size and age (table 1).

Conclusion: These results provide support for the use of adjuvant trastuzumab in this group of patients who are typically classified as very good prognosis, not routinely offered standard chemotherapy, and as such do not fit current prescribing guidelines for trastuzumab. A clinical trial to assess the benefit of adjuvant trastuzumab alone within this subgroup of HER2 patients would resolve this. These results are in keeping with those from HERA trial that suggested that patients with the best prognosis tumours (node negative and size 1-2cm) had benefit similar to the overall cohort.

The persistence of a reduction in survival in our ER positive subgroup despite endocrine therapy confirms the recent trans-ATAC analysis based on HER2 status and suggests that we cannot not rely solely on adjuvant endocrine therapy in these largely ER positive patients.

#### 408 Structure-guided design of inhibitors of the eukaryotic initiation factor 4E (eIF4E) mRNA-cap interaction as anti-cancer agents

Poster

G. Patwardhan<sup>1</sup>, N.J. Oldham<sup>2</sup>, P.M. Fischer<sup>1</sup>

<sup>1</sup>University of Nottingham, Centre for Biomolecular Sciences and School of Pharmacy, Nottingham, United Kingdom; <sup>2</sup> University of Nottingham, Chemistry, Nottingham, United Kingdom

Protein synthesis in eukaryotes is regulated by eIF4E together with the other components of the translation initiation complex eIF4F. eIF4E anchors the mRNA by recognition of the 5' cap structure m7GpppN (where N is the first transcribed nucleotide and p represents phosphates), which contains a N-7-methylated guanine base. The role of eIF4E in cell proliferation and tumour progression is well documented, thus making eIF4E an attractive cancer drug target.

eIF4E recognizes the 5' cap structure through a characteristic cation- $\pi$  interaction involving the delocalized charge of the at N-7-modified guanine in the cap and two tryptophan residues in the eIF4E binding site, along with H-bonding interactions of the guanine base and electrostatic interactions with the phosphates. Our work aims to find non-nucleotidic cap-binding antagonists, and we look mainly for moieties in such inhibitors that could maintain the critical cation- $\pi$  interaction and would avoid the need for phosphate groups that render compounds membrane-impermeable and metabolically labile.

We follow a structure-guided drug design approach that consists of defining binding site constraints and performing in silico docking of small ligands into the eIF4E cap-binding site. We find that most of the scoring functions used to rank docked ligands fail to reward for the cation- $\pi$  interaction and we have implemented a quantum mechanical (QM) scoring strategy for the scoring of docked ligands. Hits from these virtual docking and scoring approaches were screened for binding eIF4E by Electrospray Ionization mass spectrometry (ESI-MS).

Several small molecules have been identified that bind to eIF4E and the results from these binding studies, as well as the effects of hit compounds on protein translation will be discussed.

#### 409 Gene expression profiling in formalin-fixed paraffin-embedded primary melanomas

Poster

C.A. Conway<sup>1</sup>, F. Elliott<sup>1</sup>, S. Lobo<sup>1</sup>, D.T. Bishop<sup>1</sup>, J. Newton-Bishop<sup>1</sup>

<sup>1</sup>Leeds Institute of Molecular Medicine, Section of Epidemiology and Biostatistics, Leeds, United Kingdom

Melanoma is an aggressive highly metastatic disease arising from epidermal melanocytes. Diagnosis and prognosis of this disease is currently limited to histological factors such as measurement of tumour invasion by Breslow Thickness and few successful treatments are available due to our poor understanding of the metastatic phenotype. Studies investigating the molecular basis of melanoma initiation and progression have been limited in the past due to the heterogeneous nature of melanoma and limited availability of fresh primary tumour. New techniques such as whole genome gene expression profiling are improving knowledge of many human cancers, however this is often a DNA intensive method resulting in most studies of this kind on melanomas involving either small numbers of fresh tissues or melanoma cell lines. We have used a novel method for gene expression profiling of 500 cancer genes in formalin-fixed paraffin-embedded primary melanomas. For this study, we selected 27

(FFPE) primary tumours from 27 patients; 15 of who had relapsed from their primary tumour within 5 years and 12 who had not relapsed after 5 years. The deepest, most invasive part of the tumours was sampled with a core biopsy needle and total RNA was isolated for analysis with the cDNA-mediated annealing selection, extension and ligation (DASL) assay (Illumina®). This assay is designed for use on partially degraded RNA for measurement of relative gene expression levels of up to 1536 sequence targets using as little as 25ng of total RNA. We used a Cancer Panel to target 502 genes commonly altered in cancer with three probes per gene. The results were visualised with Beadstudio analysis software (Illumina®) and normalisation was carried out using cubic spline methods, prior to export of results into STATA for further analysis. All tumours produced satisfactory results for the 502 genes in the Cancer Panel after quality control tests for average fragment length and amplification ability by qPCR. Genes found to have significantly different levels of expression between the group who had relapsed and the group that had not relapsed were SKI, PAI-1, BRCA2, WT1, MLLT4, NFKBIA and FGF8 (t-test and wilcoxon p

#### 410 Diagnosis of thyroid cancer by gene expression profiling on thyroid nodule biopsy

Poster

S. Durand<sup>1</sup>, C. Ferraro-Peyret<sup>2</sup>, S. Selmi-Ruby<sup>1</sup>, C. Paulin<sup>2</sup>,

F. Borson-Chazot<sup>1</sup>, B.A. Rousset<sup>3</sup>

<sup>1</sup>INSERM UMR 664, Faculté de Médecine Laennec, Lyon, France; <sup>2</sup> Lyon

Thyroid Tumor Bank Organization, Hôpital Edouard-Herriot, Lyon, France;

<sup>3</sup> INSERM UMR 664 and Lyon Thyroid Tumor Bank Organization, Faculté de Médecine Laennec and Hôpital Edouard-Herriot, Lyon, France

The diagnosis of thyroid cancer relies on cytological examination of material collected from nodules by fine-needle aspiration biopsies (FNAB). Due to the absence of markers, it is difficult, even for experienced cytologists, to discriminate benign from malignant thyroid tumors. Thus, only 35% of patients undergoing thyroidectomy for cancer or suspicion of cancer, actually have a cancer. The diagnostic procedure must be improved to reduce the number of thyroid ablation subsequently proved to be unjustified (about 10,000 per year in France). With the aim of identifying marker genes capable to discriminate benign from malignant thyroid tumors, we designed an oligonucleotide-based nylon macroarray formed from 200-potentially informative genes. Gene expression profiles of normal and tumoral (adenomas, carcinomas) thyroid tissue were generated with the macroarray and validated by real-time PCR. In this study, we built tumor classifiers from macroarray data and we tested their performances on a series of samples corresponding to FNAB. Gene expression data deriving from samples of the Lyon Thyroid Tumor Bank, representing the "training set", were subjected to a weighted voting algorithm to generate prediction models or classifiers capable of assigning a sample to one of the two classes: benign or malignant. Three prediction models were built by considering either all thyroid carcinomas (the commun classifier) or only follicular carcinomas (the F classifier) or only papillary thyroid carcinomas (the P classifier). The classifiers were composed of 9 to 12 genes and brought into play a total of 19 "marker" genes which were used to compose a fourth predictor, the global classifier. The capacity of the 4 classifiers to discriminate benign from malignant tumors was tested on a series of FNAB (carried out on nodules after surgical resection) used as "validation set". In 23 out of 26 FNAB, the 4 classifiers gave a diagnosis similar to that of the pathologist used as "gold standard"; in the 3 other cases, the correct diagnosis was given by 3 of 4 classifiers. Thus, the combination of classifiers identified benign and malignant tumors with very high sensitivity and specificity. In conclusion, we developed a procedure of molecular diagnosis of benign versus malignant tumors applicable to the material collected by FNAB. This molecular test which complied with a pre-clinical validation stage is now subjected to a prospective, large-scale (800 patients) evaluation study.

#### 411 Identification of drug-sensitive prediction genes by an epigenetic reactivation screen of cisplatin-resistant NSCLC cell lines

Poster

I. Ibanez de Caceres<sup>1</sup>, C. Moratilla<sup>1</sup>, M. Cortes Sempere<sup>1</sup>,

R. Machado-Pinilla<sup>1</sup>, V. Rodriguez-Fanjul<sup>1</sup>, C. Manguan<sup>1</sup>,

J. de Castro Carpeño<sup>2</sup>, C. Belda-Iniesta<sup>2</sup>, P. Cejas<sup>2</sup>, R. Perona<sup>1</sup>

<sup>1</sup>Biomedical Research Institute, Translational Oncology CSIC/H. La Paz, Madrid, Spain; <sup>2</sup> Medical Oncology Division Hospital Universitario La Paz, Translational Oncology CSIC/H. La Paz, Madrid, Spain

Non-small cell lung cancer (NSCLC) shows resistance in tumors that are initially chemo-sensitive, which is a serious problem in cisplatin-based adjuvant chemotherapy. CDDP is the paradigm of cytotoxic drugs in NSCLC treatment, however, it also induces, de novo DNA hypermethylation in vivo. Histone deacetylation and aberrant promoter hypermethylation are common epigenetic mechanisms for the silencing of