

THE INCORPORATION OF 5-FLUOROURACIL INTO RNA OF NORMAL TISSUES AND AN ADENOCARCINOMA TRANSPLANTED INTO THE LIVER OF PROTEIN DEPRIVED RATS

U.Stenram, I.A.El Hag, P.-I.Christensson, C.Erichsen, B.Jakobsson and P.-E.Jönsson

Departments of Pathology, Lund, and Surgery, Lund and Helsingborg; University of Lund, Sweden

Liver metastases of adenocarcinoma are commonly treated with 5-fluorouracil (5-FUra). A main effect is due to its incorporation into RNA. At protein deprivation, common in human cancer, the incorporation of cytidine into RNA is increased in the liver in rat. We have examined the effect of dietary protein on the incorporation of 5-FUra into RNA of tumour and normal tissues in rat.

Method: An N-methyl-nitrosoguanidine-induced adenocarcinoma of the colon of rat was transplanted to the liver. The rats were given either a 25% or 0% casein diet and some of the latter amino acids parenterally for one week. The rats were given 600 nmol [³H]-5-FUra/100 g body weight in a 2 hr infusion by the hepatic artery and killed after one further hr. Amounts of labelling in the acid-soluble fraction, RNA and DNA were determined in tumour and several normal tissues.

Results: Protein deprivation increased specific RNA labelling significantly in liver, kidney and ileum. Parenteral amino acids largely eliminated this increase and increased the RNA/DNA ratio in the liver. Protein deprivation also increased the specific RNA labelling in tumour, but the increase was not statistically significant. There was no change in bone marrow.

Conclusion: Protein deprivation alters the incorporation of 5-FUra into RNA in several tissues. Trials are in progress to decrease the incorporation of 5-FUra into normal tissues.

A TUMOUR CELL-ASSOCIATED PLASMIN-INDEPENDENT PATHWAY FOR UROKINASE PROENZYME ACTIVATION

R.W.Stephens, J.J.Hargreaves, P.-S.Sim, K.-C.Leung and L.Skriver(1)

The John Curtin School of Medical Research, Canberra, Australia, and (1)The Finsen Institute, Copenhagen, Denmark

It has been shown by several authors that human tumour cells normally produce and secrete the single-chain proenzyme form of urokinase plasminogen activator, which can

be activated by plasmin to the active two-chain form. For example, using inhibitory monoclonal antibodies to plasmin we have shown that such a pathway operates in cultures of human colon tumour cells.

However we now report that a human macrophage-like cell line (RC2A) consistently secretes the DFP-sensitive two-chain form of urokinase, which is formed by a cell-associated non-plasmin pathway. The active urokinase product is also recognised by the fast-acting PAI-2 class of specific inhibitor.

ANTI-IDIOTYPIC IMMUNITY AND TUMOUR DORMANCY IN A MOUSE B-CELL LYMPHOMA

F.K.Stevenson and A.J.T.George

Lymphoma Research Unit, Tenovus Research Laboratory, General Hospital, Southampton, U.K.

Immunization of mice with purified idiotype IgM rescued from the BCL1 lymphoma specifically protects against tumour challenge with approximately 30% of mice surviving for more than 6 months. Spleens from long-term survivors with no visible tumour were examined for tumour cells using monoclonal anti-idiotypic antibody. Dormant tumour cells were detectable which on passage into naive mice gave rise to tumour in the usual time span.

In long-term survivors which eventually showed emergent tumour, the pattern of surface idiotype expression was variable, ranging from normal to completely negative, but cytoplasmic idiotype was always present. This alteration was partly due to immunoselection of a variant tumour which fails to express idiotype but has intracellular idiotype.

Host mechanisms involved in tumour suppression appear to include anti-idiotypic antibody and reactive T cells which respond *in vitro* to idiotype IgM.

SCANNING CYTOPHOTOMETRIC DETERMINATION OF DNA IN P3X63-Ag8 PLASMOCYTOMA CELLS

H.Storch(1), H.Krug(2), S.Janz(1) and Th.Kohl(1)

(1)Department of Clinical Immunology and (2)Institute of Pathology, Leipzig, G.D.R.

The IgG1 producing murine plasmacytoma cell line P3X63-AgB established from MOPC 21 has been used for fusions in hybridoma technology. Intraperitoneally transplanted cells (not fused) were investigated on smears by scanning cytophotometric measurement after Feulgen staining. Only