

supporting the notion, that lymphocytes could be used as a model for primary target tissue, e.g. lungs. However, we have not seen any significant differences in lymphocyte AHH activity and inducibility between lung cancer patients and controls.

We have also studied whether MO activities predict the *in vivo* or *in vitro* formation of benzo(a)pyrene diol-epoxide (BPDE)-DNA adducts, the model lesion for PAH-induced carcinogenesis. We have shown that the nature of P450 isozyme is of importance for the activation ability, whereas the magnitude of activity seems to be of lesser importance. This was demonstrated with the aid of monoclonal antibodies (MAb) to different P450 isozymes. For example, the MAb to rat MC-induced liver P-450 inhibited AHH in placenta, but not in liver. It readily inhibited the *in vitro* formation of BPDE-DNA adducts in placenta, but not in liver. The MAb to phenobarbital-induced isozyme did not have these effects.

THE EFFECT OF INHIBITION OF MITOCHONDRIAL PROTEIN SYNTHESIS ON THE GROWTH KINETICS OF A RAT LEUKAEMIA

Ooby van den Bogert, Bert H.J. Dantje, Trudi E. Melis and Albert M. Kroon

Laboratory of Physiological Chemistry, State University, Medical School, Bloemsingel 10, 9712 KZ Groningen, The Netherlands

Mitochondria (mt) contain DNA coding for several subunits of components of the oxidative phosphorylation system and a specific system to transcribe and translate this DNA. Inhibition of the expression of mt-genes finally diminishes the capacity for oxidative phosphorylation to an extent that cell functions (e.g. proliferation) become impaired. This has been demonstrated already in several tumour systems. Specific and continuous impairment of mt-protein synthesis by treatment with tetracyclines results also in growth inhibition of a leukaemia in the rat - it even leads to its disappearance. Cytostasis is achieved more rapidly and the rate of cytolysis is faster when tetracycline treatment is started in later stages of tumour progression. Our studies indicate that this is due to interference of tetracyclines with the cytostatic and cytolytic effects of corticosteroids on the growth of this tumour. As tetracycline treatment has lasted longer, the anti-tumour effects of (endogenous) corticosteroids become less. It is suggested that mt-protein synthesis is required for the action of corticosteroids on leukaemic cells.

PROVIRUS INTEGRATION IN [90]Sr -INDUCED OSTEOSARCOMAS OF C57BL MICE

E. Van der Raaij-van der Wal(1), J.R. Maisin(2) and J. Merregaert(1)

(1) Department of Biochemistry, Section of Biotechnology, UIA, Universiteitsplein 1, 2610 Antwerpen, Belgium; and (2) Department of Biology, SCK, Boeretang 200, 2400 Mol, Belgium

The development of [224]Ra and [90]Sr induced osteosarcomas in mice is accompanied by the expression of endogenous retroviruses in bone tissues of the treated animals in the early latency period, and later in the osteosarcomas. Using the Southern blotting procedure, we have studied the presence of somatically acquired proviruses in genomic DNA isolated from seven primary [90]Sr induced osteosarcomas and one osteosarcoma cell line (O-127a1) of the C57BL mouse strain. Newly integrated ecotropic proviruses were detected with specific hybridization probes in four primary tumours. In contrast, genomic DNA from cultured osteosarcoma cells harboured additional ecotropic recombinant (MCF-related) proviruses. No integrations were found in the vicinity (22 kbp) of c-myc. The c-myc locus is amplified in two out of eight tumour DNAs. According to our data, detectable integrations of activated retroviruses do not appear to be an essential requisite for the development of radiogenic osteosarcomas in mice, but in some cases, clonal or oligoclonal integrations might have been responsible for the deregulation of a nearby putative oncogene, allowing cells to escape normal growth control *in vivo*.

REVERTANTS OF METHIONINE-DEPENDENT H-ras-1 ONCOGENE-TRANSFORMED CELLS

Luc Vanhamme and Claude Szpirer

Département de Biologie Moléculaire, Université Libre de Bruxelles, Rue des Chevaux, 67, B1640 Rhode-St-Genèse, Belgium

Methionine-dependence is a metabolic defect reported to be exhibited by many transformed and malignant human or animal cells (Mechan *et al.*, *EBRC*, 117: 429, 1983). This defect is characterized by the inability of cells in culture to grow in a medium where methionine has been replaced by its immediate metabolic precursor, homocysteine. The biochemical basis of this phenomenon is not understood. We have shown recently that the activated H-ras-1 oncogene, derived from the EJ human carcinoma line, induces methionine-requirement after transfection in