

Trends in Cancer Mortality in the Americas, 1955–1989

Carlo La Vecchia, Franca Lucchini, Eva Negri, Peter Boyle and Fabio Levi

INTRODUCTION

DATA AND statistics on cancer mortality over the last few decades are available from the United States of America [1] and Canada [2, 3], but are more difficult to find, in a standard and comparable format, for Central and South American countries [4, 5].

The World Health Organization (WHO) mortality database includes death certification data from the early 1950s onwards for several American countries. We decided, therefore, to include in a unique publication the pattern of trends in cancer mortality between 1955 and 1989 for 14 Central and South American countries, together with—for comparative purposes—the United States of America and Canada. The tabular and graphical form of presentation are similar to those previously adopted for a companion report on cancer mortality in Europe [6–10].

The present publication has two major objectives: to provide basic documentation for epidemiology, health statistics and public health, and to offer a baseline framework for comparative inference. This may also have relevant implications for analytical epidemiology and cancer control, in terms of indicating priorities for further investigation and intervention on a public health scale.

MATERIALS AND METHODS

Although the WHO database contains some data on mortality or population for 49 American countries or territories, we excluded from consideration all countries with less than 2 000 000 population in order to avoid problems with low numbers of cases inducing instability in the rates. Out of the 24 remaining countries, only those with age-stratified mortality and population figures of sufficient detail were retained: eight countries (Bolivia, Brazil, El Salvador, Guatemala, Haiti, Honduras, Jamaica, Nicaragua) were excluded at this stage. The analysis is based on the remaining 14 Central and South American countries: Argentina, Chile, Colombia, Costa Rica, Cuba, Dominican Republic, Ecuador, Mexico, Panama, Paraguay (reporting area only), Peru, Puerto Rico (a self-governing Commonwealth, though not an independent country), Uruguay and Venezuela, plus the U.S.A. and Canada. Unfortunately, data for different ethnic groups (hispanic, white, black, etc.) was not available for the U.S.A.

During the calendar period considered (1955–1989), four different revisions of the International Classification of Diseases (ICD) were used [11–14]. Classification of cancer deaths was thus recorded for all calendar periods and countries according

to the Ninth revision of the ICD [ICD-9]. Table 1 gives the cancers or groups of cancers considered (12 sites, plus total cancer mortality), together with the corresponding ICD codes under subsequent revisions. To improve comparability of data between different countries and calendar periods, we pooled together all intestinal sites including rectum [15], all uterine cancers (cervix and endometrium) [16], and all skin neoplasms (melanomas and non-melanomatous). For most countries, no meaningful analysis was possible for rarer cancers (e.g. gallbladder, sarcomas, thyroid or lymphomas), for those sites (e.g. liver, pancreas, ovary or brain) which implicate major difficulties and uncertainties in diagnosis and certification, and also for bladder and kidney cancer.

Estimates of the resident population, generally based on official censuses, were obtained from the same WHO databank. From the matrices of certified deaths and resident populations, age-specific rates for each 5-year age group and calendar period were computed. Age-standardised rates, at all ages and truncated to 35–64 years, were based on the world standard population [17].

In a few countries, data were missing for part of one or more calendar periods (Table 2). When a single year was missing within a quinquennium, numerators and denominators were interpolated linearly for the previous and subsequent calendar year. No extrapolation was made for missing data at the beginning or the end of the calendar period considered, or when data on one or more quinquennia were not available.

PRESENTATION OF RESULTS

For each cancer site and country one table and three figures are presented.

The table includes, for each calendar year and sex, the absolute number of deaths and the overall age-standardised and truncated 35–64 years mortality rates.

The first figure includes world-standardised rates (all ages and truncated) for each 5-year calendar period in both sexes.

Age-specific rates from 30–34 to 70–74 years were plotted against the central year of birth cohort in the second figure for males and in the third for females. The points corresponding to the same age group were joined to provide more clearly spaced graphs. Thus, in this graph the cohort effect can be read in the ordinate.

Finally, to provide graphical representation of the range of variation in mortality for each cancer site, Fig. 1. gives histograms of overall age-standardised death rates in the most recent calendar quinquennium (1985–1989, whenever available).

GENERAL COMMENT

It is useful to remember that this work has essentially a descriptive interest, and comments and interpretations are consequently limited. A few general methodological comments are nonetheless included, to assist in critical reading and interpretation of patterns of rates and trends.

Correspondence to C. La Vecchia.

C. La Vecchia, F. Lucchini and F. Levi are at the Institut universitaire de médecine sociale et préventive, Bugnon 17, 1005 Lausanne, Switzerland (address for correspondence); C. La Vecchia and E. Negri are at the Istituto di Ricerche Farmacologiche "Mario Negri", Via Eritrea 62, 20157 Milano, Italy; P. Boyle is at the Division of Epidemiology and Biostatistics, European Institute of Oncology, via Ripamonti 332/10, 20141 Milan, Italy; and F. Levi is at the Registre vaudois des tumeurs, Institut universitaire de médecine sociale et préventive, CHUV–Falaises 1, 1011 Lausanne, Switzerland.

Received and accepted 7 July 1992.




Fig. 1. Histograms of geographic variation in mortality per 100 000 from major cancer sites in the Americas, in the most recent calendar period for each country.

First of all, although age-standardised rates are a useful summary statistic, accurate inspection of subsequent age-specific rates is essential before formulating any inference on trends in cancer mortality. Trends at younger ages are of utmost importance, not only because death certification and population estimates are more reliable, but also because they may contain useful indications as to the most likely future trends [18].

Due caution, moreover, should be paid to upward trends only affecting older age groups while rates in the younger middle age are stable (e.g. for cancer of the prostate, bone or leukaemias, where large diagnostic improvements have been achieved over the last two or three decades). Most of these apparent trends should in general be considered largely or totally artefactual [18].

Further, problems related to variability should always be considered, and are clearly greater in relation to smaller populations and rarer cancers.

Finally, and more complex, there are problems of death certification accuracy and reliability in various countries and calendar periods. In general, for several common cancer sites including lung, stomach, colorectum and breast, death certification data can be considered sufficiently reliable and comparable to permit meaningful inference in most American countries, particularly in relation to trends within countries and under age

65. Greater caution is required for a few countries and cancer sites whose diagnosis and certification may be largely influenced by the availability and utilisation of diagnostic techniques and accuracy of registration [4, 18, 19]. These cancers include, among others, liver, bone, prostate, brain, myeloma and other lympho-haemopoietic neoplasms. Among these, only for bone, prostate and leukaemias were there sufficient (or acceptably reliable) data to justify inclusion in the present report.

On the basis of these considerations, besides a non-uniform availability of data in various countries, trends in mortality are not given for a few rare or particularly difficult to diagnose and certify cancers including liver, gallbladder, pancreas, ovary, bladder, kidney, brain, thyroid, lymphoma and myeloma.

MOUTH OR PHARYNX

To improve comparability of data within and between countries, all oral and pharyngeal sites were grouped together.

In the 1960s, the highest rates in males were in Puerto Rico and Cuba (around 10/100 000, world standard), followed by Argentina, Paraguay and Uruguay (around 5/100 000). Overall rates of around 5/100 000 males which were stable over time were also observed in the U.S.A. and Canada. Registered mortality from oral and pharyngeal cancer was appreciably lower in most other Central and South American countries.




Fig. 1. Continued.

The pattern of trends was considerably heterogeneous in various countries. Substantial declines were observed in Cuba and Paraguay, but oral and pharyngeal cancer rates increased in Argentina and Uruguay, particularly in younger and middle-aged males, and increased in younger males in Canada. Thus, in the late 1980s Uruguay had the second highest rate of oral and pharyngeal cancer (6.9/100 000 males), after Puerto Rico (9.3/100 000). Among countries starting from lower values, appreciable rises were observed in the Dominican Republic and Mexico.

Oral and pharyngeal cancer rates in females were lower, and showed no appreciable trend in any country except Venezuela, where rates started from exceedingly high values of 4.1/100 000 in 1955–1959 but declined to 1.3/100 000 in the late 1980s.

Alcohol and tobacco are the main determinants of oral and pharyngeal cancer in the developed world, and are likely to account for a large proportion of the underlying trends in most American areas [20, 21]. A diet poor in vegetables and fresh fruit may also have some role [22, 23], as well as maté drinking in selected areas of Argentina, Brazil, Uruguay and Paraguay [24].

OESOPHAGUS

There are substantial differences in male oesophageal cancer rates between various American countries. High rates (around

or over 10/100 000, all ages) in the 1950s or 1960s were registered in Argentina, Chile, Paraguay, Uruguay and Puerto Rico, and in some of these countries, particularly Puerto Rico and Uruguay, rates were also elevated in females (around or over 5/100 000). In most of these countries, however, oesophageal cancer rates tended to decline somewhat over most recent calendar periods, particularly in middle and younger age. Still, in the late 1980s, oesophageal cancer mortality rates were over 10/100 000 males in Uruguay and Puerto Rico.

Low rates (i.e. between 1 and 3/100 000 males) were observed in several Central American countries, including Mexico, Dominican Republic, Ecuador, and also in Peru. No systematic trend over time was observed in the low mortality areas.

Over the same calendar period, oesophageal cancer rates in Canada and the U.S.A. were between 3 and 4.5/100 000 males, and tended to increase moderately but steadily, particularly at younger ages.

As for oral and pharyngeal cancers, alcohol and tobacco are the main recognised aetiological factors underlying these trends [25]. However, to explain the recent declines in several high mortality areas, the role of thermic irritation or carcinogenic substances in maté drinking [26–28] and possibly of improved diet [20, 29] on oesophageal carcinogenesis should also be considered.




Fig. 1. Continued.

STOMACH

Some Central and South American countries had exceedingly high gastric cancer rates [28]. Among males, for instance, Chile had an overall world standardised rate of 69.4/100 000 in 1950-1959, and of 34.4/100 000 in 1985-1989. Comparative figures in Costa Rica were 60.3/100 000 males in 1960-1964 and 48.6/100 000 males in 1985-1989. For comparative purpose, in the U.S.A. the range of variation over the same calendar period was between 13.9 and 5.3/100 000 males.

Other high mortality countries were Colombia, Ecuador and, to a lesser degree, Uruguay and Venezuela.

Rates for females were lower, i.e. between 22.6/100 000 in Costa Rica, 20.8/100 000 in Colombia and 2.4/100 000 in the U.S.A in the late 1980s, but the pattern of geographical differences and trends were similar.

In most American countries, stomach cancer rates have been appreciably decreasing during the last four decades, as in most developed areas of the world. The extent of the decline, however, varied appreciably from country to country, and some of them, including Colombia, Dominican Republic and Ecuador showed rises in certified mortality from gastric cancer. Stomach cancer is easy to diagnose and, in general, to certify, but it is still

possible that some of these rises are due to improved diagnosis and certification.

Although the exact causes of the largely favourable trends in stomach cancer observed in most countries are still largely undefined, it is likely that improved techniques of food storage and conservation (such as refrigeration and declined use of salt) have played a major role, together with a more affluent diet and increased availability of fresh fruit and vegetables [30]. Improved food conservation may also have reduced the prevalence of *Helicobacter pylori* (and perhaps other micro-organisms) infection in the gastric mucosa, which has been linked to elevated stomach cancer risk [31].

Most of these lines of reasoning on the causes of the decline of gastric cancer probably apply to Central and South America too, but this cannot eclipse that on a public health level several areas of exceedingly high stomach cancer rates persist in this part of the world, thus underlining the scope and importance for future monitoring and intervention.

INTESTINES, CHIEFLY COLON AND RECTUM

Within South America, the highest intestinal cancer rates in the late 1980s were registered in Uruguay (17.9/100 000 males,




Fig. 1. Continued.

15.5/100 000 females) and Argentina (13.1/100 000 males, 9.4/100 000 females). Still, these rates were comparable to those observed in the U.S.A. and Canada, around 18/100 000 males and 12-13/100 000 females. All other Central and South American countries had mortality rates around or below 10/100 000, and intestinal cancer rates for Ecuador and Mexico were around or below 3/100 000.

Trends over time varied appreciably over the calendar period and age groups, but declines were more frequent than upward trends, particularly over the more recent calendar period.

The relatively high colorectal cancer rates in Uruguay and Argentina seem to be consistent with the putative role of red meat and animal fat in the aetiology of colorectal cancer suggested by analytical studies [32, 33]. In terms of potential implications for prevention, however, the extremely low rates in most Central and South American countries are of major interest. If not due to substantial underreporting, these low rates would indicate that the diet typical of these areas is particularly favourable with regard to intestinal carcinogenesis.

LARYNX

This is another site strongly related to the consumption of alcohol and tobacco, not only in the form of cigarettes but also

of pipe and cigar tobacco [20, 21]. Not surprisingly, therefore, the high mortality areas were in Uruguay (around 7/100 000 males), Cuba and Argentina (around 5/100 000 males).

Most other Central and South American countries showed age-standardised (world population) rates between 2 and 3/100 000 males, i.e. comparable or only slightly above those registered in the U.S.A. and Canada.

Trends over time were largely heterogeneous across various countries, in the absence of generalised and systematic patterns.

With the sole exception of Cuba (with rates around 1/100 000 females), laryngeal cancer was extremely rare in American women, and the consequent patterns were unremarkable.

LUNG

In the late 1960s, Argentina had an overall world standardised lung cancer rate of over 45/100 000 males, and Uruguay and Cuba of over 40/100 000 males, i.e. some of the highest lung cancer rates in that period on a global scale. Lung cancer mortality, however, has since been stable in Cuba and has declined (39.7/100 000 males in 1985-1988) in Argentina, and is now, therefore, substantially lower than the value of 56-57/100 000 males in the U.S.A. or Canada. During the 1970s, a decline in cigarette smoking among Argentinian males was




Fig. 1. Continued.

observed [28]. Among these high mortality countries upward trends were observed only in Uruguay, where values of 54.7/100 000 males were reached in 1985-1989.

The most important observation for lung mortality rates in Central and South America is related to the noticeably low rates still registered in most countries. Although upward trends were observed in several countries, in the late 1980s overall age-adjusted (world population) rates in males were between 15 and 22/100 000 males in Chile, Costa Rica, Panama, Puerto Rico, Venezuela and Mexico, and around or below 10/100 000 males in Colombia, Dominican Republic, Ecuador, Paraguay and Peru. Some of these rates are only moderately above those registered among females in Europe or lifelong non-smokers in the U.S.A. [34], and, if not due to macroscopic under-reporting, indicate that the tobacco-related lung cancer epidemic is still in its very early phases in these populations. It has been estimated that by 1985, over 520 000 tobacco-related deaths per year occurred in the Americas, of which about 100 000 were in Latin America and the Caribbean [35].

Female lung cancer rates were very low in all Central and South American countries, with the sole exception of Cuba (14.1/100 000 females in 1985-1989). Contrary to Canada and the U.S.A., however, female lung cancer rates have not been rising in Cuba.

BONE

Bone cancer mortality rates, and the corresponding trends, are difficult to interpret, since this is another common site of metastases [18].

Rates for several countries were around 1/100 000 males, all ages, and somewhat lower in females. There was, however, a diffuse and systematic tendency towards a decline in mortality over time, which can be taken as an indication of a decline in misclassification of secondaries, and hence of improved reliability of cancer death certification over time.

SKIN, INCLUDING MELANOMA

It was not possible, on the basis of death certification alone, to distinguish melanoma from non-melanomatous (chiefly squamous cell) skin cancers. Both types of skin cancer are linked to exposure to sunshine, but the pattern of exposure chiefly linked to risk is different (acute intermittent for melanoma; long-term and cumulative for squamous cell). Also the age distribution is different, since squamous cell carcinomas are common, or restricted, to older ages [36-38].

These differences in the epidemiology of various histotypes may explain at least part of the diverging trends observed in various American countries. For instance, the substantial upward trends in the U.S.A. and Canada, restricted or greater




Fig. 1. Continued.

Table 1. Cancers or groups of cancers considered

Type of cancer	6 ICD	7 ICD	8 ICD	9 ICD
Mouth or pharynx	140–148	140–148	140–149	140–149
Oesophagus	150	150	150	150
Stomach	151	151	151	151
Intestines, chiefly colon and rectum	152–154	152–154	152–154	152–154 + 159.0
Larynx	161	161	161	161
Trachea, bronchus and lung	162	(162 + 163)–162.2	162	162
Bone	196	196	170	170
Skin including melanoma	190 + 191	190+191	172 + 173	172 + 173
Breast	170	170	174	174(M)/175(F)
Uterus (cervix and corpus)	171–174	171–174	180–182	179–182
Prostate	177	177	185	185
Leukaemias	204	204	204–207	204–208
Total, all sites, all histologies	140–239	140–239	140–239	140–239

ICD: International Classification of Diseases.

M, Male; F, female.

at younger ages and hence reflected in substantial rises in the truncated rates, are largely or totally attributable to melanoma, and probably reflect the spreading of sunbathing habits in younger generations [38]. In the U.S.A., the truncated rates reached 6.1/100 000 males and 2.8/100 000 females in the late

1980s, while the corresponding all-age rates were 3.4 and 1.6/100 000 females, respectively.

In all Central and South American countries (with the major exception of Colombia), skin cancer rates were substantially lower (around and often below 1/100 000 in both sexes), and

Table 2. (a) Calendar years for which data were missing for each country and cancer site

Country	Mouth or pharynx	Cancer sites		
		Oesophagus	Stomach	Intestines
Argentina	55-65, 71-76, 87-89	55-65, 71-76, 87-89	55-65, 71-76, 87-89	55-65, 71-76, 87-89
Canada	89	89	89	89
Chile	88-89	88-89	88-89	88-89
Colombia	70-71, 73, 78-89	70-71, 73, 78-89	70-71, 73, 78-89	70-71, 73, 78-89
Costa Rica	55-60, 89	55-60, 89	55-60, 89	55-60, 89
Cuba	55-63, 65-67, 89	55-63, 65-67, 89	55-63, 65-67, 89	55-63, 65-67, 89
Dominican Republic	55-64, 86-89	55-64, 86-89	55-64, 86-89	55-64, 86-89
Ecuador	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89
Mexico	79, 80, 84, 87-89	79, 80, 84, 87-89	79, 80, 84, 87-89	79, 80, 84, 87-89
Panama	88-89	88-89	88-89	88-89
Paraguay	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89
Peru	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89
Puerto Rico	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89
U.S.A.	89	89	89	89
Uruguay	61-62, 79, 88-89	61-62, 79, 88-89	61-62, 79, 88-89	61-62, 79, 88-89
Venezuela	84, 88-89	84, 88-89	84, 88-89	84, 88-89

have been steadily declining in both sexes and most subsequent age groups over the last four decades. Besides different genetic characteristics of the populations, in terms of skin, eye and hair colour, this probably reflects reduced long-term exposure to sunshine, with the decline of the population employed in agriculture and other outdoor activities [36-38], and includes favourable trends both for melanoma and non-melanomatous skin cancers. There is, in any case, no indication that the epidemic of skin melanoma [38] has started in most central and southern American populations.

BREAST

In South America, elevated breast cancer mortality rates were observed in Argentina (over 20/100 000 women, all ages) and Uruguay (over 25/100 000). These were comparable with the rates of 23-24/100 000 females in the U.S.A. and Canada. In these high mortality areas, no appreciable trend in overall age-standardised rates was observed over the last three decades, although these overall rates include some generally more favourable trends in younger and middle age, and some less favourable trends (particularly in Argentina and Uruguay) at older ages.

Among other countries, in the late 1980s, Cuba had an overall breast cancer rate of 15.1/100 000 females. Puerto Rico of 13.2, Costa Rica of 13.1, and Chile of 12.8/100 000 females. All other Central and South American countries had breast cancer rates below 10/100 000 and some of these (i.e. Colombia, Dominican Republic, Ecuador or Peru) around or below 5/100 000. In all these low mortality areas, however, trends over time have been substantially upwards, at all ages.

These substantial variations in breast cancer rates between various American countries probably reflect different reproductive, dietary and general lifestyle habits in various geographical areas [39-40], although it is possible that problems of death certification reliability somewhat affect breast cancer rates in some countries. Nevertheless, the elevated breast cancer mortality in Argentina and Uruguay may deserve specific attention with reference to reproductive, but also dietary habits, as well as the extremely low rates in a few other countries. The stable rates in high mortality areas, and the systematic upward trends in low mortality areas indicate that most rates may tend to level towards high values in the future, possibly reflecting a general tendency towards more uniform lifestyle habits.

UTERUS (CERVIX AND CORPUS)

On the basis of death certification, it was not possible to distinguish cancers arising from cervix and from the corpus uteri (mainly endometrium) [16], although the aetiology of the two sites is largely different [41, 42]. Still, since endometrial cancer is extremely rare at younger ages, trends in younger and middle age are largely or totally due to cervical cancer.

In comparison with recent rates in the U.S.A. and Canada (around 5/100 000 women, all ages and between 8 and 9/100 000 females, 35-64 years in the late 1980s), mortality from cancers of the uterus was extremely high in all Central and South American countries, with the sole exception of Puerto Rico (6/100 000 all ages, 11/100 000 truncated). It has been estimated that cervical cancer incidence is about six times higher in Latin America than in the U.S.A. [28]. The highest rates were

Table 2. (b) Calendar years for which data were missing for each country and cancer site

Country	Cancer sites			
	Larynx	Lung	Bone	Skin including melanoma
Argentina	55-65, 71-76, 87-89	55-65, 71-76, 87-89	55-68, 71-76, 87-89	55-65, 71-76, 87-89
Canada	89	89	56-68, 89	89
Chile	88-89	88-89	55-67, 88-89	88-89
Colombia	70-71, 73, 78-89	70-71, 73, 78-89	55-67, 70-71, 73, 78-89	70-71, 73, 78-89
Costa Rica	55-60, 89	55-60, 89	55-67, 89	55-60, 89
Cuba	55-63, 65-67, 89	55-63, 65-67, 89	55-67, 89	55-63, 65-67, 89
Dominican Republic	55-64, 86-89	55-64, 86-89	55-67, 86-89	55-64, 86-89
Ecuador	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89	55-67, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89
Mexico	79-80, 84, 87-89	79-80, 84, 87-89	55-67, 79-80, 84, 87-89	79-80, 84, 87-89
Panama	88-89	88-89	55-67, 88-89	88-89
Paraguay	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89	55-67, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89
Peru	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89	55-67, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89
Puerto Rico	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89	55-68, 78, 88-89	55-60, 67-68, 78, 88-89
U.S.A.	89	89	55-67, 89	89
Uruguay	61-62, 79, 88-89	61-62, 79, 88-89	55-67, 79, 88-89	61-62, 79, 88-89
Venezuela	84, 88-89	84, 88-89	55-67, 84, 88-89	67, 84, 88-89

in Paraguay (22/100 000 females all ages, 44/100 000 females truncated), but truncated rates above 30/100 000 females were also observed in the late 1980s in Chile, Ecuador, Mexico and Venezuela, and between 25 and 30/100 000 females in Costa Rica, Cuba, Dominican Republic and Peru. However, in most countries appreciable downward trends were observed, particularly in most recent calendar periods and younger generations. The two exceptions are Argentina, whose rates remained stable around 12/100 000 females, all ages and 24/100 000 females, truncated (and which showed upward trends in the young), and Colombia, with stable rates around 18/100 000 females, all ages and 35/100 000 females, truncated.

The major factor underlying this substantial excess of (cervix) uteri cancer mortality in Central and South America is probably venereal transmission of (viral) aetiological factor(s), due to the sexual habits of these populations and particularly male promiscuity [43-45]. Inadequate utilization of cervical smear testing certainly also has a major role [28, 46, 47].

PROSTATE

In the late 1980s, prostatic cancer rates in Canada and the U.S.A. were around 15/100 000 males (all ages) and 7/100 000 males (truncated, 35-64 years). Some upward trend over time was observed at older ages, but U.S.A. rates were totally stable below age 65. Although prostatic cancer is a disease of elderly males, death certification in these age groups can be largely influenced by recent changes and improvements in diagnostic attention and accuracy, since neoplastic foci in the prostate are

found in a large proportion of elderly men [48]. Thus, inference in trends should be essentially based on rates in middle age, although these account only for a small proportion of cases [18].

In Costa Rica, Cuba, Puerto Rico and Uruguay recent rates were similar or higher than those of the U.S.A. and Canada, and again there was some indication of upward trends at older ages. Prostatic cancer rates above 5/100 000 males truncated, or above 10/100 000 males all ages, were observed in Argentina, Chile, Panama and Venezuela, while rates were lower in other countries. Some of these low mortality areas, such as the Dominican Republic or Mexico, showed larger and more diffuse upward trends, although it is difficult to quantify how much they were affected by improved diagnosis and certification.

In comparison with most other common cancers, including not only lung and other tobacco-related neoplasms, but also stomach, colorectum and breast, the variation in prostatic cancer mortality among various American countries is relatively limited. Similarly limited is any inference on the reasons of this pattern of trends, since little is known about the causes of prostatic cancer [49].

LEUKAEMIAS

In the late 1980s, the highest overall age-standardised (world population) leukaemia rates in Latin America were around 6/100 000 males and 4/100 000 females in Costa Rica and Uruguay, followed by Argentina and Cuba. Most other Central and South American countries had leukaemia mortality rates between 3 and 4/100 000 in both sexes, although somewhat lower

Table 2. (c) Calendar years for which data were missing for each country and cancer site

Country	Cancer sites				Total, all neoplasms
	Breast	Uterus	Prostate	Leukaemias	
Argentina	55-65, 71-76, 87-89	55-65, 71-76, 87-89	55-65, 71-76, 87-89	55-65, 71-76, 87-89	55-65, 71-76, 87-89
Canada	89	89	89	89	89
Chile	88-89	88-89	88-89	88-89	88-89
Colombia	70-71, 73, 78-89	70-71, 73, 78-89	70-71, 73, 78-89	70-71, 73, 78-89	70-71, 73, 78-89
Costa Rica	55-60, 89	55-60, 89	55-60, 89	55-60, 89	55-60, 89
Cuba	55-63, 65-67, 89	55-63, 65-67, 89	55-63, 65-67, 89	55-63, 65-67, 89	55-63, 65-67, 89
Dominican Republic	55-64, 86-89	55-64, 86-89	55-64, 86-89	55-65, 86-89	55-64, 86-89
Ecuador	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89	55-60, 62, 76, 81, 83, 89
Mexico	79-80, 84, 87-89	79-80, 84, 87-89	79-80, 84, 87-89	79-80, 84, 87-89	79-80, 84, 87-89
Panama	88-89	88-89	88-89	88-89	88-89
Paraguay	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89	55-60, 64, 69, 80-81, 87-89
Peru	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89	55-65, 74-76, 79, 84-89
Puerto Rico	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89	55-60, 67-68, 78, 88-89
U.S.A.	89	89	89	89	89
Uruguay	61-62, 79, 88-89	61-62, 79, 88-89	61-62, 79, 88-89	61-62, 79, 88-89	61-62, 79, 88-89
Venezuela	84, 88-89	84, 88-89	84, 88-89	84, 87, 88-89	84, 88-89

in females. Thus, particularly for males, leukaemia mortality was lower in Latin America than in the U.S.A. and Canada, where rates in the late 1980s were over 6/100 000 males and around 4/100 000 females.

In North America, after initial rises and a peak of over 9/100 000 males and 6/100 000 females in the late 1960s, there were appreciable declines, probably reflecting therapeutic improvements, particularly for childhood acute lymphoblastic leukaemias.

Trends were less consistent in Central and South America, since several countries (including Argentina, Chile, Costa Rica, Cuba, Peru, Puerto Rico, Uruguay and Venezuela) showed some declines, though of variable degree, over recent calendar periods, but others (including the Dominican Republic, Ecuador, Mexico and Panama) showed either no clear trend or even some recent increase in mortality. It is possible that these latter trends are influenced by improved accuracy of diagnosis and certification, since the absolute values in these countries remain comparatively low. Besides the cautions on geographical and temporal differences in leukaemia death certification, and the quantitative effect of newer therapies [50, 51], no other inference is possible on leukaemia patterns and trends, since little is known of the causes of the disease, except the recognised effect of ionising radiation [52].

TOTAL, ALL NEOPLASMS

There are at least three patterns of total cancer mortality in Latin America. One includes Argentina and Uruguay, whose rates in the 1950s and 1960s were particularly elevated (between

180 and 210/100 000 males, and between 115 and 135/100 000 females). Over the last few decades, total cancer mortality has declined in Argentina to 154/100 000 males and to 100/100 000 females, and in Uruguay to 119/100 000 females, but remained over 200/100 000 males in Uruguay.

The second pattern of trends includes countries with intermediate total cancer mortality in males but relatively high mortality in females in the 1950s, with rates around 150/100 000 males in earlier calendar periods and over 100/100 000 females. These were Chile, Costa Rica and Cuba. In these countries, total cancer mortality over the last two decades declined appreciably in females and in younger and middle-aged males, but (except in Cuba) not in older males.

The third pattern includes low cancer mortality countries, with rates around or below 100/100 000 in both sexes throughout the calendar period considered, although with different trends over time. For instance, total cancer mortality rates in Peruvian males declined from 65 to 61/100 000, and in females from 71 to 65/100 000, while in Mexico rates increased from 49 to 78/100 000 males and from 76 to 80/100 000 females. In general, in these low mortality countries rates were higher in females than in males, particularly in middle age. Over the same calendar period, overall age-standardised total cancer mortality increased from 136 to 170/100 000 males in Canada, and from 141 to 165/100 000 in the USA, while remained stable in females around 110-115/100 000. Trends in North America were more favourable in middle age, particularly in females.

The patterns observed for total cancer mortality in Latin America should be related to the trends observed for the main

cancer sites. Thus, the early stage of the tobacco-related lung cancer epidemic in several Latin American areas [35] is a main determinant of current geographical differences in cancer mortality. The elevated total cancer mortalities in Argentina and Uruguay are due to the high rates of colorectal, breast and prostatic cancer in these countries, and the relatively high stomach and lung cancer rates. Several declines in intermediate mortality countries are due to the fall in stomach cancer and also, for females, in cervical cancer rates. The elevated cervical cancer rates also explain the higher total cancer rates in females than in males in several low mortality areas. The falls in stomach and cervical cancer rates are, in several countries, still larger than the recent rises in tobacco-related lung cancer in both sexes.

Inspection of truncated and age-specific rates at younger ages shows in most countries even larger falls thus indicating that overall cancer mortality is due to decline further in several areas of South America in the near future [53]. Although absolute values are still low, recent trends in the young are, however, less favourable in several countries of Central America, including Mexico, Dominican Republic and Ecuador.

- Ries LAG, Hankey BF, Miller BA, Hartman AM, Edwards BK. *Cancer Statistics Review*, 1973–88. National Cancer Institute. NIH Publ. No. 91–2789, 1991.
- Institut National du Cancer du Canada. *Statistiques canadiennes sur le cancer* 1989. Toronto, 1989 and various issues.
- Health and Welfare Canada. *Mortality Atlas of Canada*. Vol. 4: *General Mortality Patterns and Recent Trends*. Ottawa, Minister of Supply and Services of Canada, 1991.
- Parkin DM, Läärä E, Muir CS. Estimates of the worldwide frequency of sixteen major cancers in 1980. *Int J Cancer* 1988, 41, 184–197.
- Matos EL, Parkin DM, Loria DI, Vilensky M. Geographic patterns of cancer mortality in Argentina. *Int J Epidemiol* 1990, 19, 860–870.
- La Vecchia C, Lucchini F, Negri E, Boyle P, Maisonneuve P, Levi F. Trends in cancer mortality in Europe, 1955–1989: I, Digestive sites. *Eur J Cancer* 1992, 28, 132–235.
- La Vecchia C, Lucchini F, Negri E, Boyle P, Maisonneuve P, Levi F. Trends in cancer mortality in Europe, 1955–1989: II, Respiratory tract, bone, connective and soft tissues sarcomas, and skin. *Eur J Cancer* 1992, 28, 514–599.
- La Vecchia C, Lucchini F, Negri E, Boyle P, Maisonneuve P, Levi F. Trends in cancer mortality in Europe, 1955–1989: III, Breast and genital sites. *Eur J Cancer* 1992, 28, 927–998.
- La Vecchia C, Lucchini F, Negri E, Boyle P, Maisonneuve P, Levi F. Trends in cancer mortality in Europe, 1955–1989: IV, Urinary tract, eye, brain and nerves, and thyroid. *Eur J Cancer* 1992, 28, 1210–1281.
- La Vecchia C, Lucchini F, Negri E, Boyle P, Maisonneuve P, Levi F. Trends in cancer mortality in Europe, 1955–1989: V, Lymphohaemopoietic and all cancers. *Eur J Cancer* 1992, 28, 1509–1581.
- World Health Organization. *International Classification of Disease*. 6th Revision. Geneva, World Health Organization, 1950.
- World Health Organization. *International Classification of Disease*. 7th Revision. Geneva, World Health Organization, 1957.
- World Health Organization. *International Classification of Disease*. 8th Revision. Geneva, World Health Organization, 1967.
- World Health Organization. *International Classification of Disease*. 9th Revision. Geneva, World Health Organization, 1977.
- Boyle P, Zaridze DG, Smans M. Descriptive epidemiology of colorectal cancer. *Int J Epidemiol* 1985, 36, 9–18.
- Cuzick J, Boyle P. Trends in cervix cancer mortality. *Cancer Surveys* 1988, 7, 417–439.
- Doll R, Smith PG. Comparison between registries: age-standardized rates. In Waterhouse JAH, Muir CS, Shanmugaratnam K, et al., eds. *Cancer Incidence in Five Continents*, Vol. IV. IARC Scientific Publication No 42. Lyon, International Agency for Research on Cancer, 1982, 671–675.
- Doll R, Peto R. The causes of cancer: quantitative estimates of avoidable risks of cancer in the United States today. *J Natl Cancer Inst* 1981, 66, 1191–1308.
- Percy C, Staneck E, Gloeckler L. Accuracy of cancer death certificates and its effects on mortality statistics. *Am J Publ Health* 1981, 71, 242–250.
- Boyle P, Macfarlane GJ, McGinn R, et al. International epidemiology of head and neck cancer. In *Multiple Primary Tumors in the Head and Neck*. Stuttgart, George Thieme, 1990, 80–138.
- Franceschi S, Talamini R, Barra S, et al. Smoking and drinking in relation to cancers of the oral cavity, pharynx, larynx and esophagus in Northern Italy. *Cancer Res* 1990, 50, 6502–6507.
- McLaughlin JK, Gridley G, Block G, et al. Dietary factors in oral and pharyngeal cancer. *J Natl Cancer Inst* 1988, 80, 1237–1243.
- Franceschi S, Bidoli E, Baròn AE, et al. Nutrition and cancer of the oral cavity and pharynx in North-East Italy. *Int J Cancer* 1991, 47, 20–25.
- Franco EL, Kowalski LP, Oliveira BV, et al. Risk factors for oral cancer in Brazil: a case-control study. *Int J Cancer* 1989, 43, 992–1000.
- Tuyns AJ, Péquignot G, Jensen OM. Les cancers de l'oesophage en Ile-et-Vilaine en fonction des niveaux de consommation d'alcool et de tabac. Des risques qui se multiplient. *Bull Cancer (Paris)* 1977, 64, 45–60.
- Victoria CG, Munoz N, Day NE, Barcelos LB, Peccin DA, Braga NM. Hot beverages and oesophageal cancer in Southern Brazil: a case-control study. *Int J Cancer* 1987, 39, 710–716.
- IARC. *Monographs on the Evaluation of Carcinogenic Risks to Humans. Coffee, Tea, Maté, Methylxanthines and Methylglyoxal*. Vol. 51. Geneva, World Health Organization, International Agency for Research on Cancer, 1991.
- Litvak J, Blattner W, Blot W, et al. Workshop on cancer epidemiology in Latin America and the Caribbean. *J Natl Cancer Inst* 1992, 84, 484–488.
- Decarli A, Liati P, Negri E, Franceschi S, La Vecchia C. Vitamin A and other dietary factors in the etiology of esophageal cancer. *Nutr Cancer* 1987, 10, 29–37.
- Howson CP, Hiyama T, Wynder EL. The decline of gastric cancer: epidemiology of an unplanned triumph. *Epidem Rev* 1986, 8, 1–27.
- Parsonnet J, Friedman GD, Vandersteen DP, Vogelstein JH, Orentreich N, Sibley RK. Helicobacter pylori infection and the risk of gastric carcinoma. *N Engl J Med* 1991, 325, 1127–1131.
- La Vecchia C, Negri E, Decarli A, D'Avanzo B, Gallotti L, Franceschi S. A case-control study of diet and colo-rectal cancer in Northern Italy. *Int J Cancer* 1988, 41, 492–498.
- Willett WC, Stampfer MJ, Colditz GA, Rosner BA, Speizer FE. Relation of meat, fat, and fiber intake to the risk of colon cancer in a prospective study among women. *N Engl J Med* 1990, 323, 1664–1672.
- Shopland DR, Eyre HJ, Pechacek TF. Smoking-attributable cancer mortality in 1991: is lung cancer now the leading cause of death among smokers in the United States? *J Natl Cancer Inst* 1991, 83, 1142–1148.
- U.S. Department of Health and Human Services. *Smoking and Health in the Americans*. A Report of the Surgeon General of the Public Health Service, Washington, USGPO, 1992.
- Lee JAH. Melanoma and exposure to sunlight. *Epidem Rev* 1982, 4, 110–136.
- Levi F, La Vecchia C, Te VC, Mezzanotte G. Descriptive epidemiology of skin cancer in the Swiss Canton of Vaud. *Int J Cancer* 1988, 42, 811–816.
- Glass AG, Hoover RN. The emerging epidemic of melanoma and squamous cell skin cancer. *JAMA* 1989, 262, 2097–2100.
- Boyle P. Epidemiology of breast cancer. *Baillière's Clin Oncol* 1988, 2, 1–57.
- Willett WC. The search for the causes of breast and colon cancer. *Nature* 1989, 338, 389–394.
- Brinton LA, Fraumeni JF Jr. Epidemiology of uterine cervical cancer. *J Chron Dis* 1986, 39, 1051–1106.
- Parazzini F, La Vecchia C, Bocciarelli L, Franceschi S. The epidemiology of endometrial cancer. *Gynecol Oncol* 1991, 41, 1–16.
- Skegg DCG, Corwin PA, Paul C, Doll R. Importance of the male factor in cancer of the cervix. *Lancet* 1982, ii, 581–583.
- Brinton LA, Reeves WC, Brenes MM, et al. The male factor in the etiology of cervical cancer among sexually monogamous women. *Int J Cancer* 1989, 44, 199–203.
- Herrero R, Brinton LA, Reeves WC, et al. Sexual behavior, venereal

- diseases, hygiene practices, and invasive cervical cancer in a high-risk population. *Cancer* 1990, **65**, 380–386.
46. La Vecchia C. The epidemiology of cervical neoplasia. *Biomed Pharmacother* 1985, **39**, 426–433.
 47. IARC Working Group on Evaluation of Cervical Screening Programmes. Screening for squamous cervical cancer: duration of low risk after negative results of cervical cytology and its implication for screening policies. *Br Med J* 1986, **293**, 659–664.
 48. Breslow N, Chan CE, Dhondt G, *et al.* Latent carcinoma of prostate at autopsy in seven areas. *Int J Cancer* 1977, **20**, 680–688.
 49. Zaridze DG, Boyle P. Cancer of the prostate: epidemiology and aetiology. *Br J Urol* 1987, **59**, 493–502.
 50. Niemeyer CM, Hitchcock-Bryan S, Sallan SE. Comparative analysis of treatment programs for childhood acute lymphoblastic leukemia. *Sem Oncol* 1985, **12**, 122–130.
 51. La Vecchia C, Levi F, Lucchini F, Garattini S. Progress of anticancer drugs in reducing mortality from selected cancers in Europe: an assessment. *Anti-Cancer Drugs* 1991, **2**, 215–221.
 52. Linet MS. *The Leukemias*. New York, Oxford University Press, 1985, 20–65.
 53. Doll R. Are we winning the fight against cancer? An epidemiological assessment. *Eur J Cancer* 1990, **26**, 500–508.

Acknowledgements—This study has been made possible by a core grant of the Swiss (Aargau and St-Gall) League against cancer. Supports were also received by the Italian Association for Cancer Research and the Italian League against cancer.

This study was conducted within the framework of the CNR (Italian National Research Council) Applied Project “Clinical Application of Oncological Research” and of the “Europe Against Cancer” Programme of the Commission of the European Communities.

Argentina

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	1738	3.88	6.89	374	0.76	0.99
70-74	438	3.83	6.73	98	0.76	1.09
75-79	1586	3.69	7.51	335	0.68	0.94
80-84	3163	4.21	8.85	745	0.80	1.11
85-89	1268	4.11	8.92	289	0.74	1.05

*Information is missing for one or more years.

Chile

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	213	1.89	2.87	112	0.86	1.04
60-64	281	2.18	3.93	137	0.91	1.35
65-69	307	2.24	3.55	132	0.81	1.13
70-74	396	2.53	4.50	162	0.84	0.96
75-79	488	2.73	4.84	141	0.63	0.91
80-84	531	2.60	4.80	137	0.53	0.73
85-89	310	2.35	4.29	116	0.67	0.91

*Information is missing for one or more years.

Colombia

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	319	1.96	2.49	364	1.98	2.24
60-64	363	1.89	2.50	372	1.71	1.75
65-69	486	2.19	2.91	449	1.81	2.06
70-74	213	2.13	2.06	179	1.56	1.57
75-79	407	2.41	3.10	292	1.58	1.52
80-84						
85-89						





*Information is missing for one or more years.

Costa Rica

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	42	2.97	4.96	19	1.36	0.82
65-69	79	4.03	5.15	27	1.17	1.55
70-74	66	2.68	4.11	25	1.02	0.96
75-79	84	3.00	2.87	27	0.80	0.63
80-84	107	3.14	3.24	37	0.99	0.78
85-89	117	3.92	4.33	47	1.31	1.23

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	212	7.53	11.23	53	1.92	2.51
65-69	594	9.58	12.19	132	2.27	2.28
70-74	1553	7.88	11.35	359	1.98	2.49
75-79	1523	6.43	10.64	382	1.87	2.36
80-84	1522	6.08	8.40	423	1.72	2.07
85-89	1189	5.45	7.85	389	1.78	2.37

*Information is missing for one or more years.

Dominican Republic

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	59	1.24	1.19	33	0.70	0.78
70-74	83	1.51	2.01	52	0.98	1.46
75-79	106	1.64	2.22	61	1.00	1.42
80-84	187	2.59	3.13	97	1.35	1.37
85-89	45	3.06	3.50	28	1.60	2.44

*Information is missing for one or more years.

Ecuador

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	42	1.20	1.10	19	0.53	0.69
65-69	67	1.06	1.37	34	0.47	0.44
70-74	70	0.95	1.02	37	0.45	0.50
75-79	66	0.88	1.16	39	0.43	0.52
80-84	81	1.28	1.54	41	0.55	0.74
85-89	91	0.93	1.18	81	0.70	0.74





*Information is missing for one or more years.

Mexico

Deaths from cancer of the mouth or pharynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	206	1.08	1.80	134	0.67	0.75
60-64	712	1.45	2.02	324	0.61	0.89
65-69	825	1.48	2.23	378	0.63	0.76
70-74	940	1.36	1.95	459	0.59	0.80
75-79	718	1.54	2.25	359	0.68	0.90
80-84	877	1.66	2.38	399	0.62	0.81
85-89	700	1.72	2.25	363	0.75	0.97

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	40	2.79	3.47		46	3.24	4.55	
60-64	32	1.89	1.86		37	2.78	2.13	
65-69	45	2.37	2.53		32	1.73	1.77	
70-74	64	2.75	3.56		37	1.57	1.64	
75-79	82	3.10	4.89		64	2.45	2.72	
80-84	86	2.68	3.76		73	2.14	2.14	
85-89*	75	3.41	3.45		40	1.86	2.40	

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64								
65-69								
70-74	72	4.18	6.35		23	1.12	1.06	
75-79	69	3.16	5.39		29	1.07	0.67	
80-84*	32	1.70	3.56		14	0.52	0.67	
85-89*	36	2.89	5.24		11	0.70	1.49	

*Information is missing for one or more years.

Peru

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64								
65-69*	160	1.26	1.94		70	0.50	0.55	
70-74*	127	0.92	1.23		78	0.47	0.75	
75-79*	64	0.76	0.88		38	0.38	0.47	
80-84*	156	0.83	1.05		75	0.33	0.36	
85-89								




*Information is missing for one or more years.

Puerto Rico




Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64*	366	9.93	17.92		109	2.73	3.14	
65-69*	209	10.06	16.97		39	1.76	1.98	
70-74	638	10.59	18.40		173	2.60	3.59	
75-79*	498	9.34	16.06		117	1.93	2.10	
80-84	683	8.83	14.54		169	1.85	2.31	
85-89*	466	9.26	16.30		85	1.40	1.78	




*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.






Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Uruguay

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	323	5.04	9.34		73	0.98	1.17	
60-64*	204	4.90	9.77		47	0.98	1.07	
65-69	410	5.62	11.24		96	1.16	1.38	
70-74	486	6.28	12.23		81	0.91	1.05	
75-79*	402	6.12	13.15		61	0.71	0.94	
80-84	574	6.68	13.82		87	0.81	1.19	
85-89*	372	6.86	13.84		73	1.04	1.50	

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Venezuela

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	235	3.20	4.78		344	4.10	5.33	
60-64	306	3.55	5.50		309	3.13	3.68	
65-69	362	3.44	5.26		334	2.79	3.47	
70-74	374	2.95	4.55		316	2.31	2.59	
75-79	374	2.40	3.76		329	1.99	2.09	
80-84	356	2.26	3.53		295	1.72	1.85	
85-89*	314	2.24	3.51		203	1.33	1.40	

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Canada

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	1886	4.29	4.90		497	1.13	1.38	
60-64	2008	4.15	5.38		597	1.19	1.73	
65-69	2204	4.18	5.74		645	1.13	1.75	
70-74	2457	4.33	6.35		875	1.32	2.11	
75-79	2743	4.43	7.39		959	1.28	2.07	
80-84	2930	4.28	7.32		1120	1.32	2.21	
85-89*	2794	4.51	7.94		1044	1.34	2.16	

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

United States of America

Deaths from cancer of the mouth or pharynx

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	22473	4.69	7.43		6318	1.16	1.84	
60-64	24323	4.70	7.99		7734	1.29	2.23	
65-69	25933	4.75	8.50		8933	1.35	2.52	
70-74	27582	4.75	9.40		11026	1.52	2.90	
75-79	29047	4.64	8.28		12586	1.56	2.86	
80-84	29141	4.32	7.73		13394	1.49	2.60	
85-89*	21937	3.81	6.83		10812	1.37	2.24	

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Argentina

Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	5441	12.28	16.73	1978	4.04	4.08
70-74*	1478	13.14	18.11	559	4.15	3.59
75-79*	4534	10.54	14.63	1611	3.15	2.87
80-84	7167	9.30	13.87	2480	2.51	2.52
85-89*	2819	8.87	13.25	986	2.33	2.44

*Information is missing for one or more years.

Chile

Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	926	8.46	12.79	561	4.37	6.55
60-64	1134	9.28	13.59	724	4.96	6.51
65-69	1477	11.11	14.83	924	5.81	7.16
70-74	1597	10.61	13.28	913	4.89	5.61
75-79	1653	9.59	11.25	1027	4.49	5.23
80-84	1711	8.64	9.71	1016	3.80	3.86
85-89*	1084	8.36	7.98	656	3.73	3.57

*Information is missing for one or more years.

Colombia

Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	573	3.49	5.09	318	1.75	2.38
60-64	663	3.51	4.79	433	2.04	3.03
65-69	893	4.09	5.65	567	2.33	3.02
70-74	408	3.98	5.38	248	2.16	2.57
75-79*	741	4.46	5.57	483	2.66	2.92
80-84						
85-89						


*Information is missing for one or more years.

Costa Rica


Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	71	5.16	6.60	21	1.57	2.56
65-69	99	5.14	5.71	55	2.85	2.32
70-74	89	3.81	3.91	47	1.84	2.17
75-79	127	4.69	4.95	51	1.76	1.51
80-84	101	3.10	3.05	50	1.46	1.33
85-89*	120	4.40	2.81	63	1.81	1.68


*Information is missing for one or more years.




Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	161	5.72	8.35	43	1.63	1.73
65-69*	316	5.18	5.77	145	2.60	2.44
70-74	967	4.98	6.28	336	1.87	2.16
75-79	978	4.29	5.15	356	1.71	1.80
80-84	1082	4.24	5.19	401	1.66	2.03
85-89*	983	4.40	5.74	328	1.48	1.65

*Information is missing for one or more years.

Dominican Republic

Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	46	0.95	1.63	15	0.29	0.55
70-74	62	1.15	2.35	28	0.54	0.86
75-79	81	1.25	1.87	35	0.49	0.76
80-84	78	1.07	1.60	54	0.77	1.04
85-89*	21	1.36	1.51	6	0.39	0.62

*Information is missing for one or more years.

Ecuador

Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	65	1.83	1.89	19	0.55	0.42
65-69	174	2.87	3.14	38	0.54	0.60
70-74	184	2.55	3.13	53	0.64	0.60
75-79*	177	2.44	2.25	73	0.91	0.98
80-84*	172	2.80	2.11	65	0.95	0.71
85-89*	261	2.70	2.71	86	0.81	0.72


*Information is missing for one or more years.

Mexico


Deaths from cancer of the oesophagus

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	207	1.14	1.50	114	0.58	0.93
60-64	699	1.45	2.20	342	0.67	0.89
65-69	791	1.43	2.01	403	0.68	0.87
70-74	1045	1.52	2.07	562	0.75	1.01
75-79	721	1.63	2.10	380	0.76	1.01
80-84*	898	1.70	2.22	475	0.76	0.90
85-89*	746	1.85	2.01	349	0.72	0.79


*Information is missing for one or more years.




Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama*Deaths from cancer of the oesophagus*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	37	2.61	3.55	14	0.99	0.95
60-64	33	1.95	2.07	14	0.97	0.82
65-69	34	1.86	1.98	17	0.82	1.22
70-74	55	2.38	3.37	20	0.93	0.96
75-79	52	2.02	2.45	36	1.44	2.12
80-84	52	1.65	2.27	25	0.78	0.52
85-89*	40	1.89	2.95	19	0.92	1.03

*Information is missing for one or more years.

Paraguay (reporting)*Deaths from cancer of the oesophagus*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	80	4.55	5.14	26	1.20	1.21
75-79	111	5.25	7.94	25	0.94	1.14
80-84*	76	4.22	5.47	27	1.09	1.35
85-89*	55	4.45	5.98	11	0.77	0.46

*Information is missing for one or more years.

Peru*Deaths from cancer of the oesophagus*





Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	198	1.67	2.30	56	0.41	0.35
70-74*	283	2.08	2.92	50	0.34	0.32
75-79*	128	1.50	1.92	45	0.45	0.61
80-84*	245	1.37	1.14	70	0.33	0.29
85-89						

*Information is missing for one or more years.

Puerto Rico*Deaths from cancer of the oesophagus*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	590	15.99	26.69	249	6.56	10.46
65-69*	317	15.08	25.37	122	5.67	8.20
70-74	893	14.75	22.00	326	4.93	6.04
75-79*	716	13.25	19.37	271	4.45	5.41
80-84	890	10.81	14.82	261	2.72	2.69
85-89*	565	10.74	15.76	143	2.29	2.36

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay*Deaths from cancer of the oesophagus*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1009	15.89	23.50	529	6.94	6.74
60-64*	617	14.91	20.23	320	6.42	6.29
65-69	1172	16.21	22.79	433	4.95	4.66
70-74	1184	15.21	22.70	424	4.48	4.00
75-79*	1017	15.03	23.37	354	3.98	4.71
80-84	1216	13.42	21.51	459	3.62	3.86
85-89*	694	11.96	16.25	295	3.66	3.14

*Information is missing for one or more years.

Venezuela*Deaths from cancer of the oesophagus*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	248	3.45	5.47	197	2.35	3.63
60-64	335	3.93	5.73	251	2.59	3.86
65-69	399	4.12	5.35	296	2.55	3.22
70-74	418	3.51	4.48	314	2.27	2.74
75-79	513	3.52	4.32	290	1.76	1.92
80-84*	464	3.15	3.76	262	1.53	1.34
85-89*	365	2.79	3.57	185	1.25	1.39

*Information is missing for one or more years.

Canada*Deaths from cancer of the oesophagus*





Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1277	2.95	3.48	544	1.22	1.39
60-64	1466	3.09	3.83	574	1.12	1.29
65-69	1748	3.35	4.32	668	1.12	1.42
70-74	1988	3.51	4.77	815	1.16	1.43
75-79	2289	3.67	5.16	949	1.17	1.51
80-84	2694	3.86	5.42	1041	1.11	1.26
85-89*	2616	4.26	6.51	1011	1.16	1.32

*Information is missing for one or more years.

United States of America*Deaths from cancer of the oesophagus*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	17794	3.72	5.98	4976	0.90	1.33
60-64	19700	3.80	6.20	5665	0.92	1.51
65-69	21441	3.91	6.59	6844	1.01	1.78
70-74	23378	4.00	6.80	8153	1.10	1.95
75-79	27042	4.29	7.42	9713	1.16	2.02
80-84	30232	4.43	7.49	11086	1.18	1.94
85-89*	26583	4.53	7.63	9373	1.14	1.74

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	10888	24.60	31.35	5834	11.94	12.25
70-74*	2795	24.97	29.34	1447	10.93	11.34
75-79*	7206	16.81	21.45	4026	8.00	8.46
80-84	11007	14.31	18.69	6170	6.45	7.53
85-89*	4273	13.38	17.83	2346	5.69	6.64

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Chile

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	7667	69.41	103.90	5666	43.87	61.32
60-64	8180	66.55	93.56	6204	42.60	54.03
65-69	8204	61.26	81.07	5896	36.92	42.72
70-74	8894	58.78	76.85	5972	31.71	35.77
75-79	8388	48.25	60.78	5063	21.93	24.12
80-84	7766	38.73	47.78	4480	16.78	17.87
85-89*	4481	34.38	42.52	2524	14.33	15.11

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Colombia

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	3279	19.65	31.73	2801	15.12	23.34
60-64	4428	22.05	35.35	3989	18.24	27.17
65-69	6147	27.70	40.79	4963	20.14	28.52
70-74*	2920	28.00	38.93	2404	20.60	29.13
75-79*	4841	28.42	40.74	3879	20.76	27.21

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Costa Rica

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	824	60.29	79.76	477	34.13	44.13
65-69	1118	56.64	74.68	627	30.85	35.79
70-74	1198	50.33	69.29	708	28.83	33.78
75-79	1276	47.01	55.16	663	22.30	24.67
80-84	1507	45.40	49.62	732	20.57	23.11
85-89*	1405	48.59	47.25	789	22.64	23.08

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	501	17.90	20.48	230	8.83	7.38
65-69*	975	16.12	13.66	411	7.13	7.02
70-74	2363	11.83	12.43	1072	5.91	6.51
75-79	2298	10.01	10.74	1056	5.04	5.54
80-84	2190	8.41	9.93	1125	4.58	5.36
85-89*	1721	7.51	8.97	830	3.67	4.34

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Dominican Republic

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	198	4.09	5.54	90	1.81	2.53
70-74	243	4.25	6.09	100	1.80	2.64
75-79	223	3.45	4.59	130	1.98	3.00
80-84	330	4.66	6.01	198	2.76	4.17
85-89*	89	5.85	8.47	55	3.46	5.16

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Ecuador

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	544	14.76	19.20	491	13.63	14.66
65-69	1384	22.27	26.16	1003	14.58	17.48
70-74	1703	25.98	26.50	1284	16.27	17.40
75-79*	1683	23.21	26.41	1357	16.27	18.16
80-84*	1724	27.62	30.54	1338	19.10	20.83
85-89*	2660	27.11	29.18	2092	18.74	18.08

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Mexico

Deaths from cancer of the stomach

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	1496	8.04	11.92	1629	8.43	11.61
60-64	4613	9.60	14.32	4660	9.15	12.44
65-69	5782	10.40	13.39	5565	9.33	11.66
70-74	6668	9.90	13.01	6598	8.70	11.04
75-79*	4282	9.42	11.41	4349	8.55	9.96
80-84*	5133	9.60	11.24	4775	7.62	8.90
85-89*	3834	9.53	10.82	3669	7.52	8.85

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama*Deaths from cancer of the stomach*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	242	16.83	22.15	124	8.43	13.46
60-64	263	15.69	20.54	134	8.18	11.22
65-69	258	13.66	19.38	140	6.97	9.32
70-74	324	13.81	17.77	177	7.72	9.77
75-79	348	13.46	17.73	196	7.41	9.60
80-84	378	11.68	15.49	231	6.90	9.64
85-89*	260	12.10	17.38	152	7.02	8.87

*Information is missing for one or more years.

Paraguay (reporting)*Deaths from cancer of the stomach*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	271	15.69	24.79	215	10.08	14.89
75-79	333	15.59	21.20	248	9.33	14.16
80-84	189	10.03	12.03	113	4.93	7.75
85-89*	129	10.48	12.21	84	5.85	6.22

*Information is missing for one or more years.

Peru*Deaths from cancer of the stomach*




Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	2366	19.33	27.46	1994	14.34	19.08
70-74*	2697	19.81	24.70	2144	13.94	16.97
75-79*	1391	16.14	20.59	1149	11.99	14.75
80-84*	2685	14.26	16.38	2331	10.80	12.51
85-89*						

*Information is missing for one or more years.




Puerto Rico*Deaths from cancer of the stomach*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	1093	29.46	40.28	552	13.73	16.30
65-69*	576	26.95	34.09	269	12.02	15.36
70-74	1335	21.17	24.80	627	9.15	11.75
75-79*	773	13.64	14.21	497	8.17	10.30
80-84	1068	12.35	12.69	544	5.77	7.79
85-89*	685	12.39	14.33	333	5.20	5.83




*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay*Deaths from cancer of the stomach*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	2419	38.11	54.77	1463	19.56	22.03
60-64*	1454	35.61	45.60	875	17.98	17.24
65-69	2599	35.99	44.96	1441	16.62	16.56
70-74	2532	32.64	39.81	1425	15.23	14.55
75-79*	1704	24.53	29.42	1009	11.25	10.91
80-84	1691	18.29	21.98	1099	9.02	10.29
85-89*	981	16.83	20.98	562	7.16	7.60

*Information is missing for one or more years.

Venezuela*Deaths from cancer of the stomach*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	2138	28.91	44.84	1790	20.81	32.34
60-64	2747	32.16	48.82	2130	21.12	30.90
65-69	3388	33.93	46.77	2521	20.77	29.41
70-74	3743	30.87	41.70	2463	17.34	23.82
75-79	3881	25.99	35.16	2653	15.97	19.63
80-84*	3253	21.57	27.45	2223	12.87	15.50
85-89*	2334	17.56	20.68	1575	10.28	11.59

*Information is missing for one or more years.

Canada*Deaths from cancer of the stomach*




Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	9611	22.25	27.52	4894	10.92	11.84
60-64	9172	18.96	22.29	4682	8.96	9.87
65-69	8518	16.22	18.92	4513	7.38	7.83
70-74	8031	14.03	16.29	4063	6.53	7.41
75-79	7403	11.66	13.37	4144	5.08	5.79
80-84	6905	9.58	11.19	4124	4.29	4.91
85-89*	5385	8.33	9.26	3263	3.71	4.30

*Information is missing for one or more years.




United States of America*Deaths from cancer of the stomach*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	67973	13.92	16.98	39982	7.04	8.33
60-64	61202	11.44	13.81	37004	5.68	6.70
65-69	52535	9.19	11.16	33495	4.56	5.44
70-74	46883	7.72	9.70	30578	3.70	4.50
75-79	43690	6.58	8.28	29439	3.10	3.77
80-84	42482	5.86	7.47	29009	2.75	3.39
85-89*	33091	5.28	6.95	22133	2.30	2.93




*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64								
65-69*	7138	16.17	18.37		6800	13.95	15.72	
70-74*	1892	16.98	18.97		1839	13.94	16.13	
75-79*	6058	14.14	16.14		5817	11.56	13.00	
80-84	10111	13.05	15.02		9502	9.87	11.27	
85-89*	4216	13.07	14.85		3883	9.40	11.10	

*Information is missing for one or more years.

Chile

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	590	5.25	7.11		788	5.97	8.79	
60-64	677	5.41	7.79		945	6.42	8.30	
65-69	914	6.78	8.69		1204	7.48	8.93	
70-74	1101	7.16	8.63		1451	7.62	8.26	
75-79	1219	7.01	7.51		1646	7.09	8.01	
80-84	1391	6.89	7.57		1699	6.35	7.15	
85-89*	908	6.89	7.26		1185	6.64	7.02	

*Information is missing for one or more years.

Colombia

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	528	3.18	4.64		686	3.71	5.64	
60-64	602	3.00	4.23		801	3.62	5.54	
65-69	831	3.60	5.06		991	3.93	5.46	
70-74*	405	3.77	5.08		472	4.04	4.92	
75-79*	651	3.76	4.92		799	4.15	5.33	





*Information is missing for one or more years.

Costa Rica

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59	73	5.22	5.28		81	5.72	7.25	
60-64*								
65-69	107	5.24	7.26		119	5.84	7.18	
70-74	107	4.41	4.38		131	5.15	6.57	
75-79	150	5.25	5.65		179	5.89	6.71	
80-84	193	5.55	5.98		235	6.50	5.84	
85-89*	223	7.19	6.71		247	6.89	7.70	

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64*	303	10.74	12.80		327	12.38	13.73	
65-69*	681	11.14	9.84		723	12.53	11.88	
70-74	1924	9.65	10.64		1880	10.40	12.41	
75-79	2065	9.01	9.78		2209	10.61	12.19	
80-84	2335	9.00	10.69		2586	10.59	12.47	
85-89*	2170	9.42	10.62		2396	10.80	13.55	

*Information is missing for one or more years.

Dominican Republic

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64								
65-69	119	2.37	3.19		117	2.45	3.11	
70-74	145	2.64	3.36		144	2.60	3.34	
75-79	190	2.82	3.24		190	2.89	3.50	
80-84	238	3.27	4.15		240	3.31	4.37	
85-89*	59	3.98	3.10		57	3.51	4.67	

*Information is missing for one or more years.

Ecuador

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59								
60-64*	88	2.37	2.76		127	3.51	3.76	
65-69	219	3.43	3.86		265	3.80	4.54	
70-74	237	3.17	3.27		310	3.83	4.79	
75-79*	189	2.48	2.92		272	3.26	3.42	
80-84*	174	2.79	2.75		210	3.02	3.47	
85-89*	299	2.91	3.24		374	3.37	3.86	





*Information is missing for one or more years.

Mexico

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males				Females			
	Deaths No.	World Stand	Trunc 35-64		Deaths No.	World Stand	Trunc 35-64	
55-59*								
60-64	1254	2.57	3.38		1962	3.77	4.68	
65-69	1598	2.80	3.68		2181	3.62	4.75	
70-74	1878	2.64	3.43		2476	3.24	4.20	
75-79*	1232	2.61	3.41		1691	3.32	4.14	
80-84*	1322	2.39	2.93		1674	2.66	3.35	
85-89*	1258	3.02	3.61		1515	3.11	3.81	

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Panama

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	66	4.50	5.02	63	4.42	6.69
60-64	62	3.62	5.14	89	5.40	8.31
65-69	87	4.73	5.89	88	4.64	6.74
70-74	111	4.89	6.43	113	4.95	6.35
75-79	137	5.10	7.46	117	4.43	5.55
80-84	178	5.48	6.51	170	5.24	6.96
85-89*	121	5.36	5.98	110	5.06	5.08

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Paraguay (reporting)

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	123	6.92	8.35	156	7.27	8.71
75-79	119	5.28	5.31	151	5.66	7.12
80-84*	70	3.61	4.46	76	3.43	4.33
85-89*	58	4.67	4.47	59	4.11	3.55

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Peru

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	427	3.41	4.17	442	3.16	4.25
65-69*	397	2.81	3.55	487	3.14	3.83
70-74*	223	2.46	2.52	262	2.73	3.41
75-79*	425	2.21	2.26	503	2.36	2.44

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Puerto Rico

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	219	5.84	8.18	248	6.21	8.52
60-64*	149	7.03	9.00	141	6.23	7.52
65-69*	425	6.85	9.70	453	6.58	9.01
70-74*	418	7.46	9.39	411	6.79	8.41
75-79*	596	7.22	10.28	579	6.16	8.11
80-84*	457	8.74	13.31	458	7.16	8.28

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Uruguay

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	963	15.09	20.00	1201	16.09	18.90
60-64*	688	16.84	21.90	855	17.80	21.78
65-69	1440	19.99	24.28	1660	19.50	23.04
70-74	1553	19.98	24.87	1827	19.65	21.30
75-79*	1269	18.55	21.99	1415	16.24	18.04
80-84	1609	17.25	20.09	1783	14.78	17.22
85-89*	1066	17.90	19.22	1216	15.51	16.18

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Venezuela

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	369	5.07	6.21	573	6.57	10.02
60-64	437	4.98	7.14	646	6.30	9.11
65-69	550	5.35	6.72	710	5.76	7.21
70-74	626	4.89	6.56	784	5.38	7.39
75-79	799	5.17	6.11	1065	6.31	7.30
80-84*	718	4.51	5.40	896	5.16	6.20
85-89*	647	4.72	5.68	784	5.10	5.84

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Canada

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	9187	21.16	25.18	9029	20.70	27.46
60-64	10001	20.64	24.66	10054	19.75	25.72
65-69	11280	21.33	25.01	11274	19.00	24.50
70-74	12383	21.55	24.93	12620	18.53	23.99
75-79	13185	20.72	23.94	13364	16.60	20.61
80-84	13538	18.79	21.72	13238	14.17	17.68
85-89*	11781	18.33	21.14	11301	13.15	16.03

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




United States of America

Deaths from cancer of the intestines, chiefly colon and rectum

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	93153	19.11	23.75	97241	17.48	23.85
60-64	100422	18.83	23.13	106615	16.78	22.40
65-69	108340	19.01	23.01	114345	15.96	20.84
70-74	115943	19.02	22.68	123915	15.22	19.44
75-79	126259	18.87	22.00	134830	14.28	18.07
80-84	134641	18.28	21.04	140903	13.27	16.09
85-89*	112404	17.53	20.44	115588	12.45	14.98

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	244	1.69	2.15	230	1.36	1.52
80-84	2612	3.39	4.44	2692	2.85	3.64
85-89*	1477	4.63	5.87	1497	3.67	4.58

*Information is missing for one or more years.

Chile*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84	694	3.38	4.62	772	2.79	3.56
85-89*	616	4.64	5.67	662	3.85	5.13

*Information is missing for one or more years.

Colombia*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84						
85-89						

Costa Rica*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84	207	5.73	6.73	133	3.58	4.17
85-89*	248	7.75	8.72	180	5.15	6.00

*Information is missing for one or more years.

Cuba*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	71	1.46	2.01	44	0.95	1.23
80-84	354	1.39	1.60	271	1.13	1.38
85-89*	348	1.54	1.94	285	1.29	1.46

*Information is missing for one or more years.

Dominican Republic*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84	303	4.05	6.60	233	3.13	4.32
85-89*	32	1.84	4.15	8	0.42	0.87

*Information is missing for one or more years.

Ecuador*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	33	1.61	2.03	25	1.12	1.08
80-84*	100	1.60	1.86	85	1.21	1.97
85-89*	179	1.75	2.50	213	1.92	2.22

*Information is missing for one or more years.

Mexico*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84*	527	0.95	1.19	531	0.85	1.17
85-89*	547	1.32	1.69	572	1.18	1.64

*Information is missing for one or more years.

Panama*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	10	1.97	2.22	4	0.74	1.66
80-84	71	2.15	2.57	51	1.54	1.48
85-89*	43	1.70	1.91	25	1.15	1.58

*Information is missing for one or more years.

Paraguay (reporting)*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	1	0.23	0.00	2	0.40	0.59
80-84*	11	0.61	1.22	44	2.06	2.54
85-89*	2	0.17	0.26	5	0.30	0.46

*Information is missing for one or more years.

Peru*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84*	300	1.40	1.78	279	1.25	1.58
85-89						

*Information is missing for one or more years.

Puerto Rico*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	15	1.08	1.55	8	0.55	0.73
80-84	107	1.30	1.86	62	0.68	0.91
85-89*	89	1.75	2.59	28	0.46	0.55

*Information is missing for one or more years.

Uruguay*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79						
80-84	72	0.81	1.19	46	0.39	0.30
85-89*	51	0.88	1.34	37	0.49	0.49

*Information is missing for one or more years.

Venezuela*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	161	4.52	5.45	162	4.24	5.45
80-84*	408	2.56	3.56	316	1.78	2.67
85-89*	252	1.80	2.30	203	1.33	1.68

*Information is missing for one or more years.

Canada*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	178	1.41	2.28	80	0.54	0.78
80-84	907	1.32	1.86	468	0.56	0.76
85-89*	1049	1.73	2.40	501	0.63	0.76

*Information is missing for one or more years.

United States of America*Deaths from cancer of the liver*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74						
75-79*	1632	1.24	1.74	818	0.48	0.72
80-84	9058	1.32	1.94	4317	0.48	0.68
85-89*	8799	1.48	2.19	4008	0.50	0.73




*Information is missing for one or more years.

Argentina

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	2308	5.15	9.54	193	0.40	0.53
70-74	584	5.11	8.99	66	0.49	0.72
75-79	2189	5.12	10.39	197	0.40	0.51
80-84	3995	5.30	11.22	321	0.36	0.59
85-89	1551	5.00	9.89	113	0.28	0.40

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Chile

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	166	1.46	2.44	49	0.38	0.54
60-64	177	1.41	2.67	32	0.22	0.40
65-69	264	1.92	3.69	57	0.35	0.51
70-74	319	2.09	3.43	62	0.33	0.40
75-79	414	2.35	3.65	78	0.34	0.43
80-84	436	2.19	3.73	72	0.27	0.28
85-89	275	2.12	2.85	43	0.26	0.38

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Colombia

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	224	1.37	2.05	153	0.86	0.80
60-64	325	1.73	2.33	175	0.81	1.12
65-69	374	1.72	2.43	228	0.93	1.00
70-74	218	2.09	2.79	108	0.94	0.81
75-79	363	2.17	2.86	154	0.82	0.98

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Costa Rica

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	23	1.72	2.72	7	0.42	1.06
60-64						
65-69	25	1.24	1.94	4	0.22	0.00
70-74	47	2.03	2.06	9	0.37	0.25
75-79	53	2.00	2.19	13	0.42	0.33
80-84	83	2.56	2.48	21	0.60	0.51
85-89	91	3.19	2.73	21	0.63	0.65

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Cuba

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	197	7.08	9.21	32	1.21	1.61
60-64	461	7.51	9.74	91	1.58	2.40
65-69	1097	5.52	6.77	146	0.80	1.02
70-74	1232	5.48	6.99	210	1.04	1.40
75-79	1379	5.50	6.92	233	0.98	1.30
80-84	1197	5.41	6.53	210	1.02	1.66

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Dominican Republic

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	34	0.75	0.91	12	0.30	0.30
70-74	53	1.01	1.48	10	0.20	0.40
75-79	54	0.79	1.35	18	0.30	0.50
80-84	116	1.65	2.13	38	0.53	0.73
85-89	25	1.68	2.29	7	0.49	1.05

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Ecuador

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	33	0.87	1.10	6	0.18	0.16
65-69	76	1.27	0.88	16	0.21	0.28
70-74	95	1.39	1.92	20	0.24	0.19
75-79	86	1.21	1.29	20	0.24	0.35
80-84	83	1.36	1.87	18	0.26	0.37
85-89	111	1.16	1.22	24	0.22	0.21

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Mexico

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	324	1.74	2.74	135	0.68	0.77
60-64	926	1.94	2.71	373	0.72	1.09
65-69	1120	2.03	2.92	388	0.65	0.84
70-74	1411	2.05	2.94	438	0.58	0.86
75-79	990	2.20	3.02	310	0.62	0.89
80-84	1219	2.34	2.76	323	0.52	0.58
85-89	914	2.33	2.94	230	0.48	0.51

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 35-64 to 70-74; bottom panel, age-specific death rate for females 35-64 to 70-74.

Panama

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	26	1.91	2.01	6	0.43	0.83
60-64	33	2.04	2.17	4	0.25	0.17
65-69	40	2.09	1.76	5	0.26	0.17
70-74	52	2.25	3.08	14	0.59	0.90
75-79	46	1.84	3.08	12	0.40	0.43
80-84	55	1.72	2.57	11	0.35	0.51
85-89*	45	2.12	2.61	7	0.31	0.18

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	55	3.15	6.39	10	0.47	0.75
75-79	41	1.88	3.79	10	0.34	0.43
80-84*	29	1.60	2.07	7	0.34	0.36
85-89*	25	2.09	2.89	0	0.00	0.00

*Information is missing for one or more years.

Peru

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	119	1.00	1.28	28	0.18	0.28
70-74*	144	1.11	1.27	31	0.20	0.19
75-79*	63	0.75	0.98	16	0.17	0.21
80-84*	126	0.65	0.70	27	0.12	0.12
85-89						





*Information is missing for one or more years.

Puerto Rico

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	164	4.42	7.27	36	0.94	2.24
65-69*	85	4.04	6.03	23	1.08	1.56
70-74	280	4.62	7.76	63	0.95	1.23
75-79*	260	4.93	7.73	49	0.81	0.97
80-84	367	4.52	6.63	53	0.57	0.76
85-89*	213	4.12	5.85	34	0.56	0.83

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	406	6.38	11.33	17	0.23	0.30
60-64*	271	6.53	11.41	26	0.53	0.79
65-69	488	6.71	11.65	38	0.46	0.71
70-74	608	7.85	15.10	35	0.40	0.45
75-79	466	7.05	12.93	23	0.29	0.54
80-84	671	7.66	14.33	38	0.34	0.53
85-89*	391	6.96	12.47	20	0.26	0.35

*Information is missing for one or more years.

Venezuela

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	222	3.02	5.19	132	1.58	2.22
60-64	290	3.45	5.44	145	1.46	1.74
65-69	344	3.39	5.39	191	1.69	1.88
70-74	347	2.87	4.04	158	1.11	1.55
75-79	450	3.02	4.29	165	1.02	1.08
80-84*	423	2.84	3.58	135	0.80	0.86
85-89*	374	2.78	4.38	76	0.50	0.49

*Information is missing for one or more years.

Canada

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	737	1.76	2.65	118	0.27	0.40
60-64	829	1.78	2.81	123	0.25	0.31
65-69	994	1.97	3.01	133	0.24	0.38
70-74	1260	2.25	3.35	197	0.32	0.58
75-79	1444	2.35	3.72	212	0.29	0.50
80-84	1748	2.52	4.00	303	0.37	0.62
85-89*	1561	2.55	3.96	315	0.43	0.76





*Information is missing for one or more years.

United States of America

Deaths from cancer of the larynx

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	9869	2.07	3.41	1045	0.20	0.36
60-64	11067	2.15	3.68	1240	0.21	0.45
65-69	12295	2.26	3.87	1560	0.25	0.54
70-74	13548	2.29	3.85	1981	0.29	0.63
75-79	14233	2.24	3.66	2650	0.35	0.75
80-84	14624	2.12	3.42	3186	0.38	0.75
85-89*	11690	1.97	3.18	2761	0.38	0.69

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Argentina

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	20442	45.71	81.74	2754	5.74	8.26
70-74*	5481	48.13	81.71	754	5.80	8.52
75-79*	17902	41.98	70.22	2641	5.41	7.24
80-84	31214	41.30	72.60	4989	5.42	7.76
85-89*	12388	39.65	68.28	2044	5.29	8.24

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Chile

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1391	12.37	21.72	565	4.36	6.66
60-64	1739	13.80	23.34	718	4.90	7.74
65-69	2259	16.56	27.14	851	5.33	8.22
70-74	2846	18.70	29.82	945	5.10	7.79
75-79	3453	19.67	30.98	1165	5.21	7.76
80-84	4279	21.50	31.80	1414	5.56	8.03
85-89*	2810	21.72	31.48	1016	6.08	8.59

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Colombia

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	767	4.69	7.35	510	2.68	4.67
60-64	1161	6.08	8.92	608	2.79	4.31
65-69	1603	7.28	10.59	840	3.39	5.24
70-74*	873	8.38	12.26	484	4.17	6.50
75-79*	1657	9.75	13.23	929	4.85	7.63

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Costa Rica

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	87	6.43	9.81	53	3.89	6.06
60-64*	150	7.69	11.65	77	3.97	4.91
65-69	197	8.31	11.37	118	4.84	5.13
70-74	322	12.01	16.57	154	5.31	5.51
75-79	510	15.81	18.73	222	6.45	7.94
80-84	475	16.53	17.32	205	6.03	5.95

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Cuba

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1102	39.73	52.12	386	14.81	19.01
60-64*	2533	41.78	47.69	799	14.05	17.68
65-69	3844	47.24	2573	1446	18.33	18.33
70-74	5658	37.78	44.44	2772	13.56	16.63
75-79	9767	38.30	44.87	3198	13.40	17.32
80-84	8767	39.30	49.39	3049	14.10	18.77

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Dominican Republic

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	175	3.69	5.55	75	1.75	2.56
70-74	232	4.11	7.06	87	1.68	2.42
75-79	272	4.20	6.61	128	2.06	3.40
80-84	469	6.51	9.66	201	2.82	4.78
85-89*	111	7.31	10.36	31	1.99	2.79

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Ecuador

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	135	3.63	4.21	100	2.38	4.05
65-69	267	4.20	4.85	176	2.53	3.34
70-74	333	4.58	6.09	172	2.13	3.01
75-79*	332	4.84	5.89	162	2.00	2.19
80-84*	409	6.67	7.42	203	2.90	3.34
85-89*	717	7.39	8.61	326	2.85	3.41

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Mexico

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	1021	5.55	8.04	645	3.25	5.10
60-64	2996	6.20	9.74	1868	3.60	5.27
65-69	4231	7.58	11.26	2446	4.04	6.24
70-74	5615	8.14	12.01	3032	4.05	5.99
75-79*	4735	10.59	14.41	2201	4.42	6.33
80-84	6940	13.28	17.71	3127	5.12	7.11
85-89*	3893	14.95	18.12	2728	5.75	7.69

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	112	7.72	14.33	25	1.78	1.91
60-64	137	8.14	14.76	38	2.33	3.68
65-69	170	9.03	13.59	36	1.72	3.19
70-74	241	10.37	18.25	62	2.73	3.70
75-79	342	13.14	22.30	82	3.23	4.68
80-84	462	14.55	21.68	126	3.89	5.77
85-89*	318	14.77	20.05	99	4.64	6.56

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	153	8.84	17.18	49	2.44	4.99
75-79	194	9.24	16.00	62	2.37	3.36
80-84*	160	8.70	15.63	49	2.34	4.10
85-89*	141	11.68	18.35	31	2.28	2.77

*Information is missing for one or more years.

Peru

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	841	6.93	9.76	358	2.60	3.37
70-74*	953	6.99	9.53	393	2.55	3.62
75-79*	545	6.27	8.67	227	2.43	2.99
80-84*	1266	6.78	8.23	591	2.74	3.79
85-89						




*Information is missing for one or more years.

Puerto Rico




Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	428	11.72	16.97	170	4.37	6.61
65-69*	277	13.18	17.63	111	5.19	7.88
70-74	924	15.15	21.31	363	5.35	8.15
75-79*	858	15.88	22.30	346	6.02	8.84
80-84	1274	15.76	23.59	595	6.49	8.40
85-89*	927	17.60	25.52	409	6.58	7.58




*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	2111	33.18	58.27	257	3.57	4.61
60-64*	1612	39.25	64.60	193	4.07	4.69
65-69	3154	43.78	66.69	309	3.85	4.38
70-74	3463	44.52	68.61	346	3.84	5.41
75-79*	3218	48.25	76.09	328	3.99	5.71
80-84	4663	52.43	87.48	455	4.05	5.75
85-89*	3075	54.73	87.54	297	4.12	5.86

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Venezuela

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	709	9.49	15.91	382	4.49	7.35
60-64	997	11.15	19.64	529	5.25	9.41
65-69	1407	13.79	21.12	744	6.36	10.03
70-74	1899	15.37	25.44	920	6.65	11.23
75-79	2471	16.36	23.58	1184	7.08	10.76
80-84*	2420	15.77	23.52	1188	6.97	10.28
85-89*	2150	16.14	21.97	1077	7.11	10.47

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Canada

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	9291	22.72	40.15	1535	3.67	5.86
60-64	12869	28.70	47.83	17969	4.18	7.20
65-69	17898	36.04	56.33	3033	5.59	10.29
70-74	24203	43.93	66.50	4772	7.70	14.60
75-79	30602	49.64	71.73	7636	10.93	20.47
80-84	38286	55.02	77.46	12029	15.00	27.18
85-89*	34809	56.18	76.43	15851	19.19	33.31

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

United States of America

Deaths from cancer of the trachea, bronchus and lung

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	130670	27.87	52.90	22732	4.30	7.47
60-64	175669	34.65	62.17	30462	5.22	9.94
65-69	228748	42.46	72.69	46830	7.38	14.72
70-74	286198	49.41	81.48	74003	10.70	22.02
75-79	341613	53.88	85.42	110866	14.46	29.00
80-84	394356	56.74	85.78	161195	18.88	35.09
85-89*	344231	57.01	83.74	167140	22.36	38.71

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Argentina

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	259	2.23	3.32	207	1.66	2.41
70-74*	285	2.47	3.55	178	1.42	2.11
75-79*	825	1.95	3.18	521	1.12	1.64
80-84	1151	1.53	2.26	799	0.94	1.26
85-89*	444	1.41	1.92	299	0.82	1.22

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Chile

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	105	1.62	2.55	83	1.18	2.12
70-74*	322	1.87	2.54	239	1.21	1.69
75-79*	349	1.73	2.29	252	1.06	1.53
80-84	312	1.38	1.71	247	0.92	1.21
85-89*	173	1.20	1.36	139	0.78	0.80

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Colombia

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	136	1.18	1.67	115	0.98	1.42
70-74*	134	0.97	1.06	145	1.02	1.48
75-79*	224	1.05	1.50	217	0.92	1.26
80-84						
85-89						

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Costa Rica

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	11	0.99	0.35	7	0.58	1.39
70-74*	17	0.43	0.37	16	0.47	0.38
75-79*	34	0.74	0.61	25	0.63	0.94
80-84	40	0.82	1.01	26	0.61	0.85
85-89*	50	1.37	1.91	39	0.98	1.03

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Cuba

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	134	2.07	2.18	110	1.75	2.43
70-74	290	1.42	1.67	200	1.05	1.41
75-79	332	1.43	1.81	227	1.04	1.24
80-84	318	1.21	1.44	164	0.67	0.66
85-89*	246	1.09	1.28	178	0.82	1.02

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Dominican Republic

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	16	0.83	1.64	7	0.51	0.64
70-74	25	0.39	0.47	36	0.62	1.40
75-79	46	0.59	0.68	38	0.53	0.82
80-84	59	0.73	0.65	37	0.46	0.67
85-89*	7	0.37	0.69	15	0.97	1.91

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Ecuador

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	22	0.73	0.72	25	0.72	1.20
70-74	79	0.90	0.87	90	1.02	1.18
75-79*	100	1.10	1.32	86	0.92	0.94
80-84*	108	1.40	1.47	90	1.18	1.50
85-89*	184	1.50	1.43	138	1.14	1.65

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Mexico

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	248	0.87	1.07	215	0.74	1.09
70-74	647	0.77	1.03	616	0.70	1.06
75-79*	540	0.94	1.13	414	0.69	0.81
80-84*	318	0.48	0.59	282	0.38	0.48
85-89*	210	0.45	0.62	176	0.32	0.42

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	6	0.60	0.45	8	0.70	0.88
70-74	19	0.58	0.75	8	0.23	0.13
75-79	20	0.59	0.65	14	0.40	0.62
80-84	22	0.52	0.29	19	0.52	0.43
85-89*	21	0.77	0.52	10	0.33	0.41

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	57	2.68	3.56	90	4.03	7.99
75-79	54	2.14	3.63	56	1.92	2.80
80-84*	21	0.95	1.23	24	0.99	0.92
85-89*	18	1.55	1.59	16	1.08	1.35

*Information is missing for one or more years.

Peru

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	65	0.78	0.82	63	0.78	0.76
70-74*	167	0.96	1.11	129	0.72	0.93
75-79*	81	0.72	0.83	74	0.66	0.85
80-84*	189	0.74	0.76	176	0.68	1.12
85-89						





*Information is missing for one or more years.

Puerto Rico

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	11	0.95	0.78	10	1.00	1.91
70-74	89	1.38	1.79	68	1.04	1.47
75-79*	65	1.18	2.03	51	0.87	1.20
80-84	52	0.64	0.99	18	0.16	0.09
85-89*	29	0.60	1.16	6	0.11	0.13

*Information is missing for one or more years.

**Uruguay**

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	53	1.78	3.59	34	1.05	1.38
70-74	112	1.48	2.25	84	0.99	1.28
75-79*	82	1.36	1.71	65	0.89	0.68
80-84	79	0.93	1.07	67	0.68	0.92
85-89*	51	1.01	1.10	32	0.50	0.70

*Information is missing for one or more years.

Venezuela

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	84	1.34	1.73	80	1.29	2.04
70-74	206	1.14	1.57	178	1.04	1.63
75-79	195	0.88	1.08	168	0.74	0.87
80-84*	224	0.98	1.10	172	0.75	0.98
85-89*	117	0.57	0.67	109	0.52	0.71

*Information is missing for one or more years.

Canada

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	113	1.03	0.78	80	0.67	0.77
70-74	584	1.01	1.02	330	0.53	0.59
75-79	503	0.80	0.86	334	0.47	0.43
80-84	424	0.62	0.59	290	0.38	0.35
85-89*	343	0.60	0.61	229	0.35	0.30





*Information is missing for one or more years.

United States of America

Deaths from cancer of the bone

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	2087	0.96	1.09	1507	0.59	0.64
70-74	5282	0.92	1.00	3900	0.57	0.63
75-79	4651	0.75	0.81	3423	0.45	0.49
80-84	3204	0.49	0.51	2359	0.30	0.27
85-89*	2379	0.44	0.43	1875	0.28	0.27

*Information is missing for one or more years.



Top panel: total deaths, middle panel: age-specific death rate for males 30-34 to 70-74; bottom panel: age-specific death rate for females 30-34 to 70-74.

Top panel: total deaths, middle panel: age-specific death rate for males 30-34 to 70-74; bottom panel: age-specific death rate for females 30-34 to 70-74.

Top panel: total deaths, middle panel: age-specific death rate for males 30-34 to 70-74; bottom panel: age-specific death rate for females 30-34 to 70-74.




Top panel: total deaths, middle panel: age-specific death rate for males 30-34 to 70-74; bottom panel: age-specific death rate for females 30-34 to 70-74.

Argentina

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	768	1.72	2.40	490	1.00	1.23
70-74	219	1.90	2.49	128	0.96	1.26
75-79	563	1.31	1.81	354	0.72	0.93
80-84	860	1.13	1.89	542	0.62	0.98
85-89	211	0.66	1.12	171	0.46	0.80

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Chile

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	101	0.83	1.02	86	0.59	0.66
60-64	121	0.88	1.04	116	0.72	0.87
65-69	180	1.33	1.74	161	0.96	1.10
70-74	162	1.03	1.35	177	0.91	1.08
75-79	215	1.20	1.54	225	0.95	1.19
80-84	156	0.73	1.21	135	0.51	0.82
85-89	196	0.73	1.30	112	0.66	1.15

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Colombia

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	170	0.98	1.23	207	1.06	1.22
60-64	332	1.16	1.44	276	1.22	1.35
65-69	308	1.37	1.55	311	1.19	1.54
70-74	162	1.46	1.97	137	1.12	1.12
75-79	218	1.27	1.47	210	1.06	1.22

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Costa Rica

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	33	2.49	2.72	16	0.94	1.29
60-64	38	1.78	2.14	21	0.95	1.20
65-69	32	1.25	1.97	28	1.12	1.37
70-74	27	0.92	1.16	14	0.43	0.71
75-79	21	0.62	0.92	9	0.23	0.41
80-84	23	0.73	1.18	17	0.44	0.73

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	58	2.02	1.12	38	1.35	1.19
60-64	135	2.21	1.92	57	0.96	0.33
65-69	274	1.37	1.57	143	0.76	0.64
70-74	266	1.14	1.21	169	0.78	0.70
75-79	92	0.37	0.62	77	0.33	0.55
80-84	74	0.35	0.62	70	0.33	0.49

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Dominican Republic

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	19	0.39	0.51	16	0.34	0.29
70-74	17	0.30	0.46	14	0.24	0.46
75-79	24	0.33	0.62	18	0.27	0.43
80-84	5	0.07	0.19	3	0.03	0.04
85-89	0	0.00	0.00	0	0.00	0.00

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Ecuador

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	20	0.56	0.24	38	1.04	0.77
65-69	32	0.51	0.52	52	0.72	0.72
70-74	50	0.65	0.79	61	0.76	0.67
75-79	39	0.51	0.56	42	0.51	0.60
80-84	34	0.51	0.65	24	0.32	0.22
85-89	69	0.68	0.56	76	0.66	0.87

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Mexico

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	152	0.78	0.97	137	0.63	0.70
60-64	359	0.68	1.10	374	0.70	0.82
65-69	386	0.66	0.97	437	0.69	0.73
70-74	537	0.74	0.98	564	0.71	0.89
75-79	371	0.78	1.06	324	0.61	0.80
80-84	174	0.30	0.51	163	0.25	0.38
85-89	168	0.39	0.56	137	0.27	0.35

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	12	0.76	1.08	12	0.88	1.50
60-64	17	1.01	1.43	10	0.52	0.20
65-69	28	1.48	2.02	14	0.77	1.05
70-74	18	0.68	1.15	6	0.26	0.13
75-79	21	0.79	0.72	15	0.58	0.93
80-84	18	0.52	0.99	9	0.27	0.50
85-89*	13	0.58	1.14	15	0.60	0.84

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	21	1.05	0.98	29	1.37	2.25
75-79	20	0.89	1.26	22	0.83	0.80
80-84*	4	0.20	0.12	5	0.17	0.00
85-89*	8	0.71	0.30	6	0.32	0.70

*Information is missing for one or more years.

Peru

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	101	0.76	1.23	107	0.72	0.72
70-74*	142	1.01	1.30	105	0.63	0.85
75-79*	66	0.73	0.85	40	0.41	0.63
80-84*	93	0.47	0.73	62	0.28	0.38
85-89*						




*Information is missing for one or more years.

Puerto Rico




Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	44	1.19	1.72	50	1.11	0.87
65-69*	22	1.04	1.15	9	0.37	0.50
70-74	48	0.75	0.91	41	0.56	0.35
75-79*	29	0.53	0.86	27	0.42	0.34
80-84	25	0.29	0.33	20	0.22	0.21
85-89*	23	0.43	0.73	16	0.23	0.19




*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	106	1.69	1.83	69	0.91	1.37
60-64*	74	1.81	2.23	54	1.09	0.84
65-69	105	1.45	1.89	86	1.01	1.23
70-74	140	1.82	3.05	87	0.98	1.24
75-79*	123	1.88	2.95	61	0.72	0.89
80-84	71	0.80	1.40	49	0.43	0.60
85-89*	49	0.88	1.28	32	0.52	1.00

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Venezuela

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	158	2.14	2.94	177	1.98	2.82
60-64	192	2.16	2.97	214	2.07	3.03
65-69*	171	2.10	2.49	188	1.86	2.44
70-74	200	1.56	2.13	216	1.45	1.80
75-79	247	1.63	1.96	227	1.32	1.43
80-84*	133	0.83	1.00	98	0.54	0.78
85-89*	55	0.38	0.46	52	0.33	0.57

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Canada

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	760	1.73	2.22	488	1.09	1.43
60-64	856	1.74	2.17	616	1.22	1.79
65-69	991	1.87	2.63	743	1.31	2.11
70-74	985	1.72	2.59	794	1.23	2.03
75-79	1089	1.75	2.93	853	1.19	2.06
80-84	1420	2.05	3.54	1029	1.28	2.35
85-89*	1376	2.23	3.61	944	1.29	2.39

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

United States of America

Deaths from cancer of the skin, including melanoma

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	10964	2.29	3.30	7989	1.48	2.17
60-64	12466	2.41	3.70	8665	1.46	2.30
65-69	13395	2.46	4.05	9423	1.47	2.49
70-74	13726	2.40	4.01	10139	1.43	2.53
75-79	17049	2.76	4.87	12055	1.54	2.71
80-84	20587	3.06	5.41	13117	1.53	2.75
85-89*	19928	3.42	6.09	11617	1.55	2.80

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	197	0.43	0.81	9138	19.09	39.56
70-74*	61	0.51	0.98	2373	18.61	37.96
75-79*	148	0.35	0.69	9228	19.71	39.07
80-84				17062	19.76	38.40
85-89*				7373	20.21	38.71

*Information is missing for one or more years.

Chile

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	15	0.13	0.25	1184	8.83	18.75
60-64	24	0.19	0.27	1365	9.17	18.77
65-69	13	0.09	0.15	1839	11.24	22.57
70-74	36	0.24	0.40	2264	12.18	24.75
75-79	61	0.34	0.57	2559	11.63	22.54
80-84				3116	12.34	23.47
85-89*				2129	12.80	23.47

*Information is missing for one or more years.

Colombia

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	11	0.06	0.05	802	4.13	8.38
60-64	11	0.06	0.09	1194	5.09	11.16
65-69	18	0.08	0.11	1525	5.75	12.16
70-74*	9	0.09	0.14	799	6.35	13.10
75-79*	15	0.08	0.13	1272	6.21	12.93
80-84						
85-89						




*Information is missing for one or more years.

Costa Rica




Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	2	0.11	0.00	134	8.15	16.13
65-69	0	0.00	0.00	183	8.48	16.62
70-74	2	0.09	0.30	195	7.57	15.51
75-79	6	0.21	0.43	279	9.18	17.77
80-84				405	11.22	20.33
85-89*				455	13.11	23.64




*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	6	0.21	0.20	346	12.20	26.36
65-69*	8	0.13	0.10	834	13.41	26.42
70-74	41	0.20	0.29	2353	13.26	27.12
75-79	19	0.09	0.16	3057	15.34	31.34
80-84				3439	15.14	28.77
85-89*				3104	15.13	28.44

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Dominican Republic

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	4	0.07	0.04	143	2.89	6.51
70-74	3	0.05	0.17	163	2.88	6.65
75-79	8	0.11	0.21	234	3.58	7.71
80-84				405	5.38	10.22
85-89*				82	4.83	10.51

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Ecuador

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	0	0.00	0.00	121	2.98	5.93
65-69	1	0.01	0.00	247	3.49	6.17
70-74	2	0.03	0.05	314	3.90	7.00
75-79				349	4.12	8.29
80-84*				354	4.84	9.38
85-89*				632	5.53	11.32

*Information is missing for one or more years.






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Mexico

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	0	0.00	0.00	780	3.77	7.80
60-64	9	0.02	0.03	2376	4.31	8.51
65-69	69	0.12	0.22	2783	4.39	8.77
70-74	118	0.16	0.29	4011	5.22	11.17
75-79	83	0.18	0.27	2929	5.64	11.59
80-84*				4050	6.37	12.72
85-89*				3440	7.05	14.33

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	79	5.25	10.89
60-64	3	0.17	0.34	125	7.39	15.57
65-69	2	0.11	0.19	145	7.07	14.16
70-74	0	0.00	0.00	152	6.39	14.62
75-79	2	0.07	0.10	200	7.31	14.21
80-84				252	7.59	14.12
85-89*				189	8.66	15.94

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	1	0.05	0.17	182	8.82	18.27
75-79	2	0.10	0.32	291	11.61	22.59
80-84*				189	8.96	18.39
85-89*				135	9.43	19.16

*Information is missing for one or more years.

Peru

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	11	0.09	0.13	616	4.26	8.74
70-74*	7	0.04	0.06	738	4.44	9.25
75-79*	8	0.08	0.11	462	4.59	10.53
80-84*				1093	4.85	9.97
85-89						




*Information is missing for one or more years.

Puerto Rico




Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	4	0.12	0.37	264	6.80	14.27
65-69*	1	0.05	0.16	175	8.36	17.75
70-74	10	0.15	0.17	571	9.24	19.45
75-79*	3	0.06	0.20	524	9.65	20.59
80-84				906	10.88	22.47
85-89*				709	13.23	29.24




*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	10	0.15	0.24	1428	20.84	43.95
60-64*	11	0.27	0.33	945	21.36	44.37
65-69	9	0.12	0.12	1799	22.81	46.39
70-74	15	0.19	0.21	2136	25.34	51.38
75-79*	16	0.24	0.42	1808	24.12	48.49
80-84				2525	24.87	49.51
85-89*				1615	25.21	49.65

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Venezuela

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1	0.02	0.00	622	6.79	12.41
60-64	2	0.02	0.07	875	7.94	15.72
65-69	16	0.17	0.14	1130	8.71	17.95
70-74	13	0.11	0.13	1401	9.31	19.12
75-79	24	0.15	0.27	1736	9.78	19.73
80-84*				1531	8.32	17.24
85-89*				1487	9.24	19.00

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Canada

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	112	0.26	0.46	9395	22.73	45.51
60-64	85	0.18	0.28	11152	23.59	48.13
65-69	94	0.18	0.17	12843	23.88	49.95
70-74	118	0.21	0.33	14724	23.94	49.00
75-79	120	0.19	0.23	16265	23.34	47.10
80-84				18514	23.19	45.12
85-89*				17366	24.17	46.65

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

United States of America

Deaths from cancer of the breast

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1134	0.23	0.35	111552	21.53	43.74
60-64	1159	0.22	0.34	123874	21.54	44.44
65-69	1222	0.22	0.33	139740	22.27	46.41
70-74	1456	0.24	0.36	154765	22.44	46.35
75-79	1422	0.17	0.25	168487	22.03	45.05
80-84				186989	22.19	44.18
85-89*				163703	22.51	44.25

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	0	0.00	0.00	6125	12.78	24.57
70-74*	0	0.00	0.00	1669	13.01	24.86
75-79*	0	0.00	0.00	5448	11.77	22.34
80-84	0	0.00	0.00	9615	11.44	22.40
85-89*	0	0.00	0.00	4093	11.64	23.43

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Chile

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	2836	20.74	44.26
60-64	0	0.00	0.00	3052	19.94	42.90
65-69	0	0.00	0.00	3575	21.42	45.33
70-74	0	0.00	0.00	4213	22.32	45.87
75-79	0	0.00	0.00	4174	18.84	38.57
80-84	0	0.00	0.00	4144	16.29	32.11
85-89*	0	0.00	0.00	2678	15.99	30.70

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Colombia

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	3398	16.93	36.23
60-64	0	0.00	0.00	4556	19.30	41.10
65-69	0	0.00	0.00	5420	20.13	41.39
70-74*	0	0.00	0.00	2350	18.49	37.81
75-79*	0	0.00	0.00	3697	18.14	35.37
80-84						
85-89*						

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Costa Rica

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	0	0.00	0.00	433	28.39	59.51
65-69	0	0.00	0.00	542	24.89	51.00
70-74	0	0.00	0.00	452	17.46	34.00
75-79	0	0.00	0.00	501	16.26	26.96
80-84	0	0.00	0.00	520	14.38	23.66
85-89*	0	0.00	0.00	570	15.98	27.67

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	0	0.00	0.00	538	19.35	36.37
65-69*	0	0.00	0.00	1055	17.39	29.88
70-74	0	0.00	0.00	2812	15.75	28.30
75-79	0	0.00	0.00	3015	14.88	25.78
80-84	0	0.00	0.00	3227	13.99	24.85
85-89*	0	0.00	0.00	2792	13.60	25.45

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Dominican Republic

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	0	0.00	0.00	384	7.89	16.17
70-74	0	0.00	0.00	478	8.35	18.33
75-79	0	0.00	0.00	559	8.19	16.24
80-84	0	0.00	0.00	908	11.92	23.65
85-89*	0	0.00	0.00	257	15.53	25.53

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Ecuador

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	0	0.00	0.00	703	17.42	33.62
65-69	0	0.00	0.00	1445	19.30	35.96
70-74	0	0.00	0.00	1735	20.81	38.11
75-79*	0	0.00	0.00	1594	18.55	35.09
80-84*	0	0.00	0.00	1481	20.32	36.62
85-89*	0	0.00	0.00	2367	20.82	37.21

*Information is missing for one or more years.





Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Mexico

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	0	0.00	0.00	4161	20.19	40.86
60-64	0	0.00	0.00	11408	20.45	42.12
65-69	0	0.00	0.00	12791	20.03	40.78
70-74	0	0.00	0.00	14768	19.09	39.15
75-79*	0	0.00	0.00	9539	18.34	36.95
80-84*	0	0.00	0.00	11269	17.83	35.11
85-89*	0	0.00	0.00	8752	17.91	34.63

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	273	17.64	34.45
60-64	0	0.00	0.00	300	16.57	36.04
65-69	0	0.00	0.00	319	15.40	27.83
70-74	0	0.00	0.00	350	14.75	29.21
75-79	0	0.00	0.00	406	15.10	28.88
80-84	0	0.00	0.00	444	13.42	26.05
85-89*	0	0.00	0.00	286	12.58	23.02

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	735	34.98	74.98
60-64	0	0.00	0.00	766	29.82	62.76
65-69	0	0.00	0.00	476	22.19	48.83
70-74	0	0.00	0.00	325	22.09	43.70

*Information is missing for one or more years.

Peru

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	2400	16.07	33.35
60-64	0	0.00	0.00	2920	17.70	35.83
65-69*	0	0.00	0.00	1414	14.10	28.10
70-74*	0	0.00	0.00	3072	13.58	26.54






*Information is missing for one or more years.

Puerto Rico


Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	731	18.76	36.63
60-64*	0	0.00	0.00	304	14.13	25.26
65-69*	0	0.00	0.00	841	13.00	22.48
70-74*	0	0.00	0.00	509	9.02	16.16
75-79*	0	0.00	0.00	622	7.08	12.45
80-84	0	0.00	0.00	367	6.34	10.95


*Information is missing for one or more years.




Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	1122	16.27	31.36
60-64*	0	0.00	0.00	746	16.82	34.25
65-69	0	0.00	0.00	1351	17.02	33.02
70-74	0	0.00	0.00	1255	14.72	27.22
75-79*	0	0.00	0.00	990	13.47	28.01
80-84	0	0.00	0.00	1226	12.19	22.49
85-89*	0	0.00	0.00	691	11.13	20.60

*Information is missing for one or more years.

Venezuela

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	2848	30.30	63.28
60-64	0	0.00	0.00	3230	28.67	61.46
65-69	0	0.00	0.00	3542	26.46	55.23
70-74	0	0.00	0.00	3924	25.91	52.69
75-79	0	0.00	0.00	4105	22.61	43.93
80-84*	0	0.00	0.00	3576	19.41	37.16
85-89*	0	0.00	0.00	2826	17.29	33.55

*Information is missing for one or more years.

Canada

Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	5431	13.18	26.33
60-64	0	0.00	0.00	5333	11.33	22.74
65-69	0	0.00	0.00	5530	10.22	19.20
70-74	0	0.00	0.00	5289	8.48	15.47
75-79	0	0.00	0.00	4812	6.66	11.38
80-84	0	0.00	0.00	4857	5.73	8.97
85-89*	0	0.00	0.00	3826	5.02	7.72






*Information is missing for one or more years.

United States of America


Deaths from cancer of the uterus (cervix and corpus)

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	0	0.00	0.00	74136	14.45	29.19
60-64	0	0.00	0.00	71222	12.49	24.60
65-69	0	0.00	0.00	65611	10.44	19.84
70-74	0	0.00	0.00	59746	8.54	15.95
75-79	0	0.00	0.00	54970	6.92	12.53
80-84	0	0.00	0.00	53403	6.03	10.44
85-89*	0	0.00	0.00	41532	5.41	9.16


*Information is missing for one or more years.




Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina*Deaths from cancer of the prostate*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	5403	12.46	7.18	0	0.00	0.00
70-74*	1463	13.38	5.91	0	0.00	0.00
75-79*	5108	11.82	5.91	0	0.00	0.00
80-84	8850	11.08	5.63	0	0.00	0.00
85-89*	3888	11.57	6.06	0	0.00	0.00

*Information is missing for one or more years.

Chile*Deaths from cancer of the prostate*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	755	7.34	5.35	0	0.00	0.00
60-64	956	8.33	4.94	0	0.00	0.00
65-69	1269	10.18	5.62	0	0.00	0.00
70-74	1609	10.99	5.45	0	0.00	0.00
75-79	1876	11.14	4.95	0	0.00	0.00
80-84	2230	11.24	4.68	0	0.00	0.00
85-89*	1631	12.62	6.09	0	0.00	0.00

*Information is missing for one or more years.

Colombia*Deaths from cancer of the prostate*









Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	800	5.30	3.56	0	0.00	0.00
60-64	851	4.76	3.29	0	0.00	0.00
65-69	1150	5.55	3.38	0	0.00	0.00
70-74*	647	6.72	3.69	0	0.00	0.00
75-79*	1167	7.45	4.01	0	0.00	0.00
80-84						
85-89						

*Information is missing for one or more years.

Costa Rica*Deaths from cancer of the prostate*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	105	7.91	4.16	0	0.00	0.00
65-69	195	10.10	4.47	0	0.00	0.00
70-74	192	8.22	3.41	0	0.00	0.00
75-79	263	9.76	2.97	0	0.00	0.00
80-84	402	12.16	4.60	0	0.00	0.00
85-89*	442	15.99	5.50	0	0.00	0.00

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Cuba*Deaths from cancer of the prostate*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	397	14.74	7.08	0	0.00	0.00
65-69*	977	16.48	5.56	0	0.00	0.00
70-74	2848	13.90	5.61	0	0.00	0.00
75-79	3535	14.68	6.07	0	0.00	0.00
80-84	4289	14.85	6.23	0	0.00	0.00
85-89*	4329	16.65	7.26	0	0.00	0.00

*Information is missing for one or more years.

Dominican Republic*Deaths from cancer of the prostate*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	163	3.53	2.05	0	0.00	0.00
70-74	297	5.52	3.19	0	0.00	0.00
75-79	445	7.00	4.60	0	0.00	0.00
80-84	704	10.07	5.34	0	0.00	0.00
85-89*	164	11.43	3.98	0	0.00	0.00

*Information is missing for one or more years.

Ecuador*Deaths from cancer of the prostate*









Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	108	3.32	1.42	0	0.00	0.00
65-69	283	5.07	1.89	0	0.00	0.00
70-74	380	5.56	2.26	0	0.00	0.00
75-79*	477	6.83	2.40	0	0.00	0.00
80-84*	533	9.26	3.15	0	0.00	0.00
85-89*	899	9.59	3.00	0	0.00	0.00

*Information is missing for one or more years.

Mexico*Deaths from cancer of the prostate*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	672	3.67	2.54	0	0.00	0.00
60-64	1844	4.02	2.92	0	0.00	0.00
65-69	2470	4.54	2.78	0	0.00	0.00
70-74	3807	5.47	3.19	0	0.00	0.00
75-79*	2718	5.99	3.19	0	0.00	0.00
80-84*	3927	7.33	3.71	0	0.00	0.00
85-89*	3598	8.98	3.97	0	0.00	0.00

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 78-74; bottom panel, age-specific death rate for females 30-34 to 78-74.

Panama

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	84	6.00	4.30	0	0.00	0.00
60-64	122	7.85	3.82	0	0.00	0.00
65-69	156	8.75	5.36	0	0.00	0.00
70-74	188	8.34	2.94	0	0.00	0.00
75-79	250	10.11	4.37	0	0.00	0.00
80-84	356	11.31	4.87	0	0.00	0.00
85-89*	264	11.93	4.25	0	0.00	0.00



*Information is missing for one or more years.

**Paraguay (reporting)**

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	171	9.72	5.45	0	0.00	0.00
75-79	194	8.98	5.10	0	0.00	0.00
80-84*	138	7.00	3.45	0	0.00	0.00
85-89*	44	3.78	2.06	0	0.00	0.00



*Information is missing for one or more years.

**Peru**

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	499	4.43	1.86	0	0.00	0.00
65-69*	624	5.01	2.49	0	0.00	0.00
70-74*	406	5.01	2.44	0	0.00	0.00
75-79*	1198	6.76	2.47	0	0.00	0.00
80-84*						
85-89*						



*Information is missing for one or more years.

**Puerto Rico**

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	394	9.88	6.23	0	0.00	0.00
65-69*	260	11.40	4.80	0	0.00	0.00
70-74	864	12.59	5.54	0	0.00	0.00
75-79*	851	13.75	6.46	0	0.00	0.00
80-84	1405	14.33	6.69	0	0.00	0.00
85-89*	1010	16.17	7.75	0	0.00	0.00

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	908	14.38	8.09	0	0.00	0.00
60-64*	607	14.93	8.98	0	0.00	0.00
65-69	1206	16.72	7.66	0	0.00	0.00
70-74	1355	17.41	7.86	0	0.00	0.00
75-79*	1172	16.34	8.59	0	0.00	0.00
80-84	1597	16.19	7.76	0	0.00	0.00
85-89*	1166	18.30	7.84	0	0.00	0.00



*Information is missing for one or more years.

**Venezuela**

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	426	7.05	5.57	0	0.00	0.00
60-64	613	8.53	6.35	0	0.00	0.00
65-69	912	10.77	7.44	0	0.00	0.00
70-74	1061	9.71	6.78	0	0.00	0.00
75-79	1407	10.50	6.38	0	0.00	0.00
80-84*	1516	10.85	5.98	0	0.00	0.00
85-89*	1461	11.75	6.28	0	0.00	0.00



*Information is missing for one or more years.

**Canada**

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	5785	12.53	5.95	0	0.00	0.00
60-64	6739	12.85	5.68	0	0.00	0.00
65-69	7418	12.96	5.57	0	0.00	0.00
70-74	8368	13.95	5.65	0	0.00	0.00
75-79	9544	14.09	6.05	0	0.00	0.00
80-84	11088	14.19	5.97	0	0.00	0.00
85-89*	11258	15.96	7.37	0	0.00	0.00



*Information is missing for one or more years.

**United States of America**

Deaths from cancer of the prostate

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	69883	13.69	7.44	0	0.00	0.00
60-64	75560	13.27	6.91	0	0.00	0.00
65-69	81881	13.31	6.89	0	0.00	0.00
70-74	91266	13.92	7.01	0	0.00	0.00
75-79	104483	14.35	6.93	0	0.00	0.00
80-84	120638	14.85	7.10	0	0.00	0.00
85-89*	110051	15.28	7.21	0	0.00	0.00

*Information is missing for one or more years.



Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.




Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	3044	6.77	7.90	2108	4.50	5.32
70-74*	664	5.83	5.93	510	4.14	4.65
75-79*	2244	5.35	5.71	1650	3.71	4.00
80-84	4028	5.39	5.48	3018	3.64	3.88
85-89*	1587	5.00	5.15	1239	3.42	3.81

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Chile
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	584	3.84	4.47	459	2.78	3.68
60-64	760	4.40	4.57	696	3.81	4.50
65-69	947	4.97	5.63	771	3.86	3.94
70-74	851	4.04	4.50	808	3.54	4.03
75-79	926	4.02	3.76	854	3.36	4.06
80-84	1048	4.20	4.12	902	3.26	3.69
85-89*	636	3.98	4.03	628	3.52	3.78

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Colombia
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	591	2.24	2.60	475	1.78	2.16
60-64	1075	3.57	3.95	994	3.19	3.69
65-69	1403	3.65	4.55	1181	3.04	3.78
70-74*	643	3.56	4.01	553	2.77	2.70
75-79*	1068	3.53	3.63	949	3.03	3.57
80-84						
85-89						

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Costa Rica
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	152	7.73	9.14	79	3.88	4.71
65-69	151	4.68	4.26	127	3.90	3.98
70-74	165	4.06	4.15	172	4.32	5.42
75-79	230	5.67	5.69	207	4.91	5.34
80-84	279	5.58	5.05	235	4.74	5.34
85-89*	256	6.02	4.91	210	4.62	4.32

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	240	7.32	10.20	167	5.23	5.42
65-69*	411	5.57	5.52	284	3.92	4.96
70-74	1045	4.79	5.01	714	3.48	3.93
75-79	1081	4.60	5.04	805	3.66	4.41
80-84	1227	4.83	5.03	885	3.67	4.11
85-89*	1032	4.75	5.17	803	3.84	4.60

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Dominican Republic
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	83	1.35	1.94	70	1.23	2.00
65-69	138	1.51	1.80	101	1.15	1.65
70-74	176	1.69	1.63	148	1.51	1.69
75-79	253	2.25	2.68	178	1.67	2.17
80-84	58	2.49	3.76	40	1.83	2.59
85-89*						

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Ecuador
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	140	2.80	3.30	118	2.46	2.71
65-69	254	2.64	2.91	215	2.10	2.70
70-74	285	2.33	2.70	281	2.30	2.23
75-79*	309	2.44	2.24	252	2.06	2.88
80-84*	350	3.25	3.27	283	2.75	2.97
85-89*	572	3.60	3.70	489	3.10	3.71

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64






Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Mexico
Deaths from leukaemias

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59*	605	2.28	2.78	502	1.81	2.11
60-64	1932	2.57	3.12	1675	2.29	2.86
65-69	2401	2.58	2.68	2184	2.38	2.81
70-74	2877	2.41	2.57	2560	2.13	2.52
75-79*	2168	2.62	2.64	1888	2.37	2.78
80-84*	3103	3.34	3.58	2683	2.88	3.41
85-89*	2425	3.61	3.76	2043	3.02	3.65

*Information is missing for one or more years.

— Males, all ages
 - - Males, 35-64
 — Females, all ages
 - - Females, 35-64







Top panel, total deaths, middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	40	1.99	2.20	36	2.05	2.77
60-64	77	3.49	4.40	57	2.36	2.73
65-69	113	4.44	4.95	74	2.66	4.03
70-74	117	3.33	4.45	68	1.91	2.56
75-79	134	3.42	3.50	104	2.84	3.68
80-84	174	3.96	3.66	116	2.58	3.22
85-89*	127	4.34	4.49	87	3.04	2.64

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Paraguay (reporting)*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	132	4.85	5.15	104	3.67	4.60
75-79	151	4.36	3.95	114	3.10	2.58
80-84*	128	4.86	5.64	90	3.11	3.14
85-89*	68	3.56	3.16	73	3.56	3.01

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Peru*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	523	2.59	2.99	463	2.22	2.52
70-74*	505	2.12	2.26	464	1.95	2.29
75-79*	314	2.28	2.34	246	1.73	2.12
80-84*	653	2.14	2.00	572	1.79	1.96
85-89						

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Puerto Rico*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	262	6.07	8.06	212	4.62	5.23
65-69*	119	4.97	4.57	104	4.21	6.12
70-74	345	5.27	6.80	259	3.72	4.75
75-79*	298	5.29	6.54	258	4.25	5.79
80-84	375	4.43	4.45	328	3.68	4.70
85-89*	218	4.06	4.83	219	3.76	4.94

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	386	6.25	6.93	266	4.19	4.76
60-64*	335	8.35	10.36	258	5.98	6.52
65-69	545	7.80	7.71	403	5.26	6.59
70-74	476	6.31	7.04	350	4.39	4.97
75-79*	340	5.54	5.60	294	4.22	5.33
80-84	444	5.33	5.57	393	4.05	4.72
85-89*	317	6.02	5.84	250	4.07	4.94

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Venezuela*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	405	3.02	3.22	359	2.73	3.30
60-64	759	5.42	6.70	709	4.87	6.09
65-69	846	4.82	4.90	733	3.99	4.60
70-74	860	3.71	4.23	713	3.05	3.31
75-79	1075	4.06	4.46	871	3.15	3.34
80-84*	1001	3.95	4.37	835	3.20	3.59
85-89*	534	3.64	3.40	510	3.49	3.90

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Canada*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	2933	6.87	7.80	2114	5.00	5.75
60-64	4112	8.53	8.88	2895	5.92	6.77
65-69	4491	8.43	9.12	3061	5.53	6.45
70-74	3922	6.94	6.61	2804	4.52	4.82
75-79	4324	6.99	6.92	3031	4.29	4.74
80-84	4561	6.58	6.90	3446	4.15	4.72
85-89*	4072	6.63	6.66	3084	4.04	4.40

*Information is missing for one or more years.







Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

United States of America*Deaths from leukaemias*

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	38156	8.27	9.40	27775	5.57	6.60
60-64	59452	9.44	11.61	36234	6.37	7.72
65-69	48751	8.99	10.28	35939	5.68	6.69
70-74	41011	7.09	7.29	31405	4.44	4.80
75-79	43125	6.80	7.14	33393	4.12	4.66
80-84	46103	6.63	7.09	37307	4.07	4.77
85-89*	38118	6.27	6.74	31427	3.89	4.38

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Argentina

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	81824	183.85	263.10	56255	116.96	174.36
70-74	21609	191.35	258.63	15174	117.43	173.66
75-79	70220	164.31	228.59	51450	106.91	156.40
80-84	121801	159.32	229.37	92229	102.47	154.34
85-89	48950	154.10	218.44	38201	99.73	152.62

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Chile

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	17091	149.55	219.51	18361	137.94	225.91
60-64	19705	154.93	216.39	21445	143.44	224.73
65-69	22714	163.91	217.76	24318	148.44	220.53
70-74	25345	162.34	209.70	26620	140.41	209.29
75-79	27259	152.05	187.54	27701	121.86	180.17
80-84	30186	146.99	175.52	30158	115.85	167.50
85-89	19454	145.97	170.48	20172	117.83	168.09

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Colombia

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	15163	86.91	124.09	20795	106.38	179.75
60-64	19262	92.93	131.71	25936	111.87	188.00
65-69	23672	99.48	134.50	29544	111.48	181.04
70-74	11160	100.17	130.66	13470	107.88	171.42
75-79	18333	101.57	130.97	21476	106.82	165.55

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Costa Rica

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	2136	147.10	194.16	1995	133.51	210.11
65-69	3121	148.72	186.93	2891	132.79	197.91
70-74	3196	127.16	158.20	2947	113.27	162.77
75-79	3915	136.02	150.36	3339	107.19	143.24
80-84	4862	139.99	145.59	3874	105.39	136.08
85-89	4763	156.69	147.08	4085	114.31	146.38

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Cuba

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	4473	157.48	191.61	3126	113.36	166.01
65-69	9811	157.66	161.21	6696	111.14	151.81
70-74	27113	134.95	152.24	18041	99.46	147.54
75-79	29684	129.56	144.40	20227	98.29	143.59
80-84	32990	127.39	141.79	22896	96.98	142.04
85-89	29376	129.21	148.91	20814	97.90	146.10

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Dominican Republic

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69	1839	35.72	47.87	1956	39.52	68.47
70-74	2412	40.27	54.15	2301	38.95	69.57
75-79	2923	41.80	56.17	2893	41.99	70.42
80-84	4159	54.62	64.99	4203	54.37	88.39
85-89	962	60.80	66.48	928	54.79	85.07

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Ecuador

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64	2123	53.68	66.23	2849	71.63	104.81
65-69	4634	69.92	77.73	5664	77.10	111.73
70-74	5494	71.77	80.74	6769	80.50	115.52
75-79	5499	70.63	77.71	6624	76.54	111.58
80-84	5663	86.13	87.97	6309	86.18	119.99
85-89	9011	86.77	89.37	10361	89.55	123.26

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Mexico

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	9733	49.18	66.50	15735	76.40	130.07
60-64	28728	55.38	78.12	43895	79.86	134.41
65-69	35389	58.91	77.51	50327	78.98	129.40
70-74	43546	58.19	76.03	59578	75.43	126.45
75-79	31713	64.37	80.01	40951	77.20	123.64
80-84	42105	73.27	87.14	51087	78.74	123.58
85-89	33498	77.87	87.78	39862	80.02	123.40

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Panama

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	1144	74.66	94.43	1222	78.45	134.68
60-64	1372	78.03	100.27	1453	82.68	138.11
65-69	1577	79.44	96.53	1511	72.69	115.73
70-74	1910	78.53	101.88	1792	74.53	119.99
75-79	2398	88.51	108.75	2180	79.78	127.46
80-84	2980	89.54	105.97	2559	75.49	112.58
85-89*	2007	89.13	103.89	1731	76.81	107.15

*Information is missing for one or more years.

Paraguay (reporting)

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69						
70-74	1795	96.36	131.71	2355	108.98	194.74
75-79	1930	84.36	110.91	2605	97.44	168.33
80-84*	1333	66.72	83.52	1615	72.45	129.09
85-89*	1006	78.20	91.60	1124	75.46	115.97

*Information is missing for one or more years.

Peru

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64						
65-69*	8623	65.32	85.46	10480	71.12	112.89
70-74*	9575	65.50	80.51	11681	71.12	111.47
75-79*	5279	57.31	68.83	6306	62.81	98.83
80-84*	12197	60.84	64.82	14657	64.64	94.48
85-89						




*Information is missing for one or more years.

Puerto Rico




Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59						
60-64*	4978	131.04	183.63	3826	95.87	147.79
65-69*	2823	131.07	169.67	1973	89.50	133.96
70-74	7998	127.50	164.45	5734	86.90	129.06
75-79*	7184	128.80	161.21	4952	84.32	123.82
80-84	10164	120.50	150.06	7146	78.96	112.97
85-89*	6698	124.03	162.23	4629	76.81	107.45




*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Uruguay

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	11727	184.84	266.14	9443	132.36	199.10
60-64*	8052	196.59	271.29	6382	138.25	202.62
65-69	15145	210.12	276.25	11190	136.18	199.79
70-74	16365	211.57	282.07	12047	136.42	192.31
75-79*	13682	203.66	277.11	9992	124.06	185.67
80-84	17922	198.18	274.04	13143	118.86	176.31
85-89*	11717	204.14	271.54	8373	118.58	168.65

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Venezuela

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	7123	90.74	133.19	10112	110.53	189.30
60-64	9536	102.91	151.51	12341	112.82	193.62
65-69	11949	110.89	146.73	14522	111.20	183.87
70-74	14056	106.70	145.26	16130	106.91	176.97
75-79	16708	104.32	132.72	18648	103.80	159.86
80-84*	15776	97.71	121.57	16855	91.84	139.14
85-89*	8455	90.43	107.07	8951	83.52	124.39

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

Canada

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	58480	136.34	175.90	49806	117.34	184.37
60-64	67823	142.21	180.87	55886	114.37	180.64
65-69	77804	149.74	189.03	62919	112.01	177.33
70-74	90097	158.92	199.05	71979	111.16	176.48
75-79	102043	162.46	202.50	80430	108.31	168.63
80-84	117846	166.23	206.42	93836	109.38	167.24
85-89*	107548	170.06	206.67	87752	112.79	170.95

*Information is missing for one or more years.






Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.

United States of America

Deaths from all neoplasms, malignant and benign

Years of Death	Males			Females		
	Deaths No.	World Stand	Trunc 35-64	Deaths No.	World Stand	Trunc 35-64
55-59	676399	141.39	198.36	604328	114.21	187.80
60-64	763006	146.57	205.64	655784	110.39	181.92
65-69	857060	154.71	216.62	721505	109.59	181.16
70-74	952662	160.76	221.53	797426	108.74	180.97
75-79	1063464	163.78	221.16	892072	107.67	177.59
80-84	1183565	166.11	219.21	1017118	110.06	175.83
85-89*	1022413	164.79	214.97	897235	111.68	174.39

*Information is missing for one or more years.



Top panel, total deaths; middle panel, age-specific death rate for males 30-34 to 70-74; bottom panel, age-specific death rate for females 30-34 to 70-74.