

either due to late onset of adenoma development or noncarrier status, molecular testing may identify affected patients or exclude noncarriers. Since combined clinical/molecular diagnosis can identify affected relatives about twenty years earlier than symptomatic probands, sixty of our probands but none of the presymptomatic relatives presented with colorectal cancer. Restorative proctocolectomy followed by an ileoanal J-pouch procedure prevents colorectal cancer and preserves sphincter function in FAP patients. Although familial adenomatous polyposis accounts for less than 1% of all colorectal cancers, hereditary nonpolyposis colorectal cancer (HNPCC) may account for up to 15%. Presymptomatic molecular diagnosis of FAP and HNPCC and preventive surgical treatment might be effective tools to further decrease mortality due to colorectal cancer.

619

COLORECTAL CANCER AND THE NEED FOR SCREENING

J. Northover

ICRF Colorectal Cancer Unit, St Mark's Hospital, London, U.K.

Colorectal cancer is the second commonest killing cancer in Europe. However, there is some cause for hope. First, the adenoma-carcinoma sequence offers a convenient target for screening; intervention during the benign period of the sequence might allow cancer prevention. Second, surgical treatment of colorectal cancer at an early pathological stage is almost always curative. Over the past two decades a major research effort has gone into screening strategies, mainly into faecal occult blood testing. There is guarded optimism while we await the outcome of European RCTs. Another potentially important method of screening in colorectal neoplasia is flexible sigmoidoscopy which, applied once around age 60, might allow cancer prevention by adenoma removal. A very large trial of this modality, involving 200,000 people, is about to start in the U.K.

620

THE PHENOMENON OF CANCER CLUSTERING

P. Boyle^{1,2}, F.E. Alexander²

¹Division of Epidemiology and Biostatistics, European Institute of Oncology, via Ripamonti 435, 20141 Milan, Italy

²Department of Public Health, University of Edinburgh, U.K.

There is a long history of disease 'clusters'; although in many reports the term cluster had only an impressionistic meaning without indication of the numbers of cases expected in the time and place in question. A large number of apparent clusters published throughout this century have involved leukaemia and initial reports were essentially non-quantitative and based on astute clinical observations and *ad hoc* investigations. From the 1930s onwards there were a few wide ranging searches for clusters with some elementary statistical analysis. Such was the **Ashington Cluster** involving three cases of AML, one of ALL and one CLL arising in a small mining village. From around 1970 the field was extended with analysis of large computerised databases becoming a reality and statistical methodology emerging. The phenomenon of space-time interaction was developed and demonstrated in childhood leukaemia in the U.K. The most recent phase of cluster research has generally involved sophisticated statistical analysis of aggregations of cases around point sources such as nuclear power installations. Perhaps the most interesting aspect of clustering at the present time is the consensus emerging that the residences at onset of cases of childhood leukaemia show weak evidence of a general tendency for spatial clustering. This is consistent with shared exposure to localised aetiological agents and is providing a clue as to where to look for aetiological factors for childhood leukaemia.

621

CANCER AND NUCLEAR INSTALLATIONS

C. Hill

Department of Biostatistics and Epidemiology and INSERM U351 Institut Gustave-Roussy, 94805 Villejuif, France

We describe here the main results of studies of cancer mortality and morbidity in the populations residing in the vicinity of nuclear plants. Reports from the U.K. have described increases in leukaemia and lymphoma risks in children living near nuclear installations. Paternal pre-conceptional exposure to radiation had been suggested as an explanation, but this explanation has since been dismissed. An infectious aetiology has also been suggested, based on the observation of an excess risk of leukaemia and non-Hodgkin's lymphoma after population mixing both

around the sites of some nuclear facilities in the U.K., and in other circumstances.

In all other countries where the problem has been studied (U.S.A., Canada, Germany, France, Sweden), no excess morbidity or mortality has been observed in the vicinity of nuclear installations. The power of these studies were reasonable. For instance in a French study, the expected number of deaths around installations was equal to 200, and therefore the probability of detecting an increase of 25% was 95%, with a type I error of 5%.

622

CHILDHOOD CANCER INCIDENCE IN RELATION TO DISTANCE FROM THE FORMER NUCLEAR TESTING SITE IN SEMIPALATINSK, KAZAKHSTAN

D.G. Zaridze

Department of Epidemiology, Cancer Research Center RAMS, Moscow, Russian Federation

There are few data on cancer risk following atmospheric nuclear explosions. Rates of childhood cancer between 1981 and 1990 in the four administrative zones of Kazakhstan were studied to assess the relationship, if any, with distance from nuclear testing sites. Risk of various cancers among children aged 14 years and younger were estimated in relation to distance from (1) a site where testing in air was performed before 1963, (2) a site where underground testing took place thereafter, and (3) a reservoir, known as "Atom Lake", created by four nuclear explosions in 1965. Risk of acute leukaemia rose significantly with increasing proximity of residence to the testing areas, although the absolute value of the risk gradient was relatively small. The relative risk for those living less than 200 km from the air-testing site was 1.76 compared with those living 400 km or more away from the site. Similar relative risks were observed for the underground site and "Atom Lake". There was also some evidence of increased risk of brain tumours in association with proximity to the test sites. In two of the four zones studied, there was substantial regional variation in acute leukaemia rates which was not attributable to distance from the test site. The findings may be affected by potential confounders, notably urban/rural status and ethnic factors but tend to confirm an association with increased risk of childhood leukaemia and exposure to radioactivity from atmospheric nuclear explosions.

623

RISKS OF EXPOSURE TO RADON GAS

S.C. Darby

Imperial Cancer Research Fund Cancer Epidemiology Unit, Radcliffe Infirmary, Gibson Building, Oxford, U.K.

Radon is a radioactive gas that occurs naturally in the earth's crust as part of the decay chain of uranium-238. It has recently been appreciated that by far the greatest source of exposure to ionizing radiation arises from the inhalation of radon indoors. There is conclusive proof that inhaled radon and its decay products can cause lung cancer, both from animal experiments and from the study of men who have worked in mines of uranium and other igneous rocks where radon levels are exceptionally high. At the present time, estimates of the risk of lung cancer from inhaled radon indoors are based on the miners studies. If these estimates are correct, then radon would be the second most important cause of lung cancer after cigarette smoking. However, there are many uncertainties in extrapolating from the mining to the indoor environment. Direct assessment of the risk of indoor radon exposure is currently in progress.

624

CANCER AND ELECTROMAGNETIC FIELDS

E. Pukkala

Finnish Cancer Registry, Helsinki, Finland

There are several studies on the cancer risk of children exposed to electromagnetic fields. The results are controversial with a majority suggesting an increased risk of childhood cancer to be associated with exposure magnetic fields. The only population-based cohort study until now included all 135,000 Finnish children aged 0-19 years who during 1970-89 lived within 500 m of overhead power lines of 110-400 kV in magnetic fields calculated to be $\geq 0.01 \mu\text{T}$. Cancer cases of these children were picked up from the countrywide Finnish Cancer Registry, and the observed numbers of various cancers were compared with the expected ones based on national incidence rates. In the whole cohort, 140 cancers were observed and 145 expected. The only statistically significant excess was found in nervous system cancer in boys (but not in girls) who were