

ranges from 1–11%. The aim of this study was to evaluate the effectiveness of thrombolytic therapy in the management of oncologic patients with major pulmonary embolism (PE).

Methods: Nineteen cancer patients (mean age 53.5 ± 9 , M/F: 9/10) were reviewed. The initial diagnosis was gynecologic cancer ($n = 7$), lung cancer ($n = 4$), breast cancer ($n = 2$), lymphoma ($n = 2$), prostate cancer ($n = 1$), histiocytoma ($n = 1$), osteosarcoma ($n = 1$) and hepatoma ($n = 1$). The clinical suspicion of PE was set by the physical examination, the ECG, the chest X-ray, the echocardiogram and the right heart catheterization findings and confirmed by lung perfusion scan. The fibrinogen during the acute phase of PE was markedly elevated (mean value 794 ± 400 mg/dl). The therapy was initiated with IV streptokinase 250,000 IU per hour for 24–72 hours and was continued with IV heparin administration for 5–7 days. **Results:** Fourteen out of nineteen patients survived and there was improvement of the clinical and scintigraphic status. The thrombolytic therapy in one patient was stopped because of major gastrointestinal bleeding. Four patients died of respiratory failure.

Conclusion: The thrombolytic treatment in oncologic patients with major pulmonary embolism seems to be effective with relatively few hemorrhagic complications.

1244

PUBLICATION

CENTRAL VENOUS CATHETER WITH RESERVOIR IN ONCOLOGICAL PATIENTS. TWO YEARS' EXPERIENCE

A. Delgado, J. Montesdeoca, M. Marrero, A. Murias, J. Almenares, F. Molano

Medical Oncology Unit, Hospital Insular de Gran Canaria, Las Palmas, Canary Island, Spain

From Feb/93, a prospective study was initiated with oncological patients who had had a central venous catheter with reservoir (CVC) implantation, in order to evaluate tolerance and incidence of complications. A total of 218 patients were included, of which 97 were male (44.3%) and 122 female (55.7%). The median age was 52.5 years (age range 9–76 years); 80% of whom had a Karnofsky >70 . The most frequent diagnoses were breast cancer (37%), head and neck cancer (14.2%) and lymphomas (13.7%). In 85.3% of the patients the reservoir was implanted in the right hemithorax (HT) and in 14% the left HT. The vein most frequently used for implantation was the jugular vein (97.7%). The median duration of catheter implantation was 8 months (1–24). In the main, the drugs used were ADM (18%), 5-FU (17.4%), CTX (14.4%). 32.2% of the cases received 24-hours continuous infusion. The median usage of the catheter was 18.8 times (1–62). The percentage of complications was 14.4% and in order of frequency: no blood return (5%), infections (3.2%), thrombosis (2.75%), pain (1.9%), rejection (1.83%) migration (1.8%), rotation of reservoir (0.4%).

Conclusion: There have been few complications and 94% of the patients have indicated that they are content with the catheter.

1245

PUBLICATION

ERYTHROPOIETIN AND CHEMOTHERAPY: EFFECTS ON HEMOSTASIS AND FIBRINOLYSIS

C. Oberhoff, B. Krummeich, U.H. Winkler, A.E. Schindler

Department of Gynecology, University Hospital, D-45122 Essen, Germany
Data from hemodialysis patients who received erythropoietin (EPO) for anaemia report an increased incidence of thrombosis.

We studied hemostasis in 13 patients with gynecological cancer receiving 5000 I.E. EPO (Boehringer Mannheim) s.c. daily, for 12 weeks because of chemotherapy-induced anaemia ($Hb < 11$ g/dl). Blood sampling was done before and monthly on therapy. Pretreatment procoagulant activity, anticoagulation, fibrinolysis and antifibrinolysis were found within the normal range. D-dimer fibrin-split product (1122 ng/ml) were elevated. No change in the dynamic parameters of coagulation and fibrinolysis was seen on therapy except a significant decrease of protein C (50%).

Protein C deficiency is a common complication of anticancer chemotherapy in gynecology. In absence of any increased intravascular coagulation, we suggest that EPO therapy induces no additional risk for thrombosis, but further analysis might be necessary to evaluate if EPO enhances iatrogenic protein C deficiency.

1246

PUBLICATION

BACTEREMIAS IN PATIENTS (PTS) WITH HEMATOLOGIC MALIGNANCIES

K. Palla, M. Hatzizianni, M. Gilianos, G. Christakis¹, A. Katrachoura¹, A.E. Athanassiou, V. Seitanides

Hematology Dept.

¹ Microbiology Department, Metaxa Cancer Hospital, Pireas 185 37, Greece

All positive blood culture isolates from a hematology unit, between 1991 and 1992, were evaluated retrospectively. A total of 88 bacteremic episodes (105 pathogens) were recorded. Patient population consisted of 31 males and 17 females and their age ranged from 17 to 78 years (median 57). 24/48 (50%) pts had 1 bacteremic episode, 12 (25%) pts had 2 episodes, 8 (17%) pts had 3 episodes and 4 (8%) pts had 4 episodes. Nearly all pts suffered from hematologic malignancies. Fever $<38^{\circ}\text{C}$ was present in 78/88 (88.6%) cases. Severe Neutropenia was present in 64/88 (72%) cases. Gram-positive bacteria were isolated in 64 (61%) cases, Gram-negative in 37 (35.2%), and others in 4 (3.8%) [1 *Bacillus*, 1 *Listeria*, 1 *Bacteroides*, 1 *Candida*]. A total of 10 pts had died within 3 weeks of first positive blood culture. Of these fatal cases, Gram-negative bacteria were isolated in 8 episodes, Gram-positive in 5, and *Candida* in 1 (3 polymicrobial isolates). Among them, *Pseudomonas aeruginosa* (3 cases) and *E. coli* (3 cases) were the most prevalent. In conclusion, although Gram-positive bacteremias are increasing steadily (61% versus 40% during 1988–89 in the same unit), Gram-negative infections proved acting as main contributing factor for fatal bacteremias.

1247

PUBLICATION

SUPPORTIVE ERYTHROPOIETIN TREATMENT IN PATIENTS WITH OVARIAN CARCINOMA UNDER CHEMOTHERAPY WITH CIS-PLATINUM

P.S. Parakevopoulos, P. Kourtidu, P. Eliopoulos, L. Zepirides, A. Papadopoulou-Boutis¹, L. Boutis

1st Department of Clinical Oncology

¹ Biological-Chemistry Laboratory of Anticancer Institute of Thessaloniki "THEAGENIO", Greece

In a randomized prospective study we evaluated the efficacy of erythropoietin to maintain the hematocrit levels above 30% in patients with stage III ovarian carcinoma under cis-platinum containing chemotherapy. In this study 20 patients were entered divided into two groups. Group A was the under investigation group with ten patients (mean age 60.4 y) and was treated with Cis-platinum 80 mg/m² + Epirubicin 60 mg/m² + Cyclophosphamide 600 mg/m² (PECX6) every four weeks, and simultaneously received erythropoietin (Epo) 2000 U subcutaneously three times a week. Group B or control group (ten patients with mean age 61.5 y) received the same chemotherapy regimen (PECX6) without Epo supportive treatment. The oscillation of hematocrit, hemoglobin and reticuloerythrocytes were measured during the 6 cycles of PEC chemotherapy, in all 20 patients, as well as the red blood cell units transfused in each patient in order to be able to continue their chemotherapy. The analysis of the results showed an increase of hematocrit in 5 patients of group A (Epo) and stable levels of hematocrit in the other 5 patients of the same group, and only 2 patients needed transfusion with 3 red blood cell units. In the control group B hematocrit decreased in all ten patients and 7 patients needed transfusion with 15 red blood cell units. From the above results we conclude that Epo supports the hematocrit levels satisfactorily. Also requiring considerably fewer red blood cell units for transfusion with all the benefits from the avoidance of adverse effects from the blood transfusion.

1248

PUBLICATION

USE OF METHYLENE BLUE AND BICARBONATE IN IFOSFAMIDE-RELATED CNS TOXICITY. CASE REPORT

H. Serve, E. Hilgenfeld, J. Maurer, E. Thiel, W.E. Berdel

Department of Hematology and Oncology, Benjamin Franklin Hospital, Freie Universität, Berlin, Germany

Severe, sometimes fatal CNS toxicity is a rare, dose-related side effect of ifosfamide (IFO). It has been suggested to be due to toxic effects of IFO metabolites on electron transport. Methylene blue (MB) has been tested in a few patients as an antidote. We have treated 2 adults with MB, who presented with severe IFO-related CNS toxicity. The 1st patient received combination chemotherapy for Wilms-Tumor containing 3.5 g IFO i.v. for 3 days. On day 1 of IFO infusion, he developed paranoid hallucinations, agitation and coma (CTC grade 4) for 3 days. With 5 further cycles of an equal chemotherapy he received 50 mg MB 4 × /d