

Methods: C595 was reduced with 2-mercaptoethanol and labelling was performed by adding 200 MBq of ^{99m}Tc and 0.1 mL of methylene diphosphonate from a bone scanning kit. In vitro studies revealed the size, stability and immunoreactivity of the conjugate. The ability of the conjugate to bind to bladder tumours was initially tested in an ex-vivo model in which five fresh cystectomy specimens containing tumour were filled with ^{99m}Tc -C595 for 1 h. Bladders were imaged and biopsies of tumour and normal urothelium taken. The conjugate was then administered intravesically to five patients who had been diagnosed radiologically or endoscopically with bladder cancer. Patients underwent gamma camera imaging post-washout followed by Transurethral Resection of Tumour (TUR) and biopsy 2 h later.

Results: The direct reduction mediated method produced a stable conjugate of ^{99m}Tc -C595 which had an immunoreactivity exceeding 85%. Gel filtration chromatography revealed a single peak of labelled material at the expected 150 kD region. Binding to bladder tumours was confirmed in the ex-vivo model; four out of the five tumours were successfully imaged with a mean tumour:normal ratio of 5.7:1. In the patient studies, four out of the five tumours were imaged with a mean tumour:normal ratio of 22:1.

Discussion: We have successfully labelled the monoclonal antibody C595 with ^{99m}Tc . The conjugate is stable in vitro and retains its high immunoreactivity. It has demonstrated its ability to localise to bladder tumours in an ex-vivo model and in patients when given intravesically. On the basis of these results we have embarked upon a study of intravenous ^{99m}Tc -C595 immunoscintigraphy in patients with invasive bladder cancer. This reagent may also have an application in diagnostic imaging of other intra-cavity tumours such as ovarian carcinoma.

1402

POSTER

Testicular self-examination (TSE) and testicular ultrasound (TUS) assisted follow-up for the early detection of secondary testicular tumors (STT)

L. Géczi, I. Bodrogi. *National Institute of Oncology, Budapest, Hungary*

Purpose: To improve, by an analysis of the results of 3 years' experience, the early detection of STT assisted by monthly TSE and annual TUS.

Methods: Between April 1 1995 and April 1 1998 annually 702, 782, and 761 patients (pts) participated in our new follow-up program. We provided informative materials to our previously treated and controlled testicular cancer patients about the early signs of testicular cancer, the correct method of self-investigation and the importance of the early detection of STT. If any signs of tumor were palpable TUS was applied within a few days. During this period 14 STT cases were detected; 8 other STT pts were managed in our Department in the usual way. The clinical details of the two groups were compared.

Results: In the first group 4 stage I/A and 10 stage I/B tumors were detected; in 10 pts adjuvant chemotherapy was used and in 4 pts a wait and see approach was adopted. In group 2 the stage distribution was 2 I/A, 2 I/B, 1 II/A and 3 III/B. Three pts received adjuvant and 3 curative chemotherapy, 2 were managed by wait and see policies, and 1 received mediastinal irradiation with chemotherapy. Early detection helped to reduce the duration of symptoms, the number of chemotherapy cycles, the side effects and treatment costs.

Conclusion: With regular follow-up the prognosis of STT is good (CR: 100%), and a biopsy to detect CIS is unnecessary. The pts must be followed regularly for up to 5 years, must be prepared to use TSE-techniques and must recognize the importance of the early detection of STT. TUS, in combination with TSE, enhances the chance of early diagnosis; subsequently these tests can provide the basis for a cost-effective treatment with reduced side effects.

1403

POSTER

Radiotherapy (XRT) for testicular intraepithelial neoplasia (TIN)

J. Classen, K.-P. Dieckmann, M. Bamberg. *University of Tuebingen, Radiooncology, Hoppe-Seyler-Str. 3, 72076 Tuebingen, Germany*

Introduction: TIN, also known as carcinoma in situ of the testis, has been recognized as the common precursor lesion of testicular cancer except for spermatocytic seminoma. In case of TIN in a solitary testis or double sided TIN local XRT with 20 Gy is the treatment of first choice yielding 100% cure rates. However, subsequent to XRT patients are at risk for delayed Leydig-cell insufficiency eventually necessitating testosterone substitution. Therefore new treatment strategies aim at the reduction of the total dose of XRT without compromising local control of TIN.

Methods: A multicenter prospective trial was initiated for XRT of histologically proven TIN. Patients were treated with reduced total doses. A

progressive dose reduction in 2 Gy steps was performed starting with 18 Gy. All patients had complete hormonal evaluation prior to XRT and again during follow-up. At least one control biopsy after XRT was required. Acute and possible late effects of treatment were prospectively documented using the EORTC-score.

Results: To date, 13 patients have been enrolled for the trial. 11 patients were treated with 18 Gy, 8 of which have so far undergone control biopsy showing complete eradication of TIN. 2 patients were treated with 16 Gy. Acute side effects of treatment are negligible. With a median time to follow-up of 6.8 months (range: 0–18.5 mo.) so far no significant decline in serum testosterone after XRT has been observed.

Conclusion: Preliminary data of the study indicate that XRT of TIN with 18 Gy may be sufficient for local control of this premalignant lesion. Further reduction of the XRT dose in the course of the study may provide a less toxic yet highly efficient therapy for TIN.

1404

POSTER

Inhibition of telomerase activity by antisense oligonucleotides in renal cell carcinoma

R. Heicappell¹, J. Sachsinger¹, H. Krause¹, M. Müller¹, K. Müller¹. ¹Freie Universität, Urology, Berlin, Germany

With only a few exceptions, the ribonucleoprotein telomerase is found in malignant, but not in benign tissues. It has been previously shown by us and others that telomerase activity can be found in malignant tumors of bladder and kidney. Telomerase is obviously required for indefinite, malignant growth. Therefore, inhibition of telomerase activity would be a logical approach for therapy of malignant tumors such as renal cell carcinoma. Antisense RNA was chosen for telomerase inhibition in the present study. As tools for this type of therapy are not readily available, vectors were constructed allowing for expression of antisense telomerase RNA in murine RCC cells.

Murine renal cell carcinoma line RENCA, which is highly tumorigenic in mice, displays high telomerase activity, and a mean telomere length comparable to human cells (7–10 kb) was used as a model. After electroporation with eukaryotic expression vectors harboring the RNA template in antisense direction stable transfectants were isolated and tested for telomerase activity (TRAP assay), shortening of telomere length, and proliferative activity. After 35 population doublings 9 out of 13 antisense-transfected RENCA clones showed a reduced replicative potential compared to their non-treated counterparts. Fluorescence-based TRAP assay and measurement of the mean terminal restriction fragment length (TRF) gave evidence for reduced enzyme activity and resumption of telomere shortening in the transfectants. Growth inhibition was, however, of temporary nature only in some cell clones.

Antisense oligonucleotides may effectively inhibit telomerase activity, addition of telomeric DNA, and proliferation of murine renal cell carcinoma.

1405

POSTER

Gemcitabine/cisplatin in advanced transitional cell carcinoma of the urinary tract (TCC): A phase II multicenter trial

S. Mancarella, V. Lorusso, L. Manzone, M. Antimi, F. De Vita, A. Santoro, L. Maiorino, R. Mattioli, G. Luporini, D. Bilancia, M. De Lena. *Italian Cooperative Study on Bladder cancer Oncology Institute, Via Amendola 209, 70121 Bari, Italy*

Since January 1997, 54 patients (pts) affected by measurable advanced TCC were treated with gemcitabine 1000 mg/m² on days 1, 8, 15 and cisplatin 70 mg/m² on day 2 repeated every 4 weeks. Patients' median age was 67 (range 18 to 80), none had prior chemotherapy for metastatic disease, 7 had adjuvant cisplatin-chemotherapy concluded by >1 year, 52 were male and 2 female, 16 had prior cystectomy, 36 had ECOG 0–1 and 18 had ECOG 2 PS respectively. All patients entered on study have been evaluated for response and toxicity (*intent to treat*). Complete (CR) plus partial (PR) response were observed in 26/54 pts (48%) with 8 pts (15%) attaining CR. Twelve pts (22%) showed stable disease (SD) and 6 progressed (15.5%). G3–4 leukopenia was observed in 21 (39%) pts, and G4 neutropenia in 6 (11%). Grade 3–4 thrombocytopenia was recorded in 11 (20%) pts and G4 thrombocytopenia in 3 pts. The median number of cycles delivered was 4; 28 pts (52%) required dosage reduction of gemcitabine or omission to deliver it, on day 15 for neutropenia or thrombocytopenia. No other relevant side effects were recorded. Median time to progression and survival were 26 weeks and 36 weeks respectively. Out of 32 pts with disease related symptoms (haematuria, pain, weight loss), 16 (50%) improved.

Conclusions: the combination of G plus P demonstrated to be very active and tolerable in TCC ushering us in the *post-M-VAC era* for TCC pts.

1406

POSTER

Combination chemotherapy with cis-platin (CDDP), epirubicin (EPI), and docetaxel (DOC) in transitional cell urothelial cancer (TCC). A phase II study

D. Pectasides, A. Visvikis, A. Aspropotamitis, A. Halikia, A. Kalantaridou, S. Batzios, N. Karvounis, A. Athanassiou. *1st Dept. of Medical Oncology, Metaxa's Memorial Cancer Hospital, Piraeus, Greece*

Purpose: This study was conducted to evaluate the efficacy and toxicity of the combination of CDDP, EPI and DOC in locally advanced or metastatic urothelial TCC.

Methods: Thirty-two chemotherapy (CT) – naïve patients (pts) were treated with EPI 40 mg/m² I.V., DOC 75 mg/m² I.V. with standard oral steroids premedication and CDDP 75 mg/m² I.V. with pre and post-hydration. Treatment was repeated every 3 weeks. Pts were evaluated for toxicity weekly and assessed for response every 2 cycles of CT. Twenty-four pts had metastatic and 8 pts locally advanced disease. There were 6 female and 26 male with a median age of 65 y. The ECOG PS was 0 in 10 pts, 1 in 6, 2 in 2 and 3 in 4.

Results: There were 9 (30%) CRs (2, 28.5% in locally advanced and 7, 30.4% in metastatic disease) and 11 PRs (3, 42.8% and 8, 34.7%, respectively) with an ORR of 66.6% (71.4% and 65.2% respectively). The median duration of response in pts with metastatic disease was 8.5 mos and the overall Survival 14.5 mos (15 mos and 12.5 mos, respectively). Sixteen (53.3%) pts required a dose reduction and 5 (16.6%) 2 dose reductions. There were 4 episodes of febrile neutropenia and sepsis, which were successfully treated with broad spectrum antibiotics. There were no treatment delays due to myelotoxicity. G3/4 anemia and thrombocytopenia occurred in 12.5% and 9.3% of pts, respectively. G2 nephrotoxicity and G2/3 neurotoxicity occurred in 9.3% and 9.3% of pts, respectively. Alopecia was universal. G3/4 nausea/vomiting, G2/3 mucositis and G2/3 diarrhea developed 15.6%, 15.4% and 21.9% of pts, respectively. G2/3 fluid retention and G2/4 allergy occurred in 9.3% and 9.3% of pts, respectively.

Conclusion: The combination of EPI, DOC and CDDP is an active regimen for urothelial TCC. The safety profile mainly consisted of neutropenia leading to dose reduction, but with no serious infectious complications.

1407

PUBLICATION

Occupational prostate cancer risk factors in an area of coal, iron, and steel industries in Germany

T. Bandel¹, T. Reckwitz², S. Dickhut¹, P. Jedrusik^{1,2}, W. Weistenhöfer¹, H.M. Bolt¹, H. Schulze¹, K. Golka¹. *¹Institute of Occupational Physiology at the University of Dortmund, Dortmund; ²Urological Department, Städtische Kliniken Dortmund, Dortmund, Germany*

Aim of the study was to identify possible risk occupations for prostate cancer in the Ruhr-area, a former center of the German coal, iron, and steel industries.

Methods: In 238 cases with histologically proven prostate cancer and of 414 controls with benign prostatic hyperplasia all occupations ever performed for more than 6 months and life time smoking habits of were asked for with a questionnaire. Confounder-adjusted odds ratios were estimated by Fisher's exact test and logistic regression analysis.

Results: Confounder-adjusted odds ratios for both age (OR 2.56) and duration of employment (OR 2.22) were elevated in underground hard coal miners. An elevated prostate cancer risk was also observed for painters/varnishers (OR 2.84 adjusted for age, OR 2.91 adjusted for duration of employment). Steelworkers showed no increased risk (OR 0.93 adjusted for age, OR 0.94 adjusted for duration of employment). Businessmen showed a remarkably low prostate cancer risk (OR 0.38 adjusted for age, OR 0.37 adjusted for duration of employment). No differences in the smoking habits described by pack years could be found between cases and controls.

Conclusion: Coal dust components and dietary factors due to high energy expenditure must be discussed as possible risk factors for prostate cancer in hard coal miners.

1408

PUBLICATION

Salivary biochemical and immunological profile following low-dose IL-2 based immunotherapy

R.M. Nagler¹, E. Gez², R. Rubinov², D. Laufer¹, H. Ben-Aryeh¹, D. Gailini³, M. Filatov¹, A. Kuten². *¹Oral Biochemistry Laboratory; ²Dept. of Oncology; ³Dept. of Diagnostic Radiology, Rambam Medical Center and Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel*

Purpose: One of the side effects accompanying low-dose recombinant interleukin-2 (rIL-2)-based immunotherapy is salivary hypofunction. We evaluated the functional and compositional whole salivary profile at both resting and stimulated conditions in 10 renal cell carcinoma patients who received prolonged low-dose rIL-2-based immunotherapy.

Results: At three days following the termination of four weeks of the combined administration of rIL-2 and rIFN- α , 21% reduction of salivary flow rate [$p < 0.05$] at resting condition was found, accompanied by significant multiple compositional alterations, including an increase in calcium, magnesium and phosphate concentrations by 65.6% [$p < 0.01$], 50% [$p < 0.05$] and 27% [$p < 0.05$], respectively, and a 23.5% [$p < 0.05$] reduction in the total protein concentration. In contrast, no flow rate reduction was noted under stimulated condition, and the only altered compositional component was the phosphate which was increased by 29.3% [$p < 0.05$]. The concentrations of all the other salivary components analyzed, including sodium, potassium, amylase, albumin, and the immunoglobulins, IgG, IgA and secretory IgA, were not effected by the immunotherapy. At one month following the termination of the immunotherapy, no functional or compositional salivary alterations were noted.

Conclusion: We recommend salivary-supporting therapies and anticarcinogenic treatments for patients undergoing low-dose rIL-2-based immunotherapy.

1409

PUBLICATION

Immunohistochemical evaluation of the apoptotic markers Bcl2, Par-4 and p53, VEGF receptor Flk-1 and correlation with clinicopathological parameters in localised prostate adenocarcinomas

A. Giannopoulos¹, L. Nakopoulou², E. Serafetinides¹, H. Gakiopoulou², A. Rashdan¹, A. Zervas¹, M. Giannopoulos¹, C. Dimopoulos¹. *¹Athens Medical School, University of Athens, Urology, Athens; ²Athens Medical School, University of Athens, Pathology, Athens, Greece*

Purpose: Oncogenic protein Bcl2 expression is a key event in apoptosis while mutant p53 protein plays also an important role as it regulates expression of Bcl2. Prostate apoptosis response-4 (PAR-4) is specifically expressed by cells entering apoptosis. Flk-1 represents a high affinity receptor for vascular endothelial growth factor. In a retrospective study we evaluated the expression of the former markers in a cohort of 37 surgical specimens of localised prostate adenocarcinomas and the possible relationship with PSA, grade and stage.

Methods: Thirty seven formalin-fixed paraffin-embedded archival prostate cancer specimens were examined by immunohistochemistry with respect to apoptotic markers and Flk. Preoperative serum PSA, grade and pathological stage were included in statistical analysis.

Results: The expression of Par-4 and Flk-1 in malignant and normal prostatic tissue was inverse. No relation was found in expression of these markers with PSA, grade and stage. An inverse relation between p53 and Bcl2 expression with PSA was noted ($p = 0.0006$ and $p = 0.008$ respectively). Bcl2 expression in malignant tissue correlates with grade ($p = 0.044$) and stage ($p = 0.06$) of disease.

Conclusion: Bcl2 expression is considered an independent prognostic marker in early prostate cancer while correlation of the other markers needs further research.

1410

PUBLICATION

Gemcitabine and paclitaxel in previously treated patients with advanced transitional cell carcinoma

L. Marini¹, C. Sternberg¹, A. Sella², F. Calabrò¹, A. Van Rijn¹. *¹San Raffaele Scientific Institute, Medical Oncology, Rome, Italy; ²Rabin Medical Center, Medical Oncology, Petah Tikvah, Israel*

Second-line therapy is needed for patients (pts) with transitional cell carcinoma (TCC) who fail first-line M-VAC chemotherapy. Both Gemcitabine (G) and Paclitaxel (T) have demonstrated activity in TCC. Based upon the results of an Q 2 week G + T schedule (Ann Oncol 1998; 9: 733-738), 18