

patients completed treatment with a total of 107 cycles (median: 3.5 cycles, range: 1-9). Four patients are still on treatment. One complete response (breast) and 2 partial responses (larynx and lung) were observed with 7 additional stable diseases. The main toxicity is neutropenia G3-4. The following DLTs were observed: At level 3 (D:80/MS:600): febrile neutropenia G4, infection G3 leading to death, At level 4 (D:80/MS:900): stomatitis G3, dysphagia G3, fatigue G3, At level 5 (D:80/MS:1200): 2 DLT: stomatitis G3/ neutropenia G3 and stomatitis G4/fatigue G3. The MTD was reached at level 5. The recommended dose is then level 4 (D:80/MS:900). PK analysis did not demonstrate a strong PK interaction between the two compounds but at the highest dose levels, there is a trend to an increase of docetaxel AUC when this agent is given in combination with MS209. Complete data set and PK analysis will be presented.

520

A phase I pharmacokinetic study of ONT-093 in combination with paclitaxel in patients with advanced cancer

K.N. Chi¹, S.K. Chia¹, B. Sikic², R. Dixon³, M. Newman³, V.J. Wachter³, B. Toyonaga³, D. Hoth³, K.A. Gelmon¹. ¹BC Cancer Agency, Medical Oncology, Vancouver; ²Stanford University, Palo Alto, CA, USA; ³Ontogen Corporation, Carlsbad, CA, USA

Background: ONT-093 (OC144-093) is a third generation, orally bioavailable inhibitor of P-glycoprotein (P-gp). In pre-clinical studies, inhibition of P-gp and reversal of multidrug resistance (MDR) could be achieved at nM concentrations. ONT-093 does not affect paclitaxel pharmacokinetics (PK) in rodents. Phase I clinical trials of single agent ONT-093 in normal human volunteers showed no dose-limiting toxicities at biologically active serum concentrations in doses ranging from 300 to 500 mg.

Methods: We performed a Phase I pharmacokinetic trial of ONT-093 in doses from 300 to 500 mg administered orally 14 h before, 2 h before and 10 h after intravenous paclitaxel doses of 150 to 225 mg/m² repeated every 21 days. All patients received paclitaxel alone for cycle 1 to allow intra-patient comparisons of toxicity and paclitaxel pharmacokinetics.

Results: To date, 18 patients have been enrolled into 4 dose levels and have received doses of ONT-093 up to 500 mg and paclitaxel up to 175 mg/m². Toxicities have mainly been attributable to paclitaxel, and included arthralgia, myalgia, neutropenia, and peripheral neuropathy. Toxicities possibly associated with ONT-093 include grade 1-2 headaches and transient grade 1 elevation of liver transaminases in 1 patient. Three patients have had higher-grade neutropenia with cycle 2. One of these patients, who was also heavily pre-treated and had extensive hepatic metastases, had febrile neutropenia on cycle 2, dose level 4 (ONT-093 500 mg and paclitaxel 175 mg/m²). This cohort is being expanded. C_{max} concentrations of ONT-093 given at 500 mg are > 8 µM, well above that required in pre-clinical models to inhibit P-gp and completely reverse MDR. Plasma PK of paclitaxel are unchanged between cycle 1 and 2.

Conclusions: Biologically active doses of ONT-093 have been well tolerated in combination with standard doses of paclitaxel. There have been no alterations of paclitaxel PK parameters with the combination at the doses tested. These results support the continued clinical development of ONT-093 as an active, potent, non-toxic inhibitor of P-gp in conjunction with cytotoxic chemotherapy. Patient accrual continues, and final results will be presented.

521

Hyperthermia and multidrug resistance: Impact on expression and regulation of MDR genes in human cancer cells

U. Stein¹, W. Walther¹, K. Juerchott¹, S. Bergmann¹, M. Schlaefke², B. Rau², H.D. Royer³, P. Hohenberger², P.M. Schlag². ¹Max-Delbrueck-Center for Molecular Medicine, Surgery / Surgical Oncology, Berlin, Germany; ²Robert-Roessle-Hospital and Tumor Institute, Charite, Humboldt-University; ³Institute for Transplantation Diagnostics and Cell, Heinrich-Heine-Univ., Duesseldorf, Germany

Hyperthermia is used for the treatment of cancer patients in combination with chemotherapy, cytokines and/or radiotherapy. Since multidrug resistance (MDR)-associated genes are inducible by external stress factors such as heat and drugs used for chemotherapy, we investigated the influence of hyperthermia on expression, regulation and function of the MDR genes MDR1, MRP1, and MVP/LRP in *in vitro* models as well as in clinical specimens. In colon carcinoma cell lines, hyperthermia caused nuclear translocation of the transcription factor YB-1. Nuclear YB-1 interacts with the pro-

motors of the MDR1 and MRP1 genes and is associated with increased MDR1 and MRP1 gene transcription, as well as strong efflux pump activity. However, a combination of hyperthermia and drug treatment effectively reduced cell survival demonstrating that activation of MDR1 and MRP1 gene expression and increased efflux pump activity after hyperthermia did not consequently lead to an increase in a MDR phenotype. The ability of hyperthermia to abrogate drug resistance in the presence of an increase in functional MDR proteins may provide an explanation for the efficacious results seen in the clinic in colon cancer patients treated with hyperthermia and chemotherapy. We investigated MDR1 expression in colon cancer specimens of patients who were treated by radio-chemo-thermo therapy. We find that the levels of MDR1 expression in colon cancer specimens before and 6 weeks after radio-chemo-thermo-therapy were not significantly different in the majority of cases. Since induction of MDR1 gene expression by external stress factors such as heat occur directly after exposure to hyperthermia, we conclude that this increase of MDR1 gene activity has ceased after 6 weeks, the time point of resection when the tumor specimens were analyzed. We then determined the levels of MDR genes sequentially before, during, and after isolated hyperthermic isolated limb perfusion (hILP) with rTNFα/melphalan in patients with advanced soft tissue sarcoma and locoregional metastatic malignant melanoma. In the majority of patients (> 80%) MVP/LRP expression was induced during hILP, often paralleling the increase in temperature during hILP. This is the first study to investigate expression of MDR genes sequentially during hILP of patients. The result of this study demonstrates that hILP caused selective induction of MVP/LRP expression, whereas MDR1 and MRP1 expressions were rarely affected by the treatment regimen.

522

Polymorphisms of MDR1 and MRP2/cMOAT in healthy North Eastern Italian subjects

G. Toffoli, E. Cecchin, B. Basso, F. Sartor, G. Biscontin, L. Cadelli, A. Steffan, R. Sorio. C.R.O.-National Cancer Institute, Experimental and Clinical Pharmacology, Pordenone, Italy

MDR1 and MRP2 belong to the ABC transporter genes superfamily encoding for integral membrane glycoproteins that function as ATP-dependent export pumps with substrate specificity. The increased expression of these proteins in cancer cells is associated with the development of cross resistance of tumors to many antitublastic agents. Their activity in normal tissue cells has a protective role towards xenobiotics and prevents chemotherapy toxicity modulating pharmacokinetics of antineoplastic drugs. At least 15 single nucleotide polymorphisms (SNPs) have been described in a Caucasian population for MDR1, one of them resulted particularly interesting such as C3435T in exon 26, its pharmacological effect has been demonstrated and its frequency is quite variable among different ethnic groups. For MRP2 several mutations have been described in Dubin-Johnson Syndrome patients. Few of them have been detected also in Japanese and Jewish healthy subjects and described as SNPs. Their influence on the protein activity are not yet clarified but they could have a role on MRP2 expression or functionality. No population studies have been reported till now on them. We analysed the frequency of C3435T SNP for MDR1 gene and C24T SNP for MRP2 in 800 healthy blood donors from North East of Italy. Distribution of MDR1 C3435T genotype was as follows: C/C in 191 subjects (23.96%), C/T in 434 subjects (54.26%), and T/T in 175 subjects (21.78%). The allelic frequency was 48.91% for T allele, 51.09% for C allele. Distribution of MRP2 C24T genotype was as follows: C/C in 494 subjects (61.78%), C/T in 271 subjects (33.86%), and T/T in 35 patients (4.36%). The allelic frequency was 78.7% for C allele, 21.29% for T allele was. In conclusion the population we analysed showed for MDR1 a frequency quite similar to the one described in literature for the Caucasian population. For MRP2 we found out a considerable allelic frequency for C24T in our geographic area and this encourages further investigations to evaluate its impact on pharmacokinetics of drugs excreted by this transporter protein.

523

Co-dominance of cisplatin resistance in somatic cell hybrids

Y.M. Ying, D.W. Shen, X.J. Liang, M.M. Gottesman. National Cancer Institute, N.I.H., Laboratory of Cell Biology, Bethesda, USA

Intrinsic or acquired resistance to cisplatin in cancer cells remains a major obstacle to successful chemotherapy. The relevant genetic and molecular mechanisms of resistance have not yet been identified. We have isolated cisplatin-resistant human KB epidermoid carcinoma cell lines resistant to varying levels of cisplatin after single and multiple selection steps.