



Erratum

Erratum to “EUROLOX 1: Uterine safety of kilogest versus evista: preliminary results” [European Journal of Cancer, 38 Suppl 6 (2002) S83–S84]

H. Depypere, P. Neven*, T. Lunde, P. Benedetti-Panici, Tiitinen, B. Marinescu,
T. de Villiers, T. Hillard, A. Cano, E. Peer, D. Quail, B.T. Nickelsen
for the Euralox 1 Study Group

Flemish Gynecologic Oncology Group, Azalealaan 10/3, B-9100 Sint Niklaas, Belgium

The Publisher regrets that an error occurred in Fig. 1 of the published article mentioned above. This figure is correctly printed below.




Fig. 1. Gynaecological surveillance algorithm used in the Euralox 1 study. In patients who underwent a routine Pipelle biopsy at baseline (group A), this algorithm was administered in cases of repeated bleeding/spotting at the 6-month visit (visit 4) if the routine TVU showed an endometrial thickness of > 5 mm. In patients without routine Pipelle biopsy at baseline (group B) who had repeated bleeding/spotting during the study, the algorithm was administered at visit 3 or 4 as needed.

☆ PII of original article: S0959-8049(02)00299-X

* Corresponding author. Tel.: +32-1634-4634; fax: +32-1634-4629.

E-mail address: patrick.neven@uz.kuleuven.ac.be (P. Neven).