

5137

POSTER

Unsatisfactory Accuracy of Axillary Ultrasound and Ultrasound-guided Fine Needle Aspiration Cytology of Axillary Lymph Nodes in Breast Cancer Patients

M.W.H. Leenders¹, M. Broeders¹, L. Croese¹, H.L.S. Go², B.L.A.M. Langenhorst¹, W.H. Schreurs¹. ¹Medical Center Alkmaar, Surgery, Alkmaar, The Netherlands; ²Medical Center Alkmaar, Radiology, Alkmaar, The Netherlands

The purpose of our study was to evaluate the accuracy of axillary ultrasound and ultrasound-guided fine needle aspiration cytology (FNAC) in the preoperative diagnosis of axillary metastases. Between 2004 and 2009, 1151 female patients were evaluated and treated in our clinic for histologically proven breast carcinoma. Preoperative axillary ultrasound with subsequent FNAC in case of suspicious lymph nodes was performed in all patients. We analyzed the results of axillary ultrasound and FNAC retrospectively. Pathological node status was used as the reference standard (based on axillary dissection or sentinel node biopsy). 427 of 1151 patients (37.1%) had metastatic disease on final histological analysis. Axillary ultrasound showed suspicious lymph nodes in 328 patients (28.5%). Of these 328 patients, FNAC was performed. FNAC established axillary metastases in 108 patients. Final histological analysis could not find metastases in 4 of these 108 patients. Sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of axillary ultrasound alone were 43.6% (186/427), 80.4% (582/724), 56.7% (186/328) and 70.7% (582/823), respectively. When combining axillary ultrasound with FNAC of suspicious lymph nodes, sensitivity was 24.4% (104/427), specificity was 99.4% (720/724), PPV was 96.3% (104/108) and NPV was 69.0% (720/1043). Therefore, 104/427 (24.4%) node-positive patients were identified by ultrasound-guided FNAC and spared unnecessary sentinel node biopsy. Unfortunately, the percentage false negative results of ultrasound-guided FNAC (28.1%, 323/1151) was very high.

5138

POSTER

Flat Epithelial Atypia

P. Ghuijs¹, C. Boetes², F.W. van der Ent³, E.M. Heuts¹, K.B.M.I. Keymeulen¹, M.N.F. von Meyenfeldt¹, L.J.A. Strobbe⁴, K.K.B.T. van de Vijver⁵, C.A.P. Wauters⁶, M.L. Smidt¹. ¹Maastricht Medical University, Surgery, Maastricht, The Netherlands; ²Maastricht Medical University, Radiology, Maastricht, The Netherlands; ³Orbis Medisch Centrum, Surgery, Sittard-Geleen, The Netherlands; ⁴Canisius-Wilhelmina Ziekenhuis, Surgery, Nijmegen, The Netherlands; ⁵Maastricht Medical University, Pathology, Maastricht, The Netherlands; ⁶Canisius-Wilhelmina Ziekenhuis, Pathology, Nijmegen, The Netherlands

Background: Flat epithelial atypia (FEA) is a presumably neoplastic alteration of terminal duct-lobular units, that is characterized by the replacement of the native luminal epithelium by ductal cells demonstrating low-grade cytologic atypia. The architecture shows stratification of the epithelial cells. FEA is often accompanied by microcalcifications and therefore often discovered in biopsies of mammographically detected microcalcifications. FEA is frequently seen in association with atypical ductal hyperplasia (ADH), low-grade ductal carcinoma in situ (DCIS), lobular neoplasia and invasive tubular carcinomas. There is emerging evidence to suggest that FEA may even represent a precursor of DCIS. Detection of FEA in an histological biopsy should therefore be an indicator signal for possibly present (pre-)malignant lesions. The risk for subsequent breast carcinoma remains to be defined. The aim of this study is therefore to inventories the consequences of solitary FEA in histological biopsies in three Dutch teaching hospitals (university and peripheral).

Materials and Methods: Data of all FEA patient of this retrospective multicentre study were collected in a database. Patient files were detected by searching the local pathology databases for 'FEA', 'Flat Epithelial Atypia', and Dutch equivalents. Results were manually screened, only including solitary FEA.

Patient files were viewed then for information on presentation, mammography and ultrasound and chosen therapy (surgery vs follow-up). In case of excision, the definitive pathology was added.

Results: The search resulted in 161 FEA cases, of which 66 solitary FEA biopsies. The management of these patients consisted of follow-up for 41 patients (62%) and 25 (38%) underwent a lumpectomy (23) and mastectomy (2). No incidents occurred in the follow-up group so far. The definitive pathology of the lumpectomy/mastectomy showed no abnormalities or solitary FEA in both 6 patients; other findings were ADH in 3, LCIS in 2 and DCIS in 5 patients. Invasive lobular disease (ILC) was only found in one patient. Reason for choosing mastectomy were contralateral malignant disease; the definitive pathology showed no abnormalities.

Conclusions: No consistent management exists concerning solitary FEA. Lack of this study is the retrospective gathering of data, making it difficult to detect the reason for the chosen management. DCIS or ILC was discovered in 24% of all surgical patients. Therefore, FEA can be seen as a red flag, indicating the possible presence of a more malignant lesion. Additional research is warranted, preferably as a multicentre randomized controlled trial comparing surgery vs follow-up.

5139

POSTER

Lymph Node Surgery in Breast Cancer – Age-related Inequality in Surgical Procedure Use

C. Cluze¹, M. Meresse¹, M.K. BenDiane¹, D. Rey¹, R. Giorgi². ¹Ours Paca – Inserm Umr912 – Aix-Marseille Université, Marseille, France; ²Laboratoire d'Enseignement et de Recherche sur le Traitement de l'Information Médicale – Aix-Marseille Université – Assistance publique hôpitaux de Marseille Service de Santé Publique et d'Information Médicale, Marseille, France

Background: Sentinel lymph node dissection (SLND) induces lower morbidity and gives a better quality of life than axillary lymph node dissection (ALND) in breast cancer (BC) women. Furthermore, SLND is the recommended method for the axillary nodal staging of BC. However, for some patients ALND is performed without previous SLND. The aim of the study was to identify factors associated with ALND use without initial SLND in early stage BC women.

Materials and Methods: This study was performed in the cohort ELIPPSE65, a French representative cohort of women with non-metastatic BC, aged 65–80 years, and diagnosed between 2007 and 2010. The study sample was all BC women with an indication of SLND i.e. with T1 tumour without palpable axillary node(s) and who had not received neo-adjuvant chemotherapy. Characteristics of women who had SLND were compared with those who had immediately ALND without initial SLND.

Results: A total of 363 women had an indication of SLND. Among them, 6 had no axillary surgery. The axillary surgical procedure used was unknown for 6 women. Accordingly, 351 women were analyzed. ALND was performed without previous SLND in 21% of cases (n = 75). Among these women, 85% had no lymph node involvement.

After taking into account the increase of the use of SLND over time, ALND without previous SLND was more frequently performed in women older than 75 years (aOR = 2.0 [1.1–3.8]) and among those who had no child (aOR = 2.4 [1.1–5.4]). Co-morbidities, physical and psychological disabilities, living place or place of treatment did not explain these inequalities.

Conclusions: Although SLND is a widespread surgical practice, many women still have unnecessary ALND, especially oldest women and those with poor family support. This is particularly detrimental for oldest women whose quality of life and autonomy can be strongly impaired by the lymphoedema following ALND and its long term complications.

5140

POSTER

Stress and Autoaggression in Patients With Breast Cancer (BC) During the Surgical Treatment

O. Krechyk¹, I. Schepotin¹, O. Zotov¹, J. Zaielieva¹. ¹National O.O. Bohomolets Medical University, Oncology, Kiev, Ukraine

Background: Providing of successful treatment of cancer patients and their optimal quality of life requires studying of disease's psychological component and its correction. Lack of experience in providing psychological care in home oncological practice isn't compensated by taking the experience of foreign oncopsychologists and creates a necessity of investigations and elaboration proper methods of psychological support for patients with BC.

Objective: The aim of the study is to investigate the degree of the stress and autoaggression in patients with BC under radical surgery, its dependence of the extent of surgical intervention.

Materials and Methods: 75 women aged 25–65 were examined in 2010–2011 immediately after surgery. They were divided into 3 groups: I – patients with BC who underwent breast-preserving operation, II – patients with BC who underwent radical mastectomy, III – patients with benign breast tumours who underwent sectoral resection. We used Spilberger-Hanin diagnostic scale of reactive and personal anxiety, Bass-Darka diagnostic scale of indexes and forms of aggression, clinical examination and interview.

Results: From all examined patients 40 (53.4%) had signs of acute stress reaction, 51 (68%) had signs of adaptation disorder. The risk group of stress disorder consists of 31 patients with BC.

Conclusions: The degree of acute stress reaction is higher in patients with BC and doesn't depend on the extent of surgical intervention.