

for IBC. The last patient also developed distant metastases concurrently. One patient developed distant metastasis without locoregional recurrence. Of the patients with LR, one patient was treated with excision of DCIS, one patient was treated with excision combined with chemo-, hormonal, and radiation therapy. The two patients with distant metastases were treated only with hormonal therapy.

Conclusions: SSM with IBR carries no increased risk for loco regional or distant recurrence compared to a non-skin sparing mastectomy in our single institute.

5143

POSTER

Ultrasonography by Breast Surgeons in the Operating Theatre – Evaluation of the Initial Learning Phase

N. Krekel¹, B.M. Zonderhuis¹, S. Muller², H. van der Veen², A.M.F. Lopes Cardozo³, A.H.M. Taets van Amerongen⁴, E. Bergers⁴, E.S.M. de Lange de Klerk⁵, S. Meijer¹, M.P. van den Tol¹. ¹VU University Medical Center (VUmc), Surgical Oncology, Amsterdam, The Netherlands; ²Red Cross Hospital, General Surgery, Beverwijk, The Netherlands; ³Medical Center Alkmaar, General Surgery, Alkmaar, The Netherlands; ⁴VU University Medical Center (VUmc), Radiology, Amsterdam, The Netherlands; ⁵VU University Medical Center (VUmc), Epidemiology and Biostatistics, Amsterdam, The Netherlands

Background: Many studies have proven that intra-operative use of ultrasonography (US) during breast-conserving surgery is the most accurate method to achieve high rates of tumour-free margins, with small volumes excised. US, therefore, has become an increasingly popular modality to guide the excision of malignant breast tumours. This study was performed to assess the accuracy of a surgeon's learning in US-guided excision of palpable breast cancer. Difficulties and potential pitfalls were analysed.

Methods: A total of 30 female patients undergoing breast-conserving surgery for palpable T1-T2 invasive breast cancer were recruited. The US assisted three individual breast surgeons to target and excise the tumour with continuous intra-operative imaging. The main objective was to obtain adequate resection margins with optimal resection volumes. The specimen volume was measured, and the tumour diameter and histological margin status were registered from the pathology reports. An optimal resection volume was defined as the spherical tumour volume with an added 1.0-cm margin. The specimen volume was compared to the optimal resection volume. The resulting calculated resection ratio (CRR) indicated the excess tissue resection.

Results: All tumours were correctly identified during the surgery, and 29 of 30 tumours (96.7%) were removed with adequately negative resection margins; focally positive margins were found in one tumour (3.3%). Median CRR was 1.0 (range, 0.4–4.4); for all breast surgeons, CRR improved during the learning period. By the 8th procedure, all surgeons showed proficiency in performing intra-operative breast-US. The main difficulties were recognising ill-defined tumour margins, and relating the US-image to the tumour position in the breast.

Conclusions: Surgeons can easily learn the skills to perform intra-operative US for the excision of palpable breast cancer. Intra-operative surgeon-performed US is non-invasive, simple, safe and effective in obtaining adequate resection margins, as it permits a real-time localisation of the breast carcinoma and the subsequent planning of surgical margins. The resection volumes markedly improved during the learning period into optimal volume resection, thereby presumably resulting in improved cosmetic outcome. In a prospective randomised controlled clinical trial the surgical accuracy of intra-operative US guidance is compared with the traditional palpation-guided surgery for palpable breast cancer.

5144

POSTER

Comparison of Level 3 Nodal Yield in Carcinoma Breast Patients Using the Subpectoral and Interpectoral Approach of Axillary Dissection

V.S. Chauhan¹, N. Mittal², S. Raina¹. ¹Metro Hospital, Department of Surgery, Faridabad, India; ²Metro Hospital, Department of Pathology, Faridabad, India

Background: Level 3 Axillary nodal clearance during modified Radical Mastectomy in Carcinoma Breast is done conventionally by retracting the pectoralis minor muscle medially and exposing the Axillary vein. However, we have been performing the level 3 clearance by retracting the Pectoralis minor muscle laterally and entering the interpectoral groove.

The aim of the study was to compare the nodal yield of level 3 axillary nodes using the conventional Subpectoral route and the Interpectoral route.

Materials and Methods: A total of 50 females with Carcinoma Breast were enrolled in the study and were randomly allocated to the Subpectoral and

Interpectoral Axillary clearance group with 25 patients in each. Statistical analysis was done.

Results: The nodal yield was 20% higher in the interpectoral group as compared to the Subpectoral group which was statistically significant. The number of complications was lower in the subpectoral group. However, three patients required prolonged Axillary drainage for more than 3 days.

Conclusion: Level 3 Axillary Clearance using Interpectoral approach gives a better nodal yield and less complications as compared to the Subpectoral approach in Carcinoma breast patients.

5145

POSTER

Evaluation of Clinical Utility of Sentinel Lymph Node (SLN) Examination by One-step Nucleic Acid Amplification (OSNA) Assay in Breast Cancer

D. Takabatake¹, S. Noguchi², F. Akiyama³, N. Sato⁴, M. Tsujimoto⁵, H. Tsuda⁶, S. Nakamura⁷, T. Kamio⁸, K. Taniyama⁹, Y. Tokuda¹⁰. ¹Shikoku Cancer Center, Breast Surgery, Ehime, Japan; ²Osaka University Graduate School of Medicine, Breast Surgery, Osaka, Japan; ³Cancer Institute Japanese Foundation for Cancer Research, Pathology, Tokyo, Japan; ⁴Niigata Cancer Center Hospital, Breast Surgery, Niigata, Japan; ⁵Osaka Police Hospital, Pathology, Osaka, Japan; ⁶National Cancer Center Hospital, Pathology, Tokyo, Japan; ⁷Showa University School of Medicine, Breast Surgery, Tokyo, Japan; ⁸Tokyo Women's Medical University, Surgery, Tokyo, Japan; ⁹Tokyo Women's Medical University, Pathology, Hiroshima, Japan; ¹⁰Tokai University School of Medicine, Breast Surgery, Kanagawa, Japan

Background: Sentinel lymph node biopsy (SLNB) in Breast cancer has been commonly used in place of axillary lymph node dissection (ALND) for local control and nodal staging. Recently completed large scale clinical studies have suggested that ALND doesn't improve the prognosis of breast cancer patients for SLN positives cases significantly and can even increase patient morbidity. Therefore the significance of ALND in SLN positive patients will be called into question in the future and the value for nodal staging of SLNB will increase. The OSNA assay is utilized for lymph node examination by measuring the amount of CK19mRNA in lymph nodes using molecular techniques. The assay is expected to improve the accuracy and standardization of LN examination with the advantage of being able to assess a large portion of a lymph node. The OSNA assay has been in clinical use in Japan and EU. We conducted a multi central clinical study with 11 institutes in Japan to investigate the relationship between the results in SLN and non-SLN status as well as to identify the clinical and pathologic factors that show a correlation with non-SLN status.

Material and Methods: 417 clinically node negative breast cancer patients scheduled for SLNB were enrolled. After cutting off a central 1 mm thickness slice for pathological examinations from each SLN, the residual portion was applied to the OSNA assay. For patients who received ALND, non-SLN status was examined by routine pathological method at each site.

Results: Ninety-four of 417 (22.5%) patients were judged as SLN positive by the OSNA assay. 86 of these patients received ALND and 29 (33.7%) of these had metastases in the non-SLN. Among those who received ALND, 50 patients had an OSNA ++ result and 34 patients had an OSNA + result, with metastases in non-SLN in 22 and 6 patients respectively. Positive predictive value of OSNA ++ for non-SLN metastasis (22/50 = 44.0%) was significantly higher than that of OSNA + (6/34 = 17.6%) ($p = 0.01$). A strong association with non-SLN involvement was found in cT (cT1 vs cT2 $p = 0.0104$), pT (pT1 vs pT2/T3 $p = 0.0389$) and Grade (Grade 1 vs Grade 2/3 $p = 0.0066$).

Conclusions: SLN evaluation by OSNA can reflect non-SLN status appropriately. The semi-quantitative result of OSNA(++/+) has potential to indicate non-SLN status by combination with tumour size and tumour grade.

5146

POSTER

OSNA is Suitable for Intraoperative Analysis of Sentinel Lymph Node Metastasis in Breast Cancer

M. Suzuki¹, H. Nanjo², T. Sugiyama². ¹Kitamura Hospital, Breast Center, Yamagata, Japan; ²Akita Karyology and Histology Research Center, Pathology, Yurihonjo, Japan

Background: Sentinel node (SN) biopsy is the standard procedure in breast cancer patients who are clinically node-negative. However, our hospital has no pathologist, so we have not been able to perform intraoperative analysis of SNs. The one-step nucleic acid amplification (OSNA) technique is simple to perform and allows straightforward diagnosis of SN metastasis by quantitative evaluation of cytokeratin 19 m-RNA. The analysis results are available in as little as 30 minutes. This technique enables us to now perform SN analysis during the operation. In this study,