

study (Okusaka et al. Br J Cancer 2010). However, whilst statistically and clinically significant, these gains are modest and these studies serve only as a foundation on which to develop further treatments.

Advanced biliary tract cancer is relatively chemotherapy-sensitive and further chemotherapy studies are underway to expand chemotherapy options.

Attention has also turned to the targeting of cellular pathways pivotal to tumorigenesis including epithelial growth factor receptor (EGFR-), vascular endothelial growth factor (VEGF-) and mitogen-activated protein kinase (MEK-) inhibition, amongst others; no practice-changing phase III studies have yet been reported. Investigators continue to gain a better understanding of the underlying processes in the development and establishment of biliary tract cancers and the effect of targeting one or more of these cellular pathways. Identification of particularly active regimens will also lead to adjuvant and neo-adjuvant studies which are likely to have the greatest impact on patient survival.

### Special Session (Mon, 26 Sep, 13:15–14:15) Developments and Management of Lymphoedema

**293** INVITED  
**Lymphoedema: How the Prevalence and Severity of Lymphoedema Related to Cancer has Changed**

M. Sneddon<sup>1</sup>. <sup>1</sup>University of Glasgow, School of Nursing & Health Care, Glasgow, United Kingdom

This presentation will consider the impact that developments in surgical and imaging techniques are having on the incidence of cancer-related lymphoedema. It will also discuss emerging strategies for early detection of lymphoedema and identification and screening of those most at risk of developing lymphoedema following cancer treatment. These developments have the potential to minimise the severity of lymphoedema and reduce the burden on the patient and the health service.

Finally recent research and developments in the management of lymphoedema with potential to minimise progression and enhance quality of life will be identified.

**294** INVITED  
**Lymphoedema Management Options**

J. Simmons<sup>1</sup>. <sup>1</sup>British Lymphology Society, BLS, London, United Kingdom

Lymphoedema of the limbs, trunk and genitals is a common side effect following treatment for breast, gynaecological, urological, head and neck and aggressive skin cancers and in sarcomas where radical radiotherapy has been delivered.

For many years treatment for lymphoedema has centred around the 'Four cornerstones' of management namely care of the skin, exercise and positioning, lymphatic drainage and compression either in the form of hosiery of multi layered lymphoedema bandaging; and on a two phase intensive then maintenance programme.

However recent advances have been made in understanding who might be more at risk of lymphoedema together with further treatment options. There is a growing understanding of the role of low level laser therapy, intermittent pneumatic compression devices and early intervention programmes. The role of liposuction for resistant lymphoedema is becoming commonplace where conventional treatments have failed. We have moved on from the Four Cornerstones to be able to offer our patients more choice and autonomy in self management in their own home in line with the survivorship agenda.

This special session will give an overview of the current evidence for all the major therapies used in lymphoedema management.

**295** INVITED  
**The Management of Lymphoedema in Advanced Cancer**

A. Honnor<sup>1</sup>. <sup>1</sup>LOROS Hospice, Lymphoedema, Leicester, United Kingdom

Lymphoedema is a problem commonly experienced by patients with advanced cancer. There are several factors which may contribute to the onset of lymphoedema in these patients including: surgery and radiotherapy, metastatic lymphadenopathy, tumour recurrence, infection or inflammation, reduced mobility and function, effects of medication, hypoalbuminaemia and venous thrombosis.

Symptoms commonly experienced include: swelling of one or more limbs which can extend into the trunk, genitals and head or neck; ulceration;

tension of the affected tissues; heaviness of the limb causing impaired mobility, function and sensation and infection. Up to 67% of patients with advanced cancer experience pain due to oedema and lymphorrhoea (leakage of lymph fluid) can be a further complication.

Cancer patients with lymphoedema can experience a wide range of physical, psychological and social problems which can have a significant impact on their quality of life. A holistic assessment is the first step in understanding the patient's main concerns and priorities with regard to managing their swelling. However, it is important to remember that the burden of treatment on the patient should not exceed the benefit gained and, therefore, treatment will generally have to be adapted and modified to suit patients' individual needs.

This presentation will examine the management of lymphoedema in the patient with advanced cancer.

### Special Session (Mon, 26 Sep, 13:15–14:15) Controversies in the Management of Cervical Cancer

**296** INVITED  
**Neoadjuvant Chemotherapy in Locally Advanced Cervical Cancer**

N. Colombo<sup>1</sup>, M. Peiretti<sup>1</sup>, G. Parma<sup>1</sup>, R. Mancari<sup>1</sup>, A. Garbi<sup>1</sup>. <sup>1</sup>European Institute of Oncology, Department of Gynaecology, Milan, Italy

Cervical cancer is the second most common cause of female cancer mortality worldwide. In 1999 the National Cancer Institute (NCI) announcement modified the paradigm of locally advanced cervical cancer (LACC) treatment toward concurrent chemo-radiotherapy, in view of the publication of 5 randomized trials that demonstrated a survival advantage with the addition of cisplatin-based chemotherapy to radiotherapy alone. However, the lack of radiotherapy departments, especially in developing countries, the presumed high incidence of long-term complications and the poor control of metastatic disease have brought about the development of different therapeutic approaches such as neoadjuvant chemotherapy (NACT) followed by surgery.

A meta-analysis of NACT followed by radical hysterectomy showed an absolute improvement of 14% in 5 year survival compared to radiotherapy. Out of more than 800 patients evaluated in this meta-analysis, 441 were from an Italian randomized trial that compared NACT followed by surgery (group 1) with radiotherapy (group 2). Patients with FIGO stage IB2-IIB had 5-years overall and progression free survival significantly longer in group 1. To date, there are not enough data that would support the "best" neoadjuvant chemotherapy regimen. The two randomized phase II studies, SNAP01 and SNAP02, showed that TIP or TEP (epirubicin for adenocarcinoma tumours) provide an average of 42–48% optimal response rate, defined as no residual or residual disease with  $\leq 3$  mm stromal invasion. In these studies, the achievement of an optimal response rate as above described was the strongest predictor of survival and therefore this endpoint could be considered as a surrogate in trials investigating new therapeutic options. The Cochrane metanalysis indicated that trials using cycle lengths shorter than 14 days or the cisplatin dose greater than 25 mg/m<sup>2</sup> per week exhibited a greater advantage on the survival of patients with LACC. It indicates that the timing and dose intensity may greatly impact the curative effect of NACT. It seems reasonable therefore to explore dose-dense regimens in this setting.

In conclusion from the data of the literature NACT followed by radical surgery could have an important role in the treatment of LACC, but the appropriate indications and contraindications need to be better identified. Promising results have been already published, but the final answer will be given by the randomized clinical trial carried out by the European Organization for Research in Cancer Therapy (EORTC 55994), which will hopefully demonstrate whether or not NACT followed by surgery will display a better oncological outcome compared to chemo-radiotherapy for IB2-IIB cervical cancer patients.

**297** INVITED  
**Image-guided Adaptive Brachytherapy (IGABT) and External Radiotherapy in Patients With Cervix Cancer**

C. Haie-Meder<sup>1</sup>, R. Mazon<sup>1</sup>, I. Dumas<sup>2</sup>, C. Lhomme<sup>3</sup>, P. Morice<sup>4</sup>. <sup>1</sup>Institut Gustave Roussy, Service de Curietherapie, Villejuif, <sup>2</sup>Institut Gustave Roussy, Physics Department, Villejuif, <sup>3</sup>Institut Gustave Roussy, Gynaecological Service, Villejuif, <sup>4</sup>Institut Gustave Roussy, Gynaecological Surgery, Villejuif, France

Concomitant chemoradiation followed by intrauterine brachytherapy (BT) represents the standard of treatment in patients with advanced cervical

cancers. Although the addition of concomitant chemotherapy has improved survival, therapeutic challenges remain, especially in locally advanced disease. Traditionally, prescription and treatment planning in intracavitary BT for cervix cancer have used either reference points (mainly points A and B) or reference isodoses (60 Gy according to ICRU recommendations). Up until recently, when point A was not used for prescription, target volume (CTV) was assessed mainly by clinical examination. Doses to critical organs were reported at bladder and rectum ICRU points. This practice has been supported by a long-standing clinical experience that has yielded an acceptable therapeutic ratio.

The recent development of imaging has contributed to the improvement in target and organs at risk knowledge. In 2005 and 2006, GEC-ESTRO recommendations publications on image guided adaptive BT (IGABT) have defined the different volumes of interest. These recommendations have been validated with intercomparison delineation studies. IGABT is based on an adaptive 4D target concept and requires the use of imaging (preferably MRI) with the BT applicator in place. With the concomitant development of remote after-loading projectors, provided with miniaturized sources, it is now possible to plan radiation doses by adjusting dwell positions and relative dwell time values. These procedures allow better coverage of the targets while sparing OAR. Tumour visualization has also led to the development of new applicators, allowing the combination of endocavitary and interstitial brachytherapy when the CTVs cannot be properly covered without a dose increase to organs at risk. The aim of these applicators is to perform interstitial brachytherapy with a limited number of needles, with a short distance between the needles and the CTVs.

Several teams have published a significant improvement in the HR-CTV coverage without increase in doses to the organs at risk. Some authors found a D2cc dose reduction of  $7 \pm 6 \text{ Gy}/\beta 3$  for the bladder and  $7 \pm 6 \text{ Gy}/\beta 3$  for the rectum. D2cc dose reduction was also observed in the sigmoid. The recent literature data evidenced a significant improvement in local control without increase in complications. This is particularly true for tumours larger than 5 cm at diagnosis with an improvement in overall survival as high as 30%. So far, the recommended doses to the HR-CTV have not clearly defined even if some teams have found a cut off level at  $87 \text{ Gy}/\beta 10$ .

In France, between 2005 and 2007, 705 patients were included in the first prospective non-randomized multi-centre study based on a national grant: "Soutien aux Techniques Innovantes et Coûteuses" (STIC). The comparison was performed between 2D brachytherapy and IGABT. Local control at 2 years was improved from 74 to 79% and G3-4 morbidity reduced from 23 to 3% when comparing 2D brachytherapy and IGABT, without significant effect on disease free survival.

More recently, a prospective non randomized observational study, the so-called EMBRACE study registers IGABT since July 2008. So far, 20 centers have included 407 patients with cervix cancer. This study evidences the tendency to increase the indications of interstitial brachytherapy: 29% with the new applicators, allowing limited number of needles in the parametrium. The mean dose to the HR-CTV was  $88.6 \text{ Gy}/\beta 10 \pm 12.5 \text{ Gy}/\beta 10$  and the mean dose to the IR-CTV was  $68.7 \text{ Gy}/\beta 10 \pm 6.8 \text{ Gy}/\beta 10$ . These very preliminary results evidence the attempt to increase the dose to the CTVs. This study is also a very appealing approach, as it will allow the comparison of all the biologically equivalent doses. All the different teams included in this prospective study will be able to compare their results according to the doses. This prospective study will help in better defining the dose recommended in both tumour and critical organs and will provide very important information in DVH parameters.

298

INVITED

### Management Aimed at Fertility Preservation

F. Amant<sup>1</sup>. <sup>1</sup>University Hospitals Gasthuisberg, Gynaecologic Oncology, Leuven, Belgium

More than 25% of women with cervical cancer are under 40 years of age and the age of nulliparous women has increased in developed countries. Prior to assess the possibility to preserve fertility, the motivated patient should have realistic expectations. Fertility decreases with age and together with the availability of a partner, the chance to try to conceive should be part of the decision tree. A fertility sparing strategy is ideally executed in a motivated patient who is likely to try to conceive subsequently. Only few centers worldwide have sufficient experience to do these procedures on a regular basis. Most centers should call their approach experimental and time to seek for second opinion is available. Standard treatment of stage Ia1 cervical cancer (microinvasive) is treated by conisation and results in normal fertility. Standard treatment for higher stages is Wertheim-Meigs hysterectomy or radiochemotherapy. Tumour characteristics that allow consideration of fertility preservation include stage Ib1 smaller than 2 cm with negative pelvic lymph nodes. Magnetic resonance imaging enables adequate estimation of tumour diameter. When the decision is taken to try to preserve fertility, determination of lymph node status enables to identify

high risk disease that needs standard treatment. When lymph nodes are negative two options exist. Firstly, radical trachelectomy which consists of removal of the cervix, parametria and vaginal cuff. This procedure requires sufficient surgical expertise in order to be executed radically and safe. Preliminary findings of less radical procedures (ie, deep cone and simple trachelectomy) in patients with tumours less than 2 cm are comparable with the results of radical trachelectomy. Secondly, neoadjuvant chemotherapy can be administered prior to surgery in order to reduce the amount of tissue (downstaging) that needs to be resected, by a less radical trachelectomy or conisation. It appears that chemotherapy does not affect fertility and chance to conceive. Although the experience is less well reported, this latter option results in a better obstetrical outcome, with less preterm delivery and more live births.

## Special Session (Mon, 26 Sep, 13:15–14:15) Hitting the Right Pathway: New Drugs for New Targets

299

INVITED

### MET Inhibitors

Abstract not received

300

INVITED

### RAF Inhibitors

F. Di Nicolantonio<sup>1</sup>. <sup>1</sup>Institute for Cancer Research and Treatment, University of Torino School Medicine, Candiolo (Torino), Italy

In the last decade, RAF kinases have emerged as valuable druggable targets, in particular after the discovery of the prevalence of oncogenic BRAF mutations in a number of solid tumours, including cutaneous melanoma, colorectal and papillary thyroid cancers. The BRAF V600E aminoacid substitution accounts for the majority of alterations and confers constitutive kinase activity. Among RAF inhibitors undergoing preclinical or clinical development, some display some preferential selectivity towards the mutant isoform (GSK2118436, vemurafenib), while others have similar activity on both mutated and wild type BRAF or other RAF isoforms.

The first targeted drug expected to reach clinical use is vemurafenib (known also as PLX4032, RO5185426, RG7204) that has demonstrated remarkable efficacy in melanoma patients carrying BRAF V600E mutated tumours. Initial enthusiasm has been dampened by the relatively short duration of clinical response. Indeed, secondary resistance to this drug emerges in patients who relapse after a median time of 6–7 months. Acquired resistance does not seem to be mediated by additional 'gatekeeper' mutations in the BRAF target gene itself. Rather, a number of studies on cell lines and melanoma samples indicate that tumours become resistant by modulating the expression or activity of other genes (such as NRAS, CRAF, COT, MEK1, PDGFR, IGF1R, PTEN) that compensate BRAF inhibition or obviate for the need of BRAF signaling. Consequently, combinatorial strategies with MEK or PI3K pathway targeted drugs or single molecule dual RAF/MEK inhibitors (such as RO5126766) are being evaluated to delay the onset of secondary resistance and improve patient survival.

Interestingly, not all BRAF V600E mutated tumours are sensitive to vemurafenib, with 15–20% melanoma cases and the vast majority of BRAF mutant colorectal cancer patients displaying intrinsic resistance to treatment with this inhibitor. The molecular bases of these partial failures represent an area of active research investigations.

Finally, newer compounds such as PLX-PB-4 and ARQ680 (and its prodrug ARQ736) are being developed to overcome the limitations of first-generation drugs, and, in particular, their ability to promote dimerization of RAF family members and paradoxical activation of MAPK pathway in treated cells bearing oncogenic or normally activated RAS. It is likely that this could be achieved by designing inhibitors with increased potency against CRAF and/or with the ability to block RAF dimerization.

301

INVITED

### ALK in NSCLC – Paving the Way for Oncogene-directed Therapy

R. Doebele<sup>1</sup>. <sup>1</sup>University of Colorado Cancer Center, Division of Medical Oncology, Aurora CO, USA

The discovery of distinct subsets of non-small cell lung cancer (NSCLC) as defined by the presence of molecular oncogenes has greatly impacted oncogene-directed therapy for this disease. Gene fusions involving the kinase domain encoding region of the ALK gene were first reported in NSCLC in 2007 and displayed oncogenic properties. The tyrosine kinase inhibitor, crizotinib, has shown remarkable response rates and progression