

presentation is advanced or symptom burden is great. This transformation has led to some concern that contemporary palliative care has lost clarity over its own position in relation to death and dying. In this presentation, a debate on the concept of dying well in an oncology context will be offered. Using exemplars for recently conducted research in the EU and Ireland, the key questions which underpin the concept of a good death will be discussed. The international evidence on how cancer patients see a good death will be presented as well as the challenges facing clinicians who work at the acute intervention level when it becomes clear that different goals and objectives of care are indicated. Based on a recent case study, the presentation will conclude with an interpretation of how a palliative care response can be built into acute oncology practice which may be of benefit to patients with advanced disease and their families.

410 INVITED Nurses Attitudes Towards Caring for Dying Patients in Acute Settings

M. Arantzamendi Solabarrieta¹, A. Richardson², J. Addington-Hall³.

¹University of Navarra, Faculty of Nursing, Pamplona, Spain; ²Southampton University Hospitals NHS Trust and the University of Southampton, School of Health Sciences, Southampton, United Kingdom; ³University of Southampton, School of Health Sciences, Southampton, United Kingdom

There is a need to better understand nurses' experiences and factors that determine these. The evidence on nurses' views about their experiences of caring for the terminally ill in the hospital setting described them as holding largely negative views. Factors that might shape these negative views have been mentioned but have not been studied statistically. If we are able to understand the factors influencing nurses' experiences and on how they go about their daily work with terminally ill patients, we might be able to modify these and effect how they interact and care for these patients. The research reported here about Spanish hospital nurses caring for terminally ill patients, starts to tackle this subject.

A cross-sectional postal survey was conducted involving 165 hospital nurses working on acute wards in 6 hospitals of one region in Spain (65% response rate).

Nurses valued the learning experience of caring for terminally ill patients and did not transmit much discomfort regarding death and dying. They perceived the care of terminally ill patients to be more demanding and challenging than caring for other patients. They were not highly motivated to care for this group.

Focusing on the multivariate analysis of factors that influence the challenges, nurses' perceive they face aspects such as degree of discomfort with death and dying, whether they view it to be a learning experience, their competence to care emotionally and their disposition to involve relatives were among the statistically influential factors.

The results point to the complexity of nurses' experiences of caring for terminally ill patients and the factors underlying these. Nurses' perceived competence to care emotionally for terminally ill patients was revealed to be a key factor influencing nurses' experiences. Tackling nurses' perceived ability to care emotionally for terminally ill patients should form a priority if we aim to improve the care they provide to terminally ill patients. On the other hand, as nurses' inclination to involve relatives decreased, they expressed fewer challenges in caring for patients. This suggests interactions with relatives may have an significant impact on nurses' experiences and more than that expected on the basis of the literature. Further research is needed to understand the nurse-terminally ill patient's relative relationship.

Special Session (Tue, 27 Sep, 11:30–12:30) Surgical Treatment in Breast Cancer Patients With Distant Metastases

411 INVITED Surgery of the Primary Tumour

Abstract not received

412 INVITED Surgery of Liver and Pulmonary Metastases

Abstract not received

Special Session (Tue, 27 Sep, 11:30–12:30) Desmoid Fibromatosis Tumours Representing an Unmet Medical Need

413 INVITED Desmoid Type Fibromatosis: How Much Surgery and When?

A. Gronchi¹. ¹Istituto Nazionale Tumori, Department of Surgery, Milan, Italy

The first approach to desmoid type fibromatosis (DFs) today should routinely be a watchful surveillance. This policy could be the best way to select patients who really need a therapy, surgical resection included, from those who don't.

Infact DFs are made by different diseases. There seems to be a subset with favourable behaviour. Spontaneous regressions and long lasting stabilization have been repeatedly reported in up to 50% of primary cases. This subset maybe be constituted by tumours characterized by a different molecular profile, even if with an undistinguishable morphology. This hypothesis has recently been supported by the finding that a particular beta-catenin gene mutation subtype could correlate with the outcome. The results of this study, though need to be prospectively validated, look very attractive.

Beside more favourable diseases, there are also aggressive ones for which a treatment needs to be considered. Before resorting to surgery, anti-inflammatory drugs, hormonal therapies and sometimes low-dose chemotherapy are generally considered. Infact several times surgery induce more morbidity than the disease itself. This may not be true for all locations. On the one end patients affected by abdominal wall tumours may well prefer a surgical resection than a chronic therapy. Cosmesis is generally not affected and results are usually good (recurrence rate for tumours at this site being very low). On the other end, patients affected by intra-abdominal, head & neck and intrathoracic tumours are offered more often surgical resection, since the potential life-threat progression of their disease may determine.

When surgery is planned, the resection should not be accomplished with the same stricht principles of sarcoma surgery. Indeed many authors have claimed that the outcome of primary disease is quite unpredictable and not influenced by surgical margins. Therefore pursuing wide margins resection in primary surgery should always be weighted with function preservation and cosmesis.

This issue, along with the natural history of the disease, the lack of metastatic potential, the possible implication of inflammatory agents in further tumour re-growth have all become arguments in favour of a less and less aggressive primary approach.

At recurrence repeated surgical excisions are usually not the first choice and again preservations of function and cosmesis are always to be carefully taken into account.

414 INVITED Radiotherapy as Adjuvant Treatment to Surgery or as Definitive Treatment?

V. Budach¹. ¹Budach, Klinik für Strahlentherapie CCM/CKV, Berlin, Germany

Desmoid tumours need more extensive surgery than STS of the same size due to their infiltrating growth pattern needing in order to achieve clear surgical margins. Despite of adequate surgery local recurrence rates are disappointing. A specific tumour grading like for STS does not exist. However, the tumours do not de-differentiate with increasing numbers of recurrences. Lethality is low and only apply to FAP with 10%. R0-Surgery and Wait & See (W&S) for unresectable cases was the standard of care for many decades. However, during the last decade primarily wait & see policies and also neoadjuvant treatments have gained increasing importance. This conservative approach can lead to long periods of stable disease in up to 80% of all cases and is lacking lethality. The importance of the surgical margin status remains so far unclear and is the case of further debates. Many factor may be attributed to the inconsistent evidences for this disease: Most studies were small in number and therefore lack statistical significance, the margin status did not always imply meticulous pathologic work-up.

Postoperative radiotherapy might have offset the detection of the margin impact. The postoperative release of tissue-derived growth factors might be the reason for the observation, that a considerable number of desmoids recur to a certain degree and thereafter remain stable for many years. Two groups of patients are distinguishable: A slow and a relatively faster proliferation subgroup, which needs more than a W&S policy. The DFS after surgical resection and non-surgical treatment seems to be equivalent. From more recent publications RTX as a conservative treatment approach seems to be beneficial when added postoperatively or used definitively for primarily/recurrent or operable/inoperable disease. A comparative review of