

Adducts of Ferrocenylboranes and Pyridine Bases: Generation of Charge-Transfer Complexes and Reversible Coordination Polymers

Marco Fontani, Frank Peters, Wolfgang Scherer, Wolfgang Wachter, Matthias Wagner*, and Piero Zanello

Eur. J. Inorg. Chem. **1998**, 1453–1465

Table 1 contains a wrong header for the last three lines; the correct Table 1 should read as follows:

Table 1. Infrared bands [cm^{-1}] of selected compounds **1**·Do and **2**·(Do)₂ (KBr disc) in the characteristic regions 1650–1450 cm^{-1} (C=C and C=N stretch) and 1100–1050 cm^{-1} (B–N vibration)

1 ·pic	1 ·DMAP	2 ·(pic) ₂	2 ·(DMAP) ₂	1 ·bipy· 1	[2 ·bipy] _n
1628	1636	1628	1638	1622	1622
—	1544	—	1552	1540	1540
1507	—	1508	—	1493	1493
1104	1104	—	—	1100	—
1068	1076	1066	1077	1075	1075
1058	1058	1047	1060	1058	1061

pic	DMAP	bipy	bipyPr ₂ ^[a]
1606	1603	1590	1645
1563	1537	1531	1569
1497	1519	1487	1509

^[a] *N,N'*-Bis(*n*-propyl)-4,4'-bipyridinium hexafluorophosphate.

The Editor
[C 98092]