

Binary [Hydrotris(indazol-1-yl)borato]metal Complexes, $M(\text{Tp}^{4\text{Bo}})_2$ with $M = \text{Fe}, \text{Co}, \text{Ni}, \text{Cu},$ and Zn : Electronic Properties and Solvent-Dependent Framework Structures through $\text{C}-\text{H}\cdots\pi$ Interactions

Christoph Janiak,* Savaş Temizdemir, Sebastian Dechert, Werner Deck,
Frank Girgsdies, Jürgen Heinze, Mario J. Kolm, Tobias G. Scharmann,
and Oliver M. Zipffel

Eur. J. Inorg. Chem. **2000**, 1229–1241

We regret that in Table 3 on page 1232 the isomer shifts at 290 K were given incorrectly. Also $\Gamma/2$ is actually Γ . The correct Table is shown below. Changes are highlighted in **bold**.

Table 3. Parameters of ^{57}Fe -Mössbauer spectra for the iron complex **4** (cf. Figure 3)

T/K [$^{\circ}\text{C}$]	N_{lin} ^[a]	$\delta_{\alpha\text{-Fe}}$ ^[b] /mm s ⁻¹	Γ ^[c] /mm s ⁻¹	ΔE_{Q} ^[d] /mm s ⁻¹	$A_{\text{r}}/\%$ ^[e] (A_{r}/A_{Σ})
290 [17] ^[f]	3	0.35	0.28	0.15	60
		0.39	0.61	1.02	28
		1.06	0.71	2.70	13
523 [250]	2	0.20	0.79	0.00	82
		0.20	0.38	1.21	18
473 [200]	2	0.21	0.54	-0.23	62
		0.24	0.56	1.16	38
287 [14]	1	0.34	0.26	0.16	100

^[a] Number of lines used to fit the spectrum. – ^[b] Isomer shift referred to α -iron at room temperature. – ^[c] **Full width at half height**. – ^[d] Quadrupole splitting. – ^[e] Line area relative to total area. – ^[f] The spectrum was measured after heating the sample to 523 K and may contain some decomposition product.

C. Janiak
[C 199436]