

Efficient Asymmetric Synthesis of Δ - and Λ -Enantiomers of (Bipyridyl)ruthenium Complexes and Crystallographic Analysis of Δ -Bis(2,2'-bipyridine)(2,2'-bipyridine-4,4'-dicarboxylato)ruthenium: Diastereoselective Homo- and Heterochiral Ion Pairing Revisited

Régis Caspar, Hani Amouri,* Michel Gruselle, Christine Cordier, Bernard Malézieux, Raphaël Duval, and Hubert Leveque

Keywords: Ruthenium / Chiral resolution / Circular dichroism

*In our original article,^[1] the captions for Figures 2, 3 and 4 (page 501) are incorrect. We wrongly attributed the CD curves with the solid line to the Δ -enantiomers **4a**, **4a** and **5a** and those with dotted lines to the Λ -enantiomers **3b**, **4b** and **5b**. The correct captions are as follows:*

Figure 2. CD curves for **3a** (dotted line) and **3b** (solid line) in acetonitrile

Figure 3. CD curves for **5a** (dotted line) and **5b** (solid line) in acetonitrile

Figure 4. CD curves for **4a** (dotted line) and **4b** (solid line) in acetonitrile

^[1] R. Caspar, H. Amouri, M. Gruselle, C. Cordier, B. Malézieux, R. Duval, H. Leveque, *Eur. J. Inorg. Chem.* **2003**, 499–505.

Received July 3, 2003
H. Amouri