

μ - η^1 : η^1 -*N,N'*-Imidazolidine-Bridged Dicopper(II/II) Complexes of a New Dinucleating μ -Bis(tetradentate) Schiff Base Ligand: Synthesis, Structural Characterization, ^1H NMR Spectroscopy, and Magnetic Coupling

Manindranath Bera,^[a] Wing Tak Wong,^[b] Guillem Aromí,^[c] and Debashis Ray^{*[a]}

Keywords: N,O ligand / Imidazolidine / Copper complexes / Magnetic properties

The dicopper oxidation state in the title of the original article^[1] is erroneously given as (II/III); however, it should be (II/II) as shown in the correct title above.

[1] M. Bera, W. T. Wong, G. Aromí, D. Ray, *Eur. J. Inorg. Chem.* **2005**, 2526–2535.

Received: June 27, 2005
The Editors

[a] Department of Chemistry, Indian Institute of Technology, Kharagpur 721 302, India,
E-mail: dray@chem.iitkgp.ernet.in

[b] Department of Chemistry, University of Hong Kong, Pokfulam Road, Pokfulam, Hong Kong SAR, P.R. China

[c] Departament de Química Inorgànica, Universitat de Barcelona, Diagonal, 647, 08028 Barcelona, Spain