

The Hexakis(thiocyanato)ferrate(III) Ion: a Coordination Chemistry Classic Reveals an Interesting Geometry Pattern for the Thiocyanate Ligands

Anthony W. Addison,^{*,[a]} Raymond J. Butcher,^[b] Zoltán Homonnay,^[c]
Vitaly V. Pavlishchuk,^{*,[d]} Michael J. Prushan,^[e] and Laurence K. Thompson^[f]

Keywords: Iron / Thiocyanate / X-ray diffraction / Magnetic properties / IR spectroscopy / Moessbauer spectroscopy / Coordination modes / Isomers

We are grateful to Professor Ulrich Müller, of Philipps-Universität Marburg, for drawing the attention of ourselves and the referees to a prior structure report for $(\text{NMe}_4)_3\text{[Fe(NCS)}_6\text{]}$.^[1] The 1977 and 2005^[2] reports differ only in the choice of unit cell origin. The conclusions from the structure–spectra relationships discussed in the 2005 paper remain the same.

[1] U. Müller, *Acta Crystallogr.* **1977**, B33, 2197–2201.

[2] A. W. Addison, R. J. Butcher, Z. Homonnay, V. V. Pavlishchuk, M. J. Prushan, L. K. Thompson, *Eur. J. Inorg. Chem.* **2005**, 2404–2408.

Received: September 26, 2005
The Authors

[a] Chemistry Department, Drexel University,
Philadelphia, PA 19104-2875, USA

[b] Chemistry Department, Howard University,
Washington, DC 20059, USA

[c] Department of Nuclear Chemistry, Eötvös Loránd University,
Pázmány P. s 1/A, 1117 Budapest, Hungary

[d] L.V. Pisarzhevskii Institute of Physical Chemistry, Ukrainian
Academy of Sciences,
Prospekt Nauki 31, Kiev, 03028 Ukraine

[e] Department of Chemistry & Biochemistry, LaSalle University,
Philadelphia, PA 19141, USA

[f] Chemistry Department, Memorial University,
St. John's NFD, A1B 3X7 Canada