

Dioxovanadium(V) Complexes of ONO Donor Ligands Derived from Pyridoxal and Hydrazides: Models of Vanadate-Dependent Haloperoxidases

Mannar R. Maurya,^{*,[a]} Shalu Agarwal,^[a] Cerstin Bader,^[b] and Dieter Rehder^{*,[b]}

Keywords: Vanadium / Hydrazones / Catalytic bromination / Pyridoxal

Figure 1 has been depicted twice in the original article (as Figure 1 and Figure 5).^[1] For the correct Figure 5, see below.




Figure 5. Titration of $[K(H_2O)_2][VO_2(pydx-bhz)]$ (**8**) with a saturated solution of HCl in MeOH; the spectra were recorded after addition of 2-drops portions MeOH/HCl to 10 mL of ca. 10^{-4} M solution of **8** in MeOH

[1] M. R. Maurya, S. Agarwal, C. Bader, D. Rehder, *Eur. J. Inorg. Chem.* **2005**, 147–157.

Received: January 18, 2005
The Editors

^[a] Department of Chemistry, Indian Institute of Technology Roorkee, Roorkee 247667, India
E-mail: rkmanfcy@iitr.ernet.in

^[b] Institut für Anorganische und Angewandte Chemie, Universität Hamburg, Martin-Luther-King-Platz 6, 20146 Hamburg, Germany
E-mail: dieter.rehder@chemie.uni-hamburg.de