

# Phosphavinylidene(oxo)phosphorane Mes\*P(O)=C=PMes\*: A Diphosphaallene Featuring $\lambda^5\sigma^3$ - and $\lambda^3\sigma^2$ -Phosphorus Atoms

Raluca Septelean,<sup>[a]</sup> Henri Ranaivonjatovo,<sup>\*,[b]</sup> Gabriela Nemes,<sup>[a]</sup> Jean Escudié,<sup>\*,[b]</sup> Ioan Silaghi-Dumitrescu,<sup>[a]</sup> Heinz Gornitzka,<sup>[b]</sup> Luminita Silaghi-Dumitrescu,<sup>[a]</sup> and Stéphane Massou<sup>[c]</sup>

**Keywords:** Heterocumulenes / Phosphorus / Phosphaalkenes / Methylene phosphoranes

Bis(2,4,6-tri-*tert*-butylphenyl)phosphavinylidene(oxo)phosphorane (**1**) (Mes\* = 2,4,6-tri-*tert*-butylphenyl), the first  $\lambda^5\sigma^3,3\lambda^3\sigma^2$ -diphosphaallene, has been synthesised by addition of the phosphaalkenyllithium Mes\*P=C(Li)Cl to dichlorophosphane oxide Mes\*P(O)Cl<sub>2</sub>. Compound **1** is stable in the solid-state but undergoes a slow intramolecular cyclisation in solution by addition of a C–H bond of an *ortho-tert*-butyl group to the  $\sigma^3$ -P=C double bond. Water adds chemoselectively and regioselectively to the  $\lambda^5\sigma^3$ -P=C bond,


which appears to be more reactive than the  $\lambda^3\sigma^2$ -P=C bond. This leads to the hydrolysis compound **8** which has been structurally characterised. The new bis(phosphane oxide)-methylene **9** formed from the successive cyclisation of an *ortho-tert*-butyl group to the  $\lambda^5\sigma^3$ -P=C bond and addition of water to the  $\lambda^3\sigma^2$ -P=C bond has also been structurally characterised.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2006)

## Introduction

The high utility of ketenes, allenes and cumulenes has been well established in organic synthesis.<sup>[1]</sup> Phosphaallene derivatives, namely compounds with P=C=C or P=C=P skeletons, are phosphorus congeners of allenes and can thus be expected to be useful starting materials for new organophosphorus compounds.

Among the possible neutral heterocumulenes featuring the P=C=P sequence,  $1\sigma^4,3\sigma^4$ -diphosphaallenes **I**<sup>[2]</sup> and  $1\sigma^2,3\sigma^2$ -diphosphaallenes **II**<sup>[3,4]</sup> have been known for many years. More recently  $1\sigma^4,3\sigma^2$ -diphosphaallenes **III** have been isolated.<sup>[5]</sup>



Here we report the synthesis and the first aspects of the chemical behaviour of a hitherto unknown  $1\sigma^3,3\sigma^2$ -diphosphaallene **IV**. Despite continuing efforts over the past two decades to synthesise low-coordinate multiply bonded phosphorus compounds, such a functionalised molecule has

never been described even as an intermediate. Moreover, such a structure is comprised of a methylene(oxo)phosphorane unit of which only two stable –P(O)=C< representatives have been reported to date.<sup>[6]</sup>

## Results and Discussion

Mes\*P(O)=C=PMes\*, bis(2,4,6-tri-*tert*-butylphenyl)phosphavinylidene(oxo)phosphorane **1** (Mes\* = 2,4,6-tri-*tert*-butylphenyl), was obtained by addition, at –80 °C, of one equiv. of phosphaalkenyllithium **2**<sup>[7]</sup> to a solution of dichlorophosphane oxide **4**.<sup>[8]</sup> <sup>31</sup>P NMR spectroscopic analysis of the crude mixture showed the presence of **3**, **4** and **1** in a ratio of 1:1:1. The one-pot preparation of **1** probably involves the preliminary formation of phosphaalkenylphosphane oxide **5** followed by a rapid CCl/CLi exchange with phosphaalkenyllithium **2** leading to the starting dichlorophosphaalkene **3** and intermediate **6**. The latter loses LiCl to afford the title compound **1** (Scheme 1), which could be easily obtained as a yellow powder from the reaction mixture by precipitation with pentane. The structure of **1** was unambiguously determined by sophisticated <sup>1</sup>H, <sup>13</sup>C and <sup>31</sup>P NMR spectroscopic studies (1D selective <sup>31</sup>P decoupling experiments and 2D HSQC, HMBC) (see supporting information). The <sup>31</sup>P NMR spectrum of **1** displays an AX system at  $\delta$  = 261.4 ( $\sigma^2$ -P) and 117.3 ppm ( $\sigma^3$ -P) with a coupling constant <sup>2</sup>J<sub>P,P</sub> of 30.5 Hz. A large high-frequency shift of 120 ppm can be observed for the  $\lambda^3\sigma^2$ -P in comparison with the value reported for the diphosphaallene Mes\*P=C=PMes\* ( $\delta^{31}\text{P}$  = 140 ppm<sup>[4b]</sup>). The signal of the

[a] Babes-Bolyai University,

Kogalniceanu 1, 400084 Cluj-Napoca, Romania

[b] Hétérochimie Fondamentale et Appliquée, UMR 5069, Université Paul Sabatier,


118 route de Narbonne, 31062 Toulouse cedex 9, France

[c] Structure Fédérative Toulousaine en Chimie Moléculaire, FR 2599, Université Paul Sabatier,

118 route de Narbonne, 31062 Toulouse cedex 9, France

Supporting information for this article is available on the WWW under <http://www.eurjic.org> or from the author.


$\lambda^5\sigma^3$ -phosphorus atom is slightly lower in frequency in relation to those observed for  $\text{Mes}^*\text{P}(\text{O})=\text{C}(\text{R})\text{SiMe}_3$  [ $\delta^{31}\text{P}$  = 153.7 ppm ( $\text{R} = \text{Ph}$ ),<sup>[6a]</sup> 161.1 ppm ( $\text{R} = \text{Me}_3\text{Si}$ )<sup>[6b]</sup>]. The  $\text{sp}$  hybridised carbon atom gives rise to a doublet of doublets at  $\delta = 229.1$  ppm ( $^1J_{\text{C,P}(\text{O})} = 177.4$ ,  $^1J_{\text{C,P}} = 46.8$  Hz). The  $^1\text{H}$  NMR spectrum at room temperature shows slow rotation of the  $\text{Mes}^*$  group bonded to the  $\lambda^3\sigma^2$ -phosphorus atom. Dynamic  $^1\text{H}$  NMR spectroscopy was performed between 213 and 328 K and the coalescence temperatures for the *o*-*t*Bu groups and for the aromatic protons on the  $\text{Mes}^*$  group bonded to the  $\lambda^3\sigma^2$ -P are 268 and 253 K, respectively, giving a rotation barrier of 12.2 kcal mol<sup>-1</sup>. Similar values have been reported for the rotation barriers of  $\text{Mes}^*$  groups in the phospharsaallene  $\text{Mes}^*\text{P}=\text{C}=\text{AsMe}_3$  (13.6 kcal mol<sup>-1</sup>)<sup>[9]</sup> and in the diphosphaallene  $\text{Mes}^*\text{P}=\text{C}=\text{PMes}^*$  (14.0 kcal mol<sup>-1</sup>).<sup>[10]</sup> Hindered rotation can be observed for the  $\text{Mes}^*$  group bonded to the P(O) moiety even at 328 K (two singlets for the *o*-*t*Bu groups and two doublets of doublets for the aromatic protons). A molecular ion peak can be observed in the mass spectrum of **1** with the expected fragments at  $m/z = 565$  [ $\text{M} - \text{Me}$ ]<sup>+</sup> and 523 [ $\text{M} - t\text{Bu}$ ]<sup>+</sup>.



Scheme 1. Synthesis of  $\text{Mes}^*\text{P}(\text{O})=\text{C}=\text{PMes}^*$  (**1**).

Compound **1** is stable for months at room temperature in the solid-state but attempts to crystallise it in various solvents at low temperatures did not afford satisfactory single crystals for X-ray diffraction studies. Compound **1** slowly rearranges within one day at room temperature in solution. The rearrangement affords the phosphalkenylphosphane oxides **7a** and **7b** in near quantitative yield in the form of two geometric isomers in the ratio of 85:15 (Scheme 2). B3LYP/6-31G(d)//ONIOM B3LYP/6-31G(d):PM3 calculations support this easy isomerisation and **7** was found to be lower in energy than **1** by about 54 kcal mol<sup>-1</sup>.

The  $^{31}\text{P}$  NMR spectra of **7a** and **7b** exhibit signals in the typical range for phosphalkenes<sup>[11a–11d]</sup> [ $\delta^{31}\text{P}$  for P=C in **7a** at  $\delta = 339.1$  ppm (dd,  $^2J_{\text{P,P}} = 70.2$  Hz,  $^2J_{\text{P,H}} = 24.4$  Hz) and in **7b** at  $\delta = 353.0$  ppm (dd,  $^2J_{\text{P,P}} = 73.3$  Hz,  $^2J_{\text{P,H}} = 21.4$  Hz, P=C)] and phosphane oxides<sup>[11c]</sup> [ $\delta^{31}\text{P}$  for P(O) in **7a** at  $\delta = 49.3$  ppm (d,  $^2J_{\text{P,P}} = 70.2$  Hz) and in **7b** at  $\delta = 38.7$  ppm (d,  $^2J_{\text{P,P}} = 73.3$  Hz)]. Similar chemical shifts at 327 and 10 ppm have been reported for  $\text{Mes}^*\text{P}=\text{CH}-\text{P}(\text{O})-\text{HMe}_3$ .<sup>[12]</sup> The *E* structure has been tentatively assigned to the major isomer **7a** on the basis of the  $^2J_{\text{P,H}}$  coupling constant. For example, in the  $\text{Mes}^*\text{P}=\text{CHR}$  system, the larger



Scheme 2. Rearrangement reaction and hydrolysis of **1**.

coupling constant always corresponds to the *E* isomer [18 Hz (*Z*) and 24.7 Hz (*E*) in  $\text{Mes}^*\text{P}=\text{C}(\text{H})\text{SiMe}_3$ ].<sup>[7a]</sup> Moreover ONIOM B3LYP/6-31G(d):PM3 geometry optimisations of **7** place the *Z* isomer higher in energy by 3.63 kcal mol<sup>-1</sup>.

This intramolecular cyclisation results from the addition of a C–H bond of an *ortho-tert*-butyl substituent of a supermesityl group to the  $\sigma^3\text{-P}=\text{C}$  double bond.<sup>[13]</sup> This rearrangement shows that the methylene(oxo)phosphorane moiety is much less stable than the phosphane oxide one. In contrast to bis(methylene)phosphoranes  $\text{RP}(\text{C}=\text{R}')_2$  which isomerise into phosphiranes,<sup>[14]</sup> the  $\text{O}=\text{P}=\text{C}$  unit in **1** does not undergo a similar cyclisation (which would lead to the three-membered ring oxaphosphirane).

The structure of **1** was also established unambiguously from its hydrolysis products. The  $\lambda^5\sigma^3\text{-P}=\text{C}$  double bond is much more reactive than the  $\lambda^3\sigma^2\text{-P}=\text{C}$  bond since one equiv. of water adds only to this double bond to afford the two geometric isomers **8a** and **8b** (Scheme 2). Again, ONIOM B3LYP/6-31G(d):PM3 optimisations show that **8b** (*E* isomer) is lower in energy by 5.27 kcal mol<sup>-1</sup>.

The  $^{31}\text{P}$  NMR spectra of **8a** and **8b** exhibit characteristic signals of phosphalkenes [ $\delta^{31}\text{P}$  **8a** = 305.5 ppm (dd,  $^2J_{\text{P,P}} = 47.3$  Hz,  $^2J_{\text{P,H}} = 24.4$  Hz, P=C), **8b** = 335.4 ppm (dd,  $^2J_{\text{P,P}} = 68.5$  Hz,  $^2J_{\text{P,H}} = 24.4$  Hz, P=C)] and phosphinic acids [ $\delta^{31}\text{P}$  **8a** = 24.6 ppm {d,  $^2J_{\text{P,P}} = 47.3$  Hz, P(O)}; **8b** = 34.9 ppm {d,  $^2J_{\text{P,P}} = 68.5$  Hz, P(O)}].

This addition is regioselective and gives, as expected, the derivative with the hydrogen atom bonded to the central  $\text{sp}^2$  carbon atom. However, since the reaction of water with **1** is slow in solution (about one day in pentane at room temperature), the competitive cyclisation products **7a,b** are also formed. Compounds **7** and **8** are obtained in the ratio **7a** (25%), **7b** (5%), **8a** (20%) and **8b** (50%). The X-ray structure of the *Z* isomer **8b** has been determined (Figure 1) and displays standard bond lengths and angles.

Addition of water to the *Z/E* mixture of **7** afforded, after one week, the new bis(phosphane oxide)methylene **9** in the form of only one diastereoisomer, which was isolated as pale yellow crystals in 85% yield. The  $^{31}\text{P}$  NMR spectrum of **9** displays two doublets at  $\delta = 52.1$  and 4.8 ppm (small coupling constant  $^2J_{\text{P,P}}$  of 2.7 Hz). In the nondecoupled  $^{31}\text{P}$  NMR spectrum, derivative **9** exhibits a doublet of doublets




Figure 1. Structure of **8** (*E*) (50% probability level). Selected bond lengths [Å] and angles [°]: P1–C1 1.650(4), P2–C1 1.822(3), P2–C2 1.828(4), P1–C20 1.847(5), P2–O1 1.510(3), P2–O2 1.550(3), O1–C1–P2 122.3(3), O1–P2–C1 108.72(17), O2–P2–C1 107.32(18), O1–P2–C2 108.39(17), O2–P2–C2 105.60(18), C1–P2–C2 116.26(19), C20–P1–C1–P2 168.0.

of triplets for the signal at 4.8 ppm ( $^1J_{\text{P,H}} = 517.4$  Hz,  $^2J_{\text{P,H}} = 2.8$  Hz,  $^2J_{\text{P,P}} = 2.7$  Hz), the very large  $^1J_{\text{P,H}}$  coupling constant being characteristic of a hydrogen bonded to a  $\lambda^5\sigma^4$ -phosphorus atom. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra display hindered rotation of the Mes\* group. Dynamic  $^1\text{H}$  NMR spectroscopy was performed between 253 and 323 K. The coalescence temperatures for the *o*-*t*Bu groups and for the aromatic protons on the Mes\* group bonded to the P(H) atom are 288 K in both cases, corresponding to a rotation barrier of 14.2 kcal mol $^{-1}$  which is 2 kcal mol $^{-1}$  higher than that of the Mes\* group bonded to the  $\lambda^3\sigma^2$ -phosphorus atom in **1**.

The X-ray structure of **9** (Figure 2) displays standard bond lengths and angles.




Figure 2. Structure of **9** (50% probability level). Selected bond lengths [Å] and angles [°]: P1–C1 1.827(3), P2–C1 1.822(3), P2–C2 1.819(3), P2–C9 1.794(4), P1–C20 1.818(4), P1–O1 1.482(2), P2–O2 1.481(2), P1–C1–P2 115.14(18), O1–P1–C1 112.78(15), O2–P2–C1 111.13(14), O2–P2–C2 119.96(16), O1–P1–C20 120.11(14).

In conclusion, the first phosphavinylidene(oxo)phosphorane **1**, a derivative with three “cumulated formal double bonds”, was obtained as a stable compound in the solid-state in a one-pot procedure. The ability of **1** to form transition metal complexes is now under investigation.


## Experimental Section

**General:** All experiments were carried out in flame-dried glassware under an argon atmosphere using high-vacuum-line techniques.

Solvents were dried and freshly distilled from sodium benzophenone ketyl and carefully deoxygenated on a vacuum line by several “freeze-pump-thaw” cycles. NMR spectra were recorded in  $\text{CDCl}_3$  on the following spectrometers:  $^1\text{H}$ , Bruker Avance 300 (300.13 MHz) and Avance 400 (400.13 MHz);  $^{13}\text{C}\{^1\text{H}\}$ , Bruker Avance 300 (75.47 MHz) and Avance 400 (100.62 MHz) (reference TMS);  $^{31}\text{P}$ , Bruker AC200 (81.02 MHz) and Avance 400 (162.04 MHz) (reference  $\text{H}_3\text{PO}_4$ ). Melting points were determined on a Wild Leitz-Biomed apparatus. Mass spectra were obtained on a Hewlett-Packard 5989A spectrometer by EI at 70 eV.

**Synthesis of Mes\*P(O)=C=PMe\* (**1**):** To a solution of Mes\*P=CCl $_2$  **3** (3.15 g, 8.67 mmol) in THF (20 mL) cooled to  $-80^\circ\text{C}$  was added a solution of *n*-butyllithium (5.7 mL, 1.6 M in hexane, 9.1 mmol). After stirring for 1 h, the solution of Mes\*P=C(Cl)Li was transferred by cannula to a solution of Mes\*P(O)Cl $_2$  **4** (3.00 g, 8.67 mmol) in THF (30 mL) at  $-80^\circ\text{C}$ . The reaction mixture became dark brown and after 15 min stirring and subsequent warming to  $-20^\circ\text{C}$ , the solvent was removed under vacuum. Pentane (30 mL) was added to the residue and **3** and **4** dissolved. However **1** and LiCl precipitated due to both being very poorly soluble in pentane. After filtration, **1** was extracted with  $\text{CH}_2\text{Cl}_2$ . Removal of the solvent gave a pale yellow powder of **1** (2.01 g, 40%, m.p.  $125^\circ\text{C}$ , dec.).

The carbon and phosphorus atoms are numbered as follows:



observed: 33.8 (C27–29 and C35–37), 37.8 (C26 and C34), 121.9 (C22 and C24) and 153.2 ppm (C21 and C25) in the  $^{13}\text{C}$  NMR spectrum; 1.39 (s, C27–29 and C35–37) and 7.32 ppm (s, C22 and C24) in the  $^1\text{H}$  NMR spectrum.

**Synthesis of Cyclisation Products 7a and 7b:** A solution of Mes\* $\text{-P(O)=C=PMes}^*$  **1** (1.20 g, 2.01 mmol) in toluene (10 mL) was heated at 80 °C for 1 h. NMR analysis showed the conversion of **1** into **7a** and **7b** in the ratio 85:15. Removal of solvent in vacuo led to a powder of **7a,b**. Attempts at fractional crystallisation in various solvents (pentane, Et<sub>2</sub>O, CHCl<sub>3</sub>) did not allow the complete separation of the two isomers **7a** and **7b**. Selective  $^{31}\text{P}$  decoupled  $^1\text{H}$  and  $^{13}\text{C}$  NMR experiments were performed. At 283 K, signals of some  $t\text{Bu}$  groups were too broad to be observed by  $^1\text{H}$  NMR spectroscopy.

**7a [(E) isomer]:**  $^1\text{H}$  NMR (CDCl<sub>3</sub>, 400.13 MHz, 283 K):  $\delta$  = 1.31, 1.48 and 1.56 (3s, 27 H, H from  $t\text{Bu}$  groups), 2.36 [ddd,  $^2J_{\text{H,H}} = 15.0$  Hz,  $^2J_{\text{H,P2}} = 19.3$  Hz,  $^4J_{\text{H,P1}} = 3.9$  Hz, 1 H, H on C9 (carbon atom bonded to P2)], 2.50 (dd,  $^2J_{\text{H,H}} = 15.0$  Hz,  $^2J_{\text{H,P2}} = 7.1$  Hz, 1 H, H on C9), 7.16, 7.28, 7.36 and 7.40 (4 broad s,  $4 \times 1$  H, arom H), 7.58 (dd,  $^2J_{\text{H,P1}} = 22.3$  Hz,  $^2J_{\text{H,P2}} = 13.8$  Hz, 1 H, H on C1) ppm.  $^{13}\text{C}$  NMR (CDCl<sub>3</sub>, 100.62 MHz, 283 K):  $\delta$  = 31.2, 31.3 and 32.5 (C13–15, C17–19, C31–33), 34.2 (d,  $^4J_{\text{C,P1}} = 7.0$  Hz, C27–29, C35–37), 37.5 (C30), 38.3 (C26, C34), 39.0 (d,  $^2J_{\text{C,P2}} = 5.8$  Hz, C8), 40.0 (d,  $J_{\text{C,P2}} = 2.8$  Hz) and 40.8 (d,  $J_{\text{C,P2}} = 3.6$  Hz) (C12, C16), 43.7 (dd,  $^1J_{\text{C,P2}} = 75.9$  Hz,  $^3J_{\text{C,P1}} = 6.0$  Hz, C9), 118.7 (d,  $^3J_{\text{C,P2}} = 13.8$  Hz) and 122.8 (d,  $^3J_{\text{C,P2}} = 10.0$  Hz, C4, C6), 122.2 (C22, C24), 127.1 (d,  $^1J_{\text{C,P2}} = 95.6$  Hz, C2), 138.8 (dd,  $^1J_{\text{C,P1}} = 67.5$  Hz,  $^3J_{\text{C,P2}} = 17.7$  Hz, C20), 150.1 (C23), 153.4 (C21, C25), 154.2 (d,  $^2J_{\text{C,P2}} = 7.2$  Hz) and 157.0 (d,  $^2J_{\text{C,P2}} = 20.2$  Hz, C3, C7), 155.7 (d,  $^4J_{\text{C,P2}} = 2.5$  Hz, C5), 165.3 (dd,  $^1J_{\text{C,P2}} = 74.8$  Hz,  $^1J_{\text{C,P1}} = 63.2$  Hz, C1) ppm.  $^{31}\text{P}$  NMR (CDCl<sub>3</sub>, 81.02 MHz, 298 K):  $\delta$  = 49.3 (d,  $^2J_{\text{P,P}} = 70.2$  Hz, P2), 339.1 (dd,  $^2J_{\text{P,P}} = 70.2$  Hz,  $^2J_{\text{P,H}} = 24.4$  Hz P1) ppm. MS (EI = 70 eV):  $m/z$  (%) = 565 (8) [ $\text{M} - 15$ ]<sup>+</sup>, 523 (15) [ $\text{M} - t\text{Bu}$ ]<sup>+</sup>, 335 (100) [ $\text{M} - \text{Mes}^*$ ]<sup>+</sup>, 57 (90) [ $t\text{Bu}$ ]<sup>+</sup>.

**7b [(Z) isomer]:**  $^1\text{H}$  NMR (CDCl<sub>3</sub>, 400.13 MHz, 283 K):  $\delta$  = 7.82 (dd,  $^2J_{\text{H,P2}} = 212.0$  Hz,  $^2J_{\text{H,P1}} = 17.6$  Hz, 1 H, CH=P) ppm (other  $^1\text{H}$  signals and  $^{13}\text{C}$  signals of the minor isomer could not be assigned unambiguously).  $^{31}\text{P}$  NMR (CDCl<sub>3</sub>, 81.02 MHz, 298 K):  $\delta$  = 38.7 (d,  $^2J_{\text{P,P}} = 73.3$  Hz, PO), 353.0 (dd,  $^2J_{\text{P,P}} = 73.3$  Hz,  $^2J_{\text{P,H}} = 21.4$  Hz P=C) ppm.

**Hydrolysis of 7a and 7b. Synthesis of 9:** An excess of degassed water was added to the mixture of **7a,b** (0.60 g, 1.04 mmol) in Et<sub>2</sub>O. After stirring for a week at room temperature, NMR analysis showed the formation of **9**. Et<sub>2</sub>O was removed under vacuum and recrystallisation from pentane afforded pure **9** as pale yellow crystals (0.48 g, 78%, m.p. 203 °C). HSQC, HMBC, NOESY and selective  $^{31}\text{P}$  decoupled  $^1\text{H}$  and  $^{13}\text{C}$  NMR experiments were performed.  $^1\text{H}$  NMR (CDCl<sub>3</sub>, 400.13 MHz, 253 K):  $\delta$  = 0.46 (dd,  $^2J_{\text{H,P2}} = 6.2$  Hz,  $^2J_{\text{H,H}} = 15.3$  Hz, 1 H, H on C9, carbon atom bonded to P2), 0.90 and 1.21 (2s,  $2 \times 3$  H, H on C10,11), 1.30 (s, 9 H, H on C13–15), 1.38 (s, 9 H, H on C31–33), 1.52 and 1.65 (2s,  $2 \times 9$  H, H on C27–29 and C35–37), 1.53 (s, 9 H, H on C17–19), 1.54 (dd,  $^2J_{\text{H,P2}} = 20.4$  Hz,  $^2J_{\text{H,H}} = 15.3$  Hz, 1 H, H on C9), 2.43 (dddd,  $^2J_{\text{H,P1}} = 15.0$  Hz,  $^2J_{\text{H,P2}} = 7.2$  Hz,  $^2J_{\text{H,H}} = 15.3$  Hz,  $^3J_{\text{H,H}} = 7.1$  Hz, 1 H, H on C1), 3.30 (dt,  $^2J_{\text{H,P1}}$  and  $^2J_{\text{H,P2}} = 16.6$  Hz,  $^2J_{\text{H,H}} = 15.3$  Hz, 1 H, H on C1), 7.07 (dd,  $^4J_{\text{H,P2}} = 2.0$  Hz,  $^4J_{\text{H,H}} = 1.0$  Hz, 1 H, H on C4), 7.41 (dd,  $^4J_{\text{H,P2}} = 5.2$  Hz,  $^4J_{\text{H,H}} = 1.0$  Hz, 1 H, H on C6), 7.45 (dd,  $^4J_{\text{H,P1}} = 2.0$  Hz,  $^4J_{\text{H,H}} = 1.0$  Hz, 1 H, H on C22), 7.54 (dd,  $^4J_{\text{H,P1}} = 4.1$  Hz,  $^4J_{\text{H,H}} = 1.0$  Hz, 1 H, H on C24), 8.52 (ddd,  $^1J_{\text{H,P1}} = 517.4$  Hz,  $^3J_{\text{H,P2}} = 2.8$  Hz,  $^3J_{\text{H,H}} = 7.1$  Hz, 1 H, H on P1) ppm.  $^{13}\text{C}$  NMR (CDCl<sub>3</sub>, 100.62 MHz, 253 K):  $\delta$  = 31.2 (C13–15), 31.4 (C31–33), 31.9 and 33.5 (C10–11), 32.2 (C17–19), 33.6 and 34.3

(C27–29 and C35–37), 35.3 (C30), 35.4 (C12), 37.7 (C16), 38.7 (d,  $^2J_{\text{C,P2}} = 6.2$  Hz, C8), 39.2 (C26 and C34), 39.6 (d,  $^1J_{\text{C,P2}} = 72.1$  Hz, C9), 40.6 (dd,  $^1J_{\text{C,P1}} = 55.1$  Hz,  $^1J_{\text{C,P2}} = 50.5$  Hz, C1), 118.9 (d,  $^3J_{\text{C,P2}} = 14.4$  Hz, C4), 123.3 (d,  $^3J_{\text{C,P2}} = 10.2$  Hz, C6), 123.6 (d,  $^3J_{\text{C,P1}} = 11.9$  Hz) and 124.1 (d,  $^3J_{\text{C,P1}} = 13.4$  Hz) (C22 and C24), 125.2 (dd,  $^1J_{\text{C,P2}} = 96.5$  Hz,  $^3J_{\text{C,P1}} = 3.9$  Hz, C2), 128.2 (d,  $^1J_{\text{C,P1}} = 99.0$  Hz, C20), 153.6 (d,  $^4J_{\text{C,P1}} = 3.1$  Hz, C23), 154.2 (d,  $^2J_{\text{C,P2}} = 8.1$  Hz, C7), 156.0 (d,  $^4J_{\text{C,P1}} = 5.7$  Hz, C21), 156.3 (d,  $^2J_{\text{C,P2}} = 33.0$  Hz, C3), 156.6 (d,  $^4J_{\text{C,P2}} = 2.4$  Hz, C5), 157.4 (d,  $^2J_{\text{C,P1}} = 11.9$  Hz, C25) ppm.  $^{31}\text{P}$  NMR (CDCl<sub>3</sub>, 81.02 MHz, 298 K):  $\delta$  = 4.8 (ddt,  $^2J_{\text{P,P}} = 2.7$  Hz,  $^1J_{\text{P,H}} = 517.4$  Hz,  $^2J_{\text{P,H}} = 2.8$  Hz, P1), 52.1 (d,  $^2J_{\text{P,P}} = 2.7$  Hz, P2) ppm. MS (EI = 70 eV):  $m/z$  (%) = 599 (2) [ $\text{M} + 1$ ]<sup>+</sup>, 581 (1) [ $\text{M} - \text{O} - 1$ ]<sup>+</sup>, 541 (100) [ $\text{M} - t\text{Bu}$ ]<sup>+</sup>, 353 (85) [ $\text{M} - \text{Mes}^*$ ]<sup>+</sup>, 57 (70) [ $t\text{Bu}$ ]<sup>+</sup> C<sub>37</sub>H<sub>60</sub>O<sub>2</sub>P<sub>2</sub> (598.82): calcd. C 74.21, H 10.10; found C 74.56, H 9.85.

**Synthesis of 8a and 8b by Hydrolysis of the P(O)=C Double Bond of 1:** An excess of degassed water was added to a solution of **1** (1.15 g, 1.98 mmol) in pentane (20 mL). After stirring for 2 d at room temperature, NMR analysis showed the formation of a mixture of **7a,b** and **8a,b**. Attempts at crystallisation from various solvents did not allow complete separation of the two derivatives and only some single crystals of **8**, suitable for an X-ray structure determination, could be obtained by crystallisation from CHCl<sub>3</sub> and cyclohexane.  $^{31}\text{P}$  NMR (CDCl<sub>3</sub>, 81.02 MHz, 298 K) **8a** (30%),  $\delta$  = 24.6 (d,  $^2J_{\text{P,P}} = 47.3$  Hz, PO), 305.5 ppm (dd,  $^2J_{\text{P,P}} = 47.3$  Hz,  $^2J_{\text{P,H}} = 24.4$  Hz, P=C) **8b** (70%),  $\delta$  = 34.9 ( $^2J_{\text{P,P}} = 68.5$  Hz, PO), 335.4 ppm (dd,  $^2J_{\text{P,P}} = 68.5$  Hz,  $^2J_{\text{P,H}} = 24.4$  Hz, P=C).

#### X-ray Structure Determinations

**Crystal Data for 8 and 9. Compound 8:** C<sub>43</sub>H<sub>73</sub>Cl<sub>3</sub>O<sub>2</sub>P<sub>2</sub>,  $M = 790.30$  (CHCl<sub>3</sub> and cyclohexane in the lattice), triclinic,  $P\bar{1}$ ,  $a = 13.998(2)$ ,  $b = 14.106(2)$ ,  $c = 14.348(2)$  Å,  $\alpha = 91.531(4)$ ,  $\beta = 119.170(3)$ ,  $\gamma = 108.678(3)^\circ$ ,  $V = 2285.6(6)$  Å<sup>3</sup>,  $Z = 2$ ,  $T = 173(2)$  K. 9004 reflections (5491 independent,  $R_{\text{int}} = 0.0740$ ) were collected. Largest electron density residue:  $0.341 \text{ e}\cdot\text{\AA}^{-3}$ ,  $R_1$  [for  $I > 2\sigma(I)$ ] = 0.0565 and  $wR_2 = 0.1433$  (all data) with  $R_1 = \Sigma||F_o| - |F_c||/\Sigma|F_o|$  and  $wR_2 = [\Sigma w(F_o^2 - F_c^2)^2/\Sigma w(F_o^2)^2]^{0.5}$ . **Compound 9:** C<sub>37</sub>H<sub>60</sub>O<sub>2</sub>P<sub>2</sub>,  $M = 598.79$ , monoclinic,  $P2_1/c$ ,  $a = 10.275(1)$ ,  $b = 25.939(2)$ ,  $c = 14.342(1)$  Å,  $\beta = 110.884(2)^\circ$ ,  $V = 3571.4(6)$  Å<sup>3</sup>,  $Z = 4$ ,  $T = 173(2)$  K. 15676 reflections (5066 independent,  $R_{\text{int}} = 0.1213$ ) were collected. Largest electron density residue:  $0.250 \text{ e}\cdot\text{\AA}^{-3}$ ,  $R_1$  [for  $I > 2\sigma(I)$ ] = 0.0558 and  $wR_2 = 0.0943$  (all data).

All data for structures reported in this paper were collected at low temperature using an oil-coated shock-cooled crystal on a Bruker-AXS CCD 1000 diffractometer with Mo- $K_\alpha$  radiation ( $\lambda = 0.71073$  Å). The structures were solved by direct methods (SHELXS-97)<sup>[15]</sup> and all non-hydrogen atoms were refined anisotropically using the least-squares method on  $F^2$  (SHELXL-97).<sup>[16]</sup>

CCDC-605519 (for **8**) and -605520 (for **9**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

**Supporting Information:** (for details see the footnote on the first page of this article): variable temperature  $^1\text{H}$  NMR, 2D HSQC and 2D HMBC,  $^{13}\text{C}\{^1\text{H}, ^{31}\text{P}\}$  and  $^{13}\text{C}\{^1\text{H}\}$  Jmod spectra of diphosphaallene Mes\*P(O)=C=PMes\* (**1**), variable-temperature  $^1\text{H}$  NMR spectra of bis(phosphane oxide)methylene **9**.

#### Acknowledgments

We thank the CNRS and the Ministry of Foreign Affairs [AI Brancusi no. 08581TK (Fr) and 18015 (Ro)]. R. S. thanks ERASMUS/

SOCRATES and CEEX SUPRACOM and G. N. the AUF for a grant.

- [1] a) *The Chemistry of Ketenes, Allenes and Related Compounds* (Ed.: S. Patai), **1980**; b) *The Chemistry of Allenes* (Ed.: S. R. Landor), **1982**; c) *Allenenes in Organic Synthesis* (Eds.: H. F. Schuster, G. M. Coppola), **1984**.
- [2] a) F. Ramirez, N. B. Desai, B. Hansen, N. McKelvie, *J. Am. Chem. Soc.* **1961**, *83*, 3539–3540; b) H. J. Bestmann, R. Zimmermann, in: *Methoden der Organischen Chemie*, Houben–Weyl, 4<sup>th</sup> ed., vol. E1, **1982**, pp. 752–758; c) A. W. Johnson, W. C. Kaska, K. A. O. Starzewski, D. A. Dixon, in: *Ylides and Imines of Phosphorus* (Ed.: A. W. Johnson), Wiley-Interscience, New York, **1993**; d) O. I. Kolodiaznyi, in: *Phosphorus Ylides* (Ed.: O. I. Kolodiaznyi), Wiley-VCH, Weinheim, **1999**.
- [3] For reviews see: a) R. Appel, in: *Multiple Bonds and Low Coordination in Phosphorus Chemistry* (Eds.: M. Regitz, O. J. Scherer), Thieme: Stuttgart, **1990**, p. 157–219; b) J. Escudié, H. Ranaivonjatovo, L. Rigon, *Chem. Rev.* **2000**, *100*, 3639–3696; c) J. Escudié, H. Ranaivonjatovo, M. Bouslikhane, Y. El Harouch, L. Baiget, G. Cretiu-Nemes, *Russ. Chem. Bull.* **2004**, *53*, 1020–1033.
- [4] a) R. Appel, P. Fölling, B. Josten, M. Siray, V. Winkhaus, F. Knoch, *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 618–619; b) H. H. Karsch, F. H. Köhler, H.-U. Reisacher, *Tetrahedron Lett.* **1984**, *25*, 3687–3690; c) M. Yoshifuji, K. Toyota, N. Inamoto, *J. Chem. Soc., Chem. Commun.* **1984**, 689–690; d) M. Gouygou, M. Koenig, J. Escudié, C. Couret, *Heteroat. Chem.* **1991**, *2*, 221–227; e) M. Yoshifuji, S. Sasaki, N. Inamoto, *J. Chem. Soc., Chem. Commun.* **1989**, 1732–1733; f) M. Yoshifuji, T. Niitsu, D. Shiomi, N. Inamoto, *Tetrahedron Lett.* **1989**, *30*, 5433–5436; g) K. Toyota, A. Nakamura, M. Yoshifuji, *Chem. Commun.* **2002**, 3012–3013.
- [5] a) U. Krüger, H. Pritzkow, H. Grützmacher, *Chem. Ber.* **1991**, *124*, 329–331; b) H. P. Schrödel, G. Jochem, A. Schmidpeter, H. Nöth, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1853–1856; c) T. Kato, H. Gornitzka, A. Baceiredo, G. Bertrand, *Angew. Chem. Int. Ed.* **2000**, *39*, 3319–3321; d) D. Martin, H. Gornitzka, A. Baceiredo, G. Bertrand, *Eur. J. Inorg. Chem.* **2005**, 2619–2624.
- [6] a) R. Appel, C. Casser, *Tetrahedron Lett.* **1984**, *25*, 4109–4112; b) R. Appel, H. Knoch, H. Kunze, *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 157–158.
- [7] a) S. J. Goede, F. Bickelhaupt, *Chem. Ber.* **1991**, *124*, 2677–2684; b) E. Niecke, M. Nieger, O. Schmidt, D. Gudat, W. W. Schoeller, *J. Am. Chem. Soc.* **1999**, *121*, 519–522; c) M. Yoshifuji, S. Ito, *Top. Curr. Chem.* **2003**, *223*, 67–89.
- [8] S. Freeman, M. J. P. Harger, *J. Chem. Soc., Perkin Trans. 1* **1987**, 1399–1406.
- [9] H. Ranaivonjatovo, H. Ramdane, H. Gornitzka, J. Escudié, J. Satgé, *Organometallics* **1998**, *17*, 1631–1633.
- [10] M. Yoshifuji, S. Sasaki, T. Niitsu, N. Inamoto, *Tetrahedron Lett.* **1989**, *30*, 187–188.
- [11] For reviews see: a) S. Lochschmidt, A. Schmidpeter, *Phosphorus Sulfur Silicon Relat. Elem.* **1986**, *29*, 73–109; b) R. Appel, F. Knoll, *Adv. Inorg. Chem.* **1989**, *33*, 259–361; c) L. N. Markovski, V. D. Romanenko, *Tetrahedron* **1989**, *45*, 6019–6090; d) L. Weber, *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 271–288; e) *CRC Handbook of Phosphorus-31 Nuclear Magnetic Resonance Data* (Ed.: J. C. Tebbly), CRC Press, Boca Raton, Florida (USA), **1991**.
- [12] H. H. Karsch, H.-U. Reisacher, *Phosphorus Sulfur Silicon Relat. Elem.* **1988**, *37*, 241–242.
- [13] Such intramolecular cyclisations are well-known in the chemistry of phosphallenes a) H. H. Karsch, H.-U. Reisacher, *Phosphorus Sulfur Silicon Relat. Elem.* **1988**, *36*, 213–215; b) H. H. Karsch, H.-U. Reisacher, G. Müller, *Angew. Chem. Int. Ed. Engl.* **1986**, *25*, 454–455; c) M. A. David, J. B. Alexander, D. S. Glueck, G. P. A. Yap, L. M. Liable-Sands, A. L. Rheingold, *Organometallics* **1997**, *16*, 378–383; d) M. A. David, S. N. Paisner, D. S. Glueck, *Organometallics* **1995**, *14*, 17–19.
- [14] a) R. Appel, Th. Gaitzsch, F. Knoch, G. Lenz, *Chem. Ber.* **1986**, *119*, 1977–1985; b) E. Niecke, D. A. Wildbrecht, W. Schoeller, *Angew. Chem. Int. Ed. Engl.* **1981**, *20*, 131–132.
- [15] G. M. Sheldrick, *Acta Crystallogr., Sect. A* **1990**, *46*, 467–473.
- [16] G. M. Sheldrick, *Program for Crystal Structure Refinement*, University of Göttingen **1997**.

Received: April 27, 2006

Published Online: August 7, 2006