


## SHORT COMMUNICATION

### Sandwich Complex with $\eta^4\text{-Si}_4\text{R}_4$

K. Takanashi, V. Ya. Lee, M. Ichinohe,  
A. Sekiguchi\* ..... 5471–5474

( $\eta^5$ -Cyclopentadienyl)( $\eta^4$ -tetrasilal- and  $\eta^4$ -trisilagermacyclobutadiene)cobalt: Sandwich Complexes Featuring Heavy Cyclobutadiene Ligands

**Keywords:** Cobalt / Cyclobutadiene ligand / Cyclopentadienyl ligand / Germanium / Silicon



( $\eta^5$ -Cyclopentadienyl)( $\eta^4$ -tetrasilal- and  $\eta^4$ -trisilagermacyclobutadiene)cobalt sandwich complexes [( $\eta^4$ - $\text{R}_4\text{Si}_4$ )CoCp] (**2**) and [( $\eta^4$ - $\text{R}_4\text{Si}_3\text{Ge}$ )CoCp] (**4**) ( $\text{R} = \text{SiMe}_2\text{Bu}_2$ ) were synthesized by reaction of the dipotassium salts of tetrasilacyclobutadiene dianion  $\text{K}^+{}_2[\text{R}_4\text{Si}_4]^{2-}$  (**1**) and trisilagermacyclobutadiene dianion  $\text{K}^+{}_2[\text{R}_4\text{Si}_3\text{Ge}]^{2-}$  (**3**) with  $[\text{CpCoI}_2(\text{PPh}_3)]$ . The structures of the new Co complexes were confirmed by X-ray analysis.


## FULL PAPERS

### Metal–Organic Frameworks

I. Senkovska, J. Fritsch,  
S. Kaskel\* ..... 5475–5479

New Polymorphs of Magnesium-Based Metal–Organic Frameworks  $\text{Mg}_3(\text{ndc})_3$  ( $\text{ndc} = 2,6\text{-Naphthalenedicarboxylate}$ )

**Keywords:** Metal–organic frameworks / Magnesium / 2,6-Naphthalenedicarboxylic acid / Solvent effects / Gas adsorption



Two new magnesium 2,6-naphthalenedicarboxylate ( $\text{ndc}$ ) metal–organic frameworks,  $[\text{Mg}_3(\text{ndc})_3(\text{dif})_4]$  (**1**) ( $\text{dif} = N,N\text{-diisopropylformamide}$ ) and  $[\text{Mg}_3(\text{ndc})_3(\text{dmf})_2(\text{CH}_3\text{OH})(\text{H}_2\text{O})](\text{dmf})$  (**2** = TUDMOF-3) ( $\text{dmf} = N,N\text{-dimethylformamide}$ ), have been synthesised. TUDMOF-3 has a permanent porosity with a Langmuir surface area of  $632 \text{ m}^2 \text{ g}^{-1}$ , a specific pore volume of  $0.21 \text{ cm}^3 \text{ g}^{-1}$  and a hydrogen storage capacity of 1.23 wt.-% (77 K, 1 bar). Compound **1** is not porous.


### Hybrid Mesostructures

S.-R. Zhai, S. S. Park, M. Park,  
M. H. Ullah, C.-S. Ha\* ..... 5480–5488




Direct Synthesis of Zr-Containing Hybrid Periodic Mesoporous Organosilicas with Tunable Zirconium Content

**Keywords:** Silicates / Template synthesis / Organic–inorganic hybrid composites / Mesoporous materials



Highly ordered Zr-containing periodic mesoporous organosilicas (ZrPMO) were successfully synthesized, for the first time, by employing a  $\text{ZrOCl}_2/\text{NaCl}$  combination as the promoting agent; the addition of extra amounts of acid was not needed throughout the synthetic process.

The propylene bridge in the skeleton of the H<sub>3</sub>EPTPACH<sub>2</sub>OH ligand optimizes the water-exchange rate of the [Gd(EPTPACH<sub>2</sub>OH)(H<sub>2</sub>O)]<sup>2-</sup> complex, for efficiency as a MRI contrast agent, as a result of the steric compression imposed around its water binding site, as shown by solution relaxometric characterization. The hydroxymethyl substituent group is useful for conjugation of the reporter group. However, these changes, relative to those incurred by the DTPA chelate, significantly decrease its thermodynamic and kinetic stability.



S. Torres, J. A. Martins,\*  
 J. P. André,\* G. A. Pereira, R. Kiraly,  
 E. Brücher, L. Helm, É. Tóth,  
 C. F. G. C. Geraldes\* ..... 5489–5499


H<sub>3</sub>EPTPACH<sub>2</sub>OH: Synthesis, Relaxometric Characterization and <sup>1</sup>H NMR Spectroscopic Studies on the Solution Dynamics of Its Ln<sup>III</sup> Complexes



**Keywords:** Lanthanides / Contrast agents / MRI / Microscopic protonation scheme / Stability constants

### Isomerism of Copper Complexes

The preference for axial or equatorial positioning of the bridging atoms and of the amine or pyridine N centers of tridentate ligands L in LCu<sup>II</sup>(μ-X)<sub>2</sub>Cu<sup>II</sup>L complexes was probed experimentally and by DFT calculations. Their preference for either a *syn* or *anti* configuration in an equatorial/equatorial situation was also evaluated.



J. Manzur,\* A. Vega,\* A. M. García,  
 C. Acuña, M. Sieger, B. Sarkar,  
 M. Niemeyer, F. Lissner, T. Schleid,  
 W. Kaim\* ..... 5500–5510


Coordination Alternatives in Dinuclear Bis(pyridin-2-ylalkyl)benzylaminecopper(II) Complexes with OH<sup>-</sup>, RO<sup>-</sup>, F<sup>-</sup>, or Cl<sup>-</sup> Bridges: Experimental Structures and DFT Preferences



**Keywords:** Copper / EPR spectroscopy / Isomerism / N ligands / Structure

### Coordination Oligomers/Polymers

Reactions of three preformed Cu<sup>II</sup>/bdmpp complexes [bdmpp = 2,6-bis(3,5-dimethyl-1*H*-pyrazol-1-yl)pyridine] with 4,4'-bipy, dicyanamide (dca), azide, and MCl<sub>2</sub> (M = Cu, Co, Ni) lead to a series of homo- and heterometallic coordination oligomers and polymers. Their structures and redox properties were investigated.




G.-F. Liu, Z.-G. Ren, H.-X. Li,  
 Y. Chen, Q.-H. Li, Y. Zhang,  
 J.-P. Lang\* ..... 5511–5522

Homo- and Heterometallic Coordination Oligomers and Polymers Derived from the Preformed Complexes [Cu(bdmpp)(MeCN)<sub>2</sub>](ClO<sub>4</sub>)<sub>2</sub>, [Cu(bdmpp)(N<sub>3</sub>)<sub>2</sub>], and [Cu(bdmpp)(N<sub>3</sub>)(μ-N<sub>3</sub>)<sub>2</sub>] [bdmpp = 2,6-bis(3,5-dimethyl-1*H*-pyrazol-1-yl)pyridine]: Syntheses, Structures, and Redox Properties

**Keywords:** Copper / Structures / Oligomers / Polymers / Redox properties

### Water-Soluble Ruthenium Complexes

A series of [tris(pyrazolyl)borato]ruthenium complexes bearing the water-soluble cage phosphane 1,3,5-triaza-7-phosphaadamantane (PTA) and its boranyl adduct PTA(BH<sub>3</sub>), including the water-soluble hydride [TpRuH(PTA)<sub>2</sub>], have been synthesised and characterised. Cyclic voltammetric studies in MeCN and dmsO generally show reversible Ru<sup>II/III</sup> oxidation waves and the values of *E*<sub>1/2</sub><sup>ox</sup> reflect the electron-donating character of the ligands, whose Lever electrochemical parameter (*E*<sub>L</sub>) is estimated for the first time.



S. Bolaño, J. Bravo,\* J. Castro, M. Mar  
 Rodríguez-Rocha, M. F. C. Guedes da Silva,  
 A. J. L. Pombeiro,\* L. Gonsalvi,  
 M. Peruzzini\* ..... 5523–5532

Synthesis, Reactivity, X-ray Crystal Structures and Electrochemical Behaviour of Water-Soluble [Tris(pyrazolyl)borato]ruthenium(II) Complexes of 1,3,5-Triaza-7-phosphaadamantane (PTA)

**Keywords:** Electrochemistry / Water-soluble complexes / Pyrazolylborates / Ruthenium / N ligands


# CONTENTS

## 3d–4f Coordination Polymers

A. M. Madalan, K. Bernot, F. Pointillart, M. Andruh,\* A. Caneschi\* ..... 5533–5540



First  $\text{Ni}^{\text{II}}$ – $\text{Ln}^{\text{III}}$  Coordination Polymers Constructed by Using  $[\text{Ni}(\text{bpca})_2]$  as a Building Block [Hbpca = bis(2-pyridylcarbonyl)amine]: Synthesis, Crystal Structures and Magnetic Properties



Three new heterobimetallic 1D coordination polymers have been obtained by assembling the  $[\text{Ni}(\text{bpca})_2]$  complex, as a building block, with lanthanide cations:  $[\{\text{Ln}(\text{O}_2\text{NO})(\text{H}_2\text{O})_3\}\{\text{Ni}(\text{bpca})_2\}](\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$  (Ln Gd **1**; Tb **2**; Dy **3**). Their static and dynamic magnetic properties were investigated and discussed.


**Keywords:** Nickel / Lanthanides / Heterobimetallic coordination polymers / Magnetic properties

## Ditungsten(III) Complexes

O. Coutelier, R. M. Gauvin,\* G. Nowogrocki, J. Trébosc, L. Delevoye, A. Mortreux\* ..... 5541–5547



A New Donor-Stabilized Ditungsten Amido Alkoxido Species: Synthesis, Crystal Structure, Fluxionality, and Grafting onto Silica



A new dimeric tungsten(III) complex containing amido and chelating pyridine-alkoxido ligands was synthesized through protonolysis of  $[\text{W}_2(\text{NMe}_2)_6]$  by 2-(2-pyridyl)propan-2-ol and fully characterized. Grafting of this compound onto silica has been investigated by elemental analysis, DRIFTS, and solid-state NMR spectroscopy.


**Keywords:** Tungsten / Metal-metal multiple bonds / Ligand design / Grafting / Silica / Alkyne metathesis

## Copper–Thiolate Clusters

B. K. Maiti, K. Pal, S. Sarkar\* ..... 5548–5555



Flexible  $\text{Cu}^{\text{I}}$ –Thiolate Clusters with Relevance to Metallothioneins




The structural diversity in some Cu–thiol small clusters with relevance to the chemistry of Cu-containing metallothioneins was investigated. Solvent polarity and the copper/thiol ratio may change the local cluster geometry, which causes flexibility of the copper–sulfur bond. These small clusters have superposition motifs in bigger clusters such as  $\text{Cu}_8$ -yeastMT.

## Au–Au Interactions

F. Balzano, A. Cuzzola, P. Diversi,\* F. Ghiotto, G. Uccello-Barretta ..... 5556–5562


Cationic Gold(I) Phosphanyl Thiolates: Auophilic Interactions in the Solid State and in Solution



Cationic gold(I) phosphane alkanethiolates of the formula  $[\text{Au}_2(\text{S}t\text{Bu})(\text{phosphane})_2][\text{BF}_4]$  were investigated in solution by MS and NMR spectroscopy. These compounds are dimeric in the solid state and are bound through auophilic interactions; some of these compounds maintain the integrity of the Au–Au bonds in solution depending on the nature of the phosphane.

**Keywords:** Gold / Sulfur / Thiolates / Auophilicity / Aggregation / ESI MS / NMR spectroscopy

Based upon single-crystal X-ray diffraction analysis, the potassium salt of the 9-BBN hydroborate anion has a tetrameric structure when associated with toluene, methyl THF, or a mixture of toluene/THF, and THF/diethyl ether. While with the chelating solvent DME, it has a dimeric structure.




X. Chen, S. Liu, B. Du, E. A. Meyers,  
S. G. Shore\* ..... 5563–5570

The Structure of Potassium and Tetramethylammonium Salts of the Cyclic Organohydroborate Anion  $[H_2BC_8H_{14}]^-$  with Different Solvent Ligands

**Keywords:** Agostic interactions / X-ray diffraction / Organohydroborate / Potassium salts

## Hybrid Materials



Conducting polymer/cellulose fibre composites, using polypyrrole and polyaniline, have been successfully prepared. By utilising the redox active nature of the conducting polymer coatings, silver nanopar-

ticles have been deposited on the surface of the composite fibres. This additionally gives rise to fibres with antimicrobial properties.


F. M. Kelly,\* J. H. Johnston, T. Borrmann,  
M. J. Richardson ..... 5571–5577

Functionalised Hybrid Materials of Conducting Polymers with Individual Fibres of Cellulose

**Keywords:** Polypyrrole / Polyaniline / Cellulose / Fibres / Conductivity / Antimicrobial activity

## Olefin Metathesis Catalysts

The use of Ru-based NHC-arene complexes as alternatives to the Grubbs-type catalysts used in olefin metathesis is described.




N. Ledoux,\* B. Allaert,  
F. Verpoort ..... 5578–5583

Ruthenium-Based NHC-Arene Systems as Ring-Opening Metathesis Polymerisation Catalysts

**Keywords:** Carbene ligands / Homogeneous catalysis / Olefin polymerisation / Ruthenium

A series of iron, cobalt and nickel complexes bearing 2-arylimino-9-phenyl-1,10-phenanthrolines has been synthesized, characterized and evaluated for ethylene oligomerization. Most of the complexes show good catalytic activity for ethylene oligomerization, with the cobalt catalysts being more active than their iron analogues.



S. Jie, S. Zhang,  
W.-H. Sun\* ..... 5584–5598

2-Arylimino-9-phenyl-1,10-phenanthroline-iron, -cobalt and -nickel Complexes: Synthesis, Characterization and Ethylene Oligomerization Behavior

**Keywords:** Nitrogen ligands / Late transition metals / Iron / Ethylene / Oligomerization

# CONTENTS

---


## Octaruthenium Grids on HOPG

D.-L. Shieh, Y.-S. Lin, K.-B. Shiu,  
J.-L. Lin\* ..... 5599–5603



Orientation and Arrangement of Octaruthenium Supramolecules with Alkyl Chains on Graphite

**Keywords:** Supramolecular chemistry / Scanning tunneling microscopy / Ruthenium



$\{[\text{Ru}_2(\text{CO})_4(\text{NH}_2\text{C}_{16}\text{H}_{33})_2](\mu\text{-O}_2\text{CCO}_2)\}_4 / \text{HOPG}$

The self-assemblies of octaruthenium grid-type supramolecules,  $[\text{Ru}_2(\text{CO})_4(\text{NH}_2\text{C}_{16}\text{H}_{33})_2](\mu\text{-O}_2\text{CCO}_2)_4$ , on highly oriented pyrolytic graphite (HOPG) in air and in phenyloctane were studied by scanning tunneling microscopy (STM).

---

## REMOVAL

L.-Y. Li, Q. Yuan, L. Liao,\* C.-H. Yan,  
J. Chen ..... 5605

Removal: “Ultra-Simple Synthesis of Ordered Mesoporous  $\gamma$ -Alumina: High Thermal Stability and Catalytic Activity”

---

If not otherwise indicated in the article, papers in issue 34 were published online on November 20, 2007