

# SPOTLIGHTS ...


## Phosphorus Chemistry

C. A. Dyker, N. Burford\*

### catena-Phosphorus Cations

*Chem. Asian J.*

DOI: 10.1002/asia.200700229



**A new chapter:** catena-Phosphorus cations represent a new and developing area in fundamental phosphorus chemistry that complements the series of neutral and anionic polyphosphorus compounds. The picture shows examples of phosphinophosphonium frameworks (each vertex represents a phosphorus center).


## Protein Engineering

T. Leta Aboye, R. J. Clark,  
D. J. Craik, U. Göransson\*

### Ultra-Stable Peptide Scaffolds for Protein Engineering—Synthesis and Folding of the Circular Cystine Knotted Cyclotide Cycloviolacin O2

*ChemBioChem*

DOI: 10.1002/cbic.200700357



**Key to the challenge of cross-linking disulfide bonds in a circular peptide backbone.** The cyclic cystine knot motif is an attractive scaffold for protein engineering. However, the synthesis and folding of members of the most diverse and biologically active cyclo-

tide subfamily, the bracelets, has been a big challenge. This study describes the key solution to these problems and thus provides an efficient method of exploring the most potent cyclic cystine knot peptides.


## Electronic properties

M. Melucci,\* M. Zambianchi,  
A. Zanelli, N. Camaioni, M. Gazzano,  
A. Bongini, G. Barbarella

### Polyvinyl-Locked versus Free Quaterthiophene: Effect of Spatial Constraints on the Electronic Properties of *n*-Hexylquaterthiophene

*ChemPhysChem*

DOI: 10.1002/cphc.200700459



**Close chromophores:** In a triad of quaterthiophene covalently linked to a polymer backbone (see figure), intramolecular as well as intermolecular interactions between pendant quater-

thiophenes lead to a peculiar solid-state supramolecular organization which is responsible for a marked enhancement of the electron affinity.

## Kinase Inhibitors


C. M. Crane, A. K. H. Hirsch,  
M. S. Alphey, T. Sgraja, S. Lauw,  
V. Illarionova, F. Rohdich,\*  
W. Eisenreich, W. N. Hunter,\*  
A. Bacher, F. Diederich\*

### Synthesis and Characterization of Cytidine Derivatives that Inhibit the Kinase IspE of the Non-Mevalonate Pathway for Isoprenoid Biosynthesis


*ChemMedChem*

DOI: 10.1002/cmdc.200700208

**The binding modes** of two water-soluble, cytidine-based inhibitors in complex with the *A. aeolicus* kinase IspE were elucidated by co-crystal structure analysis. Because key active site residues in *A. aeolicus* IspE are identical to those of the corresponding enzymes of *M. tuberculosis* and *P. falciparum*, useful structural information was gained for future structure-based development of inhibitors of the parasite enzymes.



## ... ON OUR SISTER JOURNALS



**Rhenium clusters probe reactivity of TiO<sub>2</sub> surface-defect sites:** Graphical representation of deprotonated rhenium clusters adsorbed on defective TiO<sub>2</sub> surface interacting with surface after sites.


K. Suriye, R. J. Lobo-Lapidus,  
G. J. Yeagle, P. Praserthdam,  
R. D. Britt, B. C. Gates\*

**Probing Defect Sites on TiO<sub>2</sub> with [Re<sub>3</sub>(CO)<sub>12</sub>H<sub>3</sub>]: Spectroscopic Characterization of the Surface Species**

*Chem. Eur. J.*  
DOI: 10.1002/chem.200701514

### Surface Chemistry

H-bonds offer a convenient and versatile strategy for the design and understanding of supramolecular arrangements. Different approaches to the design of noncovalent structures leading to both calamitic and bent-core liquid crystals, as well as their characterization, are reported. A possible correlation between the promoted liquid crystal order and the structure of the noncovalent complex is proposed.




### H-Bonded Liquid Crystals

A. Pérez, N. Gimeno, F. Vera,  
M. B. Ros,\* J. L. Serrano,  
M. R. De la Fuente

**New H-Bonded Complexes and Their Supramolecular Liquid-Crystalline Organizations**

*Eur. J. Org. Chem.*  
DOI: 10.1002/ejoc.200700725

**Kissing complexes** formed on the basis of highly specific noncanonical interactions of RNA hairpins can be harnessed for the controlled assembly of DNA nanoobjects. Two DNA minicircles, each equipped with a different RNA hairpin motif (see picture), mediate a tight and specific binding of the two circular DNA nanoobjects. These interactions may offer the possibility to construct DNA nanoobjects with increased complexity.




### DNA Architectures

G. Mayer, D. Ackermann, N. Kuhn,  
M. Famulok\*

**Construction of DNA Architectures with RNA Hairpins**

*Angew. Chem. Int. Ed.*  
DOI: 10.1002/anie.704709



### Towards sustainable stockings?

Nylon 6 was efficiently converted into its monomer caprolactam, which was isolated in over 90% yield with excellent purity as the sole product of the

reaction, by treatment with supercritical secondary or tertiary alcohols. The present method opens up a new avenue in plastic recycling chemistry.

### Polymer Recycling

A. Kamimura,\* Y. Oishi, K. Kaiso,  
T. Sugimoto, K. Kashiwagi

**Supercritical Secondary Alcohols as Useful Media To Convert Polyamide into Monomeric Lactams**

*ChemSusChem*  
DOI: 10.1002/cssc.200700024