

(Benzimidazolylmethyl)cyclen: A Potential Sensitive Fluorescent PET Chemosensor for Zinc

Abir El Majzoub,^[a] Cyril Cadiou,^[a] Isabelle Déchamps-Olivier,^[a] Françoise Chuburu,^{*[a]} and Michel Aplincourt^[a]

Keywords: Zinc / Macrocyclic ligands / X-ray diffraction / UV/Vis spectroscopy / Fluorescent probes / Fluorimetric titrations

The ¹³C NMR data of ligand LH and complex [ZnLH](NO₃)₂ on page 5094 of the original article^[1] are erroneous; the correct values are given below.

LH: ¹³C NMR (62.9 MHz, D₂O): δ = 43.9, 44.3, 45.5, 51.2, 52.7 (CH₂N), 115.2, 123.3, 138.1, 153.7 (C_{Ar}) ppm.

[ZnLH](NO₃)₂: ¹³C NMR (62.9 MHz, CD₃OD): δ = 45.1, 46.2, 47.4, 52.2, 53.9 (CH₂N), 113.5, 118.5, 125.2, 125.8, 135.6, 139.3, 156.3 (C_{Ar}).

The Authors

[1] A. El Majzoub, C. Cadiou, I. Déchamps-Olivier, F. Chuburu, M. Aplincourt, *Eur. J. Inorg. Chem.* **2007**, 5087–5097.

Received: December 11, 2007

Published Online: January 7, 2008

[a] GRECI, Université de Reims Champagne-Ardenne,
B. P. 1039, 51687 Reims Cedex 2, France
E-mail: francoise.chuburu@univ-reims.fr