

Chemistry and Physical Properties of the Phosphide Telluride Zr_2PTe_2

Kristina Tschulik,^[a] Michael Ruck,^[a] Michael Binnewies,^[b] Edgar Milke,^[b]
Stefan Hoffmann,^[c] Walter Schnelle,^[c] Boniface P. T. Fokwa,^[d] Michael Gilleßen,^[d] and
Peer Schmidt*^[a]

Keywords: Phosphorus / Tellurium / Zirconium / Chemical vapor transport / Thermodynamics / Structure elucidation

The ICOHP values for the Zr–P and Zr–Te bonds reported on page 3107 (2nd column)^[1] should be multiplied by 3 and the value for the Zr–Zr bond by 9. Therefore, the correct values read: -2.420 , -1.770 and $-0.369 \text{ eV bond}^{-1}$ for Zr–P, Zr–Te and Zr–Zr, respectively. However, this correction does not influence the discussion made in the text.

The Authors

[1] K. Tschulik, M. Ruck, M. Binnewies, E. Milke, S. Hoffmann, W. Schnelle, B. P. T. Fokwa, M. Gilleßen, P. Schmidt, *Eur. J. Inorg. Chem.* **2009**, 3102–3110.

Received: March 26, 2010
Published Online: April 13, 2010

[a] Inorganic Chemistry, Dresden University of Technology,
Helmholtzstraße 10, 01069 Dresden, Germany
Fax: +49-351-463-37287
E-mail: peer.schmidt@chemie.tu-dresden.de
<http://www.peer-schmidt.de>

[b] Inorganic Chemistry, Leibniz University Hannover,
Callinstr. 9, 30167 Hannover, Germany

[c] Max Planck Institute for Chemical Physics of Solids,
Nöthnitzer Str. 40, 01187 Dresden, Germany

[d] Inorganic Chemistry, RWTH Aachen,
Landoltweg 1, 52074 Aachen, Germany