## Erratum: "Three-dimensional mixing in Stokes flow: the partitioned pipe mixer problem revisited" Eur. J. Mech. B/Fluids 18(5) (1999) 783–792

V.V. Meleshko <sup>a</sup>, O.S. Galaktionov <sup>a,b</sup>, G.W.M. Peters <sup>b</sup>, H.E.H. Meijer <sup>b</sup>

<sup>a</sup> Institute of Hydromechanics, National Academy of Sciences, 03057 Kiev, Ukraine
 <sup>b</sup> Dutch Polymer Institute, Eindhoven Polymer Laboratories, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

We have noticed a couple of unfortunate misprints in the formulae of this paper. Equation (4) should read:

$$\psi = 0, \quad \frac{\partial \psi}{\partial r} = -V \quad \text{for } r = a, \ 0 \leqslant \theta \leqslant \pi,$$

$$\psi = 0, \quad \frac{\partial \psi}{\partial \theta} = 0 \quad \text{for } 0 \leqslant r \leqslant a, \ \theta = 0, \pi.$$
(4)

Equation (11) should read

$$\Psi = 0, \quad \frac{\partial \Psi}{\partial \xi} = V \quad \text{at} \quad \xi = \frac{1}{2}\pi, \ |\eta| \le \infty,$$

$$\Psi = 0, \quad \frac{\partial \Psi}{\partial \xi} = 0 \quad \text{at} \quad \xi = \pi, \ |\eta| \le \infty.$$
(11)

The important equation (6) is correct as stated. The misprints had no influence on the results shown in figures 1–5.